

Technical Report for the Diabetes Prevention Impact Toolkit

December 2016

CONTENTS

Section	Page
1. Introduction	1
2. Input Dashboard	1
2.1 State Module Population Characteristics	1
2.2 Employer Module Population Characteristics	9
2.2.1 Assume National Average for Population Characteristics	10
2.2.2 Assume State Average for Population Characteristics	11
2.2.3 Assume Industry Average for Population Characteristics	12
2.2.4 Assume Occupation Average for Population Characteristics	15
2.2.5 Entering Employee Population Characteristics	16
2.3 Insurer Module Population Characteristics	17
2.3.1 Assume National Average for Population Characteristics	18
2.3.2 Assume State Average for Population Characteristics	18
2.3.3 Assume Industry Average for Population Characteristics	20
2.3.4 Assume Occupation Average for Population Characteristics	20
2.3.5 Entering Insured Population Characteristics	21
2.4 Diabetes Incidence Rates for Each Risk Group	21
2.4.1 Persons with Prediabetes	23
2.4.2 Persons with Prediabetes and Other Persons at Risk for Type 2 Diabetes	24
2.4.3 Persons with High-Risk Prediabetes	24
2.5 Additional Impact Toolkit Inputs	25
2.5.1 Screening	28
2.5.2 Program Enrollment and Participation	29
2.5.3 Intervention Weight Loss and Regain Schedule	31
2.5.4 Program Budget	33
2.5.5 Program Costs	33
2.5.6 Medical Costs	35
2.5.7 Productivity Costs (Employer Module Only)	37
3. Output Dashboard	38
3.1 Projected Participants	38
3.2 Cumulative Projected Cases of Diabetes	39
3.3 Cumulative Medical Costs per Participant	41

3.4	Net Costs (Program Costs minus Medical and Productivity Cost Savings) per Participant	43
3.4.1	Productivity Costs in the Employer Module	44
3.5	Cumulative Quality Adjusted Life Years (QALYs) Gained	45
3.6	Incremental Cost-Effectiveness Ratios (ICERs)	46
3.7	Cumulative Years of Life Gained.....	48

References

R-1

FIGURES

Number		Page
1.	Three Risk Groups Eligible to Participate in National DPP	2
2.	Employer Input Dashboard Collapsed.....	26
3.	Employer Input Dashboard (Expanded)	27
4.	Program Cost Inputs Showing Average Screening Cost Calculation	29
5.	Program Enrollment and Participation Inputs	30
6.	Projected Participants	39
7.	Projected Cases of Diabetes and Years with Diabetes Averted by Participating in the National DPP Lifestyle Change Program	40
8.	Cumulative Medical Costs per Participant	42
9.	Net Costs (Program Costs minus Medical and Productivity Cost Savings) per Participant.....	44
10.	Cumulative Quality Adjusted Life Years (QALYs) Gained	46
11.	Incremental Cost-Effectiveness Ratios (ICERs).....	47
12.	Cumulative Years of Life Gained	49

TABLES

Number		Page
1.	National Estimates of Prediabetes and Diabetes (NHANES 2011–2014)	4
2.	National Estimates of “Persons with Prediabetes” and BMI \geq 24 (NHANES 2011–2014)	5
3.	National Estimates of “Persons with Prediabetes,” “Persons with Prediabetes and Other Persons at Risk for Type 2 Diabetes,” and Related Estimates (NHANES 2011–2014)	5
4.	State-level Predicted Prevalence across Impact Toolkit Risk Groups (Full Adult Population)	6
5.	National and State-Level Adult Population Estimates for Calculating Risk Group Size (BRFSS 2014)	7
6.	State-level Predicted Prevalence across Impact Toolkit Risk Groups (Adult Employed Population)	11
7.	Industry Group Predicted Prevalence across Impact Toolkit Risk Groups.....	14
8.	Occupation Group Predicted Prevalence across Impact Toolkit Risk Groups.....	15
9.	State-level Predicted Prevalence across Impact Toolkit Risk Groups (Adult Privately Insured Population)	19
10.	Annual Incidence Probability of Diabetes by A1c Category	22
11.	Low, High, and Midpoint Annual Incidence Probability Estimates by A1C from Zhang et al. (2010)	22
12.	Persons with Prediabetes: Calculation of a Weighted Average Annual Probability of Diabetes by A1C Category.....	23
13.	Persons with Prediabetes and Other Persons at Risk for Type 2 Diabetes: Calculation of a Weighted Average Annual Probability of Diabetes by A1C Category	24
14.	Default Intervention Weight Loss and Weight Regain Schedule	33
15.	Estimated Diabetes Attributable Medical Costs by Year and Duration of Diabetes	36

Acknowledgments

The Diabetes Prevention Impact Toolkit and this accompanying Technical Report were developed by RTI International through Contract Number 200-2013-M-53964B, Purchase Order 200-2-14-F-61255 from the Centers for Disease Control and Prevention. The contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention, the Department of Health and Human Services, or the U.S. government.

We would like to acknowledge the following members of the Expert Advisory Panel who provided advice on the technical approach, development, and data sources to be used for the development of the Impact Toolkit: Barbara Bartman, Jonathan Brown, William Herman, Matt Petersen, and Kristi Pier.

1. INTRODUCTION

The Diabetes Prevention Impact Toolkit predicts the health and economic effects of the National Diabetes Prevention Program (National DPP) lifestyle change program on a state, employer, or insurer's population at risk for type 2 diabetes. Specifically, the Impact Toolkit allows users to estimate program costs, diabetes-related medical costs, and return on investment (ROI) along with other cost and health outcome measures. This report describes the methods, equations, and data used by the Impact Toolkit to project outcomes.

The Impact Toolkit has three modules: State, Employer, and Insurer. In general, these modules are very similar. The key difference is that, in the Employer and Insurer modules, the user can choose from a set of predefined populations or enter a customized set of population characteristics. In contrast, the State module only offers state populations for estimating the health and economic effects of the program. The differences between the modules are mainly reflected in the Population Characteristics section of the Input Dashboard. Sections 2.1 through 2.3 describe the Population Characteristics inputs for each module separately. All other sections of this technical report apply to all three modules, except for Section 2.5.7 Productivity Costs, which only applies to the Employer module.

2. INPUT DASHBOARD

For each module (State, Employer, and Insurer), the Population Characteristics section has different features and functions. Section 2.1 covers the technical details of population characteristics for the State module, Section 2.2 covers the Employer module, and Section 2.3 covers the Insurer module. Sections 2.4 and 2.5 cover the technical details associated with additional sections of the Input Dashboard that are the same in each module (e.g., screening assumptions, cost assumptions).

2.1 State Module Population Characteristics

When a state is selected from the dropdown list (or the U.S. map), the Impact Toolkit references a table of state-level data that contains the predicted prevalence for each of the risk groups eligible to participate in the National DPP lifestyle change program. The default risk group is set to "Persons with prediabetes" because the American Diabetes Association (ADA) and other organizations recommend that persons with prediabetes be referred to an intensive diet and physical activity behavioral counseling program (ADA, 2016). The National DPP is an example of such a program. You can also select a larger group ("Persons with prediabetes and other persons at risk for type 2 diabetes"), which has a slightly lower annual probability of developing diabetes, or a smaller group ("Persons with high-risk prediabetes"), which has a higher annual probability of developing diabetes. The larger group ("Persons with prediabetes and other persons at risk for type 2 diabetes") includes everyone from the group "Persons with prediabetes" and additional persons with a high risk

score on the CDC Prediabetes Screening Test (CDC, 2015a). The smaller group (“Persons with high-risk prediabetes”) only includes a high-risk subset of the group “Persons with prediabetes.” All three risk groups include only persons with a body mass index [BMI] ≥ 24 kg/m², which is the eligibility criteria for the National DPP (CDC, 2015b). The user will select one of these three risk groups as shown in Figure 1.

Figure 1. Three Risk Groups Eligible to Participate in National DPP

The screenshot shows a selection interface for a risk group. The title is "Risk Group to Participate in Program". There are three radio button options, each with an information icon (i):

- Persons with prediabetes
- Persons with prediabetes and other persons at risk for type 2 diabetes
- Persons with high-risk prediabetes

Below the options, there is a section for "Annual probability of diabetes:". It features a horizontal bar chart where the value "3.8" is displayed in a light blue box, followed by a dark blue box containing a white percentage sign "%". Below the bar, the text "Suggested Range: 1% to 7%" is shown in a smaller font.

Note: The default risk group is set to “Persons with prediabetes.” The larger group (“Persons with prediabetes and other persons at risk for type 2 diabetes”) includes everyone from the group “Persons with prediabetes” and additional persons with a high risk score on the CDC Prediabetes Screening Test (CDC, 2015a). The smaller group (“Persons with high-risk prediabetes”) only includes a high-risk subset of the group “Persons with prediabetes.” Each group has a different “Annual probability of diabetes.” See Section 2.4 for more information on how the default “Annual probability of diabetes” was calculated for each risk group.

State estimates of the number of people in each of these risk groups were derived using a prediction equation based on age, sex, race/ethnicity, and BMI characteristics. These risk groups were estimated in two steps. First, we estimated an ordered logistic regression model using National Health and Nutrition Examination Survey (NHANES) (2011–2014) data to predict the probability of having prediabetes (prediabetes was defined as fasting plasma glucose [FPG] 100–125 mg/dl or A1C 5.7%–6.4%). The ordered logistic regression approach allowed us to account for various diabetes-related groups, including normal glucose, prediabetes, undiagnosed diabetes, and diagnosed diabetes. Second, we applied

this prediction equation to the sample of state residents in the Behavioral Risk Factor Surveillance System (BRFSS) (2014) for each state. In other words, based on the age, sex, race/ethnicity, and BMI characteristics observed in each state, we predicted the number of people who would have prediabetes and BMI ≥ 24 kg/m². For Asian persons, we allowed a lower BMI threshold (BMI ≥ 22 kg/m²) per CDC's Diabetes Prevention Recognized Program criteria (CDC, 2015b). Persons with diagnosed or undiagnosed diabetes were excluded from all risk groups (persons with diabetes are not eligible for the National DPP).

For the larger risk group, "Persons with prediabetes and other persons at risk for type 2 diabetes," we calculated the CDC Prediabetes Screening Test score for each person in BRFSS (using age, BMI, and physical activity status) to ascertain how many were at risk for diabetes in each state (CDC, 2015a). Persons with a score of 9 or higher are eligible to participate in the National DPP. From this estimate of state residents with a score of 9 or higher, we then subtracted the predicted prevalence of persons with diabetes and added the predicted prevalence of persons from the prediabetes risk group that had a risk score of less than 9. This is necessary because the group of "Persons with prediabetes and other persons at risk for type 2 diabetes" should contain everyone in the prediabetes group even if they had a score of less than 9. A step-by-step description of the algorithm for estimating the state-level predicted prevalence of "Persons with prediabetes and other persons at risk for type 2 diabetes" follows:

1. Generate indicator variable for having a high risk score for each person in the sample (a score of 9 or higher on the CDC Prediabetes Screening Test qualifies as a high score). Estimate the survey weighted mean of this indicator variable for each state's sample of participants. This mean represents an estimate of the proportion of people in the sample with a high risk score.
 - a) We were able to calculate each person's screening test result based on their age, BMI, and physical activity status. We generated a variable for persons with "no physical activity" based on each person's yes or no response to a question about participating in *any* leisure time physical activity. Additional questions from the screening test regarding family history of diabetes and having had a baby that weighs more than 9 pounds at birth are only worth 1 point each and cannot change the outcome of the test (all other questions are worth at least 5 points, and these additional questions cannot increase the score to 9 or higher).
2. Generate state predictions of "Persons with prediabetes" (BMI ≥ 24 kg/m²) who have a *low* risk score (i.e., "Persons with prediabetes" who are not in the "Persons with prediabetes and other persons at risk for type 2 diabetes" group except for the fact that they have prediabetes). This was done by generating predictions of prediabetes status (using the ordered logistic regression model) among persons with a low risk score (less than 9) and a high BMI (BMI ≥ 24 kg/m²). We estimated the survey weighted mean prediction (of prediabetes) for each state's sample of participants.

3. Generate an indicator variable for persons with a high BMI ($BMI \geq 24 \text{ kg/m}^2$) and low risk score (less than 9). Estimate the survey weighted mean for the state's sample of participants.
4. Multiply (2) and (3) to get "Persons with prediabetes" ($BMI \geq 24 \text{ kg/m}^2$) who would not otherwise qualify for the "Persons with prediabetes and other persons at risk for type 2 diabetes" group.
5. Generate state predictions for total diabetes (undiagnosed and diagnosed diabetes) in persons with a high risk score and $BMI \geq 24 \text{ kg/m}^2$ (these persons are not eligible for the National DPP). Estimate the survey weighted mean of these predictions for each state's sample of participants. This mean represents an estimate of the proportion of people in the sample with a high risk score *and* diabetes.
6. Calculate the full state-level risk group, "Persons with prediabetes and other persons at risk for type 2 diabetes" as follows:
 - a. "Persons with prediabetes and other persons at risk for type 2 diabetes" = $\{(1) - [(5) * (1)]\} + (4)$

Finally, the smallest group, "Persons with high-risk prediabetes," was calculated using the predicted prevalence for the "Persons with prediabetes" risk group and the fraction of persons with prediabetes that are considered high risk (34.3%). We used NHANES data (2011–2014) to estimate this fraction, where we defined high-risk prediabetes as having an A1C of 6.0%–6.4% or an FPG of 110–125 mg/dl. As shown in Section 2.4, persons with these levels of A1C or FPG have a higher annual probability of developing diabetes.

National estimates offered in the State module ("UNITED STATES" selection in the dropdown menu) do not use a prediction equation to predict the prevalence of "Persons with prediabetes" because their prediabetes status is obtained directly from the results of the NHANES laboratory data (i.e., A1C or FPG test results). Thus, we used the NHANES data (2011–2014) alone for the U.S. population selection in the State module, whereas state-level data were based on (1) the prediabetes prediction equation estimated in NHANES (2011–2014) and (2) the state-level characteristics observed in BRFSS (2014).

Using the national data from NHANES, we demonstrate the method for parsing the Impact Toolkit's risk groups in Tables 1 through 3. Table 1 shows the division of persons between normal glucose, prediabetes, undiagnosed diabetes, and diagnosed diabetes in NHANES 2011–2014.

Table 1. National Estimates of Prediabetes and Diabetes (NHANES 2011–2014)

	Normal Glucose	Prediabetes	Undiagnosed Diabetes	Diagnosed Diabetes
Estimate	52.5%	35.6%	3.1%	8.8%

Not all persons with prediabetes are eligible for the National DPP. To be eligible, a person with prediabetes must have BMI ≥ 24 kg/m² (≥ 22 kg/m² for Asians). As shown in Table 2, 29.2% of the population has prediabetes and BMI ≥ 24 kg/m².

Table 2. National Estimates of “Persons with Prediabetes” and BMI \geq 24 (NHANES 2011–2014)

Normal Glucose	Prediabetes, BMI <24	Prediabetes, BMI \geq 24	Diagnosed and Undiagnosed Diabetes
52.5%	6.4%	29.2%	11.9%

The National DPP may also serve other persons at risk for diabetes who have BMI ≥ 24 kg/m². Persons without A1C or FPG lab values in the prediabetes range are considered “at risk for type 2 diabetes” if they have a CDC Prediabetes Screening Test score of 9 or higher (CDC, 2015a, 2015b). This additional risk group is shown in Table 3, where the normal glucose population is divided into persons at risk for type 2 diabetes with BMI ≥ 24 kg/m² (12.1%) and those who are not at risk (40.4%). The risk group in the penultimate column of Table 3, BMI ≥ 24 “Prediabetes and other persons at risk for type 2 diabetes,” includes the 12.1% with “Normal glucose, at risk for type 2 diabetes, BMI ≥ 24 ,” and the 29.2% with “Prediabetes, BMI ≥ 24 .”

Table 3. National Estimates of “Persons with Prediabetes,” “Persons with Prediabetes and Other Persons at Risk for Type 2 Diabetes,” and Related Estimates (NHANES 2011–2014)

Normal Glucose, Not at Risk for Type 2 Diabetes ^a	Normal Glucose, at Risk for Type 2 Diabetes, BMI \geq 24 ^b	Prediabetes, BMI <24	Prediabetes, BMI \geq 24	Prediabetes and Other Persons at Risk for Type 2 Diabetes, BMI \geq 24 ^c	Diagnosed and Undiagnosed Diabetes
40.4%	12.1%	6.4%	29.2%	41.3%	11.9%

Note: Persons “at risk for type 2 diabetes” are persons with a score of 9 or higher on the CDC Prediabetes Screening Test (CDC, 2015a). The percentages in this table do not add up to 100% because the “Persons with prediabetes, BMI \geq 24” are subsumed in the “Persons with prediabetes and other persons at risk, BMI \geq 24” estimate (i.e., the percentages add up to 100% if you ignore the “Prediabetes, BMI \geq 24” column and the “Normal Glucose, at Risk for Type 2 Diabetes, BMI \geq 24” column).

^a “Normal glucose, not at risk for type 2 diabetes” excludes persons eligible for the National DPP (“Prediabetes, BMI \geq 24” and “normal glucose, at risk for type 2 diabetes, BMI \geq 24”) and “diagnosed and undiagnosed type 2 diabetes.”

^b Persons with “Normal glucose, at risk for type 2 diabetes, BMI \geq 24” do not have A1C or FPG lab values in the prediabetes range; however, they do have a score of 9 or higher on the CDC Prediabetes Screening Test (CDC, 2015a) based on their diabetes risk factors (e.g., age, BMI, physical activity status).

^c The “prediabetes and other persons at risk for type 2 diabetes, BMI \geq 24” group includes persons with “Prediabetes, BMI \geq 24” and persons with “normal glucose, at risk for type 2 diabetes, BMI \geq 24” (i.e., persons with a score of 9 or higher on the CDC Prediabetes Screening Test) (CDC, 2015a).

Table 4 shows the predicted prevalence of each risk group, for each state. Table 4 is used in the State module to look up the number of people in each risk group in a particular state (i.e., people who are eligible and might participate in a National DPP). The state-level predicted prevalence estimates in Table 4 are multiplied by the total adult population (aged 18 or older) in the selected state (Table 5) to get the total number of people in a given risk group who are eligible for the National DPP. Depending on the screening and participation assumptions (see Sections 2.5.1 and 2.5.2), only a fraction of these eligible state residents will end up participating and reducing their risk of progression to diabetes.

Table 4. State-level Predicted Prevalence across Impact Toolkit Risk Groups (Full Adult Population)

National or State Abbreviation	Prediabetes	Prediabetes and Others at Risk	High-Risk Prediabetes
US (National, NHANES 2011–2014)	29.18%	41.28%	10.00%
AL (State, BRFSS 2014)	30.23%	44.28%	10.36%
AK	29.23%	40.41%	10.01%
AZ	29.71%	42.03%	10.18%
AR	30.73%	45.60%	10.53%
CA	30.04%	40.77%	10.29%
CO	26.21%	37.16%	8.98%
CT	29.17%	41.53%	9.99%
DE	31.05%	44.62%	10.64%
DC	25.49%	34.20%	8.73%
FL	29.99%	42.83%	10.27%
GA	30.08%	42.49%	10.30%
HI	30.62%	39.25%	10.49%
ID	28.48%	41.43%	9.76%
IL	28.83%	41.64%	9.88%
IN	29.51%	43.92%	10.11%
IA	29.32%	43.23%	10.05%
KS	29.53%	43.21%	10.12%
KY	29.23%	44.04%	10.01%
LA	30.98%	44.87%	10.61%
ME	28.83%	42.79%	9.88%
MD	30.65%	42.95%	10.50%
MA	27.64%	39.36%	9.47%
MI	29.49%	43.55%	10.10%
MN	28.86%	42.11%	9.89%
MS	31.64%	46.42%	10.84%
MO	29.13%	43.10%	9.98%
MT	28.12%	40.77%	9.63%

(continued)

Table 4. State-level Predicted Prevalence across Impact Toolkit Risk Groups (Full Adult Population) (continued)

National or State Abbreviation	Prediabetes	Prediabetes and Others at Risk	High-Risk Prediabetes
NE	29.21%	42.78%	10.01%
NV	29.80%	42.26%	10.21%
NH	29.02%	42.53%	9.94%
NJ	30.58%	43.30%	10.48%
NM	30.55%	42.89%	10.46%
NY	29.48%	42.30%	10.10%
NC	30.22%	43.52%	10.35%
ND	29.66%	43.73%	10.16%
OH	29.91%	44.16%	10.25%
OK	30.42%	44.64%	10.42%
OR	28.40%	39.97%	9.73%
PA	29.71%	43.70%	10.18%
RI	28.91%	42.16%	9.90%
SC	30.40%	43.77%	10.41%
SD	28.99%	42.86%	9.93%
TN	30.02%	44.12%	10.28%
TX	31.25%	44.83%	10.71%
UT	25.81%	36.72%	8.84%
VT	27.29%	40.44%	9.35%
VA	29.41%	42.25%	10.08%
WA	29.37%	41.77%	10.06%
WV	30.50%	46.19%	10.45%
WI	29.91%	43.59%	10.25%
WY	28.48%	41.81%	9.76%

Table 5. National and State-Level Adult Population Estimates for Calculating Risk Group Size (BRFSS 2014)

National or State Abbreviation	National or State Population Estimate ^a
US	245,561,099
AL	3,739,646
AK	556,360
AZ	5,091,417
AR	2,266,396
CA	29,544,655
CO	4,115,447

(continued)

Table 5. National and State-Level Adult Population Estimates for Calculating Risk Group Size (BRFSS 2014) (continued)

National or State Abbreviation	National or State Population Estimate ^a
CT	2,832,225
DE	730,755
DC	545,460
FL	15,832,660
GA	7,623,372
HI	1,112,388
ID	1,204,877
IL	9,888,842
IN	5,030,005
IA	2,386,030
KS	2,186,730
KY	3,402,842
LA	3,537,716
ME	1,068,811
MD	4,649,776
MA	5,365,728
MI	7,693,748
MN	4,191,574
MS	2,260,730
MO	4,673,556
MT	802,869
NE	1,417,407
NV	2,166,196
NH	1,061,487
NJ	6,949,942
NM	1,579,709
NY	15,519,718
NC	7,682,975
ND	583,766
OH	8,968,842
OK	2,944,523
OR	3,109,293
PA	10,099,122
RI	839,958

(continued)

Table 5. National and State-Level Adult Population Estimates for Calculating Risk Group Size (BRFSS 2014) (continued)

National or State Abbreviation	National or State Population Estimate ^a
SC	3,749,025
SD	649,956
TN	5,067,014
TX	19,900,570
UT	2,068,310
VT	506,408
VA	6,499,147
WA	5,475,871
WV	1,474,021
WI	4,459,989
WY	453,235

^a Weighted estimates from BRFSS 2014 (noninstitutionalized adult population aged 18 or older). National estimate is the sum of the state-level estimates.

In the State module, we used a table of the national and state-level adult population (noninstitutionalized, as reported in BRFSS 2014) to estimate the number of people eligible for a National DPP in each state. These population estimates are based on the survey-weighted BRFSS 2014 population counts (see Table 5). The State module assumes that the entire eligible state population is being offered the National DPP. Thus, population estimates cannot be changed in the State module. However, in the Employer module (Section 2.2) and Insurer module (Section 2.3), you can enter smaller or larger population sizes to represent the approximate size of an employee population or insured adult population.

2.2 Employer Module Population Characteristics

Before selecting a radio button to designate your population’s characteristics, you will first enter your population size (i.e., “Number of employees” in the Employer module). This number should include all employees, not just those participating in the National DPP lifestyle change program. The default number of employees is set to 1,000. Entering a number of employees less than or equal to 11 (under the default settings) will lead to a result of 0 projected participants. Under the default settings, a minimum of 12 people is required to get at least 1 participant. Having a low number of participants can lead to fractional results, such as 0.01 cases averted.

Once you have entered the total number of employees at a firm, you then will select a radio button to determine the population characteristics of your employee population. There are several predefined sets of population characteristics as well as an option to enter your own

employee population characteristics (age, race/ethnicity, sex, BMI). The proportion of the population that is eligible for the National DPP will be calculated using the set of population characteristics defined here and a prediction equation estimated in NHANES (2011–2014).

When selecting one of the first four radio buttons for predefined sets of population characteristics (“Assume national average for population characteristics” or “Assume state average for population characteristics” or “Assume industry average for population characteristics” or “Assume occupation average for population characteristics”) in the Employer module, the Impact Toolkit references a lookup table of predicted prevalence estimates for each of the risk groups eligible to participate in a National DPP. Sections 2.2.1 through 2.2.4 describe the data and methods underlying each set of predefined population characteristics. The default risk group is set to “Persons with prediabetes” because the American Diabetes Association (ADA) and other organizations recommend that persons with prediabetes be referred to an intensive diet and physical activity behavioral counseling program (ADA, 2016). The National DPP is an example of such a program. You can also select a larger group (“Persons with prediabetes and other persons at risk for type 2 diabetes”), which has a slightly lower annual probability of developing diabetes, or a smaller group (“Persons with high-risk prediabetes”), which has a higher probability of developing diabetes. All three risk groups only include persons who are eligible for the National DPP (BMI \geq 24 kg/m²).

Unlike the first four radio button selections that use lookup tables, when you select the radio button “Enter employee characteristics,” the predicted prevalence of each risk group is calculated based on a prediction equation and other assumptions. See Section 2.2.5 for the data and methods related to implementing the prediction equation when you enter your own employee population characteristics.

2.2.1 Assume National Average for Population Characteristics

National average population characteristics in the Employer module are survey-weighted estimates from the sample of U.S. employed persons in NHANES (2011–2014). National estimates offered in the Employer module do not use a prediction equation to predict the prevalence of “Persons with prediabetes” because their prediabetes status is obtained directly from the results of the NHANES laboratory data (i.e., A1C or FPG test results). Thus, we used the NHANES data (2011–2014) alone for the national selection in the employer module, whereas state-level data are based on (1) the prediabetes prediction equation estimated in NHANES (2011–2014) and (2) the state-level characteristics observed in BRFSS (2014). See Section 2.1 and Tables 1 through 3 to see how the risk groups were parsed from the national data.

2.2.2 Assume State Average for Population Characteristics

State estimates of the number of people in each risk group were estimated using the methods described in Section 2.1. The only difference in the Employer module is that risk group estimates were generated for the sample of *employed* adults in BRFSS (2014). Table 6 presents the predicted prevalence of each risk group, for each state’s population of employed adults. This lookup table is used in the Employer module to estimate the number of people in each risk group in a particular state’s employed population (i.e., people that are eligible and might participate in a National DPP). The state predicted prevalence estimates in Table 6 are multiplied by the “Number of Employees” (entered by the user) to get the total number of people in a given risk group that are eligible for the National DPP. Depending on your screening and participation assumptions (see Section 2.5.1 and 2.5.2), only a fraction of these eligible employed adults will end up participating and reducing their risk of progression to diabetes.

Table 6. State-level Predicted Prevalence across Impact Toolkit Risk Groups (Adult Employed Population)

National or State Abbreviation	Prediabetes	Prediabetes and Others at Risk	High Risk Prediabetes
US (National, NHANES)	26.95%	39.77%	9.23%
AL (State, BRFSS)	28.31%	40.95%	9.70%
AK	28.21%	38.50%	9.66%
AZ	27.76%	38.78%	9.51%
AR	29.35%	43.07%	10.06%
CA	29.26%	39.09%	10.02%
CO	25.11%	35.10%	8.60%
CT	28.40%	39.98%	9.73%
DE	29.61%	42.60%	10.14%
DC	22.27%	29.36%	7.63%
FL	28.06%	39.46%	9.61%
GA	28.94%	40.44%	9.92%
HI	30.55%	38.86%	10.47%
ID	27.19%	39.55%	9.32%
IL	27.59%	39.43%	9.45%
IN	27.92%	41.94%	9.56%
IA	27.63%	41.41%	9.47%
KS	28.35%	41.28%	9.71%
KY	26.61%	39.47%	9.12%
LA	29.63%	42.54%	10.15%
ME	26.99%	40.30%	9.25%
MD	29.90%	41.61%	10.24%
MA	26.45%	37.62%	9.06%

(continued)

Table 6. State-level Predicted Prevalence across Impact Toolkit Risk Groups (Adult Employed Population) (continued)

National or State Abbreviation	Prediabetes	Prediabetes and Others at Risk	High Risk Prediabetes
MI	27.30%	40.10%	9.35%
MN	27.32%	40.02%	9.36%
MS	30.40%	43.98%	10.42%
MO	27.40%	40.64%	9.39%
MT	26.42%	38.37%	9.05%
NE	27.91%	41.03%	9.56%
NV	28.68%	39.85%	9.83%
NH	27.43%	39.41%	9.40%
NJ	30.00%	42.33%	10.28%
NM	29.51%	41.09%	10.11%
NY	27.65%	39.36%	9.47%
NC	28.77%	40.72%	9.86%
ND	28.59%	42.61%	9.79%
OH	27.98%	41.00%	9.58%
OK	28.74%	41.61%	9.85%
OR	26.53%	36.41%	9.09%
PA	27.87%	41.18%	9.55%
RI	27.51%	39.72%	9.42%
SC	28.84%	41.00%	9.88%
SD	27.64%	40.93%	9.47%
TN	28.20%	39.83%	9.66%
TX	30.45%	43.36%	10.43%
UT	25.01%	35.40%	8.57%
VT	25.41%	37.58%	8.71%
VA	28.56%	40.58%	9.79%
WA	28.29%	39.80%	9.69%
WV	28.32%	42.59%	9.70%
WI	28.24%	41.51%	9.67%
WY	26.65%	39.02%	9.13%

2.2.3 Assume Industry Average for Population Characteristics

Industry characteristics are based on the sample of employed persons in NHANES (2011–2014) and the relative rates of obesity across industries. Industries are defined in NHANES using the U.S. Census Bureau’s 2002 Occupation and Industry coding system (U.S. Census, 2016). The U.S. Census Bureau defines industry as “the type of activity at a person’s place of work.” To estimate the number of people in each risk group, we started by estimating a baseline employed group using the mean values of each population characteristic (age, sex,

race/ethnicity, and BMI) observed in NHANES (2011–2014) for employed adults. BMI was modeled as a categorical variable with the following categories: normal weight, overweight, and obese. Overweight was defined as a BMI 24.0–29.9 kg/m² to match the National DPP criteria. Obesity was defined as BMI \geq 30 kg/m².

An analysis of National Health Interview Survey (NHIS) data by Luckhaupt and colleagues (2014) provided us with obesity prevalence ratios by industry. In other words, workers in some industries have a higher risk of obesity than the average across all industries, and Luckhaupt et al. quantified this in a prevalence ratio (controlling for sociodemographic factors, such as age and race/ethnicity). If a particular industry has a prevalence ratio of 1.0, then people in that industry have the same risk of obesity as the average person across all industries. Prevalence ratios greater than 1.0 signal a higher risk of obesity, whereas prevalence ratios less than 1.0 signal a lower risk of obesity. We used these prevalence ratios to inflate the percentage of people who are obese in a given industry according to the prevalence ratio associated with that industry from Luckhaupt et al. (2014).

To implement this approach, we also had to assume that as the percentage of obesity increased for a given industry population, the percentage overweight also increased, but to a smaller degree. We made this assumption because more than three-quarters of the employed population observed in NHANES (2011–2014) is overweight or obese, and some prevalence ratios were large enough that allowing a one to one increase in the percentage overweight (as the obesity percentage increases) led to a population with more than 100% being overweight or obese. Thus, we used NHANES data to estimate the marginal effect of the probability of obesity on the probability of overweight in the employed population (we used an ordinary least squares approach for this estimate). We found that for every one percentage point increase in the probability of being obese, there was a 0.63 percentage point increase in the probability of being overweight.

Using the prediction equation for prediabetes, we estimated the probability of having prediabetes for overweight employed people and for obese employed people. We used the mean values for other characteristics included in the prediction equation (age, sex, and race/ethnicity). The predicted probabilities of prediabetes for overweight people and obese people were then combined according to their relative weights. For instance, in the baseline employed group (prevalence ratio = 1.0), 33.9% of employed persons were obese and 41.6% were overweight (and the remaining 24.5% were normal weight or underweight). A prediction of the probability of prediabetes was generated for the obese group and the overweight group, and then these groups received weights of 33.9% and 41.6%, respectively. We did not calculate a probability of prediabetes for the 24.5% of the population that was assumed to be normal weight because they are not eligible for the National DPP. They effectively have a probability of 0.

The steps described above help us calculate the predicted prevalence of “Persons with prediabetes” (and BMI \geq 24 kg/m²), but do not help us calculate the larger risk group “Persons with prediabetes and other persons at risk for type 2 diabetes” and the smaller risk group “Persons with high-risk prediabetes.” To calculate the “Persons with prediabetes and other persons at risk for type 2 diabetes” group, we assumed that this larger group was about 1.5 times as large as the “Persons with prediabetes” group that was estimated for each industry using the methods above. We assumed that this relationship was 1.48 based on a comparison of the “Persons with prediabetes” group and the “Persons with prediabetes and other persons at risk for type 2 diabetes” group in the NHANES data (2011–2014) among employed people. Similarly, we assumed that the smaller risk group (“Persons with high-risk prediabetes”) was 0.34 times the size of the “Persons with prediabetes” group based on the relationship between these groups observed in the NHANES data (2011–2014) among employed people. The resulting predicted prevalence of each risk group is shown in Table 7.

Table 7. Industry Group Predicted Prevalence across Impact Toolkit Risk Groups

Industry Group	Prediabetes	Prediabetes and Others at Risk	High Risk Prediabetes
Agriculture, forestry, fishing, and hunting	29.06%	43.01%	9.96%
Mining	31.80%	47.07%	10.90%
Utilities	27.46%	40.64%	9.41%
Construction	29.06%	43.01%	9.96%
Manufacturing	29.75%	44.02%	10.19%
Wholesale trade	26.55%	39.29%	9.10%
Retail trade	27.69%	40.98%	9.49%
Transportation and warehousing	30.20%	44.70%	10.35%
Information	32.94%	48.76%	11.29%
Finance and insurance	27.92%	41.32%	9.56%
Real estate, rental and leasing	22.21%	32.87%	7.61%
Professional, scientific, and technical services	26.32%	38.95%	9.02%
Management of companies and enterprises	29.29%	43.35%	10.03%
Administrative, support, waste management, and remediation services	29.06%	43.01%	9.96%
Education services	30.66%	45.38%	10.50%
Healthcare and social assistance	33.40%	49.43%	11.44%
Arts, entertainment, and recreation	24.04%	35.57%	8.23%
Accommodation and food services	26.09%	38.61%	8.94%
Other services (except public administration)	26.32%	38.95%	9.02%
Public administration	35.46%	52.48%	12.15%

Note: Industry groups are defined by the U.S. Census Bureau’s 2002 Indexes of Industry and Occupation (U.S. Census, 2016).

2.2.4 Assume Occupation Average for Population Characteristics

Occupation characteristics are based on the sample of employed persons in NHANES (2011–2014) and the relative rates of obesity across occupations. Occupations are defined in NHANES using the U.S. Census Bureau’s 2002 Occupation and Industry coding system (U.S. Census, 2016). The U.S. Census Bureau defines occupation as the “kind of work a person does to earn a living.”

The methods for estimating the predicted prevalence of each risk group for an occupation are identical to the methods for the industry groups described in Section 2.2.3. The prevalence ratios associated with occupation groups from Luckhaupt et al. (2014) were used to implement the same approach, and the predicted prevalence of each risk group is shown in Table 8.

Table 8. Occupation Group Predicted Prevalence across Impact Toolkit Risk Groups

Occupation Group	Prediabetes	Prediabetes and Others at Risk	High-risk Prediabetes
Management	31.12%	46.05%	10.66%
Business and financial operations	29.75%	44.02%	10.19%
Computer and mathematical	29.29%	43.35%	10.03%
Architecture and engineering	37.51%	55.52%	12.85%
Life, physical, and social science	23.12%	34.22%	7.92%
Community and social service	36.14%	53.49%	12.38%
Legal	25.86%	38.28%	8.86%
Education, training, and library	30.66%	45.38%	10.50%
Art, design, entertainment, sports, and media	24.72%	36.59%	8.47%
Healthcare practitioners and technical	26.55%	39.29%	9.10%
Healthcare support	32.94%	48.76%	11.29%
Protective service	34.77%	51.46%	11.91%
Food preparation and serving related	25.86%	38.28%	8.86%
Building and ground cleaning and maintenance	23.35%	34.56%	8.00%
Personal care and service	32.49%	48.08%	11.13%
Sales and related	26.32%	38.95%	9.02%
Office and administrative support	31.80%	47.07%	10.90%
Farming, fishing, and forestry	29.52%	43.69%	10.11%
Construction and extraction	26.55%	39.29%	9.10%
Installation, maintenance, and repair	27.46%	40.64%	9.41%
Production	28.83%	42.67%	9.88%
Transportation and material moving	30.66%	45.38%	10.50%

Note: Occupation groups are defined by the U.S. Census Bureau’s 2002 Indexes of Industry and Occupation (U.S. Census, 2016).

2.2.5 Entering Employee Population Characteristics

For customized results based on your organization's unique population characteristics, select the "Enter employee characteristics" option. When this option is selected, you will see that the fields for each characteristic have already been filled with default values. These default values reflect national averages for the employed population in the United States. You can change these to reflect your own employee population characteristics. If you want to return to the default values at any point, click the "RESTORE DEFAULTS" link in the upper right-hand corner. This button will clear all of the data that you have entered and restore the default data.

The values entered in the employee characteristics fields are used to predict the percentage of employees in each risk group (see Section 2.4 for more information on the risk groups). The first step to predicting each of the risk groups is to predict the prevalence of the primary risk group, "Persons with prediabetes." The other two risk groups are predicted indirectly, using multipliers to inflate or deflate the prediction for "Persons with prediabetes" based on the relative size of the other two risk groups observed in NHANES data (2011–2014). To predict the percentage of employees with prediabetes, we estimated an ordered logistic regression model NHANES (2011–2014) to account for other diabetes-related outcomes, including normal glucose, prediabetes, undiagnosed diabetes, and diagnosed diabetes. The model included independent variables for age, sex, race/ethnicity, and BMI. BMI was modeled as a categorical variable with the following categories: normal weight, overweight, and obese. Overweight was defined as a BMI 24.0–29.9 kg/m² to match the National DPP criteria. Obesity was defined as BMI \geq 30 kg/m².

To limit the burden of data collection and data entry on the user, we developed an ad hoc method to apply this prediction model using the set of population characteristic averages entered by the user. Instead of asking the user to provide a person-level dataset of their employee population, we generated a predicted prevalence of "Persons with prediabetes" based on the characteristic averages with special emphasis on two key predictive variables: overweight and obesity status. Using the prediction model for prediabetes, we estimated two separate predicted prevalence estimates of "Persons with prediabetes": one for overweight persons and another for obese persons. These two predicted prevalence estimates are combined in a weighted average to estimate the predicted prevalence of "Persons with prediabetes" in your employee population. We did not calculate a predicted prevalence of "Persons with prediabetes" for the proportion of the population that is normal weight because they are not eligible for the National DPP. They effectively have a probability of 0.

The following example illustrates this method. Using the default employee population characteristics in the Employer module (age, sex, and race/ethnicity), we predicted the prevalence of "Persons with prediabetes" for overweight and obese persons separately. The

predicted prevalence of “Persons with prediabetes” for each BMI group was then combined according to their relative proportions in the employee population. In the default employee population characteristics, 33.9% of employed persons were obese, and 41.6% were overweight (and 24.5% were normal weight, which effectively have a probability of 0). A weighted average predicted prevalence of “Persons with prediabetes” was calculated based on these weights and represents the proportion of your employee population that is predicted to have prediabetes and $BMI \geq 24 \text{ kg/m}^2$.

The steps described above give us the predicted prevalence of “Persons with prediabetes” (and $BMI \geq 24 \text{ kg/m}^2$). Refer to the last paragraph in Section 2.2.3 for more information on how we calculated the larger risk group “Persons with prediabetes and other persons at risk for type 2 diabetes” and the smaller risk group “Persons with high-risk prediabetes” using the relative size of these risk groups in NHANES (2011–2014).

2.3 Insurer Module Population Characteristics

Before selecting a radio button to designate your population’s characteristics, you will first enter your population size (i.e., “Number of Adults Insured” in the Insurer module). This number should include all privately insured adults, not just those participating in the prevention program. The default number of privately insured adults is set to 1,000.

Once you have entered the total number of privately insured adults in your population, you will then select a radio button to determine the population characteristics of your population. There are several predefined sets of population characteristics as well as an option to enter your own privately insured adult population characteristics (age, race/ethnicity, sex, BMI). The proportion of the population that is eligible for the National DPP will be calculated using the set of population characteristics defined here and a prediction equation estimated in NHANES (2011–2014).

When selecting one of the first four radio buttons (“Assume national average for population characteristics” or “Assume state average for population characteristics” or “Assume industry average for population characteristics” or “Assume occupation average for population characteristics”) in the Insurer module, the Impact Toolkit references a lookup table of predicted prevalence estimates for each of the risk groups eligible to participate in a National DPP. Sections 2.3.1 through 2.3.4 describe the data and methods underlying each set of predefined population characteristics. The default risk group is set to “Persons with Prediabetes.” because the American Diabetes Association (ADA) and other organizations recommend that persons with prediabetes be referred to an intensive diet and physical activity behavioral counseling program (ADA, 2016). The National DPP is an example of such a program. You can also select a larger group (“Persons with prediabetes and other persons at risk for type 2 diabetes”), which has a slightly lower annual probability of developing diabetes, or a smaller group (“Persons with high-risk prediabetes”), which has a

higher annual probability of developing diabetes. All three risk groups only include persons who are eligible for the National DPP (BMI \geq 24 kg/m²).

Unlike the first four radio button selections that use lookup tables, when you select the radio button “Enter insured adult population characteristics,” the predicted prevalence of each risk group is calculated based on a prediction equation and other assumptions. See Section 2.3.5 for the data and methods related to implementing the prediction equation when you enter your own privately insured adult population characteristics.

2.3.1 Assume National Average for Population Characteristics

National average population characteristics in the Insurer module are survey-weighted estimates from the sample of U.S. privately insured persons in NHANES (2011–2014). National estimates offered in the Insurer module do not use a prediction equation to predict the prevalence of “Persons with prediabetes” because their prediabetes status is obtained directly from the results of the NHANES laboratory data (i.e., A1C or FPG test results). Thus, we used the observed NHANES data (2011–2014) alone for the national selection in the Insurer module, whereas state-level data are based on (1) the prediabetes prediction equation estimated in NHANES (2011–2014) and (2) the state-level characteristics among privately insured persons in BRFSS (2014). See Section 2.1 and Tables 1 through 3 to see how the risk groups were parsed from the national data.

2.3.2 Assume State Average for Population Characteristics

State estimates of the number of people in each risk group were obtained using the methods described in Section 2.1. The only difference in the Insurer module is that risk group estimates were generated for the sample of *privately insured* adults in BRFSS (2014).

Table 9 presents the predicted prevalence of each risk group, for each state’s population of privately insured adults. This lookup table is used in the Insurer module to estimate the number of people in each risk group in a particular state’s privately insured adult population (i.e., people who are eligible and might participate in a National DPP). The state predicted prevalence estimates in Table 9 are multiplied by the “Number of Adults Insured” (entered by the user) to get the total number of people in a given risk group that are eligible for the National DPP. Depending on your screening and participation assumptions (see Sections 2.5.1 and 2.5.2), only a fraction of these eligible insured adults will end up participating and reducing their risk of progression to diabetes.

Table 9. State-level Predicted Prevalence across Impact Toolkit Risk Groups (Adult Privately Insured Population)

National or State Abbreviation	Prediabetes	Prediabetes and Others at Risk	High Risk Prediabetes
US (National, NHANES)	26.56%	38.31%	9.10%
AL (State, BRFSS)	29.69%	43.17%	10.17%
AK	28.44%	39.78%	9.74%
AZ	28.00%	39.49%	9.59%
AR	29.35%	43.07%	10.06%
CA	29.26%	39.09%	10.02%
CO	25.11%	35.10%	8.60%
CT	28.04%	39.71%	9.61%
DE	29.65%	42.83%	10.16%
DC	22.30%	28.81%	7.64%
FL	30.07%	42.51%	10.30%
GA	28.94%	40.44%	9.92%
HI	30.55%	38.86%	10.47%
ID	27.09%	39.50%	9.28%
IL	28.30%	41.12%	9.69%
IN	27.92%	41.94%	9.56%
IA	27.82%	41.52%	9.53%
KS	28.35%	41.28%	9.71%
KY	27.92%	42.08%	9.57%
LA	30.63%	43.99%	10.49%
ME	26.99%	40.30%	9.25%
MD	30.96%	43.48%	10.61%
MA	25.39%	35.87%	8.70%
MI	28.71%	42.31%	9.83%
MN	27.71%	40.43%	9.49%
MS	30.92%	44.69%	10.59%
MO	27.40%	40.64%	9.39%
MT	25.99%	37.81%	8.90%
NE	27.62%	41.00%	9.46%
NV	28.80%	40.80%	9.87%
NH	27.79%	40.52%	9.52%
NJ	29.67%	42.05%	10.16%
NM	29.28%	41.19%	10.03%

(continued)

Table 9. State-level Predicted Prevalence across Impact Toolkit Risk Groups (Adult Privately Insured Population) (continued)

National or State Abbreviation	Prediabetes	Prediabetes and Others at Risk	High Risk Prediabetes
NY	29.25%	42.06%	10.02%
NC	29.17%	41.87%	9.99%
ND	28.20%	42.21%	9.66%
OH	29.06%	42.96%	9.96%
OK	28.74%	41.61%	9.85%
OR	27.49%	38.58%	9.42%
PA	28.98%	42.91%	9.93%
RI	28.03%	41.01%	9.60%
SC	29.86%	42.81%	10.23%
SD	27.64%	40.93%	9.47%
TN	30.63%	44.24%	10.49%
TX	30.45%	43.36%	10.43%
UT	24.65%	34.99%	8.44%
VT	25.57%	37.79%	8.76%
VA	28.97%	41.11%	9.93%
WA	28.35%	40.43%	9.71%
WV	29.84%	45.41%	10.22%
WI	28.47%	41.83%	9.75%
WY	26.65%	39.02%	9.13%

2.3.3 Assume Industry Average for Population Characteristics

Industry characteristics in the Insurer module are based on the sample of *employed* persons in NHANES (2011–2014) and the relative rates of obesity across industries. We used the sample of *employed* persons for this selection in the *Insurer* module because insured adults who work in a particular industry are employed by definition. Industries are defined in NHANES using the U.S. Census Bureau’s 2002 Occupation and Industry coding system.

The methods for estimating the predicted prevalence of each risk group for an industry are identical to the methods described in Section 2.2.3. The predicted prevalence of each risk group by industry is shown in Table 7.

2.3.4 Assume Occupation Average for Population Characteristics

Occupation characteristics in the Insurer module are based on the sample of *employed* persons in NHANES (2011–2014) and the relative rates of obesity across occupations. We

used the sample of *employed* persons for this selection in the *Insurer* module because insured adults who work in a particular occupation are employed by definition. Occupations are defined in NHANES using the U.S. Census Bureau's 2002 Occupation and Industry coding system.

The methods for estimating the predicted prevalence of each risk group for an industry are identical to the methods described in Section 2.2.4. The predicted prevalence of each risk group by occupation is shown in Table 8.

2.3.5 Entering Insured Population Characteristics

For customized results based on your unique insured population's characteristics, select the "Enter insured adult population characteristics" option. When this option is selected, you will see that the fields for each characteristic have already been filled with default values. These default values reflect national averages for the privately insured adult population in the United States. You can change these to reflect your own privately insured adult population characteristics. If you want to return to the default values at any point, click the "RESTORE DEFAULTS" link in the upper right-hand corner. This button will clear all of the data that you have entered and restore the default data.

The values entered in the insured adult population characteristics fields are used to predict the percentage of insured adults in each risk group. The methods for generating the predicted prevalence of each risk group are identical to the methods described in Section 2.2.5 for the Employer module. The resulting predicted prevalence of the risk group you select will be reported in the Output Dashboard.

2.4 Diabetes Incidence Rates for Each Risk Group

Defining the population characteristics allows prediction of the prevalence of each risk group at the state, employee, and privately insured adult population level. Using the radio buttons under the "Risk Group to Participate in Program" heading, you can select from the three risk groups eligible for the National DPP ("Persons with prediabetes," "Persons with prediabetes and other persons at risk for type 2 diabetes," and "Persons with high-risk prediabetes" [see Sections 2.4.1, 2.4.2, and 2.4.3, respectively]). The annual probability of developing diabetes is different for each risk group. The default annual probability for each risk group is prepopulated in the input box. You can change this number if you would like to assume a higher or lower probability of diabetes. We recommend using the default annual probabilities of diabetes unless you have generated inputs from your own data sources or identified updated data available in the scientific literature.

We determined the diabetes incidence rates for each risk group using two key sources: (1) Selvin et al.'s (2010) analysis of Atherosclerosis Risk in Communities study (ARIC) data and (2) a systematic review of annual diabetes probabilities (in the prediabetes range) by Zhang and colleagues (2010). We calculated a weighted average incidence across the A1C

categories presented by Selvin et al. and Zhang et al. (Table 10, green cells). The Zhang et al. review included studies with a combined sample size roughly 4 times larger than Selvin et al.'s ARIC sample. Thus, the Zhang et al. review received a weight of about 80%, and the Selvin et al. analysis received a weight of about 20%.

Table 10. Annual Incidence Probability of Diabetes by A1c Category

Source	Total N	<5.0%	5.0–5.5%	5.5–5.9%	6.0–6.4%
Selvin et al. (2010)	11,092	0.0046	0.0087	0.0166	0.0399
Zhang et al. (2010)	44,203	0.0030	0.0105	0.0340	0.0725
Weighted average	55,295	0.0033	0.0101	0.0305	0.0660
Rounded (%)	55,295	0.3%	1.0%	3.1%	6.6%
Unweighted average	55,295	0.0038	0.0096	0.0253	0.0562
Rounded (%)	55,295	0.4%	1.0%	2.5%	5.6%

Note: Weighted average was calculated based on the relative sample sizes of each study (see Total N column). We calculated the midpoint of the annualized incidence probabilities presented in Zhang et al. (2010) (see Table 11).

We used the midpoint of the low and high estimates (Table 11) reported for each A1C category in the Zhang et al. (2010) review. The average age was slightly older in Selvin et al. (2010) (56.7) than in Zhang et al. (2010) (53.4), and the race/ethnicity makeup differed substantially. Several studies in Zhang et al. (2010) were from Asian and American Indian populations, which were not well represented in Selvin et al.'s ARIC sample (U.S. population, 78% white, 22% black). From the data in Selvin et al. (2010) and Zhang et al. (2010), we determined that a plausible range for annual diabetes incidence was 1% to 7% for these National DPP eligible risk groups. Just under the "Annual Probability of Diabetes" entry field in the Impact Toolkit, we state this recommended range. Using values outside of this range may lead to results with low credibility.

Table 11. Low, High, and Midpoint Annual Incidence Probability Estimates by A1C from Zhang et al. (2010)

Estimate from Zhang et al. (2010)	A1C <5.0%	A1C 5.0–5.5%	A1C 5.5–5.9%	A1C 6.0–6.4%
Zhang et al. (2010) Low	*0.0030	0.0030	0.0180	0.0500
Zhang et al. (2010) High	0.0030	0.0180	0.0500	0.0950
Midpoint	0.0030	0.0105	0.0340	0.0725
Rounded (%)	0.3%	1.1%	3.4%	7.3%

Note: We assumed that the low for A1C <5.0 was the same as the high estimate. At this low value, few studies provided data. In Table 10, we can see that Selvin et al.'s estimate is actually higher than Zhang et al.'s for this low A1C category (the only A1C category in which this occurs).

The accepted range for prediabetes is 5.7% to 6.4% using the A1C blood test or 100 to 125 mg/dl using the FPG blood test (CDC National Diabetes Statistics, 2014). The two highest A1C categories reported in Zhang et al. (2010) and Selvin et al. (2010) are 5.5% to 5.9% and 6.0% to 6.4%, which cover the full range of prediabetes (5.7% to 6.4%) as measured by the A1C blood test. We used the weighted average diabetes incidence probabilities from Table 10 for 5.5% to 5.9% (3.1%) and 6.0% to 6.4% (6.6%) to calculate an annual incidence probability for each risk group. The weighted average annual probabilities of diabetes from Table 10 for <5.0% (3.3%) and 5.0% to 5.5% (1.0%) were not used to determine risk group annual incidence probabilities because these A1C values do not fall in the prediabetes range (5.7% to 6.4%). Some people qualified for the risk group “persons with prediabetes and other persons at risk for type 2 diabetes” based on the CDC Prediabetes Screening Test score and may have had an A1C value less than 5.7%. For these people, we assumed that their annual probability of diabetes was equal to 3.1% (the annual probability for persons with an A1C of 5.5%–5.9%) (see Table 10).

2.4.1 Persons with Prediabetes

This group includes persons predicted to have prediabetes (and BMI ≥ 24 kg/m²). “Persons with prediabetes” have a blood sugar level higher than normal, but not high enough for a diagnosis of diabetes (FPG 100–125 mg/dl or A1C 5.7%–6.4%). The default annual probability of diabetes for persons with prediabetes is 3.8%. This was calculated as a weighted average using (1) the distribution of the risk group (persons with prediabetes) across A1C categories as the weight (see Table 12, column 2) and (2) the weighted average annual probability for persons with an A1C of 5.5% to 5.9% (3.1%) and 6.0% to 6.4% (6.6%) (see Table 10) for the annual probabilities of diabetes that are being weighted.

Table 12. Persons with Prediabetes: Calculation of a Weighted Average Annual Probability of Diabetes by A1C Category

A1C Categories	Distribution of Risk Group Across A1C Categories	Weighted Average Annual Probability	Annual Probabilities Weighted by A1C Category
<5.7%–5.9% ^a	79.8%	3.1%	2.4%
6.0%–6.4%	20.2%	6.6%	1.3%
			Sum of Weighted Probabilities: 3.8%

Note: The distribution of the “Persons with prediabetes” risk group across A1C categories is based on a survey-weighted estimate from NHANES (2011–2014) for national estimates. The “Sum of Weighted Probabilities” is 3.8% (and not 3.7%) due to rounding of the weighted probabilities for each A1C category.

^a Some people qualified as having prediabetes based on their FPG test results, but had an A1C < 5.7%.

2.4.2 Persons with Prediabetes and Other Persons at Risk for Type 2 Diabetes

This is the largest group, as it includes all “Persons with prediabetes” and “other persons at risk for type 2 diabetes.” This group is generally about 1.5 times larger than the risk group “persons with prediabetes.” To enlarge the “persons with prediabetes” risk group, we identified the additional people in the “...other persons at risk for type 2 diabetes” group using criteria from the CDC Prediabetes Screening Test (CDC, 2015b). A score of 9 or higher on the Prediabetes Screening Test determines whether a person is at risk for diabetes. See Section 2.1 for a description of these risk groups and how they were calculated in the State module (some slight differences in methods are noted in Sections 2.2 and 2.3 for the Employer module and Insurer module, respectively). Because this group includes a broader range of persons at risk for diabetes, the default annual probability of diabetes is slightly lower at 3.6%. This probability was calculated as a weighted average using (1) the distribution of the risk group (“persons with prediabetes and other persons at risk for type 2 diabetes”) across A1C categories as the weight (see Table 13, column 2) and (2) the weighted average annual probability for persons with an A1C of 5.5%–5.9% (3.1%) and 6.0%–6.4% (6.6%) (see Table 10) for the annual probabilities of diabetes that are being weighted.

Table 13. Persons with Prediabetes and Other Persons at Risk for Type 2 Diabetes: Calculation of a Weighted Average Annual Probability of Diabetes by A1C Category

A1C Categories	Distribution of Risk Group Across A1C Categories	Weighted Average Annual Probability	Annual Probabilities Weighted by A1C Category
<5.5%–5.9% ^a	85.7%	3.1%	2.6%
6.0%–6.4%	14.3%	6.6%	0.9%
			Sum of Weighted Probabilities: 3.6%

Notes: The distribution of the “Persons with prediabetes and other persons at risk for type 2 diabetes” group across A1C categories is based on a survey-weighted estimate from NHANES (2011–2014). The “Sum of Weighted Probabilities” is 3.6% (and not 3.5%) due to rounding of the weighted probabilities for each A1C category.

^a Some people qualified for the risk group “persons with prediabetes and other persons at risk for type 2 diabetes” based on the CDC Prediabetes Screening Test score and may have had an A1C value less than 5.7%. For these people, we assumed that their annual probability of diabetes was equal to 3.1% (the annual probability for persons with an A1C of 5.5%–5.9%) (see Table 10).

2.4.3 Persons with High-Risk Prediabetes

This group is a subset of the group with prediabetes. This group is the smallest (about 34% of the prediabetes group), but it represents those with the highest risk of progressing to diabetes. “Persons with high-risk prediabetes” are defined as persons with an A1C between 6.0% and 6.4% or an FPG between 110 and 125 mg/dl and they have a default annual

probability of diabetes of 6.2%. This definition of “Persons with high-risk prediabetes” was chosen because the annual risk of developing diabetes was higher for persons with an A1C between 6.0% and 6.4% (Selvin et al., 2010) or an FPG between 110 and 125 mg/dl (Nichols et al., 2010), compared to persons with lower A1C and FPG values that are still in the prediabetes range.

This annual probability was determined by taking a weighted average of two annual probabilities. In NHANES (2011–2014), we found that about 20% of persons with prediabetes would qualify as having “high-risk prediabetes” under the A1C criteria alone (6.0%–6.4%). We assumed that this group had an annual probability of diabetes equal to 6.6% (see Table 10). When we added in the FPG criteria (110–125 mg/dl), the predicted prevalence of persons with high-risk prediabetes, among “Persons with prediabetes” increased from 20% to 34%. We assumed that the incremental persons qualifying as high-risk under the FPG criteria had a somewhat lower annual probability of diabetes based on the annual probability reported in Nichols et al. (2010) (5.6%) for persons with an FPG 110–125 mg/dl. The persons qualifying under the A1C criteria (with an associated annual probability of 6.6%) and the incremental persons qualifying under the FPG criteria (with an associated probability of 5.6%) were combined in a weighted average of roughly 60% and 40%, respectively. These weights are based on the proportion diagnosed with A1C criteria and the incremental proportion diagnosed with FPG criteria. The resulting annual probability for persons with high-risk prediabetes was 6.2%.

2.5 Additional Impact Toolkit Inputs

After selecting your population characteristics and risk group, you can customize additional inputs by clicking on the “CUSTOMIZE FURTHER” button just above the “GET RESULTS” button (Figure 2). For these additional inputs, we recommend beginning with “Screening” in the left column and ending with “Medical Costs” in the right column (your choice of screening options will affect subsequent input choice) (Figure 3). In other words, read through and customize the inputs as if you were reading a newspaper. First go down the left column, and then go down the right column (note: depending on the window or device you are using to view the Impact Toolkit, input fields may appear in a different number of columns).

Figure 2. Employer Input Dashboard Collapsed





Figure 3. Employer Input Dashboard (Expanded)



Start

Finish


Although everyone in your population can be recruited and screened for the National DPP, only a small percentage of this target population will actually be eligible and willing to participate in the program. If they do participate, they will receive the weight loss, diabetes risk reduction, and medical cost reduction benefits associated with the program. In Sections 2.5.1 through 2.5.7, we describe all assumptions related to implementing the National DPP in your population. In Section 3, we describe the benefits realized by conducting this program, as reported in the Output Dashboard.

2.5.1 Screening

In the screening section, you can choose if you would like to screen potential participants for prediabetes, if they have not already been screened. If you choose to conduct screening for the unscreened persons in your population, then screening costs will be incurred and these costs will be included in the toolkit's calculation of program costs (see Section 2.5.5).

Screening costs are shown in the screening cost calculation box in the Program Costs section (the box will appear in that section if you have chosen to conduct a screening program). This box calculates the average screening cost per person as the product of the assumed screening cost (\$12.50 in the default setting) and the average number of people screened per case detected (two in the default setting) (see Figure 4). This calculation allows us to account for the costs of negative screenings in the overall program costs. For further details on program cost calculations, see Section 2.5.5 on program costs. To assume a different number of people screened for the average screening cost calculation, edit the number in the box for "Average number of people screened for each case of prediabetes detected." An increase in this number would reflect an unscreened population with a low prevalence of prediabetes, while a decrease would reflect an unscreened population with a high prevalence of prediabetes. If you are not sure about the underlying prevalence in your unscreened population, then we recommend using the default setting of two people screened per case detected. This reflects a prevalence of prediabetes of about 50% among unscreened people that agree to participate in a screening program. To our knowledge, there are no extant data on the prevalence of prediabetes among people who participate in a screening program. While the typical prevalence of prediabetes is lower than 50%, we assumed that it would be higher among a self-selected group of screening participants (i.e., people are more likely to agree to screening if they have risk factors for prediabetes).

Figure 4. Program Cost Inputs Showing Average Screening Cost Calculation




2.5.2 Program Enrollment and Participation

Here you can enter the percentage of eligible participants that have already been screened for prediabetes. The default setting is 46%, which is based on the percentage of people at risk for diabetes that have been screened in the past 3 years (Bullard et al., 2015; Kiefer et al., 2015). Persons at risk for diabetes include adults aged 45 or older or those with physical inactivity, family histories of diabetes, high blood pressure, and other risk factors per the American Diabetes Association (ADA, 2016) and the U.S. Preventive Services Task Force (Calange et al., 2008) criteria for screening for type 2 diabetes. Other analyses have shown that only about 11% of persons with prediabetes in 2010 were screened and aware of their disease status (CDC, 2013). Thus, other values, such as 11%, can be entered as an alternative with justification.

If you have chosen to conduct additional screening for your program, then you can also input the “percentage of eligible, previously unscreened persons receiving screening” (the entry field for this input will only be shown if you have chosen to conduct screening in the screening section). The default value is 100%, which assumes that all persons with a BMI≥24 who have not been screened will receive screening. You can adjust this value downward as appropriate for your population.

Next, you enter your assumed participation rate (see Figure 5). The default setting for the participation rate is 35%, which is based on the participation rate in a demonstration of the National DPP with large employers (R. Li, R. Ackermann, personal communication, June 22, 2015). The participation rate might be higher or lower for your population based on the incentives offered for participating or the perceived benefits of participation.

Figure 5. Program Enrollment and Participation Inputs



Note: An additional input field, “Percentage of eligible, previously unscreened persons receiving screening,” will appear if you choose to conduct screening for the unscreened persons in your population in the screening section.

In the final number shown in Figure 5, the program enrollment and participation inputs are used to determine the total percentage of participation among *eligible* adults. For example, if 46% of eligible adults have previously been screened, no additional screening occurs (the default value), and the participation rate is 35%, then the total percentage of eligible adults participating will be 16% ($46\% \times 35\% = 16\%$). This percentage is then applied to the predicted prevalence of the selected risk group to get the percentage of projected participants. For example, in the default setting, the predicted prevalence of “persons with prediabetes” is 29% in the national sample. Thus, $29\% \times 16\% = 4.7\%$ of the total population that will participate in the National DPP if no additional screenings are conducted. However, in a scenario where a screening program is conducted to identify additional eligible persons, the calculation is somewhat more complex (Equation 1). Generally, the

more adults screened will result in a larger # of eligible persons and a larger # of participants.

Equation 1. Percentage of Screened Population Who Participate in the Intervention

$$\begin{aligned} \text{Percentage of screened population who participate in the intervention} &= \% \text{ of population previously screened for prediabetes} * \% \text{ of screened persons participating in intervention} + \\ &[(100\% - \% \text{ of population previously screened for prediabetes}) * \% \text{ of previously unscreened persons receiving screening} * \% \text{ of screened persons participating in intervention}] \end{aligned}$$

Similar to the example with no additional screenings, the percentage from Equation 1 (which is used when there are additional screenings) is applied to the predicted prevalence of the selected risk group (e.g., 29% for “persons with prediabetes” in the national sample) to get the percentage of projected participants. For example, if the “% of previously unscreened persons receiving screening” is 100%, then we would calculate the percentage of projected participants as $29\% * (46\% * 35\%) + [(100\% - 46\%) * 100\% * 35\%]$, which equals 16% of the total population that will participate in the National DPP if additional screenings are conducted on 100% of the previously unscreened population.

2.5.3 Intervention Weight Loss and Regain Schedule

The DPP trial demonstrated that participants in an intensive lifestyle intervention lost about 7.2% of their weight in the first year of the program (DPP, 2002; Hamman et al., 2002). Real-world adaptations of the DPP trial, such as the National DPP, resulted in a smaller weight loss effect, 4.4% on average at the end of the first year of follow-up (R. Li, personal communication, June 24, 2015). This estimate is based on results from the National DPP and represents the first year weight loss for program participants and is supported by other studies.¹ A participant was defined as someone attending at least 4 of the 16 program sessions offered during the first 6 months. Attending at least 9 sessions during the first 6 months was associated with a slightly higher weight loss at the end of Year 1 (5.1%) (R. Li, personal communication, June 24, 2015).

The DPP trial and studies of real-world interventions with additional years of follow-up show that the initial weight lost is regained in future years. Based on data from these studies, we assume that about 50% of the weight lost is regained in Year 2, and another 20% is regained in Year 3 (Barte et al., 2010). In Years 4 through 10, we assume that all the weight lost has been regained (0% weight loss relative to baseline weight). These weight loss and regain assumptions are based on group-based type 2 diabetes prevention lifestyle

¹ In 2012, Ali et al. (2012) reported a 4.1% weight loss at 12 months’ follow-up for translational programs conducted in the United States and modeled on the Diabetes Prevention Program (DPP) trial’s lifestyle intervention.

change programs with a minimum of 16 sessions over the first 6 months. A minimum of 6 sessions must be offered in the second 6 months of the program to achieve full recognition status per the criteria set forth by CDC (CDC, 2015b).

The intervention weight loss and regain schedule in the Impact Toolkit shows the average percentage difference (relative to baseline body weight) for 10 years of follow-up. Data from the DPP and Diabetes Prevention Program Outcomes Study (DPPOS) trial showed that weight loss was the greatest after 1 year (7.2%) (Hamman et al., 2002) and was associated with a 58% diabetes risk reduction over the 3-year study period. Based on these DPP trial data and the associated diabetes risk reductions observed in the trial we estimated the average diabetes risk reduction from National DPP programs by discounting the DPP trial risk reduction according to Equation 2.

Equation 2. Discount Factor for Translational DPP Program Effects for Program Participants: Weight Loss

$$(\text{NDPP weight loss})/(\text{DPP weight loss}) = (4.4\%)/(7.2\%) = 61.1\%$$

Note: This discount factor is automatically recalculated in the Toolkit if the user adjusts the weight-loss effect as described above.

Thus, the National DPP program is assumed to have 61.1% of the effect on diabetes risk that was observed in the DPP trial. In DPP trial, the 58% reduction in diabetes incidence is associated with 7.2% weight reduction. Thus the 61.1% discount factor translates to a 35.4% risk reduction in the first year of participating in the National DPP. Table 14 shows the default assumptions for weight loss and the calculated diabetes risk reduction (using Equation 2) associated with each year's weight loss estimate. The diabetes risk reduction effects in each year are automatically updated per Equation 2 when the weight loss and regain effects are changed by the user. Although these default settings are based on the best available data for the average National DPP participant, your population and program may differ from the average. You can edit these weight loss and regain assumptions according to the expectations for your program.

For the Impact Toolkit, we limit customized input average loss values to 10% of baseline weight. Larger weight losses are unlikely to occur, given the 7.2% weight loss in the original DPP trial and the 4.4% weight loss observed in translational programs. In addition, weight loss greater than 10% would lead to results with low credibility under the current model calculation of the risk reduction in diabetes incidence associated with the National DPP. This risk reduction cannot exceed 100% (i.e., you cannot reduce diabetes incidence below 0%). Given Equation 2, the 10% ceiling on weight loss would lead to a discount factor of 1.39 and—multiplied by the 58% risk reduction in the original DPP—to an 81% risk reduction in diabetes incidence.

Table 14. Default Intervention Weight Loss and Weight Regain Schedule

Year	Intervention Weight Loss (from baseline weight)	Assumed Diabetes Incidence Risk Reduction
1	4.4%	35.4%
2	2.4%	19.3%
3	1.9%	15.3%
4	0.0%	0.0%
5	0.0%	0.0%
6	0.0%	0.0%
7	0.0%	0.0%
8	0.0%	0.0%
9	0.0%	0.0%
10	0.0%	0.0%

Note: Year 1 is the year in which the intervention is delivered to participants. All weight loss percentages are based on participants’ baseline weight (before the intervention). In the default setting, weight is gradually regained with a full return to baseline weight in Year 4. This regain trend is based on evidence from a review of group-based lifestyle interventions. Weight loss and regain are associated with a reduced incidence of diabetes as described in Section 2.5.3.

2.5.4 Program Budget

In the default setting, the Impact Toolkit assumes that a state, employer, or insurer will offer the program to all eligible persons who want to participate. However, if there is a limited budget for implementing a National DPP lifestyle change program, then you can check the box in this section to set a maximum budget. Once you have checked this box, an additional entry field will appear for you to enter your maximum budget. This budget you enter will limit the number of program participants based on the size of your eligible population, your program costs, and your screening costs (if you choose to screen previously unscreened persons). Equation 3 demonstrates this calculation.

Equation 3. Determining the Proportion Participating in the Program when there is a Limited Budget

$$\text{Proportion participating in intervention} = \text{Minimum} \left[\text{Proportion completing screening} * \text{Percentage of screened persons participating in intervention,} \left(\frac{\text{Total Program Budget}}{\text{Average program cost per person with screening}} \right) \right]$$

2.5.5 Program Costs

Here you can enter the per person costs of your program. The default cost of \$417 (2013 U.S. dollars) is the per-participant cost of a group-based National DPP lifestyle change

program without screening costs included (Li et al., 2015). All costs in this section should be calculated and entered per participant. If you are conducting a screening program (i.e., selected “Screen persons for prediabetes if they have not been previously screened” in the screening section), then this section will also show an “AVERAGE SCREENING COST CALCULATION” box. Here, we assume that screening costs are \$12.50 in the default setting. Either the FPG test (\$7.22) or the hemoglobin A1c test (\$17.85) can be used to diagnose prediabetes, so we assume the average cost (\$12.50) of these two tests according to the 2015 Medicare Laboratory Fee Schedule (CMS, 2016a).

If you plan to use one of these tests or believe that your screening costs differ from these estimates, then the screening test costs can be modified in this section. If you plan to use the CDC Prediabetes Screening Test (CDC, 2015a) (a questionnaire that can also be used to determine eligibility for the National DPP) instead of a blood test, then your costs may be lower than either of the blood test costs (e.g., the cost of printing and distributing self-administered questionnaires). We also assume that there are “Other screening costs,” which include the cost of a brief follow-up visit to discuss the patient’s screening test results and receive a referral to a program such as the National DPP. The default cost of \$20 for “Other screening costs” only includes the cost a brief office visit and not the costs of recruitment for screening. “Other screening costs” are based on the 2015 Medicare Physician Fee Schedule (CMS, 2016b) associated with an evaluation and management visit of low complexity (HCPCS Code 99211) for an established patient (about 5 minutes of face-to-face time at the cost of \$20.02). This cost can be adjusted to reflect a more intensive screening program with recruitment costs. Recruitment costs for a National DPP are not widely reported in the literature yet, however a couple of sources suggest that recruitment costs are fairly low,² ranging from about \$1 to \$14.

As noted in Section 2.5.1, we also account for the costs of negative screenings by assuming that you have to screen two people to detect one case of prediabetes, on average. The number of people screened per case detected (which can be modified in the Screening Section) is used in this section to calculate the screening cost per participant (Equation 4).

Equation 4. Screening Cost per Participant (If Conducting a Screening Program)

$$\text{Average screening cost per person detected} = (\text{Screening cost per person} + \text{Other screening costs per person}) * \text{Number of persons screened per case detected}$$

² Recruitment costs vary widely across studies depending on the intensity of the recruitment strategy. Recruitment costs at an employee worksite program with 1,800 employees (just 107 were found to be eligible based on blood-test confirmed prediabetes criteria) were \$1,500 (\$1,094 for printing/ mailing and \$406 for tent cards, flyers, and posters) (Taradesh et al., 2015). Thus, recruitment costs were about \$14 per eligible person recruited. Krukowski et al. (2013) reported recruitment costs of \$1.13 per participant. This is just the cost of flyers left at senior centers to recruit participants into a lifestyle intervention. Taradesh et al.’s estimate of \$14 for recruiting costs is more consistent with data reported from the YMCA HCIA demonstrations (personal communication, Andrew Lanza, CDC).

Next, the average screening cost per participant is added to the base program cost (\$417 in the default setting) to calculate the overall program cost per person. This calculation is shown in Equation 5 and is used as the overall program cost in calculations of net costs shown in the Output Dashboard.

Equation 5. Overall Program Cost (with a Screening Program)

$$\left[\frac{\text{Average program cost per participant with screening} = \text{Program cost per participant} + \left(\frac{\text{Average screening cost per participant detected} * \left(\frac{(100\% - \% \text{ of pop previously screened for prediabetes}) * \% \text{ of previously unscreened participants receiving screening}}{\% \text{ of screened participants participating in intervention}} \right)}{\% \text{ of screened participants participating in intervention}} \right)}{\% \text{ of screened participants participating in intervention}} \right]$$

Note: In Equation 5, the term “Average screening cost per person detected” is calculated from Equation 4.

2.5.6 Medical Costs

In the year that a person is diagnosed with diabetes, he or she will have substantial medical costs associated with diagnosing and treating their diabetes. In each subsequent year, the person will continue to have medical costs for the treatment of his or her disease, although not as great as the initial year of diagnosis. These are usually called diabetes-attributable medical costs, and they are defined as the excess medical costs for a person with diabetes compared with a similar person without diabetes.

The default values that are provided in the Impact Toolkit reflect the average excess medical costs for persons with diabetes based on a CDC analysis of longitudinal medical claims data from MarketScan (2001–2013) (Shrestha et al., 2016). Based on these data and a review of other cost analyses,³ we assumed default values of \$6,424 for the first year of diagnosis and \$3,900 for the years after diagnosis (2013 U.S. dollars). First year costs are approximately 1.65 times greater than subsequent year costs. If you believe that excess medical costs associated with diabetes differ in your population, we suggest maintaining this approximate relationship between the first year costs and the subsequent year costs. We also suggest staying within the suggested range of \$3,300 to \$9,900 for first year costs and \$2,000 to \$6,000 for subsequent year costs. Using excess medical cost estimates outside of these ranges may lead to results with low credibility.

³ Studies that use cross-sectional cost data may underestimate costs in the year of diabetes onset. Previous analyses using the MEPS-NHIS linked data file have noted this limitation as well (Trogon et al., 2008). However, in studies that use *longitudinal* data to follow individuals before and after the onset of diabetes, authors find a spike in medical costs in the year of onset (Nichols et al., 2000; Shrestha et al., 2016).

We used cost equations from Zhuo et al. (2014) to account for increasing age and duration of diabetes (for persons that develop the disease) in future years. Table 15 shows how the excess medical costs increase with increasing age and duration of diabetes (for persons developing diabetes). On average, the excess medical cost associated with having diabetes was about \$6,424 in the year of diagnosis and \$3,900 in the years after diagnosis (longer durations of diabetes are associated with higher costs (see Table 15), but we only calculate one multiplier for Years 2 through 10). When the user enters a different value in the Input Dashboard, a cost multiplier is calculated based on these default excess costs of diabetes. Equation 6 shows how the cost multiplier is calculated for the year of diagnosis, and Equation 7 shows how the cost multiplier is calculated for the years after diagnosis. The user can enter his or her own “Annual diabetes attributable costs in the year of diagnosis” and “Annual diabetes attributable costs in the years after diagnosis,” which are used to calculate the multipliers in Equations 6 and 7, respectively. All of the values in column 1 of Table 15 (year of diagnosis) are increased by the multiplier calculated in Equation 6, while all of the values in columns 2 through 10 are increased or decreased by the multiplier calculated in Equation 7. The “Years after diagnosis cost multiplier” will have a larger effect on costs for longer durations of diabetes because the baseline costs in Table 15 are larger for longer durations of diabetes.

Table 15. Estimated Diabetes Attributable Medical Costs by Year and Duration of Diabetes

Year of Follow-Up	Duration with Diabetes (years)									
	<1	≥1, <2	≥2, <3	≥3, <4	≥4, <5	≥5, <6	≥6, <7	≥7, <8	≥8, <9	≥9, <10
1	\$6,424									
2	\$6,424	\$2,767								
3	\$6,424	\$2,645	\$3,092							
4	\$6,424	\$2,532	\$2,980	\$3,432						
5	\$6,424	\$2,292	\$2,740	\$3,193	\$3,650					
6	\$6,424	\$2,181	\$2,629	\$3,082	\$3,540	\$4,000				
7	\$6,424	\$2,053	\$2,500	\$2,954	\$3,412	\$3,873	\$4,336			
8	\$6,424	\$1,922	\$2,370	\$2,823	\$3,282	\$3,743	\$4,207	\$4,670		
9	\$6,424	\$1,805	\$2,251	\$2,705	\$3,163	\$3,626	\$4,090	\$4,554	\$5,017	
10	\$6,424	\$1,672	\$2,118	\$2,571	\$3,029	\$3,492	\$3,956	\$4,421	\$4,885	\$5,346

Note: All participants begin as persons without diabetes. Cells with a longer duration of diabetes than year of follow-up are left blank because a participant in Year 2 cannot have diabetes for more than 2 years if they did not have diabetes at Year 0. All costs are stated in 2013 U.S. dollars.

Equation 6. Calculating the Cost Multiplier for the Year of Diagnosis

$$\text{Year of diagnosis cost multiplier} = \frac{\text{Annual diabetes attributable costs in the year of diagnosis}}{\$6,424}$$

Equation 7. Calculating the Cost Multiplier for the Years after Diagnosis

$$\text{Years after diagnosis cost multiplier} = \frac{\text{Annual diabetes attributable costs in the years after diagnosis}}{\$3,900}$$

In the Impact Toolkit, all costs incurred in the future are discounted back to the present using a discount rate assumed in the Medical Cost section of the Input Dashboard. The discount rate input box allows us to account for the fact that the money we have today has more value than money received in the future. This accounts for future inflation, lost investment opportunity, and risk. Applying this discount value allows us to more accurately compare the money that will be spent in the future with the money that is spent today. An annual discount rate of 1.0% to 5.0% is commonly used. Our default value is 3.0%.

2.5.7 Productivity Costs (Employer Module Only)

In the Employer module only, we include productivity costs associated with diabetes. We limit productivity costs to the costs of days of work missed due to diabetes and the value of these days. The value of the days of work missed (per person) is calculated by multiplying the “Days of work missed per year due to diabetes” and the “Daily earnings for persons with diabetes.” Days of work missed per year due to diabetes are the excess days of work missed by someone with diabetes compared with a similar person without diabetes (e.g., similar age, sex, comorbidities).

We used NHIS to estimate the number of days of work missed attributable to diabetes. Pooling data from the 2009 through 2013 NHIS, we estimated days of work missed at the national level.⁴ Our final estimation used a two-part model with a logit model for the first part and a generalized linear model for the second part. We controlled for the following comorbidities: arthritis, asthma, cancer, depression, chronic bronchitis, back problems, and pregnancy. We also included the following sociodemographic controls: age, age squared, race/ethnicity, education, family income, health insurance, and occupation.

⁴ In NHIS, persons with diabetes are identified by the question “Have you ever been told that you have diabetes?” The work-loss analysis was restricted to individuals employed at any point during the year. Number of workdays lost was defined using the following NHIS question: “During the past 12 months, about how many days did you miss work at a job or business because of illness or injury (do not include maternity leave)?” To estimate workdays lost due to diabetes, we tested four different models for best fit: one-part negative binomial model, two-part truncated negative binomial model with a logit, two-part generalized linear model with a logit, and a zero-inflated negative binomial model. Based on a comparison of the model residuals, the Akaike information criterion (AIC), and the Bayesian information criterion (BIC), our final estimation used a two-part model with a logit model for the first part and a GLM for the second part.

We found that, on average, a person with diabetes will miss 3.3 more days of work each year as compared to a similar person without diabetes.⁵ The default daily earnings of \$276 was calculated as a weighted average of daily earnings for males and females aged 45 to 64.⁶ This value can be modified to reflect the average earnings for your employee population.

After filling in all the fields discussed in Sections 2.1 through 2.5, click “GET RESULTS” to view your customized results in the Output Dashboard. See Section 3 for information about calculations made in the Output Dashboard and how to interpret results.

3. OUTPUT DASHBOARD

The results in the Output Dashboard are unique to your data inputs. They predict the health and economic outcomes for your program participants as a result of implementing a National DPP lifestyle change program.


3.1 Projected Participants

Figure 6 demonstrates how the projected number of participants is calculated from the total population. The total adult population being considered for the National DPP is shown in the top bar in the figure. This is determined by the total adult population in your state (State module), the number of employees at your company (employer module), or the number of insured adults (Insurer module). In the second bar from the top, a subset of the total population is projected to be eligible for the intervention based on (1) the assumed population characteristics (Sections 2.1–2.3) and (2) the risk group selected in the Input Dashboard (Section 2.4). Next, some or all (depending on your screening inputs, see Section 2.5.1 and 2.5.2) of the eligible persons have been previously or newly screened, confirming their eligibility for the intervention (shown in the third bar from the top). Among the eligible, screened persons, some or all (depending on your participation inputs, see Section 2.5.2) will participate in the intervention. The predicted number of participants is shown in the bottom bar of this figure. This number is calculated by multiplying the number of eligible people in second bar by the “Percentage of Screened Population who Participate in the Intervention” calculated in Equation 1 in Section 2.5.2.

⁵ This estimate of 3.3 work loss days is the weighted average of the estimated work loss associated with diabetes for 45- to 64-year-old males (53.8% in the employed population, NHANES 2011–2014) and females (46.2% in the employed population, NHANES 2011–2014). We selected the 45–64 age group because the mean age of persons with prediabetes is about 52 years old.

⁶ This estimate of \$276 in daily earnings is a weighted average of the estimated daily earnings for 45- to 64-year-old males (53.8% in the employed population, NHANES 2011–2014) and females (46.2% in the employed population, NHANES 2011–2014). We selected the 45–64 age group because the mean age of persons with prediabetes is about 52 years old. To estimate the daily earnings, we used the Current Population Survey’s (CPS’) 2014 annual wage estimates by 5-year age groups, aggregated these to the 45–64 age group using 2014 population counts from the Census, and deflated to wages to 2013 U.S. dollars. We calculated daily wage as the annual wage divided by 250 work days per year.

Figure 6. Projected Participants




3.2 Cumulative Projected Cases of Diabetes

Figure 7 demonstrates the effect of the National DPP lifestyle change program on new cases of type 2 diabetes. In the chart on the left in Figure 7, the cumulative number of diabetes cases with and without the National DPP intervention is shown for your population of participants. The gray line estimates the number of new diabetes cases without the National DPP. The green line represents the number of new diabetes cases with the National DPP. The difference between the two lines represents the cases of diabetes averted at a given point in time. In the chart on the right in Figure 7, the cumulative years with diabetes averted is shown for your population of participants. The cumulative years with diabetes averted (by the National DPP lifestyle change program) is the cumulative calculation of the cases averted in each year.

The projected cases of diabetes were determined using a simplified Markov model and two key inputs: (1) the annual probability of diabetes (see Sections 2.4.1 through 2.4.3); and (2) the weight loss/regain assumptions, which reduce the probability of diabetes (see Section 2.5.3). The simplified Markov model was initially a spreadsheet model that mimics a Markov model with 1-year cycles and three states: No diabetes, Diabetes, and Dead. The model begins with the full sample of program participants (final bar from “Projected Participants” figure). During each annual cycle, a percentage of these participants progresses from prediabetes to diabetes according to the annual probability of diabetes set by the user (e.g., with an annual probability of diabetes of 3.8%, 38 out of 1,000 participants develop diabetes in the first cycle). Persons developing diabetes are removed

from the sample of persons without diabetes, and the number of cases in the next cycle is calculated based on this new sample (e.g., the number of cases in the next cycle is

Figure 7. Projected Cases of Diabetes and Years with Diabetes Averted by Participating in the National DPP Lifestyle Change Program



calculated based on a sample of 962 participants [1,000 – 38 = 962] without diabetes). The calculations made by the spreadsheet-based Markov model were programmed as the Impact Toolkit to calculate cases of diabetes over the 10-year period as well as the other outcomes described in Sections 3.3 through 3.7.

To calculate the gray line (no intervention), the model assumes no weight loss and therefore no reduction in the annual probability of diabetes. To calculate the green (National DPP intervention) line, the model assumes some weight loss and some reduction in the annual probability of diabetes—leading to a smaller number of cases. In each annual cycle, the annual probability of diabetes is only reduced if there is a weight loss effect assumed for the applicable annual cycle. In the default setting, there are weight loss effects in the first 3 years that translate to diabetes risk reduction effects (see Table 14) in each of the first 3 years. The simplified Markov model also incorporates mortality. See Section 3.7 for details regarding the death rate and relative risk of death for persons with diabetes.


Clicking on the “SHOW DATA TABLE” button will open up a table with more information. The table shows the number of “Cases Averted” in each year and the “Years with Diabetes Averted.” The “Cases Averted” are calculated as the difference between the gray and green

line. The “Years with Diabetes Averted” represents the cumulative number of person-years with diabetes that are averted with intervention. The maximum number of “Cases Averted” in a single year will usually occur in the last year assumed to have some weight loss effect (Year 3 in the default setting).

3.3 Cumulative Medical Costs per Participant

Figure 8 shows the difference in cumulative medical costs with and without the National DPP intervention. Results are displayed per participant and can be easily scaled up by multiplying results by your number of participants. The gray bar on the left indicates the cumulative medical costs without the National DPP, and the green bar on the right indicates the cumulative medical costs with participation in the National DPP. The cumulative medical cost savings is the difference between these amounts. The difference can be fairly small, however the trend indicates that the gap in cumulative medical cost between participants on the National DPP and non-participants, widens over time. Click on the “SHOW DATA TABLE” button to see more detailed information, including the medical cost-savings (i.e., the difference between the gray and green bars) with the National DPP.

Figure 8. Cumulative Medical Costs per Participant



Using the simplified Markov model described in Section 3.2, medical costs are calculated for persons with diabetes and for persons without diabetes in each annual cycle using the diabetes attributable medical costs shown in Table 15. Medical costs for the full sample of participants are summed and then divided by the number of sample participants to get the medical costs per participant. Medical costs per participant are slightly lower in the green bars (National DPP Intervention) because fewer people progress to diabetes when participating in the National DPP and thus do not incur the excess costs associated with having diabetes. We assume that diabetes onset occurs at the beginning of the period and death occurs at the end of the period for our medical cost calculations.

Results in this figure are largely affected by the medical cost assumptions (Section 2.5.6, Table 15, and Equations 6 and 7), the assumed annual probability of developing diabetes for your risk group, and the assumed weight loss effects. Even when the excess medical costs of diabetes are assumed to be large (e.g., \$6,425 in the year of diagnosis and \$3,900 in


subsequent years in the default setting), the *average* difference in medical costs can be fairly small. This is due to the fact that (1) only a fraction of participants develop diabetes each year (e.g., 3.8% each year in the default setting for persons with prediabetes), and (2) only a fraction of the projected cases of diabetes are averted among participants (this will depend on the weight loss/regain assumptions).

3.4 Net Costs (Program Costs minus Medical and Productivity Cost Savings) per Participant

Figure 9 uses the cumulative medical costs (Section 3.3) and the cumulative program costs (Section 2.5.2) to generate the cumulative net cost per participant. In each annual cycle, the cumulative medical cost savings (and productivity savings in the Employer module) is subtracted from the cumulative program cost to produce the cumulative net cost in that year. Results are displayed per participant and can be easily scaled up by multiplying results by your number of participants.

In Figure 9, program costs are represented by the “overall program cost” described in Equation 5 (see Section 2.5.5). This overall program cost (per person) includes the base cost of the program (e.g., \$417 in the default setting) and the cost of screening. The cost of screening accounts for (1) the basic screening test cost (\$12.50 in the default setting), (2) “other screening costs” (\$20 in the default setting), (3) the number of screenings per case detected (two in the default setting), and (4) the number of positive screenings that do not result in a participant (only 35% of eligible, screened persons actually participate in the default setting). See Equation 5 in Section 2.5.5 for the detailed equation that calculated the overall program cost.

Figure 9. Net Costs (Program Costs minus Medical and Productivity Cost Savings) per Participant



Note: Productivity cost savings are only included in the Employer module.

In general, the cumulative net costs decrease over time, so that by Year 10 or sooner, there may be negative cumulative net costs, indicating that the program is cost-saving (all calculations account for the time value of money using the discount rate you specify—see end of Section 2.5.6). Net costs fall because the program cost is only paid once, in Year 1, whereas medical cost savings as a result of program participation occur each year. The net cost calculation is most sensitive to your assumptions regarding the program cost, the weight loss/regain, and the medical costs. Click on the “SHOW DATA TABLE” button to see more information. It should be noted that net costs only include medical and program costs and do not otherwise reflect the health benefits to participants.

3.4.1 Productivity Costs in the Employer Module

In the Employer module, productivity cost-savings are calculated in addition to medical cost-savings. In the net costs figure (Figure 9), these productivity cost-savings are also subtracted from the overall program costs in each period to get the net costs. For each new case of diabetes, a number of days of work are missed each year due to diabetes (3.3 days in the default setting—see Section 2.5.6). Each day is valued by the average wages assumed for your employee population (\$276 per day in the default setting). For example, in the default setting, about \$911 in productivity costs are incurred each year for everyone who has developed diabetes in that cycle or a previous cycle.

The productivity cost-savings from averting cases of diabetes in the intervention scenario are averaged across all participants. Similar to the medical costs, the productivity costs with the intervention are only slightly lower than the productivity costs without the intervention due to the fact that (1) only a fraction of participants develop diabetes each year (e.g., 3.8% each year in the default setting for persons with prediabetes), and (2) only a fraction of the projected cases of diabetes are averted among participants (this will depend on the weight loss/regain assumptions). The productivity costs with the intervention, without the intervention, and the calculated cost-savings (all per person) are shown in the data table associated with Figure 9. Click on the “SHOW DATA TABLE” button to see the detailed 10 year data in the Impact Toolkit.


3.5 Cumulative Quality Adjusted Life Years (QALYs) Gained

Figure 10 shows the cumulative quality-adjusted life years (QALYs) gained for your population of participants. QALYs are a combined measure of health and time, allowing us to weight years lived by the health-related quality of life in those years. One QALY is equal to 1 year of life with a perfect quality of life.

When a person develops diabetes, we assume that their quality of life is reduced by about 5%, on average. The estimate of a 5% reduction in quality of life is based on the quality of life decrement associated with progressing from prediabetes to diabetes in the CDC/RTI model of diabetes.⁷ This reduction in quality of life is averted or delayed with each case of diabetes that is averted or delayed (due to the National DPP lifestyle change program). Similar to the method for calculating medical costs, QALYs are calculated with and without effect of the National DPP using a simplified Markov model. Using the Markov model approach the cumulative QALYs gained (shown in Figure 10) account for averting the reduction in quality of life associated with diabetes as well as the timing of when cases are averted. For instance, if weight losses are maintained throughout the whole 10-year period, then cases of diabetes will be delayed for a longer amount of time and QALY gains will be greater. If weight losses are quickly regained, then these cases of diabetes are not delayed as long and not as many QALYs will be gained.

⁷ A quality of life decrement of 0.04 relative to a baseline utility of 0.84 for persons without diabetes represents about a 5% decrease in quality of life.

Figure 10. Cumulative Quality Adjusted Life Years (QALYs) Gained




3.6 Incremental Cost-Effectiveness Ratios (ICERs)

Figure 11 shows the annual net costs per participant, QALYs gained per participant, incremental cost-effectiveness ratios (ICERs) (net costs divided by QALYs gained), and the cost per case averted (cumulative cases averted or “Years of Diabetes Averted” divided by the net costs).

The net costs are calculated per participant for each year by subtracting the cumulative medical cost-savings from the cumulative program costs (see Section 3.4 for calculation details). This information can also be seen in Figure 9. The QALYs gained are calculated per participant in Figure 11 by dividing the cumulative QALYs gained for your population of participants as in the QALYs gained figure (Figure 11) by the size of the population of participants.

Figure 11. Incremental Cost-Effectiveness Ratios (ICERs)



The ICER is a measure of the cost-effectiveness or “return on investment” associated with an intervention. It is calculated as the cumulative net costs divided by the cumulative QALYs gained. A lower ICER is better as it indicates that QALYs are gained from the intervention at a lower cost. A negative ICER indicates that the intervention is associated with QALY gains and reduced costs (i.e., a cost-saving intervention). Negative ICERs are shown in Figure 11 as “cost-saving.” To see the amount of cost-savings per person, refer to the “Net Costs” figure and accompanying table (see Section 3.4).

The Cost per Case Averted is found by dividing the cumulative cases averted (as seen in Figure 7, “Cumulative Projected Cases of Diabetes”) by the cumulative net cost. A negative cost per case averted means that the program is cost saving by that point in time. Negative costs per case averted are shown in Figure 11 as “cost-saving.”

3.7 Cumulative Years of Life Gained


Developing diabetes is associated with an increased risk of life-threatening events, such as heart attack and stroke. We assumed that persons with diabetes have twice the risk of dying in a given year (from all causes) compared with persons without diabetes, on average. Based on an unpublished CDC analysis, we observed a range of 2.0 to 4.0 for the relative risk of death for persons with diabetes (personal communication, Yiling Chen, April 11, 2015). These relative risk estimates varied by age and sex. We chose to use an estimate of 2.0.

When cases of diabetes are averted or delayed (due to the intervention), a small decrease in the number of diabetes-related deaths is achieved. These deaths averted are associated with a gain in the years of life lived during the 10-year period. Similar to other calculations discussed in Section 3, a simplified Markov model was used to calculate deaths in each annual cycle. A small percentage of people die during each cycle and are removed from the modeling sample based on an assumed mortality rate of 0.45% for persons without diabetes and 0.90% for persons with diabetes. We chose the baseline mortality rate (0.45%) from the 2010 National Vital Statistics Life Tables⁸ (Arias, 2014) based the average age (age 52) of people with prediabetes in NHANES (2011–2014).

Figure 12 shows the cumulative years of life gained for your participant population as a result of participation in the National DPP. These years of life gained do not account for the quality of life with diabetes as in the QALYs gained figure. Click on the “SHOW DATA TABLE” button to see more information. Results are calculated dynamically in the Output Dashboard and are often rounded to accommodate the best display of the results given the space restrictions. To view the unrounded results, download your results in Microsoft Excel using the green button in the upper-right corner of the Output Dashboard.

⁸ See Table 1 “Life table for the total population: United States, 2010.”

Figure 12. Cumulative Years of Life Gained



References

- Ackermann, R. T., Liss, D. T., Finch, E. A., Schmidt, K. K., Hays, L. M., Marrero, D. G., & Saha, C. (2015). A randomized comparative effectiveness trial for preventing type 2 diabetes. *American Journal of Public Health, 105*(11), 2328–2334. <http://dx.doi.org/10.2105/AJPH.2015.302641>
- Ali, M. K., Echouffo-Tcheugui, J., & Williamson, D. F. (2012). How effective were lifestyle interventions in real-world settings that were modeled on the Diabetes Prevention Program? *Health Affairs, 31*(1), 67–75. <http://dx.doi.org/10.1377/hlthaff.2011.1009>
- American Diabetes Association. (2016). Standards of medical care in diabetes—2016. *Diabetes Care, 39*(Suppl 1), S4–S5. <http://dx.doi.org/10.2337/dc16-S003>
- Arias, E. (2014). United States life tables, 2010. *National Vital Statistics Reports, 63*(7), 1–62. Retrieved from http://www.cdc.gov/nchs/data/nvsr/nvsr63/nvsr63_07.pdf
- Barte, J. C. M., ter Bogt, N. C. W., Bogers, R. P., Teixeira, P. J., Blissmer, B., Mori, T. A., & Bemelmans, W. J. E. (2010). Maintenance of weight loss after lifestyle interventions for overweight and obesity, a systematic review. *Obesity Reviews, 11*(12), 899–906. <http://dx.doi.org/10.1111/j.1467-789X.2010.00740.x>
- Bullard, K. M., Ali, M. K., Imperatore, G., Geiss, L. S., Saydah, S. H., Albu, J. B., . . . Gregg, E. W. (2015). Receipt of glucose testing and performance of two US diabetes screening guidelines, 2007-2012. *PLoS One, 10*(4), e0125249. <http://dx.doi.org/10.1371/journal.pone.0125249>
- Calonge, N., Petitti, D. B., DeWitt, T. G., Dietrich, A. J., Gordis, L., Gregory, K. D., . . . , & the U.S. Preventive Services Task Force. (2008). Screening for type 2 diabetes mellitus in adults: U.S. Preventive Services Task Force recommendation statement. *Annals of Internal Medicine, 148*(11), 846–854. <http://dx.doi.org/10.7326/0003-4819-148-11-200806030-00007>
- Centers for Disease Control and Prevention (CDC). (2013). Awareness of prediabetes—United States, 2005-2010. *Morbidity and Mortality Weekly Report, 62*(11), 209–212.
- Centers for Disease Control and Prevention (CDC). (2014a). *National diabetes statistics report: Estimates of diabetes and its burden in the United States, 2014*. Atlanta, GA: U.S. Department of Health and Human Services.
- Centers for Disease Control and Prevention (CDC). (2014b). Guide to Community Preventive Services. Diabetes prevention and control: combined diet and physical activity promotion programs to prevent type 2 diabetes among people at increased risk. Retrieved from www.thecommunityguide.org/diabetes/combineddietandpa.html
- Centers for Disease Control and Prevention (CDC). (2015a). *CDC Prediabetes Screening Test*. Retrieved from <http://www.cdc.gov/diabetes/prevention/pdf/prediabetestest.pdf>
- Centers for Disease Control and Prevention (CDC). (2015b). *Diabetes Prevention Recognition Program Standards and Operating Procedures*. Retrieved from <http://www.cdc.gov/diabetes/prevention/pdf/dprp-standards.pdf>

- Centers for Medicare and Medicaid Services (CMS). (2016a). 2015 Medicare Laboratory Fee Schedule. Retrieved February 23, 2016, from <https://www.cms.gov/apps/ama/license.asp?file=/Medicare/Medicare-Fee-for-Service-Payment/ClinicalLabFeeSched/Downloads/15CLAB.zip>
- Centers for Medicare and Medicaid Services (CMS). (2016b). 2013 Medicare Physician Fee Schedule. Retrieved February 23, 2016, from <https://www.cms.gov/apps/physician-fee-schedule/license-agreement.aspx>
- Hamman, R. F., Wing, R. R., Edelstein, S. L., Lachin, J. M., Bray, G. A., Delahanty, L., . . . Wylie-Rosett, J. (2006). Effect of weight loss with lifestyle intervention on risk of diabetes. *Diabetes Care*, *29*(9), 2102–2107. <http://dx.doi.org/10.2337/dc06-0560>
- Herman, W. H., Hoerger, T. J., Brandle, M., Hicks, K., Sorensen, S., Zhang, P., . . . Ratner, R. E., & the Diabetes Prevention Program Research Group. (2005). The cost-effectiveness of lifestyle modification or metformin in preventing type 2 diabetes in adults with impaired glucose tolerance. *Annals of Internal Medicine*, *142*(5), 323–332. <http://dx.doi.org/10.7326/0003-4819-142-5-200503010-00007>
- Hernan, W. H., Brandle, M., Zhang, P., Williamson, D. F., Matulik, M. J., Ratner, R. E., . . . Engelgau, M. M., & the Diabetes Prevention Program Research Group. (2003). Costs associated with the primary prevention of type 2 diabetes mellitus in the diabetes prevention program. *Diabetes Care*, *26*(1), 36–47. <http://dx.doi.org/10.2337/diacare.26.1.36>
- Jones, R., Jones, R. O., McCowan, C., Montgomery, A. A., & Fahey, T. (2009). The external validity of published randomized controlled trials in primary care. *BMC Family Practice*, *10*(1), 5. <http://dx.doi.org/10.1186/1471-2296-10-5>
- Kiefer, M. M., Silverman, J. B., Young, B. A., & Nelson, K. M. (2015). National patterns in diabetes screening: Data from the National Health and Nutrition Examination Survey (NHANES) 2005–2012. *Journal of General Internal Medicine*, *30*(5), 612–618. <http://dx.doi.org/10.1007/s11606-014-3147-8>
- Knowler, W. C., Barrett-Connor, E., Fowler, S. E., Hamman, R. F., Lachin, J. M., Walker, E. A., & Nathan, D. M., & the Diabetes Prevention Program Research Group. (2002). Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *The New England Journal of Medicine*, *346*(6), 393–403. <http://dx.doi.org/10.1056/NEJMoa012512>
- Knowler, W. C., Fowler, S. E., Hamman, R. F., Christophi, C. A., Hoffman, H. J., Brenneman, A. T., . . . Nathan, D. M., & the Diabetes Prevention Program Research Group. (2009). 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet*, *374*(9702), 1677–1686. [http://dx.doi.org/10.1016/S0140-6736\(09\)61457-4](http://dx.doi.org/10.1016/S0140-6736(09)61457-4)
- Krukowski, R. A., Pope, R. A., Love, S., Lensing, S., Felix, H. C., Prewitt, T. E., & West, D. (2013). Examination of costs for a lay health educator-delivered translation of the Diabetes Prevention Program in senior centers. *Preventive Medicine*, *57*(4), 400–402. <http://dx.doi.org/10.1016/j.ypmed.2013.06.027>

- Laws, R. A., St George, A. B., Rychetnik, L., & Bauman, A. E. (2012). Diabetes prevention research: A systematic review of external validity in lifestyle interventions. *American Journal of Preventive Medicine, 43*(2), 205–214. <http://dx.doi.org/10.1016/j.amepre.2012.04.017>
- Li, R., Qu, S., Zhang, P., Chattopadhyay, S., Gregg, E. W., Albright, A., . . . Pronk, N. P. (2015). Economic evaluation of combined diet and physical activity promotion programs to prevent type 2 diabetes among persons at increased risk: A systematic review for the Community Preventive Services Task Force. *Annals of Internal Medicine, 163*(6), 452–460. <http://dx.doi.org/10.7326/M15-0469>
- Li, R., Zhang, P., Chattopadhyay, S., & Hopkins, D. (2015). *Economic evaluation of translational programs to prevent type 2 diabetes among high risk adults*. Unpublished presentation by the Centers for Disease Control and Prevention Division of Diabetes Translation.
- Luckhaupt, S. E., Cohen, M. A., Li, J., & Calvert, G. M. (2014). Prevalence of obesity among U.S. workers and associations with occupational factors. *American Journal of Preventive Medicine, 46*(3), 237–248. <http://dx.doi.org/10.1016/j.amepre.2013.11.002>
- Nichols, G. A., Glauber, H. S., & Brown, J. B. (2000). Type 2 diabetes: Incremental medical care costs during the 8 years preceding diagnosis. *Diabetes Care, 23*(11), 1654–1659. <http://dx.doi.org/10.2337/diacare.23.11.1654>
- Nichols, G. A., Hillier, T. A., & Brown, J. B. (2007). Progression from newly acquired impaired fasting glucose to type 2 diabetes. *Diabetes Care, 30*(2), 228–233. <http://dx.doi.org/10.2337/dc06-1392>
- Selvin, E., Steffes, M. W., Zhu, H., Matsushita, K., Wagenknecht, L., Pankow, J., . . . Brancati, F. L. (2010). Glycated hemoglobin, diabetes, and cardiovascular risk in nondiabetic adults. *The New England Journal of Medicine, 362*(9), 800–811. <http://dx.doi.org/10.1056/NEJMoa0908359>
- Shrestha, S., Zhang, P., Hora, I., Gregg, E. (2016). Trajectory of excess medical expenditures 9 years before and after the incidence of diabetes among adults, 2001–2013. Unpublished Abstract.
- Taradash, J., Kramer, M., Molenaar, D., Arena, V., Vanderwood, K., & Kriska, A. M. (2015). Recruitment for a Diabetes Prevention Program translation effort in a worksite setting. *Contemporary Clinical Trials, 41*, 204–210. <http://dx.doi.org/10.1016/j.cct.2015.01.010>
- U.S. Census Bureau. (2016). 2002 Indexes of Industry and Occupation. Retrieved from <http://www.census.gov/people/io/>
- Zhang, X., Gregg, E. W., Williamson, D. F., Barker, L. E., Thomas, W., Bullard, K. M., . . . Albright, A. L. (2010). A1C level and future risk of diabetes: A systematic review. *Diabetes Care, 33*(7), 1665–1673. <http://dx.doi.org/10.2337/dc09-1939>
- Zhuo, X., Zhang, P., Barker, L., Albright, A., Thompson, T. J., & Gregg, E. (2014). The lifetime cost of diabetes and its implications for diabetes prevention. *Diabetes Care, 37*(9), 2557–2564. <http://dx.doi.org/10.2337/dc13-2484>