

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,515,458	100.0	807.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	596,577	23.7	191.5
2	Malignant neoplasms (C00-C97)	576,691	22.9	185.1
3	Chronic lower respiratory diseases (J40-J47)	142,943	5.7	45.9
4	Cerebrovascular diseases (I60-I69)	128,932	5.1	41.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126,438	5.0	40.6
6	Alzheimer's disease (G30)	84,974	3.4	27.3
7	Diabetes mellitus (E10-E14)	73,831	2.9	23.7
8	Influenza and pneumonia (J09-J18)	53,826	2.1	17.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45,591	1.8	14.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39,518	1.6	12.7
11	Septicemia (A40-A41)	35,748	1.4	11.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	33,642	1.3	10.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27,853	1.1	8.9
14	Parkinson's disease (G20-G21)	23,111	0.9	7.4
15	Pneumonitis due to solids and liquids (J69)	18,195	0.7	5.8
...	All other causes (Residual)	507,588	20.2	162.9
All races, both sexes, 1-4 years				
...	All causes	4,246	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,377	32.4	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	493	11.6	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	412	9.7	2.5
4	Malignant neoplasms (C00-C97)	353	8.3	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	165	3.9	1.0
6	Influenza and pneumonia (J09-J18)	112	2.6	0.7
7	Septicemia (A40-A41)	61	1.4	0.4
8	Chronic lower respiratory diseases (J40-J47)	53	1.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	42	1.0	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	34	0.8	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	22	0.5	0.1
13	Anemias (D50-D64)	14	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	14	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
...	All other causes (Residual)	1,036	24.4	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	5,401	100.0	13.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,635	30.3	4.0
2	Malignant neoplasms (C00-C97)	860	15.9	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	358	6.6	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	287	5.3	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	283	5.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	203	3.8	0.5
7	Chronic lower respiratory diseases (J40-J47)	136	2.5	0.3
8	Influenza and pneumonia (J09-J18)	118	2.2	0.3
9	Cerebrovascular diseases (I60-I69)	87	1.6	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	1.3	0.2
11	Septicemia (A40-A41)	69	1.3	0.2
12	Anemias (D50-D64)	43	0.8	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	30	0.6	0.1
14	Diabetes mellitus (E10-E14)	26	0.5	0.1
15	Pneumonitis due to solids and liquids (J69)	18	0.3	*
...	All other causes (Residual)	1,178	21.8	2.9
All races, both sexes, 15-24 years				
...	All causes	29,667	100.0	67.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,330	41.6	28.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,822	16.3	11.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,554	15.4	10.4
4	Malignant neoplasms (C00-C97)	1,611	5.4	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	998	3.4	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	432	1.5	1.0
7	Influenza and pneumonia (J09-J18)	220	0.7	0.5
8	Cerebrovascular diseases (I60-I69)	186	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	172	0.6	0.4
10	Chronic lower respiratory diseases (J40-J47)	170	0.6	0.4
11	Diabetes mellitus (E10-E14)	167	0.6	0.4
12	Septicemia (A40-A41)	134	0.5	0.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	134	0.5	0.3
14	Legal intervention (Y35,Y89.0)	102	0.3	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	99	0.3	0.2
...	All other causes (Residual)	3,536	11.9	8.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	43,748	100.0	104.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,518	35.5	37.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,100	13.9	14.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,185	9.6	10.0
4	Malignant neoplasms (C00-C97)	3,499	8.0	8.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,301	7.5	7.9
6	Diabetes mellitus (E10-E14)	686	1.6	1.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	666	1.5	1.6
8	Cerebrovascular diseases (I60-I69)	530	1.2	1.3
9	Influenza and pneumonia (J09-J18)	515	1.2	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	505	1.2	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	410	0.9	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	386	0.9	0.9
13	Septicemia (A40-A41)	351	0.8	0.8
14	Chronic lower respiratory diseases (J40-J47)	268	0.6	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	212	0.5	0.5
...	All other causes (Residual)	6,616	15.1	15.8
All races, both sexes, 35-44 years				
...	All causes	69,893	100.0	172.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,230	21.8	37.5
2	Malignant neoplasms (C00-C97)	11,717	16.8	28.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,635	15.2	26.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,599	9.4	16.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,519	3.6	6.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,449	3.5	6.0
7	Diabetes mellitus (E10-E14)	1,842	2.6	4.5
8	Cerebrovascular diseases (I60-I69)	1,718	2.5	4.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,619	2.3	4.0
10	Influenza and pneumonia (J09-J18)	859	1.2	2.1
11	Septicemia (A40-A41)	794	1.1	2.0
12	Chronic lower respiratory diseases (J40-J47)	734	1.1	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	668	1.0	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	419	0.6	1.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	411	0.6	1.0
...	All other causes (Residual)	11,680	16.7	28.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	183,247	100.0	409.8
1	Malignant neoplasms (C00-C97)	48,897	26.7	109.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,100	19.7	80.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,749	11.3	46.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	8,864	4.8	19.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,858	4.8	19.8
6	Diabetes mellitus (E10-E14)	6,012	3.3	13.4
7	Cerebrovascular diseases (I60-I69)	5,705	3.1	12.8
8	Chronic lower respiratory diseases (J40-J47)	4,634	2.5	10.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2,781	1.5	6.2
10	Septicemia (A40-A41)	2,461	1.3	5.5
11	Viral hepatitis (B15-B19)	2,317	1.3	5.2
12	Influenza and pneumonia (J09-J18)	2,224	1.2	5.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,052	1.1	4.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,984	1.1	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,375	0.8	3.1
...	All other causes (Residual)	28,234	15.4	63.1
All races, both sexes, 55-64 years				
...	All causes	323,315	100.0	849.4
1	Malignant neoplasms (C00-C97)	112,572	34.8	295.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,742	21.6	183.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,158	4.7	39.8
4	Chronic lower respiratory diseases (J40-J47)	15,044	4.7	39.5
5	Diabetes mellitus (E10-E14)	12,688	3.9	33.3
6	Cerebrovascular diseases (I60-I69)	11,205	3.5	29.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	10,749	3.3	28.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,521	2.0	17.1
9	Septicemia (A40-A41)	4,953	1.5	13.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,754	1.5	12.5
11	Influenza and pneumonia (J09-J18)	4,185	1.3	11.0
12	Viral hepatitis (B15-B19)	3,515	1.1	9.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,679	0.8	7.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,796	0.6	4.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,485	0.5	3.9
...	All other causes (Residual)	46,269	14.3	121.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	415,052	100.0	1,846.2
1	Malignant neoplasms (C00-C97)	145,593	35.1	647.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	89,708	21.6	399.0
3	Chronic lower respiratory diseases (J40-J47)	32,433	7.8	144.3
4	Cerebrovascular diseases (I60-I69)	17,584	4.2	78.2
5	Diabetes mellitus (E10-E14)	16,189	3.9	72.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,014	2.4	44.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,682	1.9	34.2
8	Septicemia (A40-A41)	6,636	1.6	29.5
9	Influenza and pneumonia (J09-J18)	6,488	1.6	28.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5,905	1.4	26.3
11	Alzheimer's disease (G30)	4,325	1.0	19.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,736	0.9	16.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,179	0.8	14.1
14	Parkinson's disease (G20-G21)	2,870	0.7	12.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,585	0.6	11.5
...	All other causes (Residual)	60,125	14.5	267.4
All races, both sexes, 75-84 years				
...	All causes	626,225	100.0	4,753.0
1	Malignant neoplasms (C00-C97)	155,348	24.8	1,179.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	149,500	23.9	1,134.7
3	Chronic lower respiratory diseases (J40-J47)	49,394	7.9	374.9
4	Cerebrovascular diseases (I60-I69)	37,596	6.0	285.4
5	Alzheimer's disease (G30)	24,224	3.9	183.9
6	Diabetes mellitus (E10-E14)	19,605	3.1	148.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,093	2.3	107.0
8	Influenza and pneumonia (J09-J18)	13,698	2.2	104.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,358	2.1	101.4
10	Parkinson's disease (G20-G21)	10,019	1.6	76.0
11	Septicemia (A40-A41)	9,798	1.6	74.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,769	1.1	51.4
13	Pneumonitis due to solids and liquids (J69)	5,139	0.8	39.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,638	0.7	35.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,859	0.6	29.3
...	All other causes (Residual)	109,187	17.4	828.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	790,545	100.0	13,779.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	235,889	29.8	4,111.6
2	Malignant neoplasms (C00-C97)	96,165	12.2	1,676.2
3	Alzheimer's disease (G30)	55,483	7.0	967.1
4	Cerebrovascular diseases (I60-I69)	54,143	6.8	943.7
5	Chronic lower respiratory diseases (J40-J47)	40,042	5.1	697.9
6	Influenza and pneumonia (J09-J18)	25,200	3.2	439.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,151	2.4	333.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,756	2.1	292.1
9	Diabetes mellitus (E10-E14)	16,608	2.1	289.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,775	1.6	222.7
11	Septicemia (A40-A41)	10,312	1.3	179.7
12	Parkinson's disease (G20-G21)	9,646	1.2	168.1
13	Pneumonitis due to solids and liquids (J69)	9,096	1.2	158.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,943	0.6	86.2
15	Atherosclerosis (I70)	3,952	0.5	68.9
...	All other causes (Residual)	180,384	22.8	3,144.1
All races, male, all ages				
...	All causes	1,254,978	100.0	818.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	308,398	24.6	201.2
2	Malignant neoplasms (C00-C97)	302,231	24.1	197.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79,257	6.3	51.7
4	Chronic lower respiratory diseases (J40-J47)	67,521	5.4	44.0
5	Cerebrovascular diseases (I60-I69)	52,335	4.2	34.1
6	Diabetes mellitus (E10-E14)	38,324	3.1	25.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31,003	2.5	20.2
8	Alzheimer's disease (G30)	25,677	2.0	16.8
9	Influenza and pneumonia (J09-J18)	25,401	2.0	16.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,649	1.8	14.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	21,621	1.7	14.1
12	Septicemia (A40-A41)	16,484	1.3	10.8
13	Parkinson's disease (G20-G21)	13,546	1.1	8.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12,745	1.0	8.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,303	0.9	7.4
...	All other causes (Residual)	226,483	18.0	147.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,405	100.0	29.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	839	34.9	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	266	11.1	3.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	239	9.9	2.9
4	Malignant neoplasms (C00-C97)	183	7.6	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	3.5	1.0
6	Influenza and pneumonia (J09-J18)	55	2.3	0.7
7	Chronic lower respiratory diseases (J40-J47)	37	1.5	0.4
8	Cerebrovascular diseases (I60-I69)	29	1.2	0.4
9	Septicemia (A40-A41)	26	1.1	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.1	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	20	0.8	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.5	*
13	Anemias (D50-D64)	9	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	8	0.3	*
15	Meningitis (G00,G03)	7	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
...	All other causes (Residual)	557	23.2	6.7
All races, male, 5-14 years				
...	All causes	3,179	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,033	32.5	4.9
2	Malignant neoplasms (C00-C97)	483	15.2	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	201	6.3	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	6.2	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	5.6	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	3.0	0.5
7	Chronic lower respiratory diseases (J40-J47)	87	2.7	0.4
8	Influenza and pneumonia (J09-J18)	63	2.0	0.3
9	Cerebrovascular diseases (I60-I69)	46	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.2	0.2
11	Septicemia (A40-A41)	34	1.1	0.2
12	Anemias (D50-D64)	27	0.8	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	16	0.5	*
14	Pneumonitis due to solids and liquids (J69)	14	0.4	*
15	Diabetes mellitus (E10-E14)	11	0.3	*
...	All other causes (Residual)	656	20.6	3.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	21,924	100.0	97.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,172	41.8	40.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,959	18.1	17.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,941	18.0	17.6
4	Malignant neoplasms (C00-C97)	966	4.4	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	684	3.1	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	275	1.3	1.2
7	Influenza and pneumonia (J09-J18)	123	0.6	0.5
8	Chronic lower respiratory diseases (J40-J47)	117	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	104	0.5	0.5
10	Diabetes mellitus (E10-E14)	101	0.5	0.5
11	Legal intervention (Y35,Y89.0)	100	0.5	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	80	0.4	0.4
13	Septicemia (A40-A41)	72	0.3	0.3
14	Anemias (D50-D64)	60	0.3	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.2	0.2
...	All other causes (Residual)	2,125	9.7	9.5
All races, male, 25-34 years				
...	All causes	30,085	100.0	143.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,426	38.0	54.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,862	16.2	23.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,499	11.6	16.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,210	7.3	10.5
5	Malignant neoplasms (C00-C97)	1,713	5.7	8.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	428	1.4	2.0
7	Diabetes mellitus (E10-E14)	388	1.3	1.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	306	1.0	1.5
9	Influenza and pneumonia (J09-J18)	290	1.0	1.4
10	Cerebrovascular diseases (I60-I69)	284	0.9	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	229	0.8	1.1
12	Septicemia (A40-A41)	188	0.6	0.9
13	Chronic lower respiratory diseases (J40-J47)	157	0.5	0.7
14	Legal intervention (Y35,Y89.0)	132	0.4	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	119	0.4	0.6
...	All other causes (Residual)	3,854	12.8	18.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	43,215	100.0	213.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,459	24.2	51.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,393	17.1	36.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,044	11.7	24.9
4	Malignant neoplasms (C00-C97)	5,034	11.6	24.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,940	4.5	9.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,572	3.6	7.8
7	Diabetes mellitus (E10-E14)	1,120	2.6	5.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,060	2.5	5.2
9	Cerebrovascular diseases (I60-I69)	906	2.1	4.5
10	Influenza and pneumonia (J09-J18)	456	1.1	2.3
11	Septicemia (A40-A41)	399	0.9	2.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	396	0.9	2.0
13	Chronic lower respiratory diseases (J40-J47)	340	0.8	1.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	258	0.6	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	0.6	1.2
...	All other causes (Residual)	6,594	15.3	32.6
All races, male, 45-54 years				
...	All causes	111,709	100.0	507.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,691	23.0	116.7
2	Malignant neoplasms (C00-C97)	24,372	21.8	110.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,917	12.5	63.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,637	5.9	30.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,868	5.3	26.6
6	Diabetes mellitus (E10-E14)	3,743	3.4	17.0
7	Cerebrovascular diseases (I60-I69)	3,201	2.9	14.5
8	Chronic lower respiratory diseases (J40-J47)	2,173	1.9	9.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2,052	1.8	9.3
10	Viral hepatitis (B15-B19)	1,637	1.5	7.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,496	1.3	6.8
12	Septicemia (A40-A41)	1,296	1.2	5.9
13	Influenza and pneumonia (J09-J18)	1,288	1.2	5.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,153	1.0	5.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	879	0.8	4.0
...	All other causes (Residual)	16,306	14.6	74.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	196,641	100.0	1,071.1
1	Malignant neoplasms (C00-C97)	62,499	31.8	340.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,388	24.6	263.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,441	5.3	56.9
4	Chronic lower respiratory diseases (J40-J47)	7,890	4.0	43.0
5	Diabetes mellitus (E10-E14)	7,624	3.9	41.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	7,589	3.9	41.3
7	Cerebrovascular diseases (I60-I69)	6,405	3.3	34.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,018	2.6	27.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,701	1.4	14.7
10	Septicemia (A40-A41)	2,648	1.3	14.4
11	Viral hepatitis (B15-B19)	2,571	1.3	14.0
12	Influenza and pneumonia (J09-J18)	2,430	1.2	13.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,657	0.8	9.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,360	0.7	7.4
15	Aortic aneurysm and dissection (I71)	975	0.5	5.3
...	All other causes (Residual)	26,445	13.4	144.1
All races, male, 65-74 years				
...	All causes	234,276	100.0	2,236.2
1	Malignant neoplasms (C00-C97)	80,223	34.2	765.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,609	24.2	540.4
3	Chronic lower respiratory diseases (J40-J47)	16,497	7.0	157.5
4	Cerebrovascular diseases (I60-I69)	9,228	3.9	88.1
5	Diabetes mellitus (E10-E14)	9,181	3.9	87.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,295	2.7	60.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,083	1.7	39.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	3,676	1.6	35.1
9	Influenza and pneumonia (J09-J18)	3,585	1.5	34.2
10	Septicemia (A40-A41)	3,379	1.4	32.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,554	1.1	24.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,956	0.8	18.7
13	Parkinson's disease (G20-G21)	1,922	0.8	18.3
14	Alzheimer's disease (G30)	1,866	0.8	17.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,502	0.6	14.3
...	All other causes (Residual)	31,720	13.5	302.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	312,693	100.0	5,610.8
1	Malignant neoplasms (C00-C97)	82,399	26.4	1,478.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	79,504	25.4	1,426.6
3	Chronic lower respiratory diseases (J40-J47)	23,736	7.6	425.9
4	Cerebrovascular diseases (I60-I69)	16,147	5.2	289.7
5	Diabetes mellitus (E10-E14)	9,676	3.1	173.6
6	Alzheimer's disease (G30)	9,128	2.9	163.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,569	2.4	135.8
8	Influenza and pneumonia (J09-J18)	6,934	2.2	124.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,832	2.2	122.6
10	Parkinson's disease (G20-G21)	6,163	2.0	110.6
11	Septicemia (A40-A41)	4,611	1.5	82.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,919	0.9	52.4
13	Pneumonitis due to solids and liquids (J69)	2,909	0.9	52.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,525	0.8	45.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,034	0.7	36.5
...	All other causes (Residual)	49,607	15.9	890.1
All races, male, 85 years and over				
...	All causes	285,425	100.0	15,069.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	87,556	30.7	4,622.7
2	Malignant neoplasms (C00-C97)	44,317	15.5	2,339.8
3	Chronic lower respiratory diseases (J40-J47)	16,464	5.8	869.3
4	Cerebrovascular diseases (I60-I69)	15,909	5.6	840.0
5	Alzheimer's disease (G30)	14,266	5.0	753.2
6	Influenza and pneumonia (J09-J18)	10,053	3.5	530.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,440	2.6	392.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,268	2.5	383.7
9	Diabetes mellitus (E10-E14)	6,476	2.3	341.9
10	Parkinson's disease (G20-G21)	5,102	1.8	269.4
11	Pneumonitis due to solids and liquids (J69)	4,383	1.5	231.4
12	Septicemia (A40-A41)	3,733	1.3	197.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,570	1.3	188.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,194	0.8	115.8
15	Atherosclerosis (I70)	1,215	0.4	64.1
...	All other causes (Residual)	55,479	19.4	2,929.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,260,480	100.0	796.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	288,179	22.9	182.0
2	Malignant neoplasms (C00-C97)	274,460	21.8	173.4
3	Cerebrovascular diseases (I60-I69)	76,597	6.1	48.4
4	Chronic lower respiratory diseases (J40-J47)	75,422	6.0	47.6
5	Alzheimer's disease (G30)	59,297	4.7	37.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47,181	3.7	29.8
7	Diabetes mellitus (E10-E14)	35,507	2.8	22.4
8	Influenza and pneumonia (J09-J18)	28,425	2.3	18.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,942	1.8	14.5
10	Septicemia (A40-A41)	19,264	1.5	12.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16,550	1.3	10.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12,021	1.0	7.6
13	Parkinson's disease (G20-G21)	9,565	0.8	6.0
14	Pneumonitis due to solids and liquids (J69)	8,519	0.7	5.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,515	0.7	5.4
...	All other causes (Residual)	278,036	22.1	175.6
All races, female, 1-4 years				
...	All causes	1,841	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	538	29.2	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	12.3	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	173	9.4	2.2
4	Malignant neoplasms (C00-C97)	170	9.2	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	4.4	1.0
6	Influenza and pneumonia (J09-J18)	57	3.1	0.7
7	Septicemia (A40-A41)	35	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	16	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	14	0.8	*
11	Cerebrovascular diseases (I60-I69)	13	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
15	Anemias (D50-D64)	5	0.3	*
...	All other causes (Residual)	472	25.6	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,222	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	602	27.1	3.0
2	Malignant neoplasms (C00-C97)	377	17.0	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	7.2	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	4.9	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	4.7	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	3.9	0.4
7	Influenza and pneumonia (J09-J18)	55	2.5	0.3
8	Chronic lower respiratory diseases (J40-J47)	49	2.2	0.2
9	Cerebrovascular diseases (I60-I69)	41	1.8	0.2
10	Septicemia (A40-A41)	35	1.6	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.5	0.2
12	Anemias (D50-D64)	16	0.7	*
13	Diabetes mellitus (E10-E14)	15	0.7	*
14	Certain conditions originating in the perinatal period (P00-P96)	14	0.6	*
15	Meningitis (G00,G03)	8	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
...	All other causes (Residual)	510	23.0	2.5
All races, female, 15-24 years				
...	All causes	7,743	100.0	36.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,158	40.8	14.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	863	11.1	4.0
3	Malignant neoplasms (C00-C97)	645	8.3	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	613	7.9	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	314	4.1	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	172	2.2	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	157	2.0	0.7
8	Influenza and pneumonia (J09-J18)	97	1.3	0.5
9	Cerebrovascular diseases (I60-I69)	82	1.1	0.4
10	Diabetes mellitus (E10-E14)	66	0.9	0.3
11	Septicemia (A40-A41)	62	0.8	0.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	54	0.7	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.7	0.3
14	Chronic lower respiratory diseases (J40-J47)	53	0.7	0.2
15	Anemias (D50-D64)	37	0.5	0.2
...	All other causes (Residual)	1,316	17.0	6.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	13,663	100.0	65.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,092	29.9	19.7
2	Malignant neoplasms (C00-C97)	1,786	13.1	8.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,238	9.1	6.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,091	8.0	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	686	5.0	3.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	386	2.8	1.9
7	Diabetes mellitus (E10-E14)	298	2.2	1.4
8	Cerebrovascular diseases (I60-I69)	246	1.8	1.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	238	1.7	1.1
10	Influenza and pneumonia (J09-J18)	225	1.6	1.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	199	1.5	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	181	1.3	0.9
13	Septicemia (A40-A41)	163	1.2	0.8
14	Chronic lower respiratory diseases (J40-J47)	111	0.8	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	0.7	0.4
...	All other causes (Residual)	2,630	19.2	12.7
All races, female, 35-44 years				
...	All causes	26,678	100.0	130.7
1	Malignant neoplasms (C00-C97)	6,683	25.1	32.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,771	17.9	23.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,242	12.2	15.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,555	5.8	7.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	877	3.3	4.3
6	Cerebrovascular diseases (I60-I69)	812	3.0	4.0
7	Diabetes mellitus (E10-E14)	722	2.7	3.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	579	2.2	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	559	2.1	2.7
10	Influenza and pneumonia (J09-J18)	403	1.5	2.0
11	Septicemia (A40-A41)	395	1.5	1.9
12	Chronic lower respiratory diseases (J40-J47)	394	1.5	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	272	1.0	1.3
14	Pregnancy, childbirth and the puerperium (O00-O99)	260	1.0	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	175	0.7	0.9
...	All other causes (Residual)	4,979	18.7	24.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	71,538	100.0	315.2
1	Malignant neoplasms (C00-C97)	24,525	34.3	108.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,409	14.6	45.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,832	9.6	30.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,996	4.2	13.2
5	Cerebrovascular diseases (I60-I69)	2,504	3.5	11.0
6	Chronic lower respiratory diseases (J40-J47)	2,461	3.4	10.8
7	Diabetes mellitus (E10-E14)	2,269	3.2	10.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,221	3.1	9.8
9	Septicemia (A40-A41)	1,165	1.6	5.1
10	Influenza and pneumonia (J09-J18)	936	1.3	4.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	831	1.2	3.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	729	1.0	3.2
13	Viral hepatitis (B15-B19)	680	1.0	3.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	556	0.8	2.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	496	0.7	2.2
...	All other causes (Residual)	11,928	16.7	52.5
All races, female, 55-64 years				
...	All causes	126,674	100.0	642.9
1	Malignant neoplasms (C00-C97)	50,073	39.5	254.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,354	16.9	108.4
3	Chronic lower respiratory diseases (J40-J47)	7,154	5.6	36.3
4	Diabetes mellitus (E10-E14)	5,064	4.0	25.7
5	Cerebrovascular diseases (I60-I69)	4,800	3.8	24.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,717	3.7	23.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,160	2.5	16.0
8	Septicemia (A40-A41)	2,305	1.8	11.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,053	1.6	10.4
10	Influenza and pneumonia (J09-J18)	1,755	1.4	8.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,503	1.2	7.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,022	0.8	5.2
13	Viral hepatitis (B15-B19)	944	0.7	4.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	650	0.5	3.3
15	Alzheimer's disease (G30)	465	0.4	2.4
...	All other causes (Residual)	19,655	15.5	99.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	180,776	100.0	1,505.8
1	Malignant neoplasms (C00-C97)	65,370	36.2	544.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,099	18.3	275.7
3	Chronic lower respiratory diseases (J40-J47)	15,936	8.8	132.7
4	Cerebrovascular diseases (I60-I69)	8,356	4.6	69.6
5	Diabetes mellitus (E10-E14)	7,008	3.9	58.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,719	2.1	31.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,599	2.0	30.0
8	Septicemia (A40-A41)	3,257	1.8	27.1
9	Influenza and pneumonia (J09-J18)	2,903	1.6	24.2
10	Alzheimer's disease (G30)	2,459	1.4	20.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,229	1.2	18.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,780	1.0	14.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,083	0.6	9.0
14	Parkinson's disease (G20-G21)	948	0.5	7.9
15	Pneumonitis due to solids and liquids (J69)	804	0.4	6.7
...	All other causes (Residual)	28,226	15.6	235.1
All races, female, 75-84 years				
...	All causes	313,532	100.0	4,124.2
1	Malignant neoplasms (C00-C97)	72,949	23.3	959.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,996	22.3	920.7
3	Chronic lower respiratory diseases (J40-J47)	25,658	8.2	337.5
4	Cerebrovascular diseases (I60-I69)	21,449	6.8	282.1
5	Alzheimer's disease (G30)	15,096	4.8	198.6
6	Diabetes mellitus (E10-E14)	9,929	3.2	130.6
7	Influenza and pneumonia (J09-J18)	6,764	2.2	89.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,526	2.1	85.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,524	2.1	85.8
10	Septicemia (A40-A41)	5,187	1.7	68.2
11	Parkinson's disease (G20-G21)	3,856	1.2	50.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,850	1.2	50.6
13	Pneumonitis due to solids and liquids (J69)	2,230	0.7	29.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,113	0.7	27.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,825	0.6	24.0
...	All other causes (Residual)	59,580	19.0	783.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	505,120	100.0	13,143.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	148,333	29.4	3,859.7
2	Malignant neoplasms (C00-C97)	51,848	10.3	1,349.1
3	Alzheimer's disease (G30)	41,217	8.2	1,072.5
4	Cerebrovascular diseases (I60-I69)	38,234	7.6	994.9
5	Chronic lower respiratory diseases (J40-J47)	23,578	4.7	613.5
6	Influenza and pneumonia (J09-J18)	15,147	3.0	394.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,711	2.3	304.7
8	Diabetes mellitus (E10-E14)	10,132	2.0	263.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,488	1.9	246.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,205	1.8	239.5
11	Septicemia (A40-A41)	6,579	1.3	171.2
12	Pneumonitis due to solids and liquids (J69)	4,713	0.9	122.6
13	Parkinson's disease (G20-G21)	4,544	0.9	118.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,749	0.5	71.5
15	Atherosclerosis (I70)	2,737	0.5	71.2
...	All other causes (Residual)	124,905	24.7	3,250.1
White, both sexes, all ages				
...	All causes	2,156,077	100.0	872.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	513,701	23.8	207.9
2	Malignant neoplasms (C00-C97)	492,348	22.8	199.3
3	Chronic lower respiratory diseases (J40-J47)	131,495	6.1	53.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109,751	5.1	44.4
5	Cerebrovascular diseases (I60-I69)	108,542	5.0	43.9
6	Alzheimer's disease (G30)	78,055	3.6	31.6
7	Diabetes mellitus (E10-E14)	57,974	2.7	23.5
8	Influenza and pneumonia (J09-J18)	46,658	2.2	18.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,268	1.7	14.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35,775	1.7	14.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	29,507	1.4	11.9
12	Septicemia (A40-A41)	28,838	1.3	11.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21,845	1.0	8.8
14	Parkinson's disease (G20-G21)	21,688	1.0	8.8
15	Pneumonitis due to solids and liquids (J69)	16,263	0.8	6.6
...	All other causes (Residual)	427,369	19.8	173.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	2,980	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,027	34.5	8.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	360	12.1	3.0
3	Malignant neoplasms (C00-C97)	275	9.2	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	237	8.0	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	3.6	0.9
6	Influenza and pneumonia (J09-J18)	66	2.2	0.5
7	Septicemia (A40-A41)	45	1.5	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.1	0.3
9	Cerebrovascular diseases (I60-I69)	23	0.8	0.2
10	Chronic lower respiratory diseases (J40-J47)	22	0.7	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	21	0.7	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	12	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.3	*
15	Meningitis (G00,G03)	8	0.3	*
...	All other causes (Residual)	721	24.2	5.9
White, both sexes, 5-14 years				
...	All causes	3,927	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,242	31.6	4.0
2	Malignant neoplasms (C00-C97)	676	17.2	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	258	6.6	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	222	5.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	4.5	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	142	3.6	0.5
7	Influenza and pneumonia (J09-J18)	83	2.1	0.3
8	Cerebrovascular diseases (I60-I69)	59	1.5	0.2
9	Septicemia (A40-A41)	55	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	49	1.2	0.2
12	Diabetes mellitus (E10-E14)	21	0.5	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	19	0.5	*
14	Anemias (D50-D64)	15	0.4	*
15	Pneumonitis due to solids and liquids (J69)	11	0.3	*
...	All other causes (Residual)	847	21.6	2.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	21,654	100.0	65.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,391	48.0	31.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,023	18.6	12.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,611	7.4	4.8
4	Malignant neoplasms (C00-C97)	1,267	5.9	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	643	3.0	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	306	1.4	0.9
7	Influenza and pneumonia (J09-J18)	165	0.8	0.5
8	Cerebrovascular diseases (I60-I69)	143	0.7	0.4
9	Diabetes mellitus (E10-E14)	115	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	102	0.5	0.3
11	Septicemia (A40-A41)	92	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	81	0.4	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.3	0.2
14	Legal intervention (Y35,Y89.0)	49	0.2	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.1
...	All other causes (Residual)	2,554	11.8	7.7
White, both sexes, 25-34 years				
...	All causes	32,715	100.0	101.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,245	40.5	41.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,227	16.0	16.2
3	Malignant neoplasms (C00-C97)	2,647	8.1	8.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,192	6.7	6.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,704	5.2	5.3
6	Diabetes mellitus (E10-E14)	432	1.3	1.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	393	1.2	1.2
8	Influenza and pneumonia (J09-J18)	390	1.2	1.2
9	Cerebrovascular diseases (I60-I69)	363	1.1	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	337	1.0	1.0
11	Septicemia (A40-A41)	237	0.7	0.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	232	0.7	0.7
13	Pregnancy, childbirth and the puerperium (O00-O99)	230	0.7	0.7
14	Chronic lower respiratory diseases (J40-J47)	163	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	128	0.4	0.4
...	All other causes (Residual)	4,795	14.7	14.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	52,998	100.0	167.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,004	24.5	41.0
2	Malignant neoplasms (C00-C97)	8,960	16.9	28.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,511	14.2	23.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,875	11.1	18.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,056	3.9	6.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,271	2.4	4.0
7	Diabetes mellitus (E10-E14)	1,257	2.4	4.0
8	Cerebrovascular diseases (I60-I69)	1,077	2.0	3.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	690	1.3	2.2
10	Influenza and pneumonia (J09-J18)	628	1.2	2.0
11	Septicemia (A40-A41)	543	1.0	1.7
12	Chronic lower respiratory diseases (J40-J47)	529	1.0	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	369	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	342	0.6	1.1
15	Viral hepatitis (B15-B19)	247	0.5	0.8
...	All other causes (Residual)	8,639	16.3	27.3
White, both sexes, 45-54 years				
...	All causes	143,426	100.0	398.0
1	Malignant neoplasms (C00-C97)	38,503	26.8	106.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,198	19.0	75.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,752	12.4	49.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,268	5.8	22.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	7,686	5.4	21.3
6	Diabetes mellitus (E10-E14)	4,307	3.0	12.0
7	Chronic lower respiratory diseases (J40-J47)	3,855	2.7	10.7
8	Cerebrovascular diseases (I60-I69)	3,747	2.6	10.4
9	Viral hepatitis (B15-B19)	1,942	1.4	5.4
10	Septicemia (A40-A41)	1,756	1.2	4.9
11	Influenza and pneumonia (J09-J18)	1,730	1.2	4.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,292	0.9	3.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,231	0.9	3.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,204	0.8	3.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	832	0.6	2.3
...	All other causes (Residual)	22,123	15.4	61.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	259,690	100.0	821.2
1	Malignant neoplasms (C00-C97)	91,865	35.4	290.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,694	21.1	173.0
3	Chronic lower respiratory diseases (J40-J47)	13,319	5.1	42.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,664	4.9	40.0
5	Diabetes mellitus (E10-E14)	9,420	3.6	29.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	9,271	3.6	29.3
7	Cerebrovascular diseases (I60-I69)	7,934	3.1	25.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,151	2.4	19.5
9	Septicemia (A40-A41)	3,667	1.4	11.6
10	Influenza and pneumonia (J09-J18)	3,338	1.3	10.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,185	1.2	10.1
12	Viral hepatitis (B15-B19)	2,782	1.1	8.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,744	0.7	5.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,225	0.5	3.9
15	Aortic aneurysm and dissection (I71)	1,057	0.4	3.3
...	All other causes (Residual)	37,374	14.4	118.2
White, both sexes, 65-74 years				
...	All causes	350,039	100.0	1,824.8
1	Malignant neoplasms (C00-C97)	124,336	35.5	648.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,927	21.1	385.4
3	Chronic lower respiratory diseases (J40-J47)	29,768	8.5	155.2
4	Cerebrovascular diseases (I60-I69)	13,812	3.9	72.0
5	Diabetes mellitus (E10-E14)	12,509	3.6	65.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,711	2.5	45.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,655	1.6	29.5
8	Influenza and pneumonia (J09-J18)	5,419	1.5	28.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	5,332	1.5	27.8
10	Septicemia (A40-A41)	5,238	1.5	27.3
11	Alzheimer's disease (G30)	3,846	1.1	20.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,995	0.9	15.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,682	0.8	14.0
14	Parkinson's disease (G20-G21)	2,614	0.7	13.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,267	0.6	11.8
...	All other causes (Residual)	50,928	14.5	265.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	551,927	100.0	4,797.1
1	Malignant neoplasms (C00-C97)	136,992	24.8	1,190.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	130,705	23.7	1,136.0
3	Chronic lower respiratory diseases (J40-J47)	46,087	8.4	400.6
4	Cerebrovascular diseases (I60-I69)	32,255	5.8	280.3
5	Alzheimer's disease (G30)	22,102	4.0	192.1
6	Diabetes mellitus (E10-E14)	15,887	2.9	138.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,871	2.3	111.9
8	Influenza and pneumonia (J09-J18)	11,920	2.2	103.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,927	2.0	95.0
10	Parkinson's disease (G20-G21)	9,391	1.7	81.6
11	Septicemia (A40-A41)	8,145	1.5	70.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,408	1.0	47.0
13	Pneumonitis due to solids and liquids (J69)	4,603	0.8	40.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,168	0.8	36.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,571	0.6	31.0
...	All other causes (Residual)	96,895	17.6	842.2
White, both sexes, 85 years and over				
...	All causes	721,162	100.0	14,031.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	216,373	30.0	4,209.8
2	Malignant neoplasms (C00-C97)	86,773	12.0	1,688.3
3	Alzheimer's disease (G30)	51,257	7.1	997.3
4	Cerebrovascular diseases (I60-I69)	49,045	6.8	954.2
5	Chronic lower respiratory diseases (J40-J47)	37,602	5.2	731.6
6	Influenza and pneumonia (J09-J18)	22,802	3.2	443.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,107	2.5	352.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,711	2.0	286.2
9	Diabetes mellitus (E10-E14)	14,019	1.9	272.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,922	1.5	212.5
11	Parkinson's disease (G20-G21)	9,173	1.3	178.5
12	Septicemia (A40-A41)	8,961	1.2	174.3
13	Pneumonitis due to solids and liquids (J69)	8,372	1.2	162.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,578	0.6	89.1
15	Atherosclerosis (I70)	3,656	0.5	71.1
...	All other causes (Residual)	164,811	22.9	3,206.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,071,966	100.0	876.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	265,596	24.8	217.1
2	Malignant neoplasms (C00-C97)	259,323	24.2	212.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68,123	6.4	55.7
4	Chronic lower respiratory diseases (J40-J47)	61,507	5.7	50.3
5	Cerebrovascular diseases (I60-I69)	43,264	4.0	35.4
6	Diabetes mellitus (E10-E14)	30,783	2.9	25.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,103	2.6	23.0
8	Alzheimer's disease (G30)	23,640	2.2	19.3
9	Influenza and pneumonia (J09-J18)	21,953	2.0	17.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	19,067	1.8	15.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,275	1.7	14.9
12	Septicemia (A40-A41)	13,371	1.2	10.9
13	Parkinson's disease (G20-G21)	12,746	1.2	10.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,698	0.8	7.1
15	Pneumonitis due to solids and liquids (J69)	8,653	0.8	7.1
...	All other causes (Residual)	188,864	17.6	154.4
White, male, 1-4 years				
...	All causes	1,689	100.0	27.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	634	37.5	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	193	11.4	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	8.5	2.3
4	Malignant neoplasms (C00-C97)	136	8.1	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.3	0.9
6	Influenza and pneumonia (J09-J18)	32	1.9	0.5
7	Septicemia (A40-A41)	20	1.2	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
10	Cerebrovascular diseases (I60-I69)	14	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
14	Meningitis (G00,G03)	6	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (Residual)	389	23.0	6.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,321	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	791	34.1	4.9
2	Malignant neoplasms (C00-C97)	379	16.3	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	157	6.8	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	141	6.1	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	107	4.6	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	2.8	0.4
7	Influenza and pneumonia (J09-J18)	43	1.9	0.3
8	Cerebrovascular diseases (I60-I69)	33	1.4	0.2
9	Septicemia (A40-A41)	29	1.2	0.2
9	Chronic lower respiratory diseases (J40-J47)	29	1.2	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.2	0.2
12	Anemias (D50-D64)	11	0.5	*
13	Pneumonitis due to solids and liquids (J69)	8	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.3	*
15	Diabetes mellitus (E10-E14)	7	0.3	*
...	All other causes (Residual)	486	20.9	3.0
White, male, 15-24 years				
...	All causes	15,839	100.0	92.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,724	48.8	45.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,307	20.9	19.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,308	8.3	7.6
4	Malignant neoplasms (C00-C97)	765	4.8	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	452	2.9	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	199	1.3	1.2
7	Influenza and pneumonia (J09-J18)	98	0.6	0.6
8	Cerebrovascular diseases (I60-I69)	83	0.5	0.5
9	Diabetes mellitus (E10-E14)	76	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	60	0.4	0.4
11	Septicemia (A40-A41)	55	0.3	0.3
12	Legal intervention (Y35,Y89.0)	48	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.1	0.1
15	Pneumonitis due to solids and liquids (J69)	21	0.1	0.1
...	All other causes (Residual)	1,590	10.0	9.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	22,675	100.0	138.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,779	43.1	59.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,181	18.4	25.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,517	6.7	9.2
4	Malignant neoplasms (C00-C97)	1,338	5.9	8.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,303	5.7	7.9
6	Diabetes mellitus (E10-E14)	258	1.1	1.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	247	1.1	1.5
8	Influenza and pneumonia (J09-J18)	226	1.0	1.4
9	Cerebrovascular diseases (I60-I69)	199	0.9	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	0.8	1.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	171	0.8	1.0
12	Septicemia (A40-A41)	139	0.6	0.8
13	Legal intervention (Y35,Y89.0)	95	0.4	0.6
14	Chronic lower respiratory diseases (J40-J47)	87	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.3	0.5
...	All other causes (Residual)	2,868	12.6	17.5
White, male, 35-44 years				
...	All causes	33,396	100.0	208.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,881	26.6	55.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,358	16.0	33.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,491	13.4	28.1
4	Malignant neoplasms (C00-C97)	4,032	12.1	25.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,356	4.1	8.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	896	2.7	5.6
7	Diabetes mellitus (E10-E14)	784	2.3	4.9
8	Cerebrovascular diseases (I60-I69)	578	1.7	3.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	532	1.6	3.3
10	Influenza and pneumonia (J09-J18)	335	1.0	2.1
11	Septicemia (A40-A41)	268	0.8	1.7
12	Chronic lower respiratory diseases (J40-J47)	248	0.7	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	229	0.7	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	200	0.6	1.3
15	Aortic aneurysm and dissection (I71)	157	0.5	1.0
...	All other causes (Residual)	5,051	15.1	31.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	88,780	100.0	495.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,861	22.4	110.8
2	Malignant neoplasms (C00-C97)	19,481	21.9	108.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,853	13.4	66.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,191	7.0	34.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,149	5.8	28.7
6	Diabetes mellitus (E10-E14)	2,791	3.1	15.6
7	Cerebrovascular diseases (I60-I69)	2,111	2.4	11.8
8	Chronic lower respiratory diseases (J40-J47)	1,815	2.0	10.1
9	Viral hepatitis (B15-B19)	1,393	1.6	7.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,042	1.2	5.8
11	Influenza and pneumonia (J09-J18)	1,010	1.1	5.6
12	Septicemia (A40-A41)	940	1.1	5.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	857	1.0	4.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	725	0.8	4.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	546	0.6	3.0
...	All other causes (Residual)	13,015	14.7	72.6
White, male, 55-64 years				
...	All causes	159,366	100.0	1,033.3
1	Malignant neoplasms (C00-C97)	51,265	32.2	332.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,820	24.4	251.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,683	5.4	56.3
4	Chronic lower respiratory diseases (J40-J47)	6,952	4.4	45.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,579	4.1	42.7
6	Diabetes mellitus (E10-E14)	5,781	3.6	37.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,740	3.0	30.7
8	Cerebrovascular diseases (I60-I69)	4,515	2.8	29.3
9	Viral hepatitis (B15-B19)	2,065	1.3	13.4
10	Septicemia (A40-A41)	1,955	1.2	12.7
11	Influenza and pneumonia (J09-J18)	1,951	1.2	12.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,833	1.2	11.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,121	0.7	7.3
14	Aortic aneurysm and dissection (I71)	807	0.5	5.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	706	0.4	4.6
...	All other causes (Residual)	21,593	13.5	140.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	198,703	100.0	2,199.7
1	Malignant neoplasms (C00-C97)	68,604	34.5	759.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,411	23.9	524.9
3	Chronic lower respiratory diseases (J40-J47)	14,986	7.5	165.9
4	Diabetes mellitus (E10-E14)	7,324	3.7	81.1
5	Cerebrovascular diseases (I60-I69)	7,253	3.7	80.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,457	2.7	60.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,305	1.7	36.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,047	1.5	33.7
9	Influenza and pneumonia (J09-J18)	3,009	1.5	33.3
10	Septicemia (A40-A41)	2,713	1.4	30.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,412	1.2	26.7
12	Parkinson's disease (G20-G21)	1,750	0.9	19.4
13	Alzheimer's disease (G30)	1,661	0.8	18.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,409	0.7	15.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,345	0.7	14.9
...	All other causes (Residual)	27,017	13.6	299.1
White, male, 75-84 years				
...	All causes	278,334	100.0	5,650.4
1	Malignant neoplasms (C00-C97)	73,162	26.3	1,485.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,648	25.4	1,434.2
3	Chronic lower respiratory diseases (J40-J47)	21,968	7.9	446.0
4	Cerebrovascular diseases (I60-I69)	13,962	5.0	283.4
5	Alzheimer's disease (G30)	8,359	3.0	169.7
6	Diabetes mellitus (E10-E14)	8,142	2.9	165.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,924	2.5	140.6
8	Influenza and pneumonia (J09-J18)	6,060	2.2	123.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,799	2.1	117.7
10	Parkinson's disease (G20-G21)	5,796	2.1	117.7
11	Septicemia (A40-A41)	3,913	1.4	79.4
12	Pneumonitis due to solids and liquids (J69)	2,608	0.9	52.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,353	0.8	47.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,320	0.8	47.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,893	0.7	38.4
...	All other causes (Residual)	44,427	16.0	901.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	262,205	100.0	15,335.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	81,292	31.0	4,754.6
2	Malignant neoplasms (C00-C97)	40,130	15.3	2,347.1
3	Chronic lower respiratory diseases (J40-J47)	15,334	5.8	896.8
4	Cerebrovascular diseases (I60-I69)	14,468	5.5	846.2
5	Alzheimer's disease (G30)	13,247	5.1	774.8
6	Influenza and pneumonia (J09-J18)	9,114	3.5	533.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,979	2.7	408.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,522	2.5	381.5
9	Diabetes mellitus (E10-E14)	5,617	2.1	328.5
10	Parkinson's disease (G20-G21)	4,875	1.9	285.1
11	Pneumonitis due to solids and liquids (J69)	4,066	1.6	237.8
12	Septicemia (A40-A41)	3,290	1.3	192.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,101	1.2	181.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,067	0.8	120.9
15	Aortic aneurysm and dissection (I71)	1,127	0.4	65.9
...	All other causes (Residual)	50,976	19.4	2,981.5
White, female, all ages				
...	All causes	1,084,111	100.0	868.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	248,105	22.9	198.8
2	Malignant neoplasms (C00-C97)	233,025	21.5	186.8
3	Chronic lower respiratory diseases (J40-J47)	69,988	6.5	56.1
4	Cerebrovascular diseases (I60-I69)	65,278	6.0	52.3
5	Alzheimer's disease (G30)	54,415	5.0	43.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41,628	3.8	33.4
7	Diabetes mellitus (E10-E14)	27,191	2.5	21.8
8	Influenza and pneumonia (J09-J18)	24,705	2.3	19.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,993	1.7	14.4
10	Septicemia (A40-A41)	15,467	1.4	12.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,147	1.2	10.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,440	1.0	8.4
13	Parkinson's disease (G20-G21)	8,942	0.8	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,672	0.7	6.1
15	Pneumonitis due to solids and liquids (J69)	7,610	0.7	6.1
...	All other causes (Residual)	238,505	22.0	191.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,291	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	393	30.4	6.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	12.9	2.8
3	Malignant neoplasms (C00-C97)	139	10.8	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	7.2	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	4.0	0.9
6	Influenza and pneumonia (J09-J18)	34	2.6	0.6
7	Septicemia (A40-A41)	25	1.9	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	10	0.8	*
10	Cerebrovascular diseases (I60-I69)	9	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
15	Diabetes mellitus (E10-E14)	4	0.3	*
...	All other causes (Residual)	330	25.6	5.6
White, female, 5-14 years				
...	All causes	1,606	100.0	10.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	451	28.1	3.0
2	Malignant neoplasms (C00-C97)	297	18.5	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	7.3	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	4.8	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	4.4	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	65	4.0	0.4
7	Influenza and pneumonia (J09-J18)	40	2.5	0.3
8	Septicemia (A40-A41)	26	1.6	0.2
8	Cerebrovascular diseases (I60-I69)	26	1.6	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.4	0.2
11	Chronic lower respiratory diseases (J40-J47)	20	1.2	0.1
12	Diabetes mellitus (E10-E14)	14	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
14	Meningitis (G00,G03)	5	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	358	22.3	2.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	5,815	100.0	36.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,667	45.9	16.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	716	12.3	4.4
3	Malignant neoplasms (C00-C97)	502	8.6	3.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	303	5.2	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	3.3	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	1.8	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	102	1.8	0.6
8	Influenza and pneumonia (J09-J18)	67	1.2	0.4
9	Cerebrovascular diseases (I60-I69)	60	1.0	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.7	0.2
11	Diabetes mellitus (E10-E14)	39	0.7	0.2
12	Septicemia (A40-A41)	37	0.6	0.2
13	Chronic lower respiratory diseases (J40-J47)	21	0.4	0.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	18	0.3	*
...	All other causes (Residual)	926	15.9	5.7
White, female, 25-34 years				
...	All causes	10,040	100.0	63.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,466	34.5	22.0
2	Malignant neoplasms (C00-C97)	1,309	13.0	8.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,046	10.4	6.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	675	6.7	4.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	401	4.0	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	230	2.3	1.5
7	Diabetes mellitus (E10-E14)	174	1.7	1.1
8	Cerebrovascular diseases (I60-I69)	164	1.6	1.0
8	Influenza and pneumonia (J09-J18)	164	1.6	1.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	146	1.5	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	1.5	0.9
12	Septicemia (A40-A41)	98	1.0	0.6
13	Chronic lower respiratory diseases (J40-J47)	76	0.8	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	61	0.6	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	0.3
...	All other causes (Residual)	1,832	18.2	11.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	19,602	100.0	124.9
1	Malignant neoplasms (C00-C97)	4,928	25.1	31.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,123	21.0	26.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,153	11.0	13.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,384	7.1	8.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	700	3.6	4.5
6	Cerebrovascular diseases (I60-I69)	499	2.5	3.2
7	Diabetes mellitus (E10-E14)	473	2.4	3.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	375	1.9	2.4
9	Influenza and pneumonia (J09-J18)	293	1.5	1.9
10	Chronic lower respiratory diseases (J40-J47)	281	1.4	1.8
11	Septicemia (A40-A41)	275	1.4	1.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	158	0.8	1.0
13	Pregnancy, childbirth and the puerperium (O00-O99)	155	0.8	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	0.7	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	0.7	0.9
...	All other causes (Residual)	3,523	18.0	22.4
White, female, 45-54 years				
...	All causes	54,646	100.0	301.8
1	Malignant neoplasms (C00-C97)	19,022	34.8	105.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,337	13.4	40.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,899	10.8	32.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,537	4.6	14.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,077	3.8	11.5
6	Chronic lower respiratory diseases (J40-J47)	2,040	3.7	11.3
7	Cerebrovascular diseases (I60-I69)	1,636	3.0	9.0
8	Diabetes mellitus (E10-E14)	1,516	2.8	8.4
9	Septicemia (A40-A41)	816	1.5	4.5
10	Influenza and pneumonia (J09-J18)	720	1.3	4.0
11	Viral hepatitis (B15-B19)	549	1.0	3.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	479	0.9	2.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	374	0.7	2.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	291	0.5	1.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	286	0.5	1.6
...	All other causes (Residual)	9,067	16.6	50.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	100,324	100.0	619.2
1	Malignant neoplasms (C00-C97)	40,600	40.5	250.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,874	15.8	98.0
3	Chronic lower respiratory diseases (J40-J47)	6,367	6.3	39.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,981	4.0	24.6
5	Diabetes mellitus (E10-E14)	3,639	3.6	22.5
6	Cerebrovascular diseases (I60-I69)	3,419	3.4	21.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,692	2.7	16.6
8	Septicemia (A40-A41)	1,712	1.7	10.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,411	1.4	8.7
10	Influenza and pneumonia (J09-J18)	1,387	1.4	8.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,352	1.3	8.3
12	Viral hepatitis (B15-B19)	717	0.7	4.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	623	0.6	3.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	519	0.5	3.2
15	Alzheimer's disease (G30)	420	0.4	2.6
...	All other causes (Residual)	15,611	15.6	96.4
White, female, 65-74 years				
...	All causes	151,336	100.0	1,491.2
1	Malignant neoplasms (C00-C97)	55,732	36.8	549.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,516	17.5	261.3
3	Chronic lower respiratory diseases (J40-J47)	14,782	9.8	145.7
4	Cerebrovascular diseases (I60-I69)	6,559	4.3	64.6
5	Diabetes mellitus (E10-E14)	5,185	3.4	51.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,254	2.2	32.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,608	1.7	25.7
8	Septicemia (A40-A41)	2,525	1.7	24.9
9	Influenza and pneumonia (J09-J18)	2,410	1.6	23.7
10	Alzheimer's disease (G30)	2,185	1.4	21.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,027	1.3	20.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,273	0.8	12.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	922	0.6	9.1
14	Parkinson's disease (G20-G21)	864	0.6	8.5
15	Pneumonitis due to solids and liquids (J69)	678	0.4	6.7
...	All other causes (Residual)	23,816	15.7	234.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	273,593	100.0	4,158.2
1	Malignant neoplasms (C00-C97)	63,830	23.3	970.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	60,057	22.0	912.8
3	Chronic lower respiratory diseases (J40-J47)	24,119	8.8	366.6
4	Cerebrovascular diseases (I60-I69)	18,293	6.7	278.0
5	Alzheimer's disease (G30)	13,743	5.0	208.9
6	Diabetes mellitus (E10-E14)	7,745	2.8	117.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,947	2.2	90.4
8	Influenza and pneumonia (J09-J18)	5,860	2.1	89.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,128	1.9	77.9
10	Septicemia (A40-A41)	4,232	1.5	64.3
11	Parkinson's disease (G20-G21)	3,595	1.3	54.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,055	1.1	46.4
13	Pneumonitis due to solids and liquids (J69)	1,995	0.7	30.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,848	0.7	28.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,678	0.6	25.5
...	All other causes (Residual)	52,468	19.2	797.4
White, female, 85 years and over				
...	All causes	458,957	100.0	13,380.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	135,081	29.4	3,938.3
2	Malignant neoplasms (C00-C97)	46,643	10.2	1,359.9
3	Alzheimer's disease (G30)	38,010	8.3	1,108.2
4	Cerebrovascular diseases (I60-I69)	34,577	7.5	1,008.1
5	Chronic lower respiratory diseases (J40-J47)	22,268	4.9	649.2
6	Influenza and pneumonia (J09-J18)	13,688	3.0	399.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,128	2.4	324.4
8	Diabetes mellitus (E10-E14)	8,402	1.8	245.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,189	1.8	238.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,821	1.7	228.0
11	Septicemia (A40-A41)	5,671	1.2	165.3
12	Pneumonitis due to solids and liquids (J69)	4,306	0.9	125.5
13	Parkinson's disease (G20-G21)	4,298	0.9	125.3
14	Atherosclerosis (I70)	2,538	0.6	74.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,511	0.5	73.2
...	All other causes (Residual)	113,826	24.8	3,318.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	290,100	100.0	679.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,372	23.6	160.2
2	Malignant neoplasms (C00-C97)	66,820	23.0	156.6
3	Cerebrovascular diseases (I60-I69)	15,853	5.5	37.1
4	Diabetes mellitus (E10-E14)	12,895	4.4	30.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,531	4.3	29.4
6	Chronic lower respiratory diseases (J40-J47)	9,086	3.1	21.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,087	2.8	18.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,858	2.7	18.4
9	Septicemia (A40-A41)	6,028	2.1	14.1
10	Alzheimer's disease (G30)	5,482	1.9	12.8
11	Influenza and pneumonia (J09-J18)	5,167	1.8	12.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,006	1.7	11.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,215	1.5	9.9
14	Certain conditions originating in the perinatal period (P00-P96)	4,055	1.4	9.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,816	1.0	6.6
...	All other causes (Residual)	55,829	19.2	130.8
Black, both sexes, 1-4 years				
...	All causes	1,052	100.0	38.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	298	28.3	10.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	15.1	5.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	10.3	3.9
4	Malignant neoplasms (C00-C97)	55	5.2	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	5.1	2.0
6	Influenza and pneumonia (J09-J18)	32	3.0	1.2
7	Chronic lower respiratory diseases (J40-J47)	25	2.4	0.9
8	Cerebrovascular diseases (I60-I69)	17	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	13	1.2	*
10	Septicemia (A40-A41)	12	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.8	*
13	Anemias (D50-D64)	6	0.6	*
14	Meningitis (G00,G03)	3	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	248	23.6	9.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,181	100.0	17.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	27.2	4.8
2	Malignant neoplasms (C00-C97)	133	11.3	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	7.6	1.3
4	Chronic lower respiratory diseases (J40-J47)	84	7.1	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	6.8	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	4.2	0.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	3.8	0.7
8	Influenza and pneumonia (J09-J18)	26	2.2	0.4
9	Anemias (D50-D64)	22	1.9	0.3
10	Cerebrovascular diseases (I60-I69)	20	1.7	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.6	*
12	Septicemia (A40-A41)	12	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.8	*
14	Pneumonitis due to solids and liquids (J69)	6	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
...	All other causes (Residual)	259	21.9	3.9
Black, both sexes, 15-24 years				
...	All causes	6,690	100.0	92.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,826	42.2	39.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,387	20.7	19.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	504	7.5	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	310	4.6	4.3
5	Malignant neoplasms (C00-C97)	260	3.9	3.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	1.6	1.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	103	1.5	1.4
8	Chronic lower respiratory diseases (J40-J47)	81	1.2	1.1
9	Anemias (D50-D64)	73	1.1	1.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	64	1.0	0.9
11	Diabetes mellitus (E10-E14)	48	0.7	0.7
12	Legal intervention (Y35,Y89.0)	47	0.7	0.6
13	Influenza and pneumonia (J09-J18)	44	0.7	0.6
14	Septicemia (A40-A41)	36	0.5	0.5
15	Cerebrovascular diseases (I60-I69)	32	0.5	0.4
...	All other causes (Residual)	771	11.5	10.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	9,068	100.0	151.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,330	25.7	38.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,657	18.3	27.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	957	10.6	16.0
4	Malignant neoplasms (C00-C97)	646	7.1	10.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	556	6.1	9.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	418	4.6	7.0
7	Diabetes mellitus (E10-E14)	228	2.5	3.8
8	Cerebrovascular diseases (I60-I69)	135	1.5	2.3
9	Anemias (D50-D64)	133	1.5	2.2
10	Pregnancy, childbirth and the puerperium (O00-O99)	126	1.4	2.1
11	Influenza and pneumonia (J09-J18)	106	1.2	1.8
12	Septicemia (A40-A41)	97	1.1	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	1.0	1.6
14	Chronic lower respiratory diseases (J40-J47)	91	1.0	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	0.7	1.0
...	All other causes (Residual)	1,433	15.8	23.9
Black, both sexes, 35-44 years				
...	All causes	13,789	100.0	248.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,718	19.7	49.0
2	Malignant neoplasms (C00-C97)	2,116	15.3	38.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,695	12.3	30.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,133	8.2	20.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	904	6.6	16.3
6	Cerebrovascular diseases (I60-I69)	513	3.7	9.2
7	Diabetes mellitus (E10-E14)	491	3.6	8.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	422	3.1	7.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	270	2.0	4.9
10	Septicemia (A40-A41)	217	1.6	3.9
11	Chronic lower respiratory diseases (J40-J47)	192	1.4	3.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	187	1.4	3.4
13	Influenza and pneumonia (J09-J18)	183	1.3	3.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	181	1.3	3.3
15	Anemias (D50-D64)	110	0.8	2.0
...	All other causes (Residual)	2,457	17.8	44.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	33,531	100.0	579.7
1	Malignant neoplasms (C00-C97)	8,398	25.0	145.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,811	23.3	135.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,410	7.2	41.7
4	Cerebrovascular diseases (I60-I69)	1,644	4.9	28.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,454	4.3	25.1
6	Diabetes mellitus (E10-E14)	1,450	4.3	25.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	802	2.4	13.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	736	2.2	12.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	727	2.2	12.6
10	Chronic lower respiratory diseases (J40-J47)	699	2.1	12.1
11	Septicemia (A40-A41)	631	1.9	10.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	503	1.5	8.7
13	Influenza and pneumonia (J09-J18)	403	1.2	7.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	356	1.1	6.2
15	Viral hepatitis (B15-B19)	278	0.8	4.8
...	All other causes (Residual)	5,229	15.6	90.4
Black, both sexes, 55-64 years				
...	All causes	53,799	100.0	1,257.7
1	Malignant neoplasms (C00-C97)	16,950	31.5	396.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,153	24.4	307.5
3	Diabetes mellitus (E10-E14)	2,771	5.2	64.8
4	Cerebrovascular diseases (I60-I69)	2,708	5.0	63.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,029	3.8	47.4
6	Chronic lower respiratory diseases (J40-J47)	1,487	2.8	34.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,422	2.6	33.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,160	2.2	27.1
9	Septicemia (A40-A41)	1,135	2.1	26.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,000	1.9	23.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	852	1.6	19.9
12	Influenza and pneumonia (J09-J18)	707	1.3	16.5
13	Viral hepatitis (B15-B19)	601	1.1	14.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	333	0.6	7.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	203	0.4	4.7
...	All other causes (Residual)	7,288	13.5	170.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	53,282	100.0	2,464.3
1	Malignant neoplasms (C00-C97)	16,995	31.9	786.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,344	25.0	617.2
3	Cerebrovascular diseases (I60-I69)	3,034	5.7	140.3
4	Diabetes mellitus (E10-E14)	3,009	5.6	139.2
5	Chronic lower respiratory diseases (J40-J47)	2,222	4.2	102.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,759	3.3	81.4
7	Septicemia (A40-A41)	1,249	2.3	57.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	955	1.8	44.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	910	1.7	42.1
10	Influenza and pneumonia (J09-J18)	853	1.6	39.5
11	Alzheimer's disease (G30)	407	0.8	18.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	407	0.8	18.8
13	Pneumonitis due to solids and liquids (J69)	257	0.5	11.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	255	0.5	11.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	225	0.4	10.4
...	All other causes (Residual)	7,401	13.9	342.3
Black, both sexes, 75-84 years				
...	All causes	57,954	100.0	5,215.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,035	25.9	1,353.1
2	Malignant neoplasms (C00-C97)	14,109	24.3	1,269.8
3	Cerebrovascular diseases (I60-I69)	3,977	6.9	357.9
4	Diabetes mellitus (E10-E14)	2,939	5.1	264.5
5	Chronic lower respiratory diseases (J40-J47)	2,517	4.3	226.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,082	3.6	187.4
7	Alzheimer's disease (G30)	1,722	3.0	155.0
8	Septicemia (A40-A41)	1,405	2.4	126.4
9	Influenza and pneumonia (J09-J18)	1,235	2.1	111.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,108	1.9	99.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	770	1.3	69.3
12	Pneumonitis due to solids and liquids (J69)	410	0.7	36.9
13	Parkinson's disease (G20-G21)	387	0.7	34.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	335	0.6	30.1
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	198	0.3	17.8
...	All other causes (Residual)	9,725	16.8	875.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	52,443	100.0	12,870.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,835	28.3	3,640.6
2	Malignant neoplasms (C00-C97)	7,142	13.6	1,752.7
3	Cerebrovascular diseases (I60-I69)	3,724	7.1	913.9
4	Alzheimer's disease (G30)	3,276	6.2	804.0
5	Diabetes mellitus (E10-E14)	1,955	3.7	479.8
6	Chronic lower respiratory diseases (J40-J47)	1,676	3.2	411.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,669	3.2	409.6
8	Influenza and pneumonia (J09-J18)	1,495	2.9	366.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,386	2.6	340.1
10	Septicemia (A40-A41)	1,169	2.2	286.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	627	1.2	153.9
12	Pneumonitis due to solids and liquids (J69)	507	1.0	124.4
13	Parkinson's disease (G20-G21)	274	0.5	67.2
14	Atherosclerosis (I70)	243	0.5	59.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	238	0.5	58.4
...	All other causes (Residual)	12,227	23.3	3,000.6
Black, male, all ages				
...	All causes	146,884	100.0	719.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,913	23.8	171.0
2	Malignant neoplasms (C00-C97)	33,978	23.1	166.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,387	5.7	41.1
4	Cerebrovascular diseases (I60-I69)	7,039	4.8	34.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,739	4.6	33.0
6	Diabetes mellitus (E10-E14)	6,048	4.1	29.6
7	Chronic lower respiratory diseases (J40-J47)	4,702	3.2	23.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,756	2.6	18.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2,744	1.9	13.4
10	Septicemia (A40-A41)	2,700	1.8	13.2
11	Influenza and pneumonia (J09-J18)	2,417	1.6	11.8
12	Certain conditions originating in the perinatal period (P00-P96)	2,252	1.5	11.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,200	1.5	10.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,828	1.2	9.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,775	1.2	8.7
...	All other causes (Residual)	25,406	17.3	124.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	596	100.0	42.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	29.0	12.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	14.6	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	9.9	4.2
4	Malignant neoplasms (C00-C97)	31	5.2	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.7	2.0
6	Influenza and pneumonia (J09-J18)	18	3.0	*
7	Chronic lower respiratory diseases (J40-J47)	16	2.7	*
8	Cerebrovascular diseases (I60-I69)	14	2.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.5	*
10	Septicemia (A40-A41)	6	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.0	*
13	Anemias (D50-D64)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Meningitis (G00,G03)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	135	22.7	9.7
Black, male, 5-14 years				
...	All causes	701	100.0	20.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	28.8	5.9
2	Malignant neoplasms (C00-C97)	77	11.0	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	8.8	1.8
4	Chronic lower respiratory diseases (J40-J47)	57	8.1	1.7
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	6.3	1.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	4.9	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.7	0.8
8	Influenza and pneumonia (J09-J18)	16	2.3	*
9	Anemias (D50-D64)	14	2.0	*
10	Cerebrovascular diseases (I60-I69)	11	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.4	*
12	Pneumonitis due to solids and liquids (J69)	5	0.7	*
13	Septicemia (A40-A41)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
...	All other causes (Residual)	131	18.7	3.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,124	100.0	140.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,536	49.5	69.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,037	20.2	28.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	421	8.2	11.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	200	3.9	5.5
5	Malignant neoplasms (C00-C97)	151	2.9	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	64	1.2	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	1.2	1.8
8	Chronic lower respiratory diseases (J40-J47)	54	1.1	1.5
9	Legal intervention (Y35,Y89.0)	46	0.9	1.3
10	Anemias (D50-D64)	45	0.9	1.2
11	Diabetes mellitus (E10-E14)	23	0.4	0.6
12	Influenza and pneumonia (J09-J18)	20	0.4	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
14	Cerebrovascular diseases (I60-I69)	16	0.3	*
15	Septicemia (A40-A41)	13	0.3	*
...	All other causes (Residual)	417	8.1	11.4
Black, male, 25-34 years				
...	All causes	6,103	100.0	212.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,086	34.2	72.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,192	19.5	41.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	577	9.5	20.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	454	7.4	15.8
5	Malignant neoplasms (C00-C97)	264	4.3	9.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	245	4.0	8.5
7	Diabetes mellitus (E10-E14)	113	1.9	3.9
8	Anemias (D50-D64)	75	1.2	2.6
9	Cerebrovascular diseases (I60-I69)	67	1.1	2.3
10	Chronic lower respiratory diseases (J40-J47)	58	1.0	2.0
11	Influenza and pneumonia (J09-J18)	55	0.9	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	0.9	1.8
13	Septicemia (A40-A41)	40	0.7	1.4
14	Legal intervention (Y35,Y89.0)	32	0.5	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	0.5	1.0
...	All other causes (Residual)	764	12.5	26.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	7,962	100.0	304.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,734	21.8	66.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,202	15.1	46.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	953	12.0	36.5
4	Malignant neoplasms (C00-C97)	754	9.5	28.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	510	6.4	19.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	338	4.2	12.9
7	Diabetes mellitus (E10-E14)	276	3.5	10.6
8	Cerebrovascular diseases (I60-I69)	266	3.3	10.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	149	1.9	5.7
10	Septicemia (A40-A41)	116	1.5	4.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	112	1.4	4.3
12	Influenza and pneumonia (J09-J18)	95	1.2	3.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	93	1.2	3.6
14	Chronic lower respiratory diseases (J40-J47)	87	1.1	3.3
15	Aortic aneurysm and dissection (I71)	67	0.8	2.6
...	All other causes (Residual)	1,210	15.2	46.3
Black, male, 45-54 years				
...	All causes	19,193	100.0	706.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,002	26.1	184.1
2	Malignant neoplasms (C00-C97)	3,956	20.6	145.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,651	8.6	60.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	979	5.1	36.0
5	Cerebrovascular diseases (I60-I69)	920	4.8	33.9
6	Diabetes mellitus (E10-E14)	801	4.2	29.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	577	3.0	21.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	488	2.5	18.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	400	2.1	14.7
10	Septicemia (A40-A41)	321	1.7	11.8
11	Chronic lower respiratory diseases (J40-J47)	317	1.7	11.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	303	1.6	11.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	290	1.5	10.7
14	Influenza and pneumonia (J09-J18)	220	1.1	8.1
15	Viral hepatitis (B15-B19)	181	0.9	6.7
...	All other causes (Residual)	2,787	14.5	102.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	31,599	100.0	1,623.4
1	Malignant neoplasms (C00-C97)	9,312	29.5	478.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,276	26.2	425.2
3	Cerebrovascular diseases (I60-I69)	1,584	5.0	81.4
4	Diabetes mellitus (E10-E14)	1,548	4.9	79.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,442	4.6	74.1
6	Chronic lower respiratory diseases (J40-J47)	815	2.6	41.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	795	2.5	40.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	787	2.5	40.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	710	2.2	36.5
10	Septicemia (A40-A41)	617	2.0	31.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	482	1.5	24.8
12	Viral hepatitis (B15-B19)	411	1.3	21.1
13	Influenza and pneumonia (J09-J18)	399	1.3	20.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	279	0.9	14.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	0.5	8.0
...	All other causes (Residual)	3,987	12.6	204.8
Black, male, 65-74 years				
...	All causes	28,908	100.0	3,127.2
1	Malignant neoplasms (C00-C97)	9,282	32.1	1,004.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,631	26.4	825.5
3	Cerebrovascular diseases (I60-I69)	1,590	5.5	172.0
4	Diabetes mellitus (E10-E14)	1,494	5.2	161.6
5	Chronic lower respiratory diseases (J40-J47)	1,260	4.4	136.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	894	3.1	96.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	615	2.1	66.5
8	Septicemia (A40-A41)	586	2.0	63.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	475	1.6	51.4
10	Influenza and pneumonia (J09-J18)	442	1.5	47.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	263	0.9	28.5
12	Alzheimer's disease (G30)	185	0.6	20.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	180	0.6	19.5
14	Pneumonitis due to solids and liquids (J69)	150	0.5	16.2
15	Parkinson's disease (G20-G21)	122	0.4	13.2
...	All other causes (Residual)	3,739	12.9	404.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	26,367	100.0	6,358.9
1	Malignant neoplasms (C00-C97)	7,072	26.8	1,705.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,967	26.4	1,680.2
3	Cerebrovascular diseases (I60-I69)	1,595	6.0	384.7
4	Chronic lower respiratory diseases (J40-J47)	1,311	5.0	316.2
5	Diabetes mellitus (E10-E14)	1,182	4.5	285.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	877	3.3	211.5
7	Alzheimer's disease (G30)	621	2.4	149.8
8	Septicemia (A40-A41)	592	2.2	142.8
9	Influenza and pneumonia (J09-J18)	586	2.2	141.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	465	1.8	112.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	405	1.5	97.7
12	Parkinson's disease (G20-G21)	222	0.8	53.5
12	Pneumonitis due to solids and liquids (J69)	222	0.8	53.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	133	0.5	32.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	91	0.3	21.9
...	All other causes (Residual)	4,026	15.3	971.0
Black, male, 85 years and over				
...	All causes	16,254	100.0	14,159.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,416	27.2	3,846.9
2	Malignant neoplasms (C00-C97)	3,071	18.9	2,675.3
3	Cerebrovascular diseases (I60-I69)	949	5.8	826.7
4	Alzheimer's disease (G30)	723	4.4	629.8
5	Chronic lower respiratory diseases (J40-J47)	718	4.4	625.5
6	Diabetes mellitus (E10-E14)	608	3.7	529.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	560	3.4	487.8
8	Influenza and pneumonia (J09-J18)	521	3.2	453.9
9	Septicemia (A40-A41)	365	2.2	318.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	331	2.0	288.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	254	1.6	221.3
12	Pneumonitis due to solids and liquids (J69)	203	1.2	176.8
13	Parkinson's disease (G20-G21)	129	0.8	112.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.4	62.7
14	Atherosclerosis (I70)	72	0.4	62.7
...	All other causes (Residual)	3,262	20.1	2,841.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	143,216	100.0	643.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,459	23.4	150.3
2	Malignant neoplasms (C00-C97)	32,842	22.9	147.5
3	Cerebrovascular diseases (I60-I69)	8,814	6.2	39.6
4	Diabetes mellitus (E10-E14)	6,847	4.8	30.8
5	Chronic lower respiratory diseases (J40-J47)	4,384	3.1	19.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,331	3.0	19.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,144	2.9	18.6
8	Alzheimer's disease (G30)	3,914	2.7	17.6
9	Septicemia (A40-A41)	3,328	2.3	15.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,806	2.0	12.6
11	Influenza and pneumonia (J09-J18)	2,750	1.9	12.4
12	Certain conditions originating in the perinatal period (P00-P96)	1,803	1.3	8.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,471	1.0	6.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,119	0.8	5.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,041	0.7	4.7
...	All other causes (Residual)	30,163	21.1	135.5
Black, female, 1-4 years				
...	All causes	456	100.0	33.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	27.4	9.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	15.8	5.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	10.7	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.7	1.9
5	Malignant neoplasms (C00-C97)	24	5.3	1.8
6	Influenza and pneumonia (J09-J18)	14	3.1	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.0	*
8	Septicemia (A40-A41)	6	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
11	Anemias (D50-D64)	3	0.7	*
11	Cerebrovascular diseases (I60-I69)	3	0.7	*
13	Meningitis (G00,G03)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Atherosclerosis (I70)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	106	23.2	7.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	480	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	24.8	3.6
2	Malignant neoplasms (C00-C97)	56	11.7	1.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	7.5	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	5.8	0.8
5	Chronic lower respiratory diseases (J40-J47)	27	5.6	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.0	0.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	2.3	*
8	Influenza and pneumonia (J09-J18)	10	2.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.9	*
9	Cerebrovascular diseases (I60-I69)	9	1.9	*
11	Septicemia (A40-A41)	8	1.7	*
11	Anemias (D50-D64)	8	1.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.0	*
14	Meningitis (G00,G03)	3	0.6	*
15	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (Residual)	125	26.0	3.8
Black, female, 15-24 years				
...	All causes	1,566	100.0	43.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	350	22.3	9.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	290	18.5	8.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	110	7.0	3.1
4	Malignant neoplasms (C00-C97)	109	7.0	3.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	5.3	2.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	64	4.1	1.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	2.6	1.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	39	2.5	1.1
9	Anemias (D50-D64)	28	1.8	0.8
10	Chronic lower respiratory diseases (J40-J47)	27	1.7	0.8
11	Diabetes mellitus (E10-E14)	25	1.6	0.7
12	Influenza and pneumonia (J09-J18)	24	1.5	0.7
13	Septicemia (A40-A41)	23	1.5	0.6
14	Cerebrovascular diseases (I60-I69)	16	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	*
...	All other causes (Residual)	327	20.9	9.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	2,965	100.0	94.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	465	15.7	14.9
2	Malignant neoplasms (C00-C97)	382	12.9	12.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	380	12.8	12.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	244	8.2	7.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	173	5.8	5.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	126	4.2	4.0
7	Diabetes mellitus (E10-E14)	115	3.9	3.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	102	3.4	3.3
9	Cerebrovascular diseases (I60-I69)	68	2.3	2.2
10	Anemias (D50-D64)	58	2.0	1.9
11	Septicemia (A40-A41)	57	1.9	1.8
12	Influenza and pneumonia (J09-J18)	51	1.7	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.4	1.3
14	Chronic lower respiratory diseases (J40-J47)	33	1.1	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	1.1	1.0
...	All other causes (Residual)	637	21.5	20.4
Black, female, 35-44 years				
...	All causes	5,827	100.0	198.5
1	Malignant neoplasms (C00-C97)	1,362	23.4	46.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	984	16.9	33.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	493	8.5	16.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	394	6.8	13.4
5	Cerebrovascular diseases (I60-I69)	247	4.2	8.4
6	Diabetes mellitus (E10-E14)	215	3.7	7.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	180	3.1	6.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	121	2.1	4.1
9	Chronic lower respiratory diseases (J40-J47)	105	1.8	3.6
10	Septicemia (A40-A41)	101	1.7	3.4
11	Influenza and pneumonia (J09-J18)	88	1.5	3.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	88	1.5	3.0
13	Pregnancy, childbirth and the puerperium (O00-O99)	85	1.5	2.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	1.4	2.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	75	1.3	2.6
...	All other causes (Residual)	1,205	20.7	41.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	14,338	100.0	467.6
1	Malignant neoplasms (C00-C97)	4,442	31.0	144.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,809	19.6	91.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	759	5.3	24.8
4	Cerebrovascular diseases (I60-I69)	724	5.0	23.6
5	Diabetes mellitus (E10-E14)	649	4.5	21.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	475	3.3	15.5
7	Chronic lower respiratory diseases (J40-J47)	382	2.7	12.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	327	2.3	10.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	314	2.2	10.2
10	Septicemia (A40-A41)	310	2.2	10.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	200	1.4	6.5
12	Influenza and pneumonia (J09-J18)	183	1.3	6.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	1.1	5.2
14	Viral hepatitis (B15-B19)	97	0.7	3.2
15	Anemias (D50-D64)	85	0.6	2.8
...	All other causes (Residual)	2,423	16.9	79.0
Black, female, 55-64 years				
...	All causes	22,200	100.0	952.3
1	Malignant neoplasms (C00-C97)	7,638	34.4	327.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,877	22.0	209.2
3	Diabetes mellitus (E10-E14)	1,223	5.5	52.5
4	Cerebrovascular diseases (I60-I69)	1,124	5.1	48.2
5	Chronic lower respiratory diseases (J40-J47)	672	3.0	28.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	635	2.9	27.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	587	2.6	25.2
8	Septicemia (A40-A41)	518	2.3	22.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	370	1.7	15.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	365	1.6	15.7
11	Influenza and pneumonia (J09-J18)	308	1.4	13.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	290	1.3	12.4
13	Viral hepatitis (B15-B19)	190	0.9	8.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	107	0.5	4.6
15	Pneumonitis due to solids and liquids (J69)	74	0.3	3.2
...	All other causes (Residual)	3,222	14.5	138.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	24,374	100.0	1,969.2
1	Malignant neoplasms (C00-C97)	7,713	31.6	623.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,713	23.4	461.6
3	Diabetes mellitus (E10-E14)	1,515	6.2	122.4
4	Cerebrovascular diseases (I60-I69)	1,444	5.9	116.7
5	Chronic lower respiratory diseases (J40-J47)	962	3.9	77.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	865	3.5	69.9
7	Septicemia (A40-A41)	663	2.7	53.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	435	1.8	35.1
9	Influenza and pneumonia (J09-J18)	411	1.7	33.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	340	1.4	27.5
11	Alzheimer's disease (G30)	222	0.9	17.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	144	0.6	11.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	120	0.5	9.7
14	Pneumonitis due to solids and liquids (J69)	107	0.4	8.6
15	Aortic aneurysm and dissection (I71)	83	0.3	6.7
...	All other causes (Residual)	3,637	14.9	293.8
Black, female, 75-84 years				
...	All causes	31,587	100.0	4,535.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,068	25.5	1,158.4
2	Malignant neoplasms (C00-C97)	7,037	22.3	1,010.4
3	Cerebrovascular diseases (I60-I69)	2,382	7.5	342.0
4	Diabetes mellitus (E10-E14)	1,757	5.6	252.3
5	Chronic lower respiratory diseases (J40-J47)	1,206	3.8	173.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,205	3.8	173.0
7	Alzheimer's disease (G30)	1,101	3.5	158.1
8	Septicemia (A40-A41)	813	2.6	116.7
9	Influenza and pneumonia (J09-J18)	649	2.1	93.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	643	2.0	92.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	365	1.2	52.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	202	0.6	29.0
13	Pneumonitis due to solids and liquids (J69)	188	0.6	27.0
14	Parkinson's disease (G20-G21)	165	0.5	23.7
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	120	0.4	17.2
...	All other causes (Residual)	5,686	18.0	816.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	36,189	100.0	12,364.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,419	28.8	3,559.7
2	Malignant neoplasms (C00-C97)	4,071	11.2	1,390.9
3	Cerebrovascular diseases (I60-I69)	2,775	7.7	948.1
4	Alzheimer's disease (G30)	2,553	7.1	872.3
5	Diabetes mellitus (E10-E14)	1,347	3.7	460.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,109	3.1	378.9
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,055	2.9	360.4
8	Influenza and pneumonia (J09-J18)	974	2.7	332.8
9	Chronic lower respiratory diseases (J40-J47)	958	2.6	327.3
10	Septicemia (A40-A41)	804	2.2	274.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	373	1.0	127.4
12	Pneumonitis due to solids and liquids (J69)	304	0.8	103.9
13	Atherosclerosis (I70)	171	0.5	58.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	166	0.5	56.7
15	Parkinson's disease (G20-G21)	145	0.4	49.5
...	All other causes (Residual)	8,965	24.8	3,063.0
American Indian, both sexes, all ages				
...	All causes	15,945	100.0	366.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,918	18.3	67.1
2	Malignant neoplasms (C00-C97)	2,810	17.6	64.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,829	11.5	42.1
4	Diabetes mellitus (E10-E14)	927	5.8	21.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	821	5.1	18.9
6	Chronic lower respiratory diseases (J40-J47)	657	4.1	15.1
7	Cerebrovascular diseases (I60-I69)	600	3.8	13.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	459	2.9	10.6
9	Influenza and pneumonia (J09-J18)	349	2.2	8.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	287	1.8	6.6
11	Septicemia (A40-A41)	269	1.7	6.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	254	1.6	5.8
13	Alzheimer's disease (G30)	226	1.4	5.2
14	Certain conditions originating in the perinatal period (P00-P96)	126	0.8	2.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	124	0.8	2.9
...	All other causes (Residual)	3,289	20.6	75.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	85	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	36.5	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	10.6	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	8.2	*
4	Malignant neoplasms (C00-C97)	5	5.9	*
5	Influenza and pneumonia (J09-J18)	3	3.5	*
6	Septicemia (A40-A41)	2	2.4	*
6	Cerebrovascular diseases (I60-I69)	2	2.4	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.4	*
9	Meningococcal infection (A39)	1	1.2	*
9	Infections of kidney (N10-N12,N13.6,N15.1)	1	1.2	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.2	*
...	All other causes (Residual)	21	24.7	6.7
American Indian, both sexes, 5-14 years				
...	All causes	96	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	33.3	4.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	11.5	*
3	Malignant neoplasms (C00-C97)	9	9.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	9.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.2	*
6	Cerebrovascular diseases (I60-I69)	4	4.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.1	*
8	Septicemia (A40-A41)	1	1.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
8	Influenza and pneumonia (J09-J18)	1	1.0	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
8	Pneumonitis due to solids and liquids (J69)	1	1.0	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.0	*
...	All other causes (Residual)	17	17.7	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	589	100.0	77.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	47.4	36.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	22.9	17.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	9.0	6.9
4	Malignant neoplasms (C00-C97)	14	2.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.2	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.0	*
7	Influenza and pneumonia (J09-J18)	5	0.8	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.7	*
8	Legal intervention (Y35,Y89.0)	4	0.7	*
10	Cerebrovascular diseases (I60-I69)	3	0.5	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.5	*
13	Diabetes mellitus (E10-E14)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	57	9.7	7.5
American Indian, both sexes, 25-34 years				
...	All causes	841	100.0	126.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	350	41.6	52.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	11.2	14.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	60	7.1	9.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	7.1	9.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	5.0	6.3
6	Malignant neoplasms (C00-C97)	34	4.0	5.1
7	Diabetes mellitus (E10-E14)	17	2.0	*
8	Cerebrovascular diseases (I60-I69)	10	1.2	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	10	1.2	*
10	Influenza and pneumonia (J09-J18)	8	1.0	*
11	Septicemia (A40-A41)	7	0.8	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.8	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.5	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
15	Legal intervention (Y35,Y89.0)	3	0.4	*
...	All other causes (Residual)	125	14.9	18.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,201	100.0	208.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	293	24.4	50.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	169	14.1	29.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	133	11.1	23.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	8.9	18.6
5	Malignant neoplasms (C00-C97)	79	6.6	13.7
6	Diabetes mellitus (E10-E14)	62	5.2	10.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	3.9	8.2
8	Cerebrovascular diseases (I60-I69)	21	1.7	3.6
9	Influenza and pneumonia (J09-J18)	20	1.7	3.5
10	Septicemia (A40-A41)	14	1.2	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.7	*
13	Viral hepatitis (B15-B19)	8	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.4	*
...	All other causes (Residual)	218	18.2	37.9
American Indian, both sexes, 45-54 years				
...	All causes	2,277	100.0	417.5
1	Malignant neoplasms (C00-C97)	379	16.6	69.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	360	15.8	66.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	350	15.4	64.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	268	11.8	49.1
5	Diabetes mellitus (E10-E14)	124	5.4	22.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	2.8	11.7
7	Cerebrovascular diseases (I60-I69)	59	2.6	10.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	1.9	7.9
9	Septicemia (A40-A41)	42	1.8	7.7
9	Chronic lower respiratory diseases (J40-J47)	42	1.8	7.7
11	Influenza and pneumonia (J09-J18)	38	1.7	7.0
12	Viral hepatitis (B15-B19)	35	1.5	6.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	1.1	4.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.4	*
...	All other causes (Residual)	424	18.6	77.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-64 years				
...	All causes	2,782	100.0	748.3
1	Malignant neoplasms (C00-C97)	663	23.8	178.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	569	20.5	153.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	7.6	57.0
4	Diabetes mellitus (E10-E14)	209	7.5	56.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	196	7.0	52.7
6	Chronic lower respiratory diseases (J40-J47)	125	4.5	33.6
7	Cerebrovascular diseases (I60-I69)	89	3.2	23.9
8	Septicemia (A40-A41)	65	2.3	17.5
9	Influenza and pneumonia (J09-J18)	54	1.9	14.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.8	13.4
11	Viral hepatitis (B15-B19)	48	1.7	12.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	1.1	8.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.7	5.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	0.6	*
15	Pneumonitis due to solids and liquids (J69)	12	0.4	*
...	All other causes (Residual)	423	15.2	113.8
American Indian, both sexes, 65-74 years				
...	All causes	2,856	100.0	1,625.2
1	Malignant neoplasms (C00-C97)	782	27.4	445.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	598	20.9	340.3
3	Diabetes mellitus (E10-E14)	218	7.6	124.1
4	Chronic lower respiratory diseases (J40-J47)	201	7.0	114.4
5	Cerebrovascular diseases (I60-I69)	115	4.0	65.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	3.7	60.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	2.8	46.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	78	2.7	44.4
9	Influenza and pneumonia (J09-J18)	58	2.0	33.0
10	Septicemia (A40-A41)	52	1.8	29.6
11	Alzheimer's disease (G30)	28	1.0	15.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.8	12.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.7	11.4
14	Parkinson's disease (G20-G21)	16	0.6	*
15	Aortic aneurysm and dissection (I71)	15	0.5	*
...	All other causes (Residual)	465	16.3	264.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	2,793	100.0	3,696.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	653	23.4	864.2
2	Malignant neoplasms (C00-C97)	596	21.3	788.8
3	Diabetes mellitus (E10-E14)	190	6.8	251.5
4	Chronic lower respiratory diseases (J40-J47)	178	6.4	235.6
5	Cerebrovascular diseases (I60-I69)	176	6.3	232.9
6	Alzheimer's disease (G30)	81	2.9	107.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	2.9	105.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	2.4	88.7
9	Influenza and pneumonia (J09-J18)	64	2.3	84.7
10	Septicemia (A40-A41)	48	1.7	63.5
11	Parkinson's disease (G20-G21)	37	1.3	49.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	34	1.2	45.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	1.1	41.0
14	Pneumonitis due to solids and liquids (J69)	24	0.9	31.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.7	*
...	All other causes (Residual)	515	18.4	681.6
American Indian, both sexes, 85 years and over				
...	All causes	2,059	100.0	8,698.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	546	26.5	2,306.7
2	Malignant neoplasms (C00-C97)	248	12.0	1,047.7
3	Cerebrovascular diseases (I60-I69)	120	5.8	507.0
4	Alzheimer's disease (G30)	113	5.5	477.4
5	Diabetes mellitus (E10-E14)	105	5.1	443.6
6	Influenza and pneumonia (J09-J18)	97	4.7	409.8
7	Chronic lower respiratory diseases (J40-J47)	96	4.7	405.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	2.7	232.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	2.5	215.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.8	156.3
11	Septicemia (A40-A41)	33	1.6	139.4
12	Pneumonitis due to solids and liquids (J69)	32	1.6	135.2
13	Parkinson's disease (G20-G21)	19	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.6	*
15	Atherosclerosis (I70)	10	0.5	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.5	*
...	All other causes (Residual)	474	23.0	2,002.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	8,638	100.0	395.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,680	19.4	76.9
2	Malignant neoplasms (C00-C97)	1,486	17.2	68.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,227	14.2	56.1
4	Diabetes mellitus (E10-E14)	453	5.2	20.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	453	5.2	20.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	352	4.1	16.1
7	Chronic lower respiratory diseases (J40-J47)	304	3.5	13.9
8	Cerebrovascular diseases (I60-I69)	257	3.0	11.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	2.3	9.0
10	Influenza and pneumonia (J09-J18)	177	2.0	8.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	1.5	5.9
12	Septicemia (A40-A41)	109	1.3	5.0
13	Certain conditions originating in the perinatal period (P00-P96)	73	0.8	3.3
14	Viral hepatitis (B15-B19)	71	0.8	3.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	0.8	3.0
...	All other causes (Residual)	1,605	18.6	73.4
American Indian, male, 1-4 years				
...	All causes	46	100.0	28.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	45.7	13.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	10.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	8.7	*
4	Malignant neoplasms (C00-C97)	3	6.5	*
5	Chronic lower respiratory diseases (J40-J47)	2	4.3	*
6	Cerebrovascular diseases (I60-I69)	1	2.2	*
6	Influenza and pneumonia (J09-J18)	1	2.2	*
...	All other causes (Residual)	9	19.6	*
American Indian, male, 5-14 years				
...	All causes	50	100.0	13.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	30.0	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	14.0	*
3	Malignant neoplasms (C00-C97)	6	12.0	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	10.0	*
5	Cerebrovascular diseases (I60-I69)	2	4.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.0	*
7	Influenza and pneumonia (J09-J18)	1	2.0	*
7	Chronic lower respiratory diseases (J40-J47)	1	2.0	*
7	Pneumonitis due to solids and liquids (J69)	1	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	2.0	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.0	*
...	All other causes (Residual)	8	16.0	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 15-24 years				
...	All causes	422	100.0	107.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	45.5	48.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	109	25.8	27.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	10.4	11.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.1	*
5	Malignant neoplasms (C00-C97)	8	1.9	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.9	*
6	Legal intervention (Y35,Y89.0)	4	0.9	*
8	Influenza and pneumonia (J09-J18)	3	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
9	Pneumonitis due to solids and liquids (J69)	2	0.5	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
12	Anemias (D50-D64)	1	0.2	*
12	Diabetes mellitus (E10-E14)	1	0.2	*
12	Aortic aneurysm and dissection (I71)	1	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	37	8.8	9.4
American Indian, male, 25-34 years				
...	All causes	558	100.0	161.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	43.2	69.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	13.3	21.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	9.0	14.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	5.9	9.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	28	5.0	8.1
6	Malignant neoplasms (C00-C97)	17	3.0	*
7	Diabetes mellitus (E10-E14)	12	2.2	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.9	*
8	Cerebrovascular diseases (I60-I69)	5	0.9	*
10	Septicemia (A40-A41)	4	0.7	*
11	Influenza and pneumonia (J09-J18)	3	0.5	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.5	*
11	Legal intervention (Y35,Y89.0)	3	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
15	Meningococcal infection (A39)	1	0.2	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (Residual)	71	12.7	20.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 35-44 years				
...	All causes	734	100.0	250.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	201	27.4	68.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	12.0	30.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	87	11.9	29.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	10.9	27.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	5.2	13.0
6	Diabetes mellitus (E10-E14)	37	5.0	12.6
7	Malignant neoplasms (C00-C97)	33	4.5	11.3
8	Influenza and pneumonia (J09-J18)	12	1.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.1	*
10	Cerebrovascular diseases (I60-I69)	7	1.0	*
11	Viral hepatitis (B15-B19)	5	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
13	Septicemia (A40-A41)	4	0.5	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.5	*
15	Pneumonitis due to solids and liquids (J69)	3	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
...	All other causes (Residual)	119	16.2	40.7
American Indian, male, 45-54 years				
...	All causes	1,347	100.0	500.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247	18.3	91.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	246	18.3	91.4
3	Malignant neoplasms (C00-C97)	190	14.1	70.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	146	10.8	54.3
5	Diabetes mellitus (E10-E14)	62	4.6	23.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	3.3	16.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	2.7	13.8
8	Cerebrovascular diseases (I60-I69)	28	2.1	10.4
9	Septicemia (A40-A41)	22	1.6	8.2
10	Viral hepatitis (B15-B19)	21	1.6	7.8
10	Influenza and pneumonia (J09-J18)	21	1.6	7.8
12	Chronic lower respiratory diseases (J40-J47)	19	1.4	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.0	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.4	*
...	All other causes (Residual)	224	16.6	83.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 55-64 years				
...	All causes	1,606	100.0	894.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	378	23.5	210.6
2	Malignant neoplasms (C00-C97)	354	22.0	197.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	9.1	81.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	118	7.3	65.7
5	Diabetes mellitus (E10-E14)	115	7.2	64.1
6	Chronic lower respiratory diseases (J40-J47)	56	3.5	31.2
7	Cerebrovascular diseases (I60-I69)	52	3.2	29.0
8	Viral hepatitis (B15-B19)	36	2.2	20.1
9	Influenza and pneumonia (J09-J18)	32	2.0	17.8
10	Septicemia (A40-A41)	27	1.7	15.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	1.5	13.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	1.4	12.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.9	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.7	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
15	Peptic ulcer (K25-K28)	5	0.3	*
...	All other causes (Residual)	210	13.1	117.0
American Indian, male, 65-74 years				
...	All causes	1,570	100.0	1,908.1
1	Malignant neoplasms (C00-C97)	431	27.5	523.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	387	24.6	470.4
3	Diabetes mellitus (E10-E14)	100	6.4	121.5
4	Chronic lower respiratory diseases (J40-J47)	96	6.1	116.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	4.6	87.5
6	Cerebrovascular diseases (I60-I69)	55	3.5	66.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	51	3.2	62.0
8	Influenza and pneumonia (J09-J18)	38	2.4	46.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	2.3	43.8
10	Septicemia (A40-A41)	24	1.5	29.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	*
12	Parkinson's disease (G20-G21)	10	0.6	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.6	*
12	Aortic aneurysm and dissection (I71)	10	0.6	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	0.6	*
...	All other causes (Residual)	230	14.6	279.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 75-84 years				
...	All causes	1,349	100.0	4,240.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	340	25.2	1,068.8
2	Malignant neoplasms (C00-C97)	324	24.0	1,018.5
3	Diabetes mellitus (E10-E14)	88	6.5	276.6
4	Chronic lower respiratory diseases (J40-J47)	86	6.4	270.3
5	Cerebrovascular diseases (I60-I69)	72	5.3	226.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	3.6	154.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	2.1	88.0
8	Alzheimer's disease (G30)	27	2.0	84.9
9	Influenza and pneumonia (J09-J18)	26	1.9	81.7
10	Parkinson's disease (G20-G21)	20	1.5	62.9
11	Septicemia (A40-A41)	16	1.2	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.0	*
13	Pneumonitis due to solids and liquids (J69)	12	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.8	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.7	*
...	All other causes (Residual)	227	16.8	713.6
American Indian, male, 85 years and over				
...	All causes	754	100.0	9,169.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	197	26.1	2,395.7
2	Malignant neoplasms (C00-C97)	120	15.9	1,459.3
3	Influenza and pneumonia (J09-J18)	40	5.3	486.4
4	Diabetes mellitus (E10-E14)	38	5.0	462.1
4	Chronic lower respiratory diseases (J40-J47)	38	5.0	462.1
6	Cerebrovascular diseases (I60-I69)	35	4.6	425.6
7	Alzheimer's disease (G30)	32	4.2	389.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.8	255.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	2.8	255.4
10	Pneumonitis due to solids and liquids (J69)	15	2.0	*
11	Septicemia (A40-A41)	9	1.2	*
11	Parkinson's disease (G20-G21)	9	1.2	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.2	*
14	Atherosclerosis (I70)	6	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.7	*
...	All other causes (Residual)	154	20.4	1,872.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, all ages				
...	All causes	7,307	100.0	338.1
1	Malignant neoplasms (C00-C97)	1,324	18.1	61.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,238	16.9	57.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	602	8.2	27.9
4	Diabetes mellitus (E10-E14)	474	6.5	21.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	368	5.0	17.0
6	Chronic lower respiratory diseases (J40-J47)	353	4.8	16.3
7	Cerebrovascular diseases (I60-I69)	343	4.7	15.9
8	Influenza and pneumonia (J09-J18)	172	2.4	8.0
9	Alzheimer's disease (G30)	161	2.2	7.4
10	Septicemia (A40-A41)	160	2.2	7.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	2.2	7.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	1.5	5.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.0	3.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	0.8	2.7
15	Certain conditions originating in the perinatal period (P00-P96)	53	0.7	2.5
...	All other causes (Residual)	1,664	22.8	77.0
American Indian, female, 1-4 years				
...	All causes	39	100.0	25.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	25.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	10.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	7.7	*
4	Septicemia (A40-A41)	2	5.1	*
4	Malignant neoplasms (C00-C97)	2	5.1	*
4	Influenza and pneumonia (J09-J18)	2	5.1	*
7	Meningococcal infection (A39)	1	2.6	*
7	Cerebrovascular diseases (I60-I69)	1	2.6	*
7	Infections of kidney (N10-N12,N13.6,N15.1)	1	2.6	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.6	*
...	All other causes (Residual)	12	30.8	*
American Indian, female, 5-14 years				
...	All causes	46	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	37.0	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	13.0	*
3	Malignant neoplasms (C00-C97)	3	6.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	6.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.3	*
5	Cerebrovascular diseases (I60-I69)	2	4.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.3	*
8	Septicemia (A40-A41)	1	2.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.2	*
...	All other causes (Residual)	9	19.6	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 15-24 years				
...	All causes	167	100.0	45.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	52.1	23.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	15.6	7.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	5.4	*
4	Malignant neoplasms (C00-C97)	6	3.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.4	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.4	*
7	Cerebrovascular diseases (I60-I69)	3	1.8	*
8	Influenza and pneumonia (J09-J18)	2	1.2	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.2	*
10	Diabetes mellitus (E10-E14)	1	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
10	Peptic ulcer (K25-K28)	1	0.6	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.6	*
...	All other causes (Residual)	20	12.0	5.4
American Indian, female, 25-34 years				
...	All causes	283	100.0	88.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	38.5	33.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	32	11.3	10.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	7.1	6.2
4	Malignant neoplasms (C00-C97)	17	6.0	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	10	3.5	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	3.5	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.2	*
8	Diabetes mellitus (E10-E14)	5	1.8	*
8	Cerebrovascular diseases (I60-I69)	5	1.8	*
8	Influenza and pneumonia (J09-J18)	5	1.8	*
11	Septicemia (A40-A41)	3	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
...	All other causes (Residual)	50	17.7	15.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 35-44 years				
...	All causes	467	100.0	164.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	19.7	32.5
2	Chronic liver disease and cirrhosis (K70,K73-K74)	82	17.6	29.0
3	Malignant neoplasms (C00-C97)	46	9.9	16.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	9.6	15.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	5.8	9.5
6	Diabetes mellitus (E10-E14)	25	5.4	8.8
7	Cerebrovascular diseases (I60-I69)	14	3.0	*
8	Septicemia (A40-A41)	10	2.1	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	1.9	*
10	Influenza and pneumonia (J09-J18)	8	1.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.9	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.9	*
13	Viral hepatitis (B15-B19)	3	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
...	All other causes (Residual)	92	19.7	32.5
American Indian, female, 45-54 years				
...	All causes	930	100.0	336.6
1	Malignant neoplasms (C00-C97)	189	20.3	68.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	122	13.1	44.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	12.2	40.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	11.2	37.6
5	Diabetes mellitus (E10-E14)	62	6.7	22.4
6	Cerebrovascular diseases (I60-I69)	31	3.3	11.2
7	Chronic lower respiratory diseases (J40-J47)	23	2.5	8.3
8	Septicemia (A40-A41)	20	2.2	7.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	2.0	*
10	Influenza and pneumonia (J09-J18)	17	1.8	*
11	Viral hepatitis (B15-B19)	14	1.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.2	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.6	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	5	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.3	*
...	All other causes (Residual)	191	20.5	69.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 55-64 years				
...	All causes	1,176	100.0	611.5
1	Malignant neoplasms (C00-C97)	309	26.3	160.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	16.2	99.3
3	Diabetes mellitus (E10-E14)	94	8.0	48.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	78	6.6	40.6
5	Chronic lower respiratory diseases (J40-J47)	69	5.9	35.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	5.6	34.3
7	Septicemia (A40-A41)	38	3.2	19.8
8	Cerebrovascular diseases (I60-I69)	37	3.1	19.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	2.2	13.5
10	Influenza and pneumonia (J09-J18)	22	1.9	11.4
11	Viral hepatitis (B15-B19)	12	1.0	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	0.7	*
13	Pneumonitis due to solids and liquids (J69)	7	0.6	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.4	*
...	All other causes (Residual)	203	17.3	105.6
American Indian, female, 65-74 years				
...	All causes	1,286	100.0	1,376.1
1	Malignant neoplasms (C00-C97)	351	27.3	375.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	211	16.4	225.8
3	Diabetes mellitus (E10-E14)	118	9.2	126.3
4	Chronic lower respiratory diseases (J40-J47)	105	8.2	112.4
5	Cerebrovascular diseases (I60-I69)	60	4.7	64.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	3.5	48.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	2.7	37.5
8	Septicemia (A40-A41)	28	2.2	30.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	27	2.1	28.9
10	Alzheimer's disease (G30)	22	1.7	23.5
11	Influenza and pneumonia (J09-J18)	20	1.6	21.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	8	0.6	*
15	Nutritional deficiencies (E40-E64)	6	0.5	*
15	Parkinson's disease (G20-G21)	6	0.5	*
...	All other causes (Residual)	223	17.3	238.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 75-84 years				
...	All causes	1,444	100.0	3,300.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	313	21.7	715.5
2	Malignant neoplasms (C00-C97)	272	18.8	621.8
3	Cerebrovascular diseases (I60-I69)	104	7.2	237.7
4	Diabetes mellitus (E10-E14)	102	7.1	233.2
5	Chronic lower respiratory diseases (J40-J47)	92	6.4	210.3
6	Alzheimer's disease (G30)	54	3.7	123.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	2.7	89.1
8	Influenza and pneumonia (J09-J18)	38	2.6	86.9
9	Septicemia (A40-A41)	32	2.2	73.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	2.1	70.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	1.5	50.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20	1.4	45.7
13	Parkinson's disease (G20-G21)	17	1.2	*
14	Pneumonitis due to solids and liquids (J69)	12	0.8	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	12	0.8	*
...	All other causes (Residual)	284	19.7	649.2
American Indian, female, 85 years and over				
...	All causes	1,305	100.0	8,448.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	349	26.7	2,259.3
2	Malignant neoplasms (C00-C97)	128	9.8	828.6
3	Cerebrovascular diseases (I60-I69)	85	6.5	550.3
4	Alzheimer's disease (G30)	81	6.2	524.4
5	Diabetes mellitus (E10-E14)	67	5.1	433.7
6	Chronic lower respiratory diseases (J40-J47)	58	4.4	375.5
7	Influenza and pneumonia (J09-J18)	57	4.4	369.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	2.6	220.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	2.3	194.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	2.1	181.3
11	Septicemia (A40-A41)	24	1.8	155.4
12	Pneumonitis due to solids and liquids (J69)	17	1.3	*
13	Parkinson's disease (G20-G21)	10	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	*
15	Aortic aneurysm and dissection (I71)	6	0.5	*
...	All other causes (Residual)	323	24.8	2,091.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	53,336	100.0	305.3
1	Malignant neoplasms (C00-C97)	14,713	27.6	84.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,586	21.7	66.3
3	Cerebrovascular diseases (I60-I69)	3,937	7.4	22.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,327	4.4	13.3
5	Diabetes mellitus (E10-E14)	2,035	3.8	11.6
6	Chronic lower respiratory diseases (J40-J47)	1,705	3.2	9.8
7	Influenza and pneumonia (J09-J18)	1,652	3.1	9.5
8	Alzheimer's disease (G30)	1,211	2.3	6.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,043	2.0	6.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	949	1.8	5.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	878	1.6	5.0
12	Septicemia (A40-A41)	613	1.1	3.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	498	0.9	2.9
14	Certain conditions originating in the perinatal period (P00-P96)	482	0.9	2.8
15	Parkinson's disease (G20-G21)	460	0.9	2.6
...	All other causes (Residual)	9,247	17.3	52.9
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	129	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	16.3	2.2
2	Malignant neoplasms (C00-C97)	18	14.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	12.4	*
4	Influenza and pneumonia (J09-J18)	11	8.5	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	7.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.1	*
6	Chronic lower respiratory diseases (J40-J47)	4	3.1	*
8	Septicemia (A40-A41)	2	1.6	*
8	Anemias (D50-D64)	2	1.6	*
10	Tuberculosis (A16-A19)	1	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
...	All other causes (Residual)	40	31.0	4.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	197	100.0	8.5
1	Malignant neoplasms (C00-C97)	42	21.3	1.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	20.3	1.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	5.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	5.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	4.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	4.6	*
7	Influenza and pneumonia (J09-J18)	8	4.1	*
8	Anemias (D50-D64)	6	3.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	2.5	*
10	Cerebrovascular diseases (I60-I69)	4	2.0	*
11	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.0	*
13	Salmonella infections (A01-A02)	1	0.5	*
13	Septicemia (A40-A41)	1	0.5	*
13	Diabetes mellitus (E10-E14)	1	0.5	*
13	Diseases of appendix (K35-K38)	1	0.5	*
...	All other causes (Residual)	45	22.8	1.9
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	734	100.0	29.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	37.2	10.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	21.8	6.4
3	Malignant neoplasms (C00-C97)	70	9.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	8.7	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	4.4	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	2.6	*
7	Cerebrovascular diseases (I60-I69)	8	1.1	*
8	Septicemia (A40-A41)	6	0.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
8	Influenza and pneumonia (J09-J18)	6	0.8	*
11	Aortic aneurysm and dissection (I71)	5	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
13	Diabetes mellitus (E10-E14)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.3	*
13	Legal intervention (Y35,Y89.0)	2	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	68	9.3	2.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	1,124	100.0	38.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	266	23.7	9.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	223	19.8	7.7
3	Malignant neoplasms (C00-C97)	172	15.3	5.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	110	9.8	3.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	91	8.1	3.1
6	Cerebrovascular diseases (I60-I69)	22	2.0	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	20	1.8	0.7
8	Influenza and pneumonia (J09-J18)	11	1.0	*
9	Septicemia (A40-A41)	10	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	10	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.8	*
11	Diabetes mellitus (E10-E14)	9	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.7	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
...	All other causes (Residual)	151	13.4	5.2
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	1,905	100.0	67.4
1	Malignant neoplasms (C00-C97)	562	29.5	19.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	273	14.3	9.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	12.5	8.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	195	10.2	6.9
5	Cerebrovascular diseases (I60-I69)	107	5.6	3.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	3.6	2.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	43	2.3	1.5
8	Diabetes mellitus (E10-E14)	32	1.7	1.1
9	Influenza and pneumonia (J09-J18)	28	1.5	1.0
10	Septicemia (A40-A41)	20	1.0	0.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.0	0.7
12	Viral hepatitis (B15-B19)	17	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.8	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	16	0.8	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	15	0.8	*
...	All other causes (Residual)	255	13.4	9.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	4,013	100.0	170.8
1	Malignant neoplasms (C00-C97)	1,617	40.3	68.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	731	18.2	31.1
3	Cerebrovascular diseases (I60-I69)	255	6.4	10.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	237	5.9	10.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	4.2	7.2
6	Diabetes mellitus (E10-E14)	131	3.3	5.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	108	2.7	4.6
8	Viral hepatitis (B15-B19)	62	1.5	2.6
9	Influenza and pneumonia (J09-J18)	53	1.3	2.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	1.0	1.8
11	Chronic lower respiratory diseases (J40-J47)	38	0.9	1.6
12	Septicemia (A40-A41)	32	0.8	1.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	0.8	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.7	1.2
15	Aortic aneurysm and dissection (I71)	20	0.5	0.9
...	All other causes (Residual)	457	11.4	19.5
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	7,044	100.0	393.8
1	Malignant neoplasms (C00-C97)	3,094	43.9	173.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,326	18.8	74.1
3	Cerebrovascular diseases (I60-I69)	474	6.7	26.5
4	Diabetes mellitus (E10-E14)	288	4.1	16.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	253	3.6	14.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	2.1	8.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	122	1.7	6.8
8	Chronic lower respiratory diseases (J40-J47)	113	1.6	6.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	97	1.4	5.4
10	Septicemia (A40-A41)	86	1.2	4.8
10	Influenza and pneumonia (J09-J18)	86	1.2	4.8
12	Viral hepatitis (B15-B19)	84	1.2	4.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	63	0.9	3.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.7	2.7
15	Aortic aneurysm and dissection (I71)	44	0.6	2.5
...	All other causes (Residual)	721	10.2	40.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	8,875	100.0	922.6
1	Malignant neoplasms (C00-C97)	3,480	39.2	361.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,839	20.7	191.2
3	Cerebrovascular diseases (I60-I69)	623	7.0	64.8
4	Diabetes mellitus (E10-E14)	453	5.1	47.1
5	Chronic lower respiratory diseases (J40-J47)	242	2.7	25.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	2.7	25.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	187	2.1	19.4
8	Influenza and pneumonia (J09-J18)	158	1.8	16.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	122	1.4	12.7
10	Septicemia (A40-A41)	97	1.1	10.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	88	1.0	9.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.8	7.6
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	0.8	7.6
14	Parkinson's disease (G20-G21)	60	0.7	6.2
15	Viral hepatitis (B15-B19)	52	0.6	5.4
...	All other causes (Residual)	1,087	12.2	113.0
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	13,551	100.0	2,805.3
1	Malignant neoplasms (C00-C97)	3,651	26.9	755.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,107	22.9	643.2
3	Cerebrovascular diseases (I60-I69)	1,188	8.8	245.9
4	Chronic lower respiratory diseases (J40-J47)	612	4.5	126.7
5	Diabetes mellitus (E10-E14)	589	4.3	121.9
6	Influenza and pneumonia (J09-J18)	479	3.5	99.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	372	2.7	77.0
8	Alzheimer's disease (G30)	319	2.4	66.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	282	2.1	58.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	222	1.6	46.0
11	Parkinson's disease (G20-G21)	204	1.5	42.2
12	Septicemia (A40-A41)	200	1.5	41.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	116	0.9	24.0
14	Pneumonitis due to solids and liquids (J69)	102	0.8	21.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	89	0.7	18.4
...	All other causes (Residual)	2,019	14.9	418.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	14,881	100.0	8,948.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,135	27.8	2,486.5
2	Malignant neoplasms (C00-C97)	2,002	13.5	1,203.9
3	Cerebrovascular diseases (I60-I69)	1,254	8.4	754.1
4	Alzheimer's disease (G30)	837	5.6	503.3
5	Influenza and pneumonia (J09-J18)	806	5.4	484.7
6	Chronic lower respiratory diseases (J40-J47)	668	4.5	401.7
7	Diabetes mellitus (E10-E14)	529	3.6	318.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	430	2.9	258.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	362	2.4	217.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	325	2.2	195.4
11	Pneumonitis due to solids and liquids (J69)	185	1.2	111.2
12	Parkinson's disease (G20-G21)	180	1.2	108.2
13	Septicemia (A40-A41)	149	1.0	89.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	114	0.8	68.6
15	Aortic aneurysm and dissection (I71)	72	0.5	43.3
...	All other causes (Residual)	2,833	19.0	1,703.6
Asian or Pacific Islander, male, all ages				
...	All causes	27,490	100.0	328.6
1	Malignant neoplasms (C00-C97)	7,444	27.1	89.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,209	22.6	74.2
3	Cerebrovascular diseases (I60-I69)	1,775	6.5	21.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,520	5.5	18.2
5	Diabetes mellitus (E10-E14)	1,040	3.8	12.4
6	Chronic lower respiratory diseases (J40-J47)	1,008	3.7	12.0
7	Influenza and pneumonia (J09-J18)	854	3.1	10.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	720	2.6	8.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	489	1.8	5.8
10	Alzheimer's disease (G30)	404	1.5	4.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	353	1.3	4.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	326	1.2	3.9
13	Septicemia (A40-A41)	304	1.1	3.6
14	Certain conditions originating in the perinatal period (P00-P96)	278	1.0	3.3
15	Parkinson's disease (G20-G21)	263	1.0	3.1
...	All other causes (Residual)	4,503	16.4	53.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	74	100.0	15.4
1	Malignant neoplasms (C00-C97)	13	17.6	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	14.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	12.2	*
4	Influenza and pneumonia (J09-J18)	4	5.4	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	5.4	*
6	Anemias (D50-D64)	2	2.7	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.4	*
...	All other causes (Residual)	27	36.5	5.6
Asian or Pacific Islander, male, 5-14 years				
...	All causes	107	100.0	9.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	23.4	2.2
2	Malignant neoplasms (C00-C97)	21	19.6	1.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	6.5	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	5.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	4.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.7	*
7	Influenza and pneumonia (J09-J18)	3	2.8	*
7	Certain conditions originating in the perinatal period (P00-P96)	3	2.8	*
9	Anemias (D50-D64)	2	1.9	*
10	Salmonella infections (A01-A02)	1	0.9	*
10	Septicemia (A40-A41)	1	0.9	*
10	Diabetes mellitus (E10-E14)	1	0.9	*
...	All other causes (Residual)	28	26.2	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-24 years				
...	All causes	539	100.0	42.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	40.6	17.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	22.6	9.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	9.8	4.2
4	Malignant neoplasms (C00-C97)	42	7.8	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.3	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.9	*
7	Cerebrovascular diseases (I60-I69)	5	0.9	*
7	Aortic aneurysm and dissection (I71)	5	0.9	*
9	Septicemia (A40-A41)	4	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
11	Influenza and pneumonia (J09-J18)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
11	Legal intervention (Y35,Y89.0)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	39	7.2	3.1
Asian or Pacific Islander, male, 25-34 years				
...	All causes	749	100.0	53.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	214	28.6	15.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	153	20.4	11.0
3	Malignant neoplasms (C00-C97)	94	12.6	6.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	11.1	6.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	8.0	4.3
6	Cerebrovascular diseases (I60-I69)	13	1.7	*
7	Chronic lower respiratory diseases (J40-J47)	10	1.3	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.9	*
9	Influenza and pneumonia (J09-J18)	6	0.8	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.8	*
11	Septicemia (A40-A41)	5	0.7	*
11	Diabetes mellitus (E10-E14)	5	0.7	*
13	Viral hepatitis (B15-B19)	4	0.5	*
14	Tuberculosis (A16-A19)	3	0.4	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.4	*
...	All other causes (Residual)	77	10.3	5.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,123	100.0	84.2
1	Malignant neoplasms (C00-C97)	215	19.1	16.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	213	19.0	16.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	175	15.6	13.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	12.0	10.1
5	Cerebrovascular diseases (I60-I69)	55	4.9	4.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	4.7	4.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	36	3.2	2.7
8	Diabetes mellitus (E10-E14)	23	2.0	1.7
9	Viral hepatitis (B15-B19)	14	1.2	*
9	Influenza and pneumonia (J09-J18)	14	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.2	*
12	Septicemia (A40-A41)	11	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.9	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.8	*
...	All other causes (Residual)	137	12.2	10.3
Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,389	100.0	216.8
1	Malignant neoplasms (C00-C97)	745	31.2	67.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	581	24.3	52.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	7.0	15.2
4	Cerebrovascular diseases (I60-I69)	142	5.9	12.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	111	4.6	10.1
6	Diabetes mellitus (E10-E14)	89	3.7	8.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	85	3.6	7.7
8	Viral hepatitis (B15-B19)	42	1.8	3.8
9	Influenza and pneumonia (J09-J18)	37	1.5	3.4
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	1.0	2.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	1.0	2.2
12	Chronic lower respiratory diseases (J40-J47)	22	0.9	2.0
13	Aortic aneurysm and dissection (I71)	18	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.6	*
...	All other causes (Residual)	269	11.3	24.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-64 years				
...	All causes	4,070	100.0	502.8
1	Malignant neoplasms (C00-C97)	1,568	38.5	193.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	914	22.5	112.9
3	Cerebrovascular diseases (I60-I69)	254	6.2	31.4
4	Diabetes mellitus (E10-E14)	180	4.4	22.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	4.2	21.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	2.5	12.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	97	2.4	12.0
8	Chronic lower respiratory diseases (J40-J47)	67	1.6	8.3
9	Viral hepatitis (B15-B19)	59	1.4	7.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	1.4	7.0
11	Septicemia (A40-A41)	49	1.2	6.1
12	Influenza and pneumonia (J09-J18)	48	1.2	5.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	40	1.0	4.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.7	3.6
15	Aortic aneurysm and dissection (I71)	28	0.7	3.5
...	All other causes (Residual)	409	10.0	50.5
Asian or Pacific Islander, male, 65-74 years				
...	All causes	5,095	100.0	1,167.1
1	Malignant neoplasms (C00-C97)	1,906	37.4	436.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,180	23.2	270.3
3	Cerebrovascular diseases (I60-I69)	330	6.5	75.6
4	Diabetes mellitus (E10-E14)	263	5.2	60.2
5	Chronic lower respiratory diseases (J40-J47)	155	3.0	35.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	3.0	34.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	106	2.1	24.3
8	Influenza and pneumonia (J09-J18)	96	1.9	22.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	62	1.2	14.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	57	1.1	13.1
11	Septicemia (A40-A41)	56	1.1	12.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	1.0	11.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.8	9.4
14	Parkinson's disease (G20-G21)	40	0.8	9.2
15	Aortic aneurysm and dissection (I71)	28	0.5	6.4
...	All other causes (Residual)	575	11.3	131.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-84 years				
...	All causes	6,643	100.0	3,311.2
1	Malignant neoplasms (C00-C97)	1,841	27.7	917.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,549	23.3	772.1
3	Cerebrovascular diseases (I60-I69)	518	7.8	258.2
4	Chronic lower respiratory diseases (J40-J47)	371	5.6	184.9
5	Diabetes mellitus (E10-E14)	264	4.0	131.6
6	Influenza and pneumonia (J09-J18)	262	3.9	130.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	2.9	95.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	128	1.9	63.8
9	Parkinson's disease (G20-G21)	125	1.9	62.3
10	Alzheimer's disease (G30)	121	1.8	60.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	92	1.4	45.9
12	Septicemia (A40-A41)	90	1.4	44.9
13	Pneumonitis due to solids and liquids (J69)	67	1.0	33.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.9	30.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	36	0.5	17.9
...	All other causes (Residual)	927	14.0	462.1
Asian or Pacific Islander, male, 85 years and over				
...	All causes	6,212	100.0	10,143.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,651	26.6	2,695.9
2	Malignant neoplasms (C00-C97)	996	16.0	1,626.3
3	Cerebrovascular diseases (I60-I69)	457	7.4	746.2
4	Influenza and pneumonia (J09-J18)	378	6.1	617.2
5	Chronic lower respiratory diseases (J40-J47)	374	6.0	610.7
6	Alzheimer's disease (G30)	264	4.2	431.1
7	Diabetes mellitus (E10-E14)	213	3.4	347.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	3.0	303.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	165	2.7	269.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	129	2.1	210.6
11	Pneumonitis due to solids and liquids (J69)	99	1.6	161.7
12	Parkinson's disease (G20-G21)	89	1.4	145.3
13	Septicemia (A40-A41)	69	1.1	112.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.8	81.6
15	Aortic aneurysm and dissection (I71)	39	0.6	63.7
...	All other causes (Residual)	1,053	17.0	1,719.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	25,846	100.0	283.8
1	Malignant neoplasms (C00-C97)	7,269	28.1	79.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,377	20.8	59.1
3	Cerebrovascular diseases (I60-I69)	2,162	8.4	23.7
4	Diabetes mellitus (E10-E14)	995	3.8	10.9
5	Alzheimer's disease (G30)	807	3.1	8.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	807	3.1	8.9
7	Influenza and pneumonia (J09-J18)	798	3.1	8.8
8	Chronic lower respiratory diseases (J40-J47)	697	2.7	7.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	525	2.0	5.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	460	1.8	5.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	323	1.2	3.5
12	Septicemia (A40-A41)	309	1.2	3.4
13	Certain conditions originating in the perinatal period (P00-P96)	204	0.8	2.2
14	Parkinson's disease (G20-G21)	197	0.8	2.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	191	0.7	2.1
...	All other causes (Residual)	4,725	18.3	51.9
Asian or Pacific Islander, female, 1-4 years				
...	All causes	55	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	18.2	*
2	Influenza and pneumonia (J09-J18)	7	12.7	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	12.7	*
4	Malignant neoplasms (C00-C97)	5	9.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	9.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	5.5	*
7	Septicemia (A40-A41)	2	3.6	*
7	Chronic lower respiratory diseases (J40-J47)	2	3.6	*
9	Tuberculosis (A16-A19)	1	1.8	*
...	All other causes (Residual)	13	23.6	*
Asian or Pacific Islander, female, 5-14 years				
...	All causes	90	100.0	7.9
1	Malignant neoplasms (C00-C97)	21	23.3	1.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	16.7	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.6	*
3	Influenza and pneumonia (J09-J18)	5	5.6	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	5.6	*
6	Anemias (D50-D64)	4	4.4	*
6	Cerebrovascular diseases (I60-I69)	4	4.4	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	4.4	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.3	*
10	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	2	2.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	2	2.2	*
13	Diseases of appendix (K35-K38)	1	1.1	*
...	All other causes (Residual)	17	18.9	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	195	100.0	15.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	27.7	4.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	19.5	3.1
3	Malignant neoplasms (C00-C97)	28	14.4	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	5.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	4.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	4.6	*
7	Influenza and pneumonia (J09-J18)	4	2.1	*
7	Chronic lower respiratory diseases (J40-J47)	4	2.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.5	*
9	Cerebrovascular diseases (I60-I69)	3	1.5	*
11	Septicemia (A40-A41)	2	1.0	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.0	*
13	Diabetes mellitus (E10-E14)	1	0.5	*
13	Meningitis (G00,G03)	1	0.5	*
13	Pneumonitis due to solids and liquids (J69)	1	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
...	All other causes (Residual)	24	12.3	2.0
Asian or Pacific Islander, female, 25-34 years				
...	All causes	375	100.0	24.6
1	Malignant neoplasms (C00-C97)	78	20.8	5.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	18.7	4.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	13.9	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	8.3	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	7.2	1.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	20	5.3	1.3
7	Cerebrovascular diseases (I60-I69)	9	2.4	*
8	Septicemia (A40-A41)	5	1.3	*
8	Influenza and pneumonia (J09-J18)	5	1.3	*
10	Anemias (D50-D64)	4	1.1	*
10	Diabetes mellitus (E10-E14)	4	1.1	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
...	All other causes (Residual)	58	15.5	3.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-44 years				
...	All causes	782	100.0	52.5
1	Malignant neoplasms (C00-C97)	347	44.4	23.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	8.1	4.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	7.7	4.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	7.7	4.0
5	Cerebrovascular diseases (I60-I69)	52	6.6	3.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	16	2.0	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	1.9	*
8	Influenza and pneumonia (J09-J18)	14	1.8	*
9	Septicemia (A40-A41)	9	1.2	*
9	Diabetes mellitus (E10-E14)	9	1.2	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.6	*
14	Aortic aneurysm and dissection (I71)	5	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.6	*
...	All other causes (Residual)	97	12.4	6.5
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,624	100.0	130.2
1	Malignant neoplasms (C00-C97)	872	53.7	69.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	150	9.2	12.0
3	Cerebrovascular diseases (I60-I69)	113	7.0	9.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	4.3	5.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	3.6	4.7
6	Diabetes mellitus (E10-E14)	42	2.6	3.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	23	1.4	1.8
8	Viral hepatitis (B15-B19)	20	1.2	1.6
9	Septicemia (A40-A41)	19	1.2	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	1.0	*
11	Influenza and pneumonia (J09-J18)	16	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	16	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.4	*
...	All other causes (Residual)	177	10.9	14.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-64 years				
...	All causes	2,974	100.0	303.7
1	Malignant neoplasms (C00-C97)	1,526	51.3	155.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	412	13.9	42.1
3	Cerebrovascular diseases (I60-I69)	220	7.4	22.5
4	Diabetes mellitus (E10-E14)	108	3.6	11.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	2.8	8.5
6	Chronic lower respiratory diseases (J40-J47)	46	1.5	4.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	1.5	4.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	1.3	4.1
9	Influenza and pneumonia (J09-J18)	38	1.3	3.9
10	Septicemia (A40-A41)	37	1.2	3.8
11	Viral hepatitis (B15-B19)	25	0.8	2.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	25	0.8	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	0.8	2.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.6	*
15	Aortic aneurysm and dissection (I71)	16	0.5	*
...	All other causes (Residual)	312	10.5	31.9
Asian or Pacific Islander, female, 65-74 years				
...	All causes	3,780	100.0	719.4
1	Malignant neoplasms (C00-C97)	1,574	41.6	299.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	659	17.4	125.4
3	Cerebrovascular diseases (I60-I69)	293	7.8	55.8
4	Diabetes mellitus (E10-E14)	190	5.0	36.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	2.4	17.1
6	Chronic lower respiratory diseases (J40-J47)	87	2.3	16.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	2.1	15.4
8	Influenza and pneumonia (J09-J18)	62	1.6	11.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	1.6	11.4
10	Septicemia (A40-A41)	41	1.1	7.8
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.8	6.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	31	0.8	5.9
13	Alzheimer's disease (G30)	30	0.8	5.7
14	Viral hepatitis (B15-B19)	26	0.7	4.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	0.6	4.6
...	All other causes (Residual)	500	13.2	95.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-84 years				
...	All causes	6,908	100.0	2,446.0
1	Malignant neoplasms (C00-C97)	1,810	26.2	640.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,558	22.6	551.7
3	Cerebrovascular diseases (I60-I69)	670	9.7	237.2
4	Diabetes mellitus (E10-E14)	325	4.7	115.1
5	Chronic lower respiratory diseases (J40-J47)	241	3.5	85.3
6	Influenza and pneumonia (J09-J18)	217	3.1	76.8
7	Alzheimer's disease (G30)	198	2.9	70.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	2.6	64.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	2.2	54.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	130	1.9	46.0
11	Septicemia (A40-A41)	110	1.6	38.9
12	Parkinson's disease (G20-G21)	79	1.1	28.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.8	19.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	53	0.8	18.8
15	Aortic aneurysm and dissection (I71)	35	0.5	12.4
...	Pneumonitis due to solids and liquids (J69)	35	0.5	12.4
...	All other causes (Residual)	1,057	15.3	374.3
Asian or Pacific Islander, female, 85 years and over				
...	All causes	8,669	100.0	8,251.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,484	28.7	2,364.5
2	Malignant neoplasms (C00-C97)	1,006	11.6	957.6
3	Cerebrovascular diseases (I60-I69)	797	9.2	758.7
4	Alzheimer's disease (G30)	573	6.6	545.4
5	Influenza and pneumonia (J09-J18)	428	4.9	407.4
6	Diabetes mellitus (E10-E14)	316	3.6	300.8
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	301	3.5	286.5
8	Chronic lower respiratory diseases (J40-J47)	294	3.4	279.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	176	2.0	167.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	160	1.8	152.3
11	Parkinson's disease (G20-G21)	91	1.0	86.6
12	Pneumonitis due to solids and liquids (J69)	86	1.0	81.9
13	Septicemia (A40-A41)	80	0.9	76.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	64	0.7	60.9
15	Aortic aneurysm and dissection (I71)	33	0.4	31.4
...	All other causes (Residual)	1,780	20.5	1,694.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2011