

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,426,264	100.0	810.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	631,636	26.0	211.0
2	Malignant neoplasms (C00-C97)	559,888	23.1	187.0
3	Cerebrovascular diseases (I60-I69)	137,119	5.7	45.8
4	Chronic lower respiratory diseases (J40-J47)	124,583	5.1	41.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121,599	5.0	40.6
6	Diabetes mellitus (E10-E14)	72,449	3.0	24.2
7	Alzheimer's disease (G30)	72,432	3.0	24.2
8	Influenza and pneumonia (J10-J18)	56,326	2.3	18.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45,344	1.9	15.1
10	Septicemia (A40-A41)	34,234	1.4	11.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,300	1.4	11.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27,555	1.1	9.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23,855	1.0	8.0
14	Parkinson's disease (G20-G21)	19,566	0.8	6.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18,573	0.8	6.2
...	All other causes (Residual)	447,805	18.5	149.6
All races, both sexes, 1-4 years				
...	All causes	4,631	100.0	28.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,610	34.8	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	515	11.1	3.2
3	Malignant neoplasms (C00-C97)	377	8.1	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	366	7.9	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	161	3.5	1.0
6	Influenza and pneumonia (J10-J18)	125	2.7	0.8
7	Septicemia (A40-A41)	88	1.9	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	65	1.4	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	1.3	0.4
10	Cerebrovascular diseases (I60-I69)	54	1.2	0.3
11	Chronic lower respiratory diseases (J40-J47)	43	0.9	0.3
12	Anemias (D50-D64)	26	0.6	0.2
13	Complications of medical and surgical care (Y40-Y84,Y88)	22	0.5	0.1
14	Meningitis (G00,G03)	20	0.4	0.1
15	Meningococcal infection (A39)	14	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	14	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.3	*
...	All other causes (Residual)	1,057	22.8	6.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex: United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	6,149	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,258	36.7	5.6
2	Malignant neoplasms (C00-C97)	907	14.8	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	390	6.3	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	344	5.6	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	253	4.1	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	219	3.6	0.5
7	Chronic lower respiratory diseases (J40-J47)	115	1.9	0.3
8	Cerebrovascular diseases (I60-I69)	95	1.5	0.2
9	Septicemia (A40-A41)	84	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	1.2	0.2
11	Influenza and pneumonia (J10-J18)	68	1.1	0.2
12	Anemias (D50-D64)	34	0.6	0.1
13	Diabetes mellitus (E10-E14)	28	0.5	0.1
14	Certain conditions originating in the perinatal period (P00-P96)	24	0.4	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	19	0.3	*
...	All other causes (Residual)	1,235	20.1	3.1
All races, both sexes, 15-24 years				
...	All causes	34,887	100.0	82.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,229	46.5	38.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,717	16.4	13.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,189	12.0	9.9
4	Malignant neoplasms (C00-C97)	1,644	4.7	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,076	3.1	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	460	1.3	1.1
7	Cerebrovascular diseases (I60-I69)	210	0.6	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	206	0.6	0.5
9	Influenza and pneumonia (J10-J18)	184	0.5	0.4
10	Pregnancy, childbirth and the puerperium (O00-099)	179	0.5	0.4
11	Diabetes mellitus (E10-E14)	165	0.5	0.4
12	Chronic lower respiratory diseases (J40-J47)	151	0.4	0.4
13	Septicemia (A40-A41)	139	0.4	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	100	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	0.3	0.2
...	All other causes (Residual)	4,139	11.9	9.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-44 years				
...	All causes	125,995	100.0	149.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,488	25.8	38.6
2	Malignant neoplasms (C00-C97)	17,573	13.9	20.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,646	12.4	18.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,576	9.2	13.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,745	6.1	9.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,192	4.1	6.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,867	2.3	3.4
8	Diabetes mellitus (E10-E14)	2,767	2.2	3.3
9	Cerebrovascular diseases (I60-I69)	2,748	2.2	3.3
10	Influenza and pneumonia (J10-J18)	1,176	0.9	1.4
11	Septicemia (A40-A41)	1,161	0.9	1.4
12	Chronic lower respiratory diseases (J40-J47)	1,096	0.9	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,061	0.8	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	905	0.7	1.1
15	Viral hepatitis (B15-B19)	595	0.5	0.7
...	All other causes (Residual)	21,399	17.0	25.4
All races, both sexes, 45-64 years				
...	All causes	466,432	100.0	623.0
1	Malignant neoplasms (C00-C97)	151,788	32.5	202.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	103,572	22.2	138.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31,121	6.7	41.6
4	Diabetes mellitus (E10-E14)	17,124	3.7	22.9
5	Cerebrovascular diseases (I60-I69)	16,859	3.6	22.5
6	Chronic lower respiratory diseases (J40-J47)	16,299	3.5	21.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	14,929	3.2	19.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12,009	2.6	16.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,613	1.4	8.8
10	Septicemia (A40-A41)	6,292	1.3	8.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	6,099	1.3	8.1
12	Influenza and pneumonia (J10-J18)	5,161	1.1	6.9
13	Viral hepatitis (B15-B19)	5,038	1.1	6.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,477	0.7	4.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,221	0.7	4.3
...	All other causes (Residual)	66,830	14.3	89.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65 years and over				
...	All causes	1,759,423	100.0	4,722.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	510,542	29.0	1,370.2
2	Malignant neoplasms (C00-C97)	387,515	22.0	1,040.0
3	Cerebrovascular diseases (I60-I69)	117,010	6.7	314.0
4	Chronic lower respiratory diseases (J40-J47)	106,845	6.1	286.8
5	Alzheimer's disease (G30)	71,660	4.1	192.3
6	Diabetes mellitus (E10-E14)	52,351	3.0	140.5
7	Influenza and pneumonia (J10-J18)	49,346	2.8	132.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37,377	2.1	100.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36,689	2.1	98.5
10	Septicemia (A40-A41)	26,201	1.5	70.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19,858	1.1	53.3
12	Parkinson's disease (G20-G21)	19,080	1.1	51.2
13	Pneumonitis due to solids and liquids (J69)	15,288	0.9	41.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11,253	0.6	30.2
15	Aortic aneurysm and dissection (I71)	10,578	0.6	28.4
...	All other causes (Residual)	287,830	16.4	772.5
All races, male, all ages				
...	All causes	1,201,942	100.0	814.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	315,706	26.3	214.0
2	Malignant neoplasms (C00-C97)	290,069	24.1	196.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78,941	6.6	53.5
4	Chronic lower respiratory diseases (J40-J47)	59,260	4.9	40.2
5	Cerebrovascular diseases (I60-I69)	54,524	4.5	37.0
6	Diabetes mellitus (E10-E14)	36,006	3.0	24.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,308	2.2	17.8
8	Influenza and pneumonia (J10-J18)	25,650	2.1	17.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,094	1.8	15.0
10	Alzheimer's disease (G30)	21,151	1.8	14.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17,866	1.5	12.1
12	Septicemia (A40-A41)	15,522	1.3	10.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14,717	1.2	10.0
14	Parkinson's disease (G20-G21)	11,300	0.9	7.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,415	0.8	6.4
...	All other causes (Residual)	203,413	16.9	137.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,541	100.0	30.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	927	36.5	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	9.6	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	208	8.2	2.5
4	Malignant neoplasms (C00-C97)	207	8.1	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	3.5	1.1
6	Influenza and pneumonia (J10-J18)	63	2.5	0.8
7	Septicemia (A40-A41)	43	1.7	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	37	1.5	0.4
9	Chronic lower respiratory diseases (J40-J47)	29	1.1	0.3
10	Cerebrovascular diseases (I60-I69)	26	1.0	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.9	0.3
12	Anemias (D50-D64)	15	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
14	Meningococcal infection (A39)	10	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
...	All other causes (Residual)	599	23.6	7.2
All races, male, 5-14 years				
...	All causes	3,625	100.0	17.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,385	38.2	6.7
2	Malignant neoplasms (C00-C97)	507	14.0	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	246	6.8	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	181	5.0	0.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	153	4.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	140	3.9	0.7
7	Chronic lower respiratory diseases (J40-J47)	78	2.2	0.4
8	Cerebrovascular diseases (I60-I69)	56	1.5	0.3
9	Septicemia (A40-A41)	52	1.4	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	1.1	0.2
11	Influenza and pneumonia (J10-J18)	35	1.0	0.2
12	Anemias (D50-D64)	16	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	14	0.4	*
14	Diabetes mellitus (E10-E14)	13	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.3	*
...	All other causes (Residual)	696	19.2	3.4

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	26,070	100.0	119.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,234	46.9	56.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,991	19.1	22.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,528	13.5	16.2
4	Malignant neoplasms (C00-C97)	1,007	3.9	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	711	2.7	3.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	279	1.1	1.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	121	0.5	0.6
8	Cerebrovascular diseases (I60-I69)	115	0.4	0.5
9	Influenza and pneumonia (J10-J18)	101	0.4	0.5
10	Diabetes mellitus (E10-E14)	96	0.4	0.4
11	Chronic lower respiratory diseases (J40-J47)	93	0.4	0.4
12	Legal intervention (Y35,Y89.0)	82	0.3	0.4
13	Septicemia (A40-A41)	68	0.3	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.2	0.3
15	Anemias (D50-D64)	55	0.2	0.3
...	All other causes (Residual)	2,533	9.7	11.6
All races, male, 25-44 years				
...	All causes	82,338	100.0	194.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23,897	29.0	56.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,923	13.3	25.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,118	11.1	21.5
4	Malignant neoplasms (C00-C97)	7,749	9.4	18.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,252	7.6	14.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,558	4.3	8.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,881	2.3	4.4
8	Diabetes mellitus (E10-E14)	1,655	2.0	3.9
9	Cerebrovascular diseases (I60-I69)	1,453	1.8	3.4
10	Influenza and pneumonia (J10-J18)	637	0.8	1.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	615	0.7	1.4
12	Septicemia (A40-A41)	604	0.7	1.4
13	Chronic lower respiratory diseases (J40-J47)	533	0.6	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	528	0.6	1.2
15	Viral hepatitis (B15-B19)	409	0.5	1.0
...	All other causes (Residual)	12,526	15.2	29.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-64 years				
...	All causes	284,159	100.0	778.2
1	Malignant neoplasms (C00-C97)	80,689	28.4	221.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,610	25.6	198.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,685	7.6	59.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	10,612	3.7	29.1
5	Diabetes mellitus (E10-E14)	10,185	3.6	27.9
6	Cerebrovascular diseases (I60-I69)	9,379	3.3	25.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,039	3.2	24.8
8	Chronic lower respiratory diseases (J40-J47)	8,340	2.9	22.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	4,615	1.6	12.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,780	1.3	10.4
11	Viral hepatitis (B15-B19)	3,604	1.3	9.9
12	Septicemia (A40-A41)	3,391	1.2	9.3
13	Influenza and pneumonia (J10-J18)	3,063	1.1	8.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,366	0.8	6.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,086	0.7	5.7
...	All other causes (Residual)	38,715	13.6	106.0
All races, male, 65 years and over				
...	All causes	787,058	100.0	5,026.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	231,016	29.4	1,475.5
2	Malignant neoplasms (C00-C97)	199,866	25.4	1,276.5
3	Chronic lower respiratory diseases (J40-J47)	50,164	6.4	320.4
4	Cerebrovascular diseases (I60-I69)	43,412	5.5	277.3
5	Diabetes mellitus (E10-E14)	24,049	3.1	153.6
6	Influenza and pneumonia (J10-J18)	21,586	2.7	137.9
7	Alzheimer's disease (G30)	20,786	2.6	132.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,098	2.3	115.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,521	2.2	111.9
10	Septicemia (A40-A41)	11,221	1.4	71.7
11	Parkinson's disease (G20-G21)	10,979	1.4	70.1
12	Pneumonitis due to solids and liquids (J69)	7,917	1.0	50.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,008	0.9	44.8
14	Aortic aneurysm and dissection (I71)	5,732	0.7	36.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,437	0.7	34.7
...	All other causes (Residual)	112,266	14.3	717.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,224,322	100.0	806.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	315,930	25.8	208.0
2	Malignant neoplasms (C00-C97)	269,819	22.0	177.6
3	Cerebrovascular diseases (I60-I69)	82,595	6.7	54.4
4	Chronic lower respiratory diseases (J40-J47)	65,323	5.3	43.0
5	Alzheimer's disease (G30)	51,281	4.2	33.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42,658	3.5	28.1
7	Diabetes mellitus (E10-E14)	36,443	3.0	24.0
8	Influenza and pneumonia (J10-J18)	30,676	2.5	20.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,250	1.9	15.3
10	Septicemia (A40-A41)	18,712	1.5	12.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,440	1.2	9.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,689	0.8	6.4
13	Parkinson's disease (G20-G21)	8,266	0.7	5.4
14	Pneumonitis due to solids and liquids (J69)	7,971	0.7	5.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7,138	0.6	4.7
...	All other causes (Residual)	240,131	19.6	158.1
All races, female, 1-4 years				
...	All causes	2,090	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	683	32.7	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	271	13.0	3.4
3	Malignant neoplasms (C00-C97)	170	8.1	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	158	7.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	72	3.4	0.9
6	Influenza and pneumonia (J10-J18)	62	3.0	0.8
7	Septicemia (A40-A41)	45	2.2	0.6
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	1.7	0.5
9	Cerebrovascular diseases (I60-I69)	28	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	28	1.3	0.4
11	Meningitis (G00,G03)	14	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.7	*
13	Anemias (D50-D64)	11	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
...	All other causes (Residual)	479	22.9	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,524	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	873	34.6	4.4
2	Malignant neoplasms (C00-C97)	400	15.8	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	6.5	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	5.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	4.5	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	2.6	0.3
7	Cerebrovascular diseases (I60-I69)	39	1.5	0.2
8	Chronic lower respiratory diseases (J40-J47)	37	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.4	0.2
10	Influenza and pneumonia (J10-J18)	33	1.3	0.2
11	Septicemia (A40-A41)	32	1.3	0.2
12	Anemias (D50-D64)	18	0.7	*
13	Diabetes mellitus (E10-E14)	15	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	10	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	538	21.3	2.7
All races, female, 15-24 years				
...	All causes	8,817	100.0	42.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,995	45.3	19.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	726	8.2	3.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	661	7.5	3.2
4	Malignant neoplasms (C00-C97)	637	7.2	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	365	4.1	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	181	2.1	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	179	2.0	0.9
8	Cerebrovascular diseases (I60-I69)	95	1.1	0.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	85	1.0	0.4
10	Influenza and pneumonia (J10-J18)	83	0.9	0.4
11	Septicemia (A40-A41)	71	0.8	0.3
12	Diabetes mellitus (E10-E14)	69	0.8	0.3
13	Chronic lower respiratory diseases (J40-J47)	58	0.7	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.6	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.5	0.2
...	All other causes (Residual)	1,520	17.2	7.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-44 years				
...	All causes	43,657	100.0	104.8
1	Malignant neoplasms (C00-C97)	9,824	22.5	23.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,591	19.7	20.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,723	10.8	11.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,458	5.6	5.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,634	3.7	3.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,493	3.4	3.6
7	Cerebrovascular diseases (I60-I69)	1,295	3.0	3.1
8	Diabetes mellitus (E10-E14)	1,112	2.5	2.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	986	2.3	2.4
10	Chronic lower respiratory diseases (J40-J47)	563	1.3	1.4
11	Septicemia (A40-A41)	557	1.3	1.3
12	Influenza and pneumonia (J10-J18)	539	1.2	1.3
13	Pregnancy, childbirth and the puerperium (O00-099)	516	1.2	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	446	1.0	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	377	0.9	0.9
...	All other causes (Residual)	8,543	19.6	20.5
All races, female, 45-64 years				
...	All causes	182,273	100.0	475.3
1	Malignant neoplasms (C00-C97)	71,099	39.0	185.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,962	17.0	80.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,436	5.2	24.6
4	Chronic lower respiratory diseases (J40-J47)	7,959	4.4	20.8
5	Cerebrovascular diseases (I60-I69)	7,480	4.1	19.5
6	Diabetes mellitus (E10-E14)	6,939	3.8	18.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,317	2.4	11.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,970	1.6	7.7
9	Septicemia (A40-A41)	2,901	1.6	7.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,833	1.6	7.4
11	Influenza and pneumonia (J10-J18)	2,098	1.2	5.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,484	0.8	3.9
13	Viral hepatitis (B15-B19)	1,434	0.8	3.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,391	0.8	3.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	945	0.5	2.5
...	All other causes (Residual)	28,025	15.4	73.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65 years and over				
...	All causes	972,365	100.0	4,501.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	279,526	28.7	1,293.9
2	Malignant neoplasms (C00-C97)	187,649	19.3	868.6
3	Cerebrovascular diseases (I60-I69)	73,598	7.6	340.7
4	Chronic lower respiratory diseases (J40-J47)	56,681	5.8	262.4
5	Alzheimer's disease (G30)	50,874	5.2	235.5
6	Diabetes mellitus (E10-E14)	28,302	2.9	131.0
7	Influenza and pneumonia (J10-J18)	27,760	2.9	128.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,856	2.0	91.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,591	1.9	86.1
10	Septicemia (A40-A41)	14,980	1.5	69.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,850	1.3	59.5
12	Parkinson's disease (G20-G21)	8,101	0.8	37.5
13	Pneumonitis due to solids and liquids (J69)	7,371	0.8	34.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,816	0.6	26.9
15	Atherosclerosis (I70)	5,094	0.5	23.6
...	All other causes (Residual)	175,316	18.0	811.5
White, both sexes, all ages				
...	All causes	2,077,549	100.0	858.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	545,974	26.3	225.5
2	Malignant neoplasms (C00-C97)	482,575	23.2	199.3
3	Cerebrovascular diseases (I60-I69)	115,864	5.6	47.9
4	Chronic lower respiratory diseases (J40-J47)	114,993	5.5	47.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103,853	5.0	42.9
6	Alzheimer's disease (G30)	67,088	3.2	27.7
7	Diabetes mellitus (E10-E14)	57,204	2.8	23.6
8	Influenza and pneumonia (J10-J18)	49,401	2.4	20.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,793	1.7	14.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,138	1.5	12.4
11	Septicemia (A40-A41)	27,373	1.3	11.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	24,191	1.2	10.0
13	Parkinson's disease (G20-G21)	18,517	0.9	7.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,264	0.9	7.5
15	Pneumonitis due to solids and liquids (J69)	15,136	0.7	6.3
...	All other causes (Residual)	371,185	17.9	153.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	3,233	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,170	36.2	9.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	376	11.6	3.0
3	Malignant neoplasms (C00-C97)	286	8.8	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	190	5.9	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	110	3.4	0.9
6	Influenza and pneumonia (J10-J18)	83	2.6	0.7
7	Septicemia (A40-A41)	59	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	1.6	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	40	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	39	1.2	0.3
11	Chronic lower respiratory diseases (J40-J47)	26	0.8	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.5	*
13	Meningitis (G00,G03)	15	0.5	*
14	Meningococcal infection (A39)	13	0.4	*
14	Anemias (D50-D64)	13	0.4	*
...	All other causes (Residual)	745	23.0	5.9
White, both sexes, 5-14 years				
...	All causes	4,460	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,668	37.4	5.3
2	Malignant neoplasms (C00-C97)	720	16.1	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	259	5.8	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	227	5.1	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	3.8	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	3.7	0.5
7	Cerebrovascular diseases (I60-I69)	67	1.5	0.2
8	Septicemia (A40-A41)	63	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	1.3	0.2
10	Influenza and pneumonia (J10-J18)	53	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	42	0.9	0.1
12	Diabetes mellitus (E10-E14)	19	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	18	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.3	*
15	Anemias (D50-D64)	13	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
...	All other causes (Residual)	888	19.9	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	25,871	100.0	78.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,824	53.4	41.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,477	13.4	10.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,213	8.6	6.7
4	Malignant neoplasms (C00-C97)	1,294	5.0	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	730	2.8	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	343	1.3	1.0
7	Cerebrovascular diseases (I60-I69)	159	0.6	0.5
8	Influenza and pneumonia (J10-J18)	138	0.5	0.4
9	Pregnancy, childbirth and the puerperium (O00-099)	116	0.4	0.3
10	Diabetes mellitus (E10-E14)	106	0.4	0.3
11	Septicemia (A40-A41)	90	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.3	0.2
13	Chronic lower respiratory diseases (J40-J47)	69	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	0.3	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.2	0.2
...	All other causes (Residual)	3,117	12.0	9.4
White, both sexes, 25-44 years				
...	All causes	93,187	100.0	139.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,120	29.1	40.7
2	Malignant neoplasms (C00-C97)	13,554	14.5	20.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,022	11.8	16.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,204	11.0	15.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,471	3.7	5.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,386	2.6	3.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2,165	2.3	3.2
8	Diabetes mellitus (E10-E14)	1,847	2.0	2.8
9	Cerebrovascular diseases (I60-I69)	1,734	1.9	2.6
10	Influenza and pneumonia (J10-J18)	866	0.9	1.3
11	Septicemia (A40-A41)	758	0.8	1.1
12	Chronic lower respiratory diseases (J40-J47)	732	0.8	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	730	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	569	0.6	0.9
15	Viral hepatitis (B15-B19)	468	0.5	0.7
...	All other causes (Residual)	15,561	16.7	23.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-64 years				
...	All causes	369,077	100.0	590.7
1	Malignant neoplasms (C00-C97)	124,216	33.7	198.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	80,177	21.7	128.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25,759	7.0	41.2
4	Chronic lower respiratory diseases (J40-J47)	14,236	3.9	22.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	12,895	3.5	20.6
6	Diabetes mellitus (E10-E14)	12,541	3.4	20.1
7	Cerebrovascular diseases (I60-I69)	11,552	3.1	18.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,246	3.0	18.0
9	Septicemia (A40-A41)	4,438	1.2	7.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,209	1.1	6.7
11	Influenza and pneumonia (J10-J18)	4,049	1.1	6.5
12	Viral hepatitis (B15-B19)	4,041	1.1	6.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2,607	0.7	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,054	0.6	3.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,939	0.5	3.1
...	All other causes (Residual)	53,118	14.4	85.0
White, both sexes, 65 years and over				
...	All causes	1,563,139	100.0	4,797.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	453,523	29.0	1,391.9
2	Malignant neoplasms (C00-C97)	342,435	21.9	1,051.0
3	Cerebrovascular diseases (I60-I69)	102,223	6.5	313.7
4	Chronic lower respiratory diseases (J40-J47)	99,870	6.4	306.5
5	Alzheimer's disease (G30)	66,386	4.2	203.7
6	Influenza and pneumonia (J10-J18)	44,048	2.8	135.2
7	Diabetes mellitus (E10-E14)	42,680	2.7	131.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33,532	2.1	102.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30,812	2.0	94.6
10	Septicemia (A40-A41)	21,809	1.4	66.9
11	Parkinson's disease (G20-G21)	18,093	1.2	55.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,988	1.0	49.1
13	Pneumonitis due to solids and liquids (J69)	13,831	0.9	42.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10,233	0.7	31.4
15	Aortic aneurysm and dissection (I71)	9,573	0.6	29.4
...	All other causes (Residual)	258,103	16.5	792.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,022,328	100.0	852.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	272,117	26.6	226.9
2	Malignant neoplasms (C00-C97)	250,322	24.5	208.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66,843	6.5	55.7
4	Chronic lower respiratory diseases (J40-J47)	54,043	5.3	45.1
5	Cerebrovascular diseases (I60-I69)	45,198	4.4	37.7
6	Diabetes mellitus (E10-E14)	29,060	2.8	24.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23,767	2.3	19.8
8	Influenza and pneumonia (J10-J18)	22,310	2.2	18.6
9	Alzheimer's disease (G30)	19,654	1.9	16.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,715	1.7	14.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,739	1.5	13.1
12	Septicemia (A40-A41)	12,450	1.2	10.4
13	Parkinson's disease (G20-G21)	10,706	1.0	8.9
14	Pneumonitis due to solids and liquids (J69)	7,984	0.8	6.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,990	0.7	5.8
...	All other causes (Residual)	167,430	16.4	139.6
White, male, 1-4 years				
...	All causes	1,781	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	675	37.9	10.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	10.7	2.9
3	Malignant neoplasms (C00-C97)	162	9.1	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	6.1	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	3.4	0.9
6	Influenza and pneumonia (J10-J18)	43	2.4	0.7
7	Septicemia (A40-A41)	29	1.6	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	26	1.5	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
11	Cerebrovascular diseases (I60-I69)	16	0.9	*
12	Meningococcal infection (A39)	9	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.4	*
15	Anemias (D50-D64)	5	0.3	*
...	All other causes (Residual)	404	22.7	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,639	100.0	16.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,017	38.5	6.3
2	Malignant neoplasms (C00-C97)	401	15.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	148	5.6	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	4.9	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	117	4.4	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	3.8	0.6
7	Cerebrovascular diseases (I60-I69)	46	1.7	0.3
8	Septicemia (A40-A41)	38	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.3	0.2
10	Chronic lower respiratory diseases (J40-J47)	32	1.2	0.2
11	Influenza and pneumonia (J10-J18)	31	1.2	0.2
12	Certain conditions originating in the perinatal period (P00-P96)	12	0.5	*
13	Diabetes mellitus (E10-E14)	11	0.4	*
14	Pneumonitis due to solids and liquids (J69)	8	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.3	*
...	All other causes (Residual)	500	18.9	3.1
White, male, 15-24 years				
...	All causes	19,175	100.0	111.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,423	54.4	60.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,939	15.3	17.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,824	9.5	10.6
4	Malignant neoplasms (C00-C97)	805	4.2	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	505	2.6	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	206	1.1	1.2
7	Cerebrovascular diseases (I60-I69)	91	0.5	0.5
8	Influenza and pneumonia (J10-J18)	75	0.4	0.4
9	Diabetes mellitus (E10-E14)	60	0.3	0.3
10	Legal intervention (Y35,Y89.0)	50	0.3	0.3
11	Septicemia (A40-A41)	43	0.2	0.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.2	0.3
13	Chronic lower respiratory diseases (J40-J47)	40	0.2	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	36	0.2	0.2
...	All other causes (Residual)	1,996	10.4	11.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-44 years				
...	All causes	61,853	100.0	181.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,957	32.3	58.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,989	12.9	23.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,974	12.9	23.4
4	Malignant neoplasms (C00-C97)	6,122	9.9	18.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,651	4.3	7.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,732	2.8	5.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,598	2.6	4.7
8	Diabetes mellitus (E10-E14)	1,118	1.8	3.3
9	Cerebrovascular diseases (I60-I69)	944	1.5	2.8
10	Influenza and pneumonia (J10-J18)	477	0.8	1.4
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	430	0.7	1.3
12	Septicemia (A40-A41)	405	0.7	1.2
13	Chronic lower respiratory diseases (J40-J47)	352	0.6	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	347	0.6	1.0
15	Viral hepatitis (B15-B19)	321	0.5	0.9
...	All other causes (Residual)	9,436	15.3	27.7
White, male, 45-64 years				
...	All causes	226,842	100.0	736.4
1	Malignant neoplasms (C00-C97)	66,232	29.2	215.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,565	25.4	186.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,863	7.9	58.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	9,236	4.1	30.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,451	3.7	27.4
6	Diabetes mellitus (E10-E14)	7,677	3.4	24.9
7	Chronic lower respiratory diseases (J40-J47)	7,199	3.2	23.4
8	Cerebrovascular diseases (I60-I69)	6,389	2.8	20.7
9	Viral hepatitis (B15-B19)	2,924	1.3	9.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,479	1.1	8.0
11	Septicemia (A40-A41)	2,405	1.1	7.8
12	Influenza and pneumonia (J10-J18)	2,396	1.1	7.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2,204	1.0	7.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,368	0.6	4.4
15	Aortic aneurysm and dissection (I71)	1,286	0.6	4.2
...	All other causes (Residual)	31,168	13.7	101.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65 years and over				
...	All causes	699,551	100.0	5,072.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	205,776	29.4	1,492.2
2	Malignant neoplasms (C00-C97)	176,563	25.2	1,280.4
3	Chronic lower respiratory diseases (J40-J47)	46,391	6.6	336.4
4	Cerebrovascular diseases (I60-I69)	37,664	5.4	273.1
5	Diabetes mellitus (E10-E14)	20,186	2.9	146.4
6	Alzheimer's disease (G30)	19,319	2.8	140.1
7	Influenza and pneumonia (J10-J18)	19,183	2.7	139.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,444	2.4	119.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,758	2.1	107.0
10	Parkinson's disease (G20-G21)	10,421	1.5	75.6
11	Septicemia (A40-A41)	9,446	1.4	68.5
12	Pneumonitis due to solids and liquids (J69)	7,166	1.0	52.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,562	0.8	40.3
14	Aortic aneurysm and dissection (I71)	5,260	0.8	38.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,962	0.7	36.0
...	All other causes (Residual)	100,450	14.4	728.4
White, female, all ages				
...	All causes	1,055,221	100.0	863.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	273,857	26.0	224.2
2	Malignant neoplasms (C00-C97)	232,253	22.0	190.1
3	Cerebrovascular diseases (I60-I69)	70,666	6.7	57.9
4	Chronic lower respiratory diseases (J40-J47)	60,950	5.8	49.9
5	Alzheimer's disease (G30)	47,434	4.5	38.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37,010	3.5	30.3
7	Diabetes mellitus (E10-E14)	28,144	2.7	23.0
8	Influenza and pneumonia (J10-J18)	27,091	2.6	22.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,078	1.7	14.8
10	Septicemia (A40-A41)	14,923	1.4	12.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,274	1.1	9.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,452	0.8	6.9
13	Parkinson's disease (G20-G21)	7,811	0.7	6.4
14	Pneumonitis due to solids and liquids (J69)	7,152	0.7	5.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,371	0.6	5.2
...	All other causes (Residual)	203,755	19.3	166.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,452	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	495	34.1	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	12.7	3.0
3	Malignant neoplasms (C00-C97)	124	8.5	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	5.6	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	3.4	0.8
6	Influenza and pneumonia (J10-J18)	40	2.8	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	2.2	0.5
8	Septicemia (A40-A41)	30	2.1	0.5
9	Cerebrovascular diseases (I60-I69)	23	1.6	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	14	1.0	*
11	Meningitis (G00,G03)	11	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.6	*
14	Anemias (D50-D64)	8	0.6	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (Residual)	331	22.8	5.4
White, female, 5-14 years				
...	All causes	1,821	100.0	11.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	651	35.7	4.3
2	Malignant neoplasms (C00-C97)	319	17.5	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	6.1	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	5.3	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	71	3.9	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	2.7	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.4	0.2
8	Septicemia (A40-A41)	25	1.4	0.2
9	Influenza and pneumonia (J10-J18)	22	1.2	0.1
10	Cerebrovascular diseases (I60-I69)	21	1.2	0.1
11	Chronic lower respiratory diseases (J40-J47)	10	0.5	*
12	Diabetes mellitus (E10-E14)	8	0.4	*
13	Anemias (D50-D64)	7	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (Residual)	391	21.5	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	6,696	100.0	41.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,401	50.8	21.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	538	8.0	3.4
3	Malignant neoplasms (C00-C97)	489	7.3	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	389	5.8	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	3.4	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	2.0	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	116	1.7	0.7
8	Cerebrovascular diseases (I60-I69)	68	1.0	0.4
9	Influenza and pneumonia (J10-J18)	63	0.9	0.4
10	Septicemia (A40-A41)	47	0.7	0.3
11	Diabetes mellitus (E10-E14)	46	0.7	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.5	0.2
13	Chronic lower respiratory diseases (J40-J47)	29	0.4	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.3	0.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.3	0.1
...	All other causes (Residual)	1,071	16.0	6.7
White, female, 25-44 years				
...	All causes	31,334	100.0	96.0
1	Malignant neoplasms (C00-C97)	7,432	23.7	22.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,163	22.9	21.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,048	9.7	9.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,215	7.1	6.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	820	2.6	2.5
6	Cerebrovascular diseases (I60-I69)	790	2.5	2.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	788	2.5	2.4
8	Diabetes mellitus (E10-E14)	729	2.3	2.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	433	1.4	1.3
10	Influenza and pneumonia (J10-J18)	389	1.2	1.2
11	Chronic lower respiratory diseases (J40-J47)	380	1.2	1.2
12	Septicemia (A40-A41)	353	1.1	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	300	1.0	0.9
14	Pregnancy, childbirth and the puerperium (O00-099)	293	0.9	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	222	0.7	0.7
...	All other causes (Residual)	5,979	19.1	18.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-64 years				
...	All causes	142,235	100.0	449.1
1	Malignant neoplasms (C00-C97)	57,984	40.8	183.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,612	15.9	71.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,896	5.6	24.9
4	Chronic lower respiratory diseases (J40-J47)	7,037	4.9	22.2
5	Cerebrovascular diseases (I60-I69)	5,163	3.6	16.3
6	Diabetes mellitus (E10-E14)	4,864	3.4	15.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,659	2.6	11.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,795	2.0	8.8
9	Septicemia (A40-A41)	2,033	1.4	6.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,730	1.2	5.5
11	Influenza and pneumonia (J10-J18)	1,653	1.2	5.2
12	Viral hepatitis (B15-B19)	1,117	0.8	3.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	769	0.5	2.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	757	0.5	2.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	571	0.4	1.8
...	All other causes (Residual)	21,595	15.2	68.2
White, female, 65 years and over				
...	All causes	863,588	100.0	4,595.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,747	28.7	1,318.4
2	Malignant neoplasms (C00-C97)	165,872	19.2	882.7
3	Cerebrovascular diseases (I60-I69)	64,559	7.5	343.5
4	Chronic lower respiratory diseases (J40-J47)	53,479	6.2	284.6
5	Alzheimer's disease (G30)	47,067	5.5	250.5
6	Influenza and pneumonia (J10-J18)	24,865	2.9	132.3
7	Diabetes mellitus (E10-E14)	22,494	2.6	119.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,088	2.0	90.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,054	1.9	85.4
10	Septicemia (A40-A41)	12,363	1.4	65.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,426	1.2	55.5
12	Parkinson's disease (G20-G21)	7,672	0.9	40.8
13	Pneumonitis due to solids and liquids (J69)	6,665	0.8	35.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,271	0.6	28.0
15	Atherosclerosis (I70)	4,675	0.5	24.9
...	All other causes (Residual)	157,291	18.2	837.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	289,971	100.0	733.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,253	24.9	182.6
2	Malignant neoplasms (C00-C97)	63,082	21.8	159.5
3	Cerebrovascular diseases (I60-I69)	17,045	5.9	43.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,917	4.8	35.2
5	Diabetes mellitus (E10-E14)	12,813	4.4	32.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9,032	3.1	22.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,397	2.9	21.2
8	Chronic lower respiratory diseases (J40-J47)	7,730	2.7	19.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,854	2.4	17.3
10	Septicemia (A40-A41)	6,108	2.1	15.4
11	Influenza and pneumonia (J10-J18)	5,311	1.8	13.4
12	Certain conditions originating in the perinatal period (P00-P96)	5,121	1.8	12.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,900	1.7	12.4
14	Alzheimer's disease (G30)	4,455	1.5	11.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,349	0.8	5.9
...	All other causes (Residual)	50,604	17.5	127.9
Black, both sexes, 1-4 years				
...	All causes	1,144	100.0	43.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	32.4	14.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	13.5	5.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	9.4	4.1
4	Malignant neoplasms (C00-C97)	65	5.7	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	3.8	1.7
6	Influenza and pneumonia (J10-J18)	34	3.0	1.3
7	Certain conditions originating in the perinatal period (P00-P96)	22	1.9	0.8
8	Septicemia (A40-A41)	21	1.8	0.8
9	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
10	Anemias (D50-D64)	12	1.0	*
10	Cerebrovascular diseases (I60-I69)	12	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
14	Meningitis (G00,G03)	4	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	265	23.2	10.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,394	100.0	21.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	471	33.8	7.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	149	10.7	2.3
3	Malignant neoplasms (C00-C97)	145	10.4	2.2
4	Chronic lower respiratory diseases (J40-J47)	69	4.9	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	4.8	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	4.8	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	2.7	0.6
8	Cerebrovascular diseases (I60-I69)	21	1.5	0.3
9	Septicemia (A40-A41)	20	1.4	0.3
9	Anemias (D50-D64)	20	1.4	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.8	*
13	Diabetes mellitus (E10-E14)	9	0.6	*
13	Influenza and pneumonia (J10-J18)	9	0.6	*
15	Meningitis (G00,G03)	8	0.6	*
...	All other causes (Residual)	276	19.8	4.2
Black, both sexes, 15-24 years				
...	All causes	7,483	100.0	112.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,295	44.0	49.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,750	23.4	26.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	416	5.6	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	308	4.1	4.6
5	Malignant neoplasms (C00-C97)	264	3.5	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	144	1.9	2.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	1.2	1.3
8	Chronic lower respiratory diseases (J40-J47)	75	1.0	1.1
9	Anemias (D50-D64)	64	0.9	1.0
10	Pregnancy, childbirth and the puerperium (O00-099)	58	0.8	0.9
11	Diabetes mellitus (E10-E14)	55	0.7	0.8
12	Cerebrovascular diseases (I60-I69)	44	0.6	0.7
13	Septicemia (A40-A41)	41	0.5	0.6
13	Influenza and pneumonia (J10-J18)	41	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.4	0.4
...	All other causes (Residual)	810	10.8	12.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-44 years				
...	All causes	27,975	100.0	246.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,280	15.3	37.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,111	14.7	36.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,017	14.4	35.4
4	Malignant neoplasms (C00-C97)	3,154	11.3	27.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,959	10.6	26.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	892	3.2	7.9
7	Cerebrovascular diseases (I60-I69)	848	3.0	7.5
8	Diabetes mellitus (E10-E14)	823	2.9	7.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	443	1.6	3.9
10	Septicemia (A40-A41)	362	1.3	3.2
11	Chronic lower respiratory diseases (J40-J47)	334	1.2	2.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	278	1.0	2.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	273	1.0	2.4
14	Influenza and pneumonia (J10-J18)	263	0.9	2.3
15	Anemias (D50-D64)	258	0.9	2.3
...	All other causes (Residual)	4,680	16.7	41.2
Black, both sexes, 45-64 years				
...	All causes	83,815	100.0	996.3
1	Malignant neoplasms (C00-C97)	22,971	27.4	273.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,662	24.7	245.6
3	Cerebrovascular diseases (I60-I69)	4,532	5.4	53.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,413	5.3	52.5
5	Diabetes mellitus (E10-E14)	3,918	4.7	46.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	3,425	4.1	40.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,187	2.6	26.0
8	Chronic lower respiratory diseases (J40-J47)	1,826	2.2	21.7
9	Septicemia (A40-A41)	1,677	2.0	19.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,514	1.8	18.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,310	1.6	15.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,134	1.4	13.5
13	Influenza and pneumonia (J10-J18)	968	1.2	11.5
14	Viral hepatitis (B15-B19)	792	0.9	9.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	466	0.6	5.5
...	All other causes (Residual)	12,020	14.3	142.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65 years and over				
...	All causes	159,270	100.0	4,965.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,937	29.5	1,463.3
2	Malignant neoplasms (C00-C97)	36,473	22.9	1,137.1
3	Cerebrovascular diseases (I60-I69)	11,539	7.2	359.7
4	Diabetes mellitus (E10-E14)	8,005	5.0	249.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,690	3.6	177.4
6	Chronic lower respiratory diseases (J40-J47)	5,400	3.4	168.3
7	Alzheimer's disease (G30)	4,390	2.8	136.9
8	Influenza and pneumonia (J10-J18)	3,906	2.5	121.8
9	Septicemia (A40-A41)	3,885	2.4	121.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,307	2.1	103.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,269	1.4	70.7
12	Pneumonitis due to solids and liquids (J69)	1,174	0.7	36.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	772	0.5	24.1
14	Aortic aneurysm and dissection (I71)	726	0.5	22.6
15	Parkinson's disease (G20-G21)	637	0.4	19.9
...	All other causes (Residual)	24,160	15.2	753.2
Black, male, all ages				
...	All causes	148,602	100.0	786.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,230	24.4	191.8
2	Malignant neoplasms (C00-C97)	32,556	21.9	172.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,605	6.5	50.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,677	5.2	40.6
5	Cerebrovascular diseases (I60-I69)	7,424	5.0	39.3
6	Diabetes mellitus (E10-E14)	5,772	3.9	30.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,443	3.0	23.5
8	Chronic lower respiratory diseases (J40-J47)	4,136	2.8	21.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,812	2.6	20.2
10	Certain conditions originating in the perinatal period (P00-P96)	2,811	1.9	14.9
11	Septicemia (A40-A41)	2,717	1.8	14.4
12	Influenza and pneumonia (J10-J18)	2,486	1.7	13.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,126	1.4	11.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,669	1.1	8.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,519	1.0	8.0
...	All other causes (Residual)	23,619	15.9	125.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	633	100.0	47.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	214	33.8	15.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	13.3	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	6.6	3.1
4	Malignant neoplasms (C00-C97)	35	5.5	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.3	2.0
6	Influenza and pneumonia (J10-J18)	16	2.5	*
7	Septicemia (A40-A41)	12	1.9	*
7	Chronic lower respiratory diseases (J40-J47)	12	1.9	*
9	Anemias (D50-D64)	9	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.4	*
11	Cerebrovascular diseases (I60-I69)	8	1.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
15	Meningococcal infection (A39)	1	0.2	*
15	Syphilis (A50-A53)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	149	23.5	11.1
Black, male, 5-14 years				
...	All causes	831	100.0	24.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	35.5	8.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	12.6	3.1
3	Malignant neoplasms (C00-C97)	83	10.0	2.5
4	Chronic lower respiratory diseases (J40-J47)	44	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.2	1.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.6	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	3.5	0.9
8	Septicemia (A40-A41)	13	1.6	*
9	Anemias (D50-D64)	9	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
10	Cerebrovascular diseases (I60-I69)	7	0.8	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
13	Meningitis (G00,G03)	3	0.4	*
13	Influenza and pneumonia (J10-J18)	3	0.4	*
13	Pneumonitis due to solids and liquids (J69)	3	0.4	*
...	All other causes (Residual)	159	19.1	4.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,793	100.0	171.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,982	51.5	88.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,331	23.0	39.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	358	6.2	10.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	183	3.2	5.4
5	Malignant neoplasms (C00-C97)	153	2.6	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	84	1.5	2.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	0.9	1.6
8	Chronic lower respiratory diseases (J40-J47)	47	0.8	1.4
9	Anemias (D50-D64)	37	0.6	1.1
10	Diabetes mellitus (E10-E14)	32	0.6	0.9
11	Legal intervention (Y35,Y89.0)	26	0.4	0.8
12	Influenza and pneumonia (J10-J18)	23	0.4	0.7
13	Septicemia (A40-A41)	21	0.4	0.6
14	Cerebrovascular diseases (I60-I69)	20	0.3	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
...	All other causes (Residual)	421	7.3	12.5
Black, male, 25-44 years				
...	All causes	17,440	100.0	322.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,414	19.6	63.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,172	18.2	58.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,555	14.7	47.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,776	10.2	32.9
5	Malignant neoplasms (C00-C97)	1,252	7.2	23.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	773	4.4	14.3
7	Diabetes mellitus (E10-E14)	483	2.8	8.9
8	Cerebrovascular diseases (I60-I69)	417	2.4	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	240	1.4	4.4
10	Septicemia (A40-A41)	179	1.0	3.3
11	Chronic lower respiratory diseases (J40-J47)	164	0.9	3.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	164	0.9	3.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	163	0.9	3.0
14	Influenza and pneumonia (J10-J18)	135	0.8	2.5
15	Anemias (D50-D64)	120	0.7	2.2
...	All other causes (Residual)	2,433	14.0	45.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-64 years				
...	All causes	49,310	100.0	1,285.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,079	26.5	340.9
2	Malignant neoplasms (C00-C97)	12,174	24.7	317.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,177	6.4	82.8
4	Cerebrovascular diseases (I60-I69)	2,567	5.2	66.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,355	4.8	61.4
6	Diabetes mellitus (E10-E14)	2,110	4.3	55.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,197	2.4	31.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,023	2.1	26.7
9	Chronic lower respiratory diseases (J40-J47)	1,009	2.0	26.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	895	1.8	23.3
11	Septicemia (A40-A41)	888	1.8	23.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	736	1.5	19.2
13	Influenza and pneumonia (J10-J18)	582	1.2	15.2
14	Viral hepatitis (B15-B19)	542	1.1	14.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	379	0.8	9.9
...	All other causes (Residual)	6,597	13.4	172.0
Black, male, 65 years and over				
...	All causes	69,687	100.0	5,650.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,283	29.1	1,644.6
2	Malignant neoplasms (C00-C97)	18,855	27.1	1,528.8
3	Cerebrovascular diseases (I60-I69)	4,372	6.3	354.5
4	Diabetes mellitus (E10-E14)	3,145	4.5	255.0
5	Chronic lower respiratory diseases (J40-J47)	2,854	4.1	231.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,337	3.4	189.5
7	Influenza and pneumonia (J10-J18)	1,672	2.4	135.6
8	Septicemia (A40-A41)	1,550	2.2	125.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,221	1.8	99.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,199	1.7	97.2
11	Alzheimer's disease (G30)	1,161	1.7	94.1
12	Pneumonitis due to solids and liquids (J69)	596	0.9	48.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	354	0.5	28.7
13	Parkinson's disease (G20-G21)	354	0.5	28.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	328	0.5	26.6
...	All other causes (Residual)	9,406	13.5	762.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	141,369	100.0	684.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,023	25.5	174.3
2	Malignant neoplasms (C00-C97)	30,526	21.6	147.7
3	Cerebrovascular diseases (I60-I69)	9,621	6.8	46.5
4	Diabetes mellitus (E10-E14)	7,041	5.0	34.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,585	3.2	22.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,312	3.1	20.9
7	Chronic lower respiratory diseases (J40-J47)	3,594	2.5	17.4
8	Septicemia (A40-A41)	3,391	2.4	16.4
9	Alzheimer's disease (G30)	3,265	2.3	15.8
10	Influenza and pneumonia (J10-J18)	2,825	2.0	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,774	2.0	13.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,411	1.7	11.7
13	Certain conditions originating in the perinatal period (P00-P96)	2,310	1.6	11.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,355	1.0	6.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	861	0.6	4.2
...	All other causes (Residual)	26,475	18.7	128.1
Black, female, 1-4 years				
...	All causes	511	100.0	39.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	30.7	12.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	13.7	5.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	12.7	5.0
4	Malignant neoplasms (C00-C97)	30	5.9	2.3
5	Influenza and pneumonia (J10-J18)	18	3.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.3	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.5	*
8	Septicemia (A40-A41)	9	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
9	Cerebrovascular diseases (I60-I69)	4	0.8	*
11	Anemias (D50-D64)	3	0.6	*
11	Meningitis (G00,G03)	3	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (Residual)	111	21.7	8.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	563	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	176	31.3	5.4
2	Malignant neoplasms (C00-C97)	62	11.0	1.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	7.8	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.7	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	5.7	1.0
6	Chronic lower respiratory diseases (J40-J47)	25	4.4	0.8
7	Cerebrovascular diseases (I60-I69)	14	2.5	*
8	Anemias (D50-D64)	11	2.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
10	Septicemia (A40-A41)	7	1.2	*
10	Diabetes mellitus (E10-E14)	7	1.2	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	1.2	*
13	Influenza and pneumonia (J10-J18)	6	1.1	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.9	*
14	Meningitis (G00,G03)	5	0.9	*
...	All other causes (Residual)	116	20.6	3.6
Black, female, 15-24 years				
...	All causes	1,690	100.0	51.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	419	24.8	12.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	313	18.5	9.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	7.4	3.8
4	Malignant neoplasms (C00-C97)	111	6.6	3.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	60	3.6	1.8
6	Pregnancy, childbirth and the puerperium (O00-099)	58	3.4	1.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	3.4	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	2.1	1.1
9	Chronic lower respiratory diseases (J40-J47)	28	1.7	0.9
10	Anemias (D50-D64)	27	1.6	0.8
11	Cerebrovascular diseases (I60-I69)	24	1.4	0.7
12	Diabetes mellitus (E10-E14)	23	1.4	0.7
13	Septicemia (A40-A41)	20	1.2	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.1	*
15	Influenza and pneumonia (J10-J18)	18	1.1	*
...	All other causes (Residual)	352	20.8	10.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-44 years				
...	All causes	10,535	100.0	176.9
1	Malignant neoplasms (C00-C97)	1,902	18.1	31.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,556	14.8	26.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,183	11.2	19.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,108	10.5	18.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	603	5.7	10.1
6	Cerebrovascular diseases (I60-I69)	431	4.1	7.2
7	Diabetes mellitus (E10-E14)	340	3.2	5.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	1.9	3.4
9	Septicemia (A40-A41)	183	1.7	3.1
10	Pregnancy, childbirth and the puerperium (O00-099)	179	1.7	3.0
11	Chronic lower respiratory diseases (J40-J47)	170	1.6	2.9
12	Anemias (D50-D64)	138	1.3	2.3
13	Influenza and pneumonia (J10-J18)	128	1.2	2.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	1.1	2.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	114	1.1	1.9
...	All other causes (Residual)	2,178	20.7	36.6
Black, female, 45-64 years				
...	All causes	34,505	100.0	754.0
1	Malignant neoplasms (C00-C97)	10,797	31.3	235.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,583	22.0	165.7
3	Cerebrovascular diseases (I60-I69)	1,965	5.7	42.9
4	Diabetes mellitus (E10-E14)	1,808	5.2	39.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,236	3.6	27.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,070	3.1	23.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	990	2.9	21.6
8	Chronic lower respiratory diseases (J40-J47)	817	2.4	17.9
9	Septicemia (A40-A41)	789	2.3	17.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	574	1.7	12.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	491	1.4	10.7
12	Influenza and pneumonia (J10-J18)	386	1.1	8.4
13	Viral hepatitis (B15-B19)	250	0.7	5.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	239	0.7	5.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	157	0.5	3.4
...	All other causes (Residual)	5,353	15.5	117.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65 years and over				
...	All causes	89,583	100.0	4,537.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,654	29.8	1,350.0
2	Malignant neoplasms (C00-C97)	17,618	19.7	892.3
3	Cerebrovascular diseases (I60-I69)	7,167	8.0	363.0
4	Diabetes mellitus (E10-E14)	4,860	5.4	246.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,353	3.7	169.8
6	Alzheimer's disease (G30)	3,229	3.6	163.5
7	Chronic lower respiratory diseases (J40-J47)	2,546	2.8	129.0
8	Septicemia (A40-A41)	2,335	2.6	118.3
9	Influenza and pneumonia (J10-J18)	2,234	2.5	113.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,086	2.3	105.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,070	1.2	54.2
12	Pneumonitis due to solids and liquids (J69)	578	0.6	29.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	418	0.5	21.2
14	Aortic aneurysm and dissection (I71)	416	0.5	21.1
15	Atherosclerosis (I70)	357	0.4	18.1
...	All other causes (Residual)	14,662	16.4	742.6
American Indian, both sexes, all ages				
...	All causes	14,037	100.0	438.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,736	19.5	85.5
2	Malignant neoplasms (C00-C97)	2,447	17.4	76.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,704	12.1	53.2
4	Diabetes mellitus (E10-E14)	811	5.8	25.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	596	4.2	18.6
6	Cerebrovascular diseases (I60-I69)	548	3.9	17.1
7	Chronic lower respiratory diseases (J40-J47)	508	3.6	15.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	395	2.8	12.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	288	2.1	9.0
10	Influenza and pneumonia (J10-J18)	267	1.9	8.3
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	254	1.8	7.9
12	Septicemia (A40-A41)	207	1.5	6.5
13	Alzheimer's disease (G30)	169	1.2	5.3
14	Certain conditions originating in the perinatal period (P00-P96)	137	1.0	4.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	0.8	3.7
...	All other causes (Residual)	2,853	20.3	89.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	94	100.0	54.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	34.0	18.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	11.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	10.6	*
4	Malignant neoplasms (C00-C97)	7	7.4	*
5	Influenza and pneumonia (J10-J18)	4	4.3	*
6	Septicemia (A40-A41)	2	2.1	*
6	Cerebrovascular diseases (I60-I69)	2	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
...	All other causes (Residual)	25	26.6	14.5
American Indian, both sexes, 5-14 years				
...	All causes	91	100.0	16.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	47.3	8.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	13.2	*
3	Malignant neoplasms (C00-C97)	5	5.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	5.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.3	*
6	Cerebrovascular diseases (I60-I69)	2	2.2	*
6	Influenza and pneumonia (J10-J18)	2	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
6	Diseases of appendix (K35-K38)	2	2.2	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.1	*
...	All other causes (Residual)	11	12.1	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	664	100.0	110.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	328	49.4	54.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	136	20.5	22.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	11.6	12.9
4	Malignant neoplasms (C00-C97)	16	2.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.7	*
7	Septicemia (A40-A41)	3	0.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
7	Cerebrovascular diseases (I60-I69)	3	0.5	*
7	Influenza and pneumonia (J10-J18)	3	0.5	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
7	Legal intervention (Y35,Y89.0)	3	0.5	*
14	Pregnancy, childbirth and the puerperium (O00-099)	2	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	53	8.0	8.8
American Indian, both sexes, 25-44 years				
...	All causes	1,960	100.0	206.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	605	30.9	63.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	176	9.0	18.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	163	8.3	17.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	152	7.8	16.0
5	Malignant neoplasms (C00-C97)	125	6.4	13.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	115	5.9	12.1
7	Diabetes mellitus (E10-E14)	54	2.8	5.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	39	2.0	4.1
9	Cerebrovascular diseases (I60-I69)	36	1.8	3.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	1.2	2.5
11	Influenza and pneumonia (J10-J18)	23	1.2	2.4
12	Septicemia (A40-A41)	19	1.0	*
12	Viral hepatitis (B15-B19)	19	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	13	0.7	*
15	Pregnancy, childbirth and the puerperium (O00-099)	7	0.4	*
...	All other causes (Residual)	390	19.9	41.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 45-64 years				
...	All causes	4,163	100.0	612.6
1	Malignant neoplasms (C00-C97)	851	20.4	125.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	824	19.8	121.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	470	11.3	69.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	336	8.1	49.4
5	Diabetes mellitus (E10-E14)	293	7.0	43.1
6	Chronic lower respiratory diseases (J40-J47)	119	2.9	17.5
7	Cerebrovascular diseases (I60-I69)	118	2.8	17.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	85	2.0	12.5
9	Septicemia (A40-A41)	77	1.8	11.3
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	1.8	10.7
11	Viral hepatitis (B15-B19)	69	1.7	10.2
12	Influenza and pneumonia (J10-J18)	50	1.2	7.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	1.0	5.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	26	0.6	3.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	0.6	3.4
...	All other causes (Residual)	709	17.0	104.3
American Indian, both sexes, 65 years and over				
...	All causes	6,670	100.0	3,047.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,712	25.7	782.2
2	Malignant neoplasms (C00-C97)	1,443	21.6	659.3
3	Diabetes mellitus (E10-E14)	463	6.9	211.5
4	Cerebrovascular diseases (I60-I69)	385	5.8	175.9
5	Chronic lower respiratory diseases (J40-J47)	371	5.6	169.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	2.8	86.4
7	Influenza and pneumonia (J10-J18)	182	2.7	83.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	174	2.6	79.5
9	Alzheimer's disease (G30)	168	2.5	76.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	105	1.6	48.0
11	Septicemia (A40-A41)	103	1.5	47.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	78	1.2	35.6
13	Parkinson's disease (G20-G21)	55	0.8	25.1
14	Pneumonitis due to solids and liquids (J69)	54	0.8	24.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.7	20.1
...	All other causes (Residual)	1,144	17.2	522.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	7,630	100.0	477.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,532	20.1	95.8
2	Malignant neoplasms (C00-C97)	1,217	16.0	76.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,184	15.5	74.0
4	Diabetes mellitus (E10-E14)	362	4.7	22.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	330	4.3	20.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	309	4.0	19.3
7	Chronic lower respiratory diseases (J40-J47)	234	3.1	14.6
8	Cerebrovascular diseases (I60-I69)	231	3.0	14.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	206	2.7	12.9
10	Influenza and pneumonia (J10-J18)	137	1.8	8.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	1.8	8.4
12	Septicemia (A40-A41)	99	1.3	6.2
13	Certain conditions originating in the perinatal period (P00-P96)	91	1.2	5.7
14	Viral hepatitis (B15-B19)	66	0.9	4.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	0.9	4.1
...	All other causes (Residual)	1,432	18.8	89.6
American Indian, male, 1-4 years				
...	All causes	51	100.0	58.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	29.4	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	17.6	*
3	Influenza and pneumonia (J10-J18)	3	5.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	5.9	*
5	Malignant neoplasms (C00-C97)	2	3.9	*
5	Cerebrovascular diseases (I60-I69)	2	3.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	2.0	*
...	All other causes (Residual)	16	31.4	*
American Indian, male, 5-14 years				
...	All causes	47	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	55.3	9.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	8.5	*
3	Malignant neoplasms (C00-C97)	3	6.4	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.1	*
6	Cerebrovascular diseases (I60-I69)	1	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.1	*
6	Diseases of appendix (K35-K38)	1	2.1	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	2.1	*
...	All other causes (Residual)	5	10.6	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 15-24 years				
...	All causes	478	100.0	156.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	237	49.6	77.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	110	23.0	35.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	14.4	22.5
4	Malignant neoplasms (C00-C97)	10	2.1	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.0	*
7	Legal intervention (Y35,Y89.0)	3	0.6	*
8	Septicemia (A40-A41)	2	0.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
10	Diabetes mellitus (E10-E14)	1	0.2	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
10	Cerebrovascular diseases (I60-I69)	1	0.2	*
10	Influenza and pneumonia (J10-J18)	1	0.2	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
...	All other causes (Residual)	25	5.2	8.2
American Indian, male, 25-44 years				
...	All causes	1,276	100.0	263.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	435	34.1	89.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	10.0	26.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	9.9	26.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	6.8	17.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	80	6.3	16.5
6	Malignant neoplasms (C00-C97)	48	3.8	9.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	26	2.0	5.4
8	Diabetes mellitus (E10-E14)	25	2.0	5.2
9	Cerebrovascular diseases (I60-I69)	24	1.9	4.9
10	Viral hepatitis (B15-B19)	17	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.1	*
12	Influenza and pneumonia (J10-J18)	12	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	8	0.6	*
14	Legal intervention (Y35,Y89.0)	6	0.5	*
15	Septicemia (A40-A41)	5	0.4	*
...	All other causes (Residual)	236	18.5	48.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-64 years				
...	All causes	2,493	100.0	759.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	564	22.6	171.8
2	Malignant neoplasms (C00-C97)	431	17.3	131.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	343	13.8	104.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	201	8.1	61.2
5	Diabetes mellitus (E10-E14)	159	6.4	48.4
6	Cerebrovascular diseases (I60-I69)	77	3.1	23.5
7	Chronic lower respiratory diseases (J40-J47)	59	2.4	18.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	2.4	18.0
9	Viral hepatitis (B15-B19)	45	1.8	13.7
10	Septicemia (A40-A41)	42	1.7	12.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.6	11.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	1.2	9.1
13	Influenza and pneumonia (J10-J18)	25	1.0	7.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.8	6.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.6	*
...	All other causes (Residual)	382	15.3	116.4
American Indian, male, 65 years and over				
...	All causes	3,042	100.0	3,178.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	832	27.4	869.4
2	Malignant neoplasms (C00-C97)	723	23.8	755.5
3	Diabetes mellitus (E10-E14)	177	5.8	185.0
4	Chronic lower respiratory diseases (J40-J47)	163	5.4	170.3
5	Cerebrovascular diseases (I60-I69)	125	4.1	130.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	3.5	111.8
7	Influenza and pneumonia (J10-J18)	96	3.2	100.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	2.6	81.5
9	Alzheimer's disease (G30)	62	2.0	64.8
10	Septicemia (A40-A41)	49	1.6	51.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	48	1.6	50.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	1.1	34.5
13	Parkinson's disease (G20-G21)	32	1.1	33.4
14	Pneumonitis due to solids and liquids (J69)	30	1.0	31.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.7	20.9
...	All other causes (Residual)	467	15.4	488.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, all ages				
...	All causes	6,407	100.0	399.9
1	Malignant neoplasms (C00-C97)	1,230	19.2	76.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,204	18.8	75.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	520	8.1	32.5
4	Diabetes mellitus (E10-E14)	449	7.0	28.0
5	Cerebrovascular diseases (I60-I69)	317	4.9	19.8
6	Chronic lower respiratory diseases (J40-J47)	274	4.3	17.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	266	4.2	16.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	2.4	9.5
9	Influenza and pneumonia (J10-J18)	130	2.0	8.1
10	Septicemia (A40-A41)	108	1.7	6.7
11	Alzheimer's disease (G30)	107	1.7	6.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	1.3	5.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	0.8	3.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	0.8	3.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	0.7	3.0
...	All other causes (Residual)	1,409	22.0	87.9
American Indian, female, 1-4 years				
...	All causes	43	100.0	50.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	39.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	16.3	*
3	Malignant neoplasms (C00-C97)	5	11.6	*
4	Septicemia (A40-A41)	2	4.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.7	*
6	Influenza and pneumonia (J10-J18)	1	2.3	*
...	All other causes (Residual)	9	20.9	*
American Indian, female, 5-14 years				
...	All causes	44	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	38.6	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	18.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.8	*
4	Malignant neoplasms (C00-C97)	2	4.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.5	*
4	Influenza and pneumonia (J10-J18)	2	4.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.3	*
7	Cerebrovascular diseases (I60-I69)	1	2.3	*
7	Chronic lower respiratory diseases (J40-J47)	1	2.3	*
7	Diseases of appendix (K35-K38)	1	2.3	*
...	All other causes (Residual)	6	13.6	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 15-24 years				
...	All causes	186	100.0	63.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	48.9	31.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	14.0	8.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	4.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	4.3	*
5	Malignant neoplasms (C00-C97)	6	3.2	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.6	*
7	Cerebrovascular diseases (I60-I69)	2	1.1	*
7	Influenza and pneumonia (J10-J18)	2	1.1	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.1	*
7	Pregnancy, childbirth and the puerperium (O00-099)	2	1.1	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.1	*
12	Septicemia (A40-A41)	1	0.5	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
12	Aortic aneurysm and dissection (I71)	1	0.5	*
12	Pneumonitis due to solids and liquids (J69)	1	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
...	All other causes (Residual)	28	15.1	9.6
American Indian, female, 25-44 years				
...	All causes	684	100.0	147.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	24.9	36.7
2	Malignant neoplasms (C00-C97)	77	11.3	16.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	72	10.5	15.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	7.2	10.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	5.4	8.0
6	Diabetes mellitus (E10-E14)	29	4.2	6.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	4.1	6.0
8	Septicemia (A40-A41)	14	2.0	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.9	*
10	Cerebrovascular diseases (I60-I69)	12	1.8	*
11	Influenza and pneumonia (J10-J18)	11	1.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.5	*
13	Pregnancy, childbirth and the puerperium (O00-099)	7	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
...	All other causes (Residual)	147	21.5	31.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-64 years				
...	All causes	1,670	100.0	475.4
1	Malignant neoplasms (C00-C97)	420	25.1	119.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	260	15.6	74.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	135	8.1	38.4
4	Diabetes mellitus (E10-E14)	134	8.0	38.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	7.6	36.2
6	Chronic lower respiratory diseases (J40-J47)	60	3.6	17.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	2.8	13.1
8	Cerebrovascular diseases (I60-I69)	41	2.5	11.7
9	Septicemia (A40-A41)	35	2.1	10.0
10	Influenza and pneumonia (J10-J18)	25	1.5	7.1
11	Viral hepatitis (B15-B19)	24	1.4	6.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	0.8	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.6	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.3	*
...	All other causes (Residual)	327	19.6	93.1
American Indian, female, 65 years and over				
...	All causes	3,628	100.0	2,945.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	880	24.3	714.5
2	Malignant neoplasms (C00-C97)	720	19.8	584.6
3	Diabetes mellitus (E10-E14)	286	7.9	232.2
4	Cerebrovascular diseases (I60-I69)	260	7.2	211.1
5	Chronic lower respiratory diseases (J40-J47)	208	5.7	168.9
6	Alzheimer's disease (G30)	106	2.9	86.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	2.6	77.9
8	Influenza and pneumonia (J10-J18)	86	2.4	69.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	2.3	66.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	57	1.6	46.3
11	Septicemia (A40-A41)	54	1.5	43.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	1.2	36.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.7	19.5
13	Pneumonitis due to solids and liquids (J69)	24	0.7	19.5
15	Parkinson's disease (G20-G21)	23	0.6	18.7
...	All other causes (Residual)	677	18.7	549.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	44,707	100.0	307.4
1	Malignant neoplasms (C00-C97)	11,784	26.4	81.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,673	23.9	73.4
3	Cerebrovascular diseases (I60-I69)	3,662	8.2	25.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,125	4.8	14.6
5	Diabetes mellitus (E10-E14)	1,621	3.6	11.1
6	Chronic lower respiratory diseases (J40-J47)	1,352	3.0	9.3
7	Influenza and pneumonia (J10-J18)	1,347	3.0	9.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	866	1.9	6.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	813	1.8	5.6
10	Alzheimer's disease (G30)	720	1.6	5.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	585	1.3	4.0
12	Septicemia (A40-A41)	546	1.2	3.8
13	Certain conditions originating in the perinatal period (P00-P96)	454	1.0	3.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	427	1.0	2.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	419	0.9	2.9
...	All other causes (Residual)	7,313	16.4	50.3
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	160	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	23.1	4.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	13.8	2.7
3	Malignant neoplasms (C00-C97)	19	11.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	6.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	4.4	*
6	Septicemia (A40-A41)	6	3.8	*
7	Influenza and pneumonia (J10-J18)	4	2.5	*
8	Pneumonitis due to solids and liquids (J69)	3	1.9	*
9	Chronic lower respiratory diseases (J40-J47)	2	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	2	1.3	*
11	Tuberculosis (A16-A19)	1	0.6	*
11	Anemias (D50-D64)	1	0.6	*
11	Meningitis (G00,G03)	1	0.6	*
11	Cerebrovascular diseases (I60-I69)	1	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	1	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (Residual)	41	25.6	5.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	204	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	37.3	4.0
2	Malignant neoplasms (C00-C97)	37	18.1	2.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	6.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	6.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	5.9	*
6	Cerebrovascular diseases (I60-I69)	5	2.5	*
7	Influenza and pneumonia (J10-J18)	4	2.0	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.0	*
11	Septicemia (A40-A41)	1	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
11	Anemias (D50-D64)	1	0.5	*
11	Diseases of appendix (K35-K38)	1	0.5	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	32	15.7	1.7
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	869	100.0	44.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	37.6	16.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	18.4	8.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	132	15.2	6.7
4	Malignant neoplasms (C00-C97)	70	8.1	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	2.8	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	2.1	*
7	Chronic lower respiratory diseases (J40-J47)	6	0.7	*
8	Septicemia (A40-A41)	5	0.6	*
9	Anemias (D50-D64)	4	0.5	*
9	Cerebrovascular diseases (I60-I69)	4	0.5	*
11	Diabetes mellitus (E10-E14)	3	0.3	*
11	Pregnancy, childbirth and the puerperium (O00-099)	3	0.3	*
11	Legal intervention (Y35,Y89.0)	3	0.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Viral hepatitis (B15-B19)	2	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.2	*
15	Influenza and pneumonia (J10-J18)	2	0.2	*
...	All other causes (Residual)	99	11.4	5.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-44 years				
...	All causes	2,873	100.0	56.3
1	Malignant neoplasms (C00-C97)	740	25.8	14.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	483	16.8	9.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	337	11.7	6.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	317	11.0	6.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	142	4.9	2.8
6	Cerebrovascular diseases (I60-I69)	130	4.5	2.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	51	1.8	1.0
8	Diabetes mellitus (E10-E14)	43	1.5	0.8
9	Pregnancy, childbirth and the puerperium (O00-099)	37	1.3	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	29	1.0	0.6
11	Viral hepatitis (B15-B19)	25	0.9	0.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.9	0.5
13	Influenza and pneumonia (J10-J18)	24	0.8	0.5
14	Septicemia (A40-A41)	22	0.8	0.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.8	0.4
...	All other causes (Residual)	446	15.5	8.7
Asian or Pacific Islander, both sexes, 45-64 years				
...	All causes	9,377	100.0	284.4
1	Malignant neoplasms (C00-C97)	3,750	40.0	113.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,909	20.4	57.9
3	Cerebrovascular diseases (I60-I69)	657	7.0	19.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	479	5.1	14.5
5	Diabetes mellitus (E10-E14)	372	4.0	11.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	224	2.4	6.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	184	2.0	5.6
8	Viral hepatitis (B15-B19)	136	1.5	4.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	1.4	4.0
10	Chronic lower respiratory diseases (J40-J47)	118	1.3	3.6
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	108	1.2	3.3
12	Septicemia (A40-A41)	100	1.1	3.0
13	Influenza and pneumonia (J10-J18)	94	1.0	2.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	1.0	2.7
15	Aortic aneurysm and dissection (I71)	51	0.5	1.5
...	All other causes (Residual)	973	10.4	29.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65 years and over				
...	All causes	30,344	100.0	2,424.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,370	27.6	668.7
2	Malignant neoplasms (C00-C97)	7,164	23.6	572.4
3	Cerebrovascular diseases (I60-I69)	2,863	9.4	228.7
4	Influenza and pneumonia (J10-J18)	1,210	4.0	96.7
5	Chronic lower respiratory diseases (J40-J47)	1,204	4.0	96.2
6	Diabetes mellitus (E10-E14)	1,203	4.0	96.1
7	Alzheimer's disease (G30)	716	2.4	57.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	701	2.3	56.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	699	2.3	55.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	485	1.6	38.7
11	Septicemia (A40-A41)	404	1.3	32.3
12	Parkinson's disease (G20-G21)	295	1.0	23.6
13	Aortic aneurysm and dissection (I71)	249	0.8	19.9
14	Pneumonitis due to solids and liquids (J69)	229	0.8	18.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	204	0.7	16.3
...	All other causes (Residual)	4,348	14.3	347.4
Asian or Pacific Islander, male, all ages				
...	All causes	23,382	100.0	330.6
1	Malignant neoplasms (C00-C97)	5,974	25.5	84.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,827	24.9	82.4
3	Cerebrovascular diseases (I60-I69)	1,671	7.1	23.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,309	5.6	18.5
5	Chronic lower respiratory diseases (J40-J47)	847	3.6	12.0
6	Diabetes mellitus (E10-E14)	812	3.5	11.5
7	Influenza and pneumonia (J10-J18)	717	3.1	10.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	563	2.4	8.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	432	1.8	6.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	320	1.4	4.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	278	1.2	3.9
12	Certain conditions originating in the perinatal period (P00-P96)	257	1.1	3.6
13	Septicemia (A40-A41)	256	1.1	3.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	247	1.1	3.5
15	Alzheimer's disease (G30)	245	1.0	3.5
...	All other causes (Residual)	3,627	15.5	51.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 1-4 years				
...	All causes	76	100.0	18.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	30.3	5.5
2	Malignant neoplasms (C00-C97)	8	10.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	10.5	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	7.9	*
5	Septicemia (A40-A41)	2	2.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.6	*
5	Pneumonitis due to solids and liquids (J69)	2	2.6	*
8	Tuberculosis (A16-A19)	1	1.3	*
8	Anemias (D50-D64)	1	1.3	*
8	Meningitis (G00,G03)	1	1.3	*
8	Influenza and pneumonia (J10-J18)	1	1.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.3	*
...	All other causes (Residual)	20	26.3	4.8
Asian or Pacific Islander, male, 5-14 years				
...	All causes	108	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	43.5	4.9
2	Malignant neoplasms (C00-C97)	20	18.5	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	8.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	4.6	*
5	Cerebrovascular diseases (I60-I69)	2	1.9	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.9	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	1.9	*
8	Septicemia (A40-A41)	1	0.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
8	Anemias (D50-D64)	1	0.9	*
8	Influenza and pneumonia (J10-J18)	1	0.9	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
8	Diseases of appendix (K35-K38)	1	0.9	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.9	*
...	All other causes (Residual)	13	12.0	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-24 years				
...	All causes	624	100.0	61.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	243	38.9	24.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	121	19.4	12.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	116	18.6	11.5
4	Malignant neoplasms (C00-C97)	39	6.3	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.8	*
7	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
8	Diabetes mellitus (E10-E14)	3	0.5	*
8	Cerebrovascular diseases (I60-I69)	3	0.5	*
8	Legal intervention (Y35,Y89.0)	3	0.5	*
11	Septicemia (A40-A41)	2	0.3	*
11	Viral hepatitis (B15-B19)	2	0.3	*
11	Anemias (D50-D64)	2	0.3	*
11	Influenza and pneumonia (J10-J18)	2	0.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	52	8.3	5.1
Asian or Pacific Islander, male, 25-44 years				
...	All causes	1,769	100.0	70.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	333	18.8	13.3
2	Malignant neoplasms (C00-C97)	327	18.5	13.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	267	15.1	10.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	230	13.0	9.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	5.7	4.0
6	Cerebrovascular diseases (I60-I69)	68	3.8	2.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	39	2.2	1.6
8	Diabetes mellitus (E10-E14)	29	1.6	1.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	24	1.4	1.0
10	Viral hepatitis (B15-B19)	22	1.2	0.9
11	Aortic aneurysm and dissection (I71)	16	0.9	*
12	Septicemia (A40-A41)	15	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.8	*
14	Influenza and pneumonia (J10-J18)	13	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	0.7	*
...	All other causes (Residual)	259	14.6	10.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-64 years				
...	All causes	5,514	100.0	357.3
1	Malignant neoplasms (C00-C97)	1,852	33.6	120.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,402	25.4	90.9
3	Cerebrovascular diseases (I60-I69)	346	6.3	22.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	302	5.5	19.6
5	Diabetes mellitus (E10-E14)	239	4.3	15.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	152	2.8	9.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	150	2.7	9.7
8	Viral hepatitis (B15-B19)	93	1.7	6.0
9	Chronic lower respiratory diseases (J40-J47)	73	1.3	4.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	1.3	4.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	1.2	4.2
12	Influenza and pneumonia (J10-J18)	60	1.1	3.9
13	Septicemia (A40-A41)	56	1.0	3.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	49	0.9	3.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.6	2.3
15	Aortic aneurysm and dissection (I71)	35	0.6	2.3
...	All other causes (Residual)	532	9.6	34.5
Asian or Pacific Islander, male, 65 years and over				
...	All causes	14,778	100.0	2,748.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,125	27.9	767.1
2	Malignant neoplasms (C00-C97)	3,725	25.2	692.7
3	Cerebrovascular diseases (I60-I69)	1,251	8.5	232.6
4	Chronic lower respiratory diseases (J40-J47)	756	5.1	140.6
5	Influenza and pneumonia (J10-J18)	635	4.3	118.1
6	Diabetes mellitus (E10-E14)	541	3.7	100.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	348	2.4	64.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	348	2.4	64.7
9	Alzheimer's disease (G30)	244	1.7	45.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	192	1.3	35.7
11	Septicemia (A40-A41)	176	1.2	32.7
12	Parkinson's disease (G20-G21)	172	1.2	32.0
13	Aortic aneurysm and dissection (I71)	147	1.0	27.3
14	Pneumonitis due to solids and liquids (J69)	125	0.8	23.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	101	0.7	18.8
...	All other causes (Residual)	1,892	12.8	351.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	21,325	100.0	285.6
1	Malignant neoplasms (C00-C97)	5,810	27.2	77.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,846	22.7	64.9
3	Cerebrovascular diseases (I60-I69)	1,991	9.3	26.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	816	3.8	10.9
5	Diabetes mellitus (E10-E14)	809	3.8	10.8
6	Influenza and pneumonia (J10-J18)	630	3.0	8.4
7	Chronic lower respiratory diseases (J40-J47)	505	2.4	6.8
8	Alzheimer's disease (G30)	475	2.2	6.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	434	2.0	5.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	338	1.6	4.5
11	Septicemia (A40-A41)	290	1.4	3.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	250	1.2	3.3
13	Certain conditions originating in the perinatal period (P00-P96)	197	0.9	2.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	0.7	2.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	141	0.7	1.9
...	All other causes (Residual)	3,647	17.1	48.8
Asian or Pacific Islander, female, 1-4 years				
...	All causes	84	100.0	21.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	16.7	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	16.7	*
3	Malignant neoplasms (C00-C97)	11	13.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	6.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	6.0	*
6	Septicemia (A40-A41)	4	4.8	*
7	Influenza and pneumonia (J10-J18)	3	3.6	*
8	Chronic lower respiratory diseases (J40-J47)	2	2.4	*
9	Cerebrovascular diseases (I60-I69)	1	1.2	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	1.2	*
9	Pneumonitis due to solids and liquids (J69)	1	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.2	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.2	*
...	All other causes (Residual)	21	25.0	5.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-14 years				
...	All causes	96	100.0	10.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	30.2	3.1
2	Malignant neoplasms (C00-C97)	17	17.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	11.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	8.3	*
5	Cerebrovascular diseases (I60-I69)	3	3.1	*
5	Influenza and pneumonia (J10-J18)	3	3.1	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.1	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.0	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	1.0	*
...	All other causes (Residual)	19	19.8	*
Asian or Pacific Islander, female, 15-24 years				
...	All causes	245	100.0	25.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	34.3	8.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	15.9	4.0
3	Malignant neoplasms (C00-C97)	31	12.7	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	6.5	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	2.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.4	*
7	Septicemia (A40-A41)	3	1.2	*
7	Pregnancy, childbirth and the puerperium (O00-099)	3	1.2	*
9	Anemias (D50-D64)	2	0.8	*
10	Syphilis (A50-A53)	1	0.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
10	Cerebrovascular diseases (I60-I69)	1	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
10	Hernia (K40-K46)	1	0.4	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
10	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.4	*
...	All other causes (Residual)	46	18.8	4.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-44 years				
...	All causes	1,104	100.0	42.4
1	Malignant neoplasms (C00-C97)	413	37.4	15.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	13.6	5.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	7.9	3.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	6.3	2.7
5	Cerebrovascular diseases (I60-I69)	62	5.6	2.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	3.8	1.6
7	Pregnancy, childbirth and the puerperium (O00-099)	37	3.4	1.4
8	Diabetes mellitus (E10-E14)	14	1.3	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	12	1.1	*
10	Influenza and pneumonia (J10-J18)	11	1.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.0	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
15	Septicemia (A40-A41)	7	0.6	*
...	All other causes (Residual)	163	14.8	6.3
Asian or Pacific Islander, female, 45-64 years				
...	All causes	3,863	100.0	220.3
1	Malignant neoplasms (C00-C97)	1,898	49.1	108.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	507	13.1	28.9
3	Cerebrovascular diseases (I60-I69)	311	8.1	17.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	4.6	10.1
5	Diabetes mellitus (E10-E14)	133	3.4	7.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	1.9	4.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	1.7	3.8
8	Chronic lower respiratory diseases (J40-J47)	45	1.2	2.6
9	Septicemia (A40-A41)	44	1.1	2.5
10	Viral hepatitis (B15-B19)	43	1.1	2.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	1.1	2.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	0.9	2.0
13	Influenza and pneumonia (J10-J18)	34	0.9	1.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	32	0.8	1.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.7	1.6
...	All other causes (Residual)	394	10.2	22.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and sex:  
United States, 2006

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65 years and over				
...	All causes	15,566	100.0	2,180.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,245	27.3	594.6
2	Malignant neoplasms (C00-C97)	3,439	22.1	481.7
3	Cerebrovascular diseases (I60-I69)	1,612	10.4	225.8
4	Diabetes mellitus (E10-E14)	662	4.3	92.7
5	Influenza and pneumonia (J10-J18)	575	3.7	80.5
6	Alzheimer's disease (G30)	472	3.0	66.1
7	Chronic lower respiratory diseases (J40-J47)	448	2.9	62.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	353	2.3	49.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	351	2.3	49.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	293	1.9	41.0
11	Septicemia (A40-A41)	228	1.5	31.9
12	Parkinson's disease (G20-G21)	123	0.8	17.2
13	Pneumonitis due to solids and liquids (J69)	104	0.7	14.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	103	0.7	14.4
15	Aortic aneurysm and dissection (I71)	102	0.7	14.3
...	All other causes (Residual)	2,456	15.8	344.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable