

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	139,241	100.0	296.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,951	20.8	61.7
2	Malignant neoplasms (C00-C97)	28,851	20.7	61.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,080	8.0	23.6
4	Cerebrovascular diseases (I60-I69)	7,121	5.1	15.2
5	Diabetes mellitus (E10-E14)	6,544	4.7	13.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,091	2.9	8.7
7	Chronic lower respiratory diseases (J40-J47)	3,949	2.8	8.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,331	2.4	7.1
9	Influenza and pneumonia (J09-J18)	3,176	2.3	6.8
10	Alzheimer's disease (G30)	3,005	2.2	6.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,903	2.1	6.2
12	Certain conditions originating in the perinatal period (P00-P96)	2,873	2.1	6.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,345	1.7	5.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,011	1.4	4.3
15	Septicemia (A40-A41)	2,004	1.4	4.3
...	All other causes (Residual)	27,006	19.4	57.5
Hispanic, both sexes, 1-4 years				
...	All causes	1,029	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	329	32.0	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	12.6	3.1
3	Malignant neoplasms (C00-C97)	105	10.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	7.7	1.9
5	Influenza and pneumonia (J09-J18)	31	3.0	0.7
6	Septicemia (A40-A41)	28	2.7	0.7
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	2.6	0.6
8	Certain conditions originating in the perinatal period (P00-P96)	17	1.7	*
9	Cerebrovascular diseases (I60-I69)	15	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
12	Anemias (D50-D64)	4	0.4	*
13	Meningitis (G00,G03)	3	0.3	*
13	Pneumonitis due to solids and liquids (J69)	3	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	240	23.3	5.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	480	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	133	27.7	3.0
2	Malignant neoplasms (C00-C97)	94	19.6	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	8.3	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	5.2	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	5.0	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	2.1	*
6	Influenza and pneumonia (J09-J18)	10	2.1	*
8	Cerebrovascular diseases (I60-I69)	8	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.5	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.3	*
11	Septicemia (A40-A41)	3	0.6	*
11	Anemias (D50-D64)	3	0.6	*
13	Meningococcal infection (A39)	2	0.4	*
13	Meningitis (G00,G03)	2	0.4	*
13	Diseases of appendix (K35-K38)	2	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
...	All other causes (Residual)	109	22.7	2.4
Hispanic, both sexes, 10-14 years				
...	All causes	540	100.0	13.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	31.7	4.3
2	Malignant neoplasms (C00-C97)	92	17.0	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	7.2	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	5.2	0.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	3.9	0.5
6	Cerebrovascular diseases (I60-I69)	18	3.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.4	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.7	*
9	Septicemia (A40-A41)	8	1.5	*
10	Diabetes mellitus (E10-E14)	6	1.1	*
10	Influenza and pneumonia (J09-J18)	6	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
13	Anemias (D50-D64)	3	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
15	Tuberculosis (A16-A19)	2	0.4	*
15	Meningitis (G00,G03)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	113	20.9	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	2,035	100.0	52.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	801	39.4	20.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	504	24.8	13.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	203	10.0	5.3
4	Malignant neoplasms (C00-C97)	165	8.1	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	2.0	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	1.7	0.9
7	Cerebrovascular diseases (I60-I69)	16	0.8	*
8	Influenza and pneumonia (J09-J18)	14	0.7	*
9	Septicemia (A40-A41)	9	0.4	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	9	0.4	*
11	Anemias (D50-D64)	8	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
13	Diabetes mellitus (E10-E14)	6	0.3	*
13	Legal intervention (Y35,Y89.0)	6	0.3	*
...	All other causes (Residual)	205	10.1	5.3
Hispanic, both sexes, 20-24 years				
...	All causes	3,168	100.0	86.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,351	42.6	36.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	713	22.5	19.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	308	9.7	8.4
4	Malignant neoplasms (C00-C97)	185	5.8	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	2.7	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	1.0	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	30	0.9	0.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	29	0.9	0.8
9	Cerebrovascular diseases (I60-I69)	17	0.5	*
10	Diabetes mellitus (E10-E14)	15	0.5	*
10	Influenza and pneumonia (J09-J18)	15	0.5	*
10	Legal intervention (Y35,Y89.0)	15	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	12	0.4	*
14	Septicemia (A40-A41)	10	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.3	*
...	All other causes (Residual)	342	10.8	9.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	3,218	100.0	77.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,226	38.1	29.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	561	17.4	13.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	320	9.9	7.7
4	Malignant neoplasms (C00-C97)	262	8.1	6.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	146	4.5	3.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	65	2.0	1.6
7	Cerebrovascular diseases (I60-I69)	44	1.4	1.1
8	Pregnancy, childbirth and the puerperium (O00-O99)	38	1.2	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	1.1	0.8
10	Diabetes mellitus (E10-E14)	22	0.7	0.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	22	0.7	0.5
12	Influenza and pneumonia (J09-J18)	20	0.6	0.5
13	Septicemia (A40-A41)	17	0.5	*
14	Legal intervention (Y35,Y89.0)	16	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.4	*
...	All other causes (Residual)	411	12.8	9.9
Hispanic, both sexes, 30-34 years				
...	All causes	3,178	100.0	78.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,034	32.5	25.6
2	Malignant neoplasms (C00-C97)	381	12.0	9.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	380	12.0	9.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	231	7.3	5.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	216	6.8	5.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	110	3.5	2.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	80	2.5	2.0
8	Cerebrovascular diseases (I60-I69)	73	2.3	1.8
9	Diabetes mellitus (E10-E14)	40	1.3	1.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	36	1.1	0.9
11	Influenza and pneumonia (J09-J18)	35	1.1	0.9
12	Septicemia (A40-A41)	20	0.6	0.5
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.6	0.5
14	Chronic lower respiratory diseases (J40-J47)	19	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.6	*
...	All other causes (Residual)	484	15.2	12.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,672	100.0	98.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	881	24.0	23.6
2	Malignant neoplasms (C00-C97)	559	15.2	15.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	350	9.5	9.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	305	8.3	8.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	246	6.7	6.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	151	4.1	4.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	150	4.1	4.0
8	Cerebrovascular diseases (I60-I69)	110	3.0	2.9
9	Diabetes mellitus (E10-E14)	79	2.2	2.1
10	Influenza and pneumonia (J09-J18)	48	1.3	1.3
11	Septicemia (A40-A41)	37	1.0	1.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.9	0.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	0.9	0.9
14	Chronic lower respiratory diseases (J40-J47)	25	0.7	0.7
15	Viral hepatitis (B15-B19)	23	0.6	0.6
15	Pregnancy, childbirth and the puerperium (O00-O99)	23	0.6	0.6
...	All other causes (Residual)	619	16.9	16.6
Hispanic, both sexes, 40-44 years				
...	All causes	4,949	100.0	150.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	952	19.2	29.0
2	Malignant neoplasms (C00-C97)	941	19.0	28.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	535	10.8	16.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	338	6.8	10.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	255	5.2	7.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	253	5.1	7.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	230	4.6	7.0
8	Cerebrovascular diseases (I60-I69)	191	3.9	5.8
9	Diabetes mellitus (E10-E14)	146	3.0	4.5
10	Viral hepatitis (B15-B19)	86	1.7	2.6
11	Septicemia (A40-A41)	56	1.1	1.7
12	Influenza and pneumonia (J09-J18)	54	1.1	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.9	1.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	0.7	1.0
15	Chronic lower respiratory diseases (J40-J47)	31	0.6	0.9
...	All other causes (Residual)	803	16.2	24.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	6,548	100.0	234.3
1	Malignant neoplasms (C00-C97)	1,531	23.4	54.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	939	14.3	33.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	893	13.6	31.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	512	7.8	18.3
5	Cerebrovascular diseases (I60-I69)	280	4.3	10.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	275	4.2	9.8
7	Diabetes mellitus (E10-E14)	232	3.5	8.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	209	3.2	7.5
9	Viral hepatitis (B15-B19)	168	2.6	6.0
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	2.5	5.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	1.5	3.4
12	Septicemia (A40-A41)	77	1.2	2.8
13	Influenza and pneumonia (J09-J18)	74	1.1	2.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	0.7	1.7
15	Chronic lower respiratory diseases (J40-J47)	44	0.7	1.6
...	All other causes (Residual)	1,006	15.4	36.0
Hispanic, both sexes, 50-54 years				
...	All causes	7,822	100.0	357.6
1	Malignant neoplasms (C00-C97)	2,019	25.8	92.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,372	17.5	62.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	670	8.6	30.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	648	8.3	29.6
5	Diabetes mellitus (E10-E14)	369	4.7	16.9
6	Cerebrovascular diseases (I60-I69)	358	4.6	16.4
7	Viral hepatitis (B15-B19)	244	3.1	11.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	192	2.5	8.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	176	2.3	8.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	113	1.4	5.2
11	Influenza and pneumonia (J09-J18)	111	1.4	5.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	103	1.3	4.7
13	Septicemia (A40-A41)	102	1.3	4.7
14	Chronic lower respiratory diseases (J40-J47)	95	1.2	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	64	0.8	2.9
...	All other causes (Residual)	1,186	15.2	54.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	9,012	100.0	546.0
1	Malignant neoplasms (C00-C97)	2,736	30.4	165.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,788	19.8	108.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	552	6.1	33.4
4	Diabetes mellitus (E10-E14)	530	5.9	32.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	525	5.8	31.8
6	Cerebrovascular diseases (I60-I69)	403	4.5	24.4
7	Viral hepatitis (B15-B19)	258	2.9	15.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	212	2.4	12.8
9	Septicemia (A40-A41)	134	1.5	8.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	125	1.4	7.6
11	Chronic lower respiratory diseases (J40-J47)	123	1.4	7.5
12	Influenza and pneumonia (J09-J18)	117	1.3	7.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	111	1.2	6.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	95	1.1	5.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	0.8	4.3
...	All other causes (Residual)	1,232	13.7	74.6
Hispanic, both sexes, 60-64 years				
...	All causes	9,490	100.0	788.5
1	Malignant neoplasms (C00-C97)	3,038	32.0	252.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,079	21.9	172.7
3	Diabetes mellitus (E10-E14)	664	7.0	55.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	474	5.0	39.4
5	Cerebrovascular diseases (I60-I69)	392	4.1	32.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	3.6	28.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	262	2.8	21.8
8	Chronic lower respiratory diseases (J40-J47)	229	2.4	19.0
9	Influenza and pneumonia (J09-J18)	173	1.8	14.4
10	Viral hepatitis (B15-B19)	150	1.6	12.5
11	Septicemia (A40-A41)	130	1.4	10.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	100	1.1	8.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	75	0.8	6.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	0.7	5.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.4	3.0
...	All other causes (Residual)	1,287	13.6	106.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	10,117	100.0	1,186.0
1	Malignant neoplasms (C00-C97)	3,235	32.0	379.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,254	22.3	264.2
3	Diabetes mellitus (E10-E14)	738	7.3	86.5
4	Cerebrovascular diseases (I60-I69)	512	5.1	60.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	388	3.8	45.5
6	Chronic lower respiratory diseases (J40-J47)	320	3.2	37.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	291	2.9	34.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	254	2.5	29.8
9	Influenza and pneumonia (J09-J18)	195	1.9	22.9
10	Septicemia (A40-A41)	172	1.7	20.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	124	1.2	14.5
12	Viral hepatitis (B15-B19)	78	0.8	9.1
13	Alzheimer's disease (G30)	72	0.7	8.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.6	7.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	0.5	6.0
...	All other causes (Residual)	1,373	13.6	161.0
Hispanic, both sexes, 70-74 years				
...	All causes	11,826	100.0	1,812.1
1	Malignant neoplasms (C00-C97)	3,451	29.2	528.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,849	24.1	436.6
3	Diabetes mellitus (E10-E14)	847	7.2	129.8
4	Cerebrovascular diseases (I60-I69)	647	5.5	99.1
5	Chronic lower respiratory diseases (J40-J47)	417	3.5	63.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	351	3.0	53.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	327	2.8	50.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	267	2.3	40.9
9	Influenza and pneumonia (J09-J18)	253	2.1	38.8
10	Septicemia (A40-A41)	220	1.9	33.7
11	Alzheimer's disease (G30)	161	1.4	24.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	127	1.1	19.5
13	Parkinson's disease (G20-G21)	96	0.8	14.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.6	11.2
15	Pneumonitis due to solids and liquids (J69)	70	0.6	10.7
...	All other causes (Residual)	1,670	14.1	255.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	14,455	100.0	2,912.1
1	Malignant neoplasms (C00-C97)	3,718	25.7	749.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,560	24.6	717.2
3	Cerebrovascular diseases (I60-I69)	950	6.6	191.4
4	Diabetes mellitus (E10-E14)	930	6.4	187.4
5	Chronic lower respiratory diseases (J40-J47)	650	4.5	131.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	402	2.8	81.0
7	Influenza and pneumonia (J09-J18)	393	2.7	79.2
8	Alzheimer's disease (G30)	342	2.4	68.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	306	2.1	61.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	282	2.0	56.8
11	Septicemia (A40-A41)	231	1.6	46.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	187	1.3	37.7
13	Parkinson's disease (G20-G21)	156	1.1	31.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.6	16.5
15	Pneumonitis due to solids and liquids (J69)	71	0.5	14.3
...	All other causes (Residual)	2,195	15.2	442.2
Hispanic, both sexes, 80-84 years				
...	All causes	15,807	100.0	4,567.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,209	26.6	1,216.2
2	Malignant neoplasms (C00-C97)	3,235	20.5	934.8
3	Cerebrovascular diseases (I60-I69)	1,183	7.5	341.8
4	Diabetes mellitus (E10-E14)	884	5.6	255.4
5	Chronic lower respiratory diseases (J40-J47)	760	4.8	219.6
6	Alzheimer's disease (G30)	650	4.1	187.8
7	Influenza and pneumonia (J09-J18)	515	3.3	148.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	433	2.7	125.1
9	Septicemia (A40-A41)	292	1.8	84.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	292	1.8	84.4
11	Parkinson's disease (G20-G21)	238	1.5	68.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	225	1.4	65.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	150	0.9	43.3
14	Pneumonitis due to solids and liquids (J69)	123	0.8	35.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	96	0.6	27.7
...	All other causes (Residual)	2,522	16.0	728.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	13,700	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,068	29.7	---
2	Malignant neoplasms (C00-C97)	2,037	14.9	---
3	Cerebrovascular diseases (I60-I69)	1,023	7.5	---
4	Alzheimer's disease (G30)	829	6.1	---
5	Chronic lower respiratory diseases (J40-J47)	684	5.0	---
6	Diabetes mellitus (E10-E14)	632	4.6	---
7	Influenza and pneumonia (J09-J18)	488	3.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	344	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	269	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	236	1.7	---
11	Septicemia (A40-A41)	220	1.6	---
12	Parkinson's disease (G20-G21)	176	1.3	---
13	Pneumonitis due to solids and liquids (J69)	110	0.8	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	93	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.5	---
...	All other causes (Residual)	2,425	17.7	---
Hispanic, both sexes, 90-94 years				
...	All causes	7,961	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,694	33.8	---
2	Malignant neoplasms (C00-C97)	776	9.7	---
3	Alzheimer's disease (G30)	585	7.3	---
4	Cerebrovascular diseases (I60-I69)	567	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	353	4.4	---
6	Influenza and pneumonia (J09-J18)	336	4.2	---
7	Diabetes mellitus (E10-E14)	284	3.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	158	2.0	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	1.8	---
11	Septicemia (A40-A41)	108	1.4	---
12	Pneumonitis due to solids and liquids (J69)	90	1.1	---
13	Parkinson's disease (G20-G21)	83	1.0	---
14	Atherosclerosis (I70)	37	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.4	---
...	All other causes (Residual)	1,532	19.2	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	3,572	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,312	36.7	---
2	Alzheimer's disease (G30)	271	7.6	---
3	Cerebrovascular diseases (I60-I69)	243	6.8	---
4	Malignant neoplasms (C00-C97)	240	6.7	---
5	Influenza and pneumonia (J09-J18)	179	5.0	---
6	Chronic lower respiratory diseases (J40-J47)	113	3.2	---
7	Diabetes mellitus (E10-E14)	104	2.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	77	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	1.7	---
11	Septicemia (A40-A41)	46	1.3	---
12	Pneumonitis due to solids and liquids (J69)	39	1.1	---
13	Atherosclerosis (I70)	29	0.8	---
14	Parkinson's disease (G20-G21)	19	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.5	---
...	All other causes (Residual)	745	20.9	---
Hispanic, both sexes, 100 years and over				
...	All causes	763	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	296	38.8	---
2	Influenza and pneumonia (J09-J18)	63	8.3	---
3	Alzheimer's disease (G30)	48	6.3	---
4	Cerebrovascular diseases (I60-I69)	44	5.8	---
5	Malignant neoplasms (C00-C97)	33	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	27	3.5	---
7	Septicemia (A40-A41)	15	2.0	---
7	Diabetes mellitus (E10-E14)	15	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.4	---
12	Atherosclerosis (I70)	7	0.9	---
12	Pneumonitis due to solids and liquids (J69)	7	0.9	---
14	Parkinson's disease (G20-G21)	6	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	---
...	All other causes (Residual)	160	21.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	76,861	100.0	316.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,498	20.2	63.9
2	Malignant neoplasms (C00-C97)	15,283	19.9	63.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,363	10.9	34.5
4	Cerebrovascular diseases (I60-I69)	3,370	4.4	13.9
5	Diabetes mellitus (E10-E14)	3,314	4.3	13.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,850	3.7	11.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,777	3.6	11.4
8	Chronic lower respiratory diseases (J40-J47)	2,019	2.6	8.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,955	2.5	8.1
10	Certain conditions originating in the perinatal period (P00-P96)	1,645	2.1	6.8
11	Influenza and pneumonia (J09-J18)	1,544	2.0	6.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,456	1.9	6.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,061	1.4	4.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,037	1.3	4.3
15	Septicemia (A40-A41)	973	1.3	4.0
...	All other causes (Residual)	13,716	17.8	56.6
Hispanic, male, 1-4 years				
...	All causes	587	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	36.1	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	11.1	3.0
3	Malignant neoplasms (C00-C97)	52	8.9	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	6.0	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.1	*
6	Septicemia (A40-A41)	14	2.4	*
6	Influenza and pneumonia (J09-J18)	14	2.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	10	1.7	*
9	Cerebrovascular diseases (I60-I69)	7	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
12	Anemias (D50-D64)	2	0.3	*
12	Meningitis (G00,G03)	2	0.3	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	145	24.7	6.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	253	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	31.2	3.5
2	Malignant neoplasms (C00-C97)	49	19.4	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	7.9	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	5.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	4.0	*
6	Influenza and pneumonia (J09-J18)	6	2.4	*
7	Cerebrovascular diseases (I60-I69)	4	1.6	*
8	Septicemia (A40-A41)	3	1.2	*
8	Anemias (D50-D64)	3	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.2	*
11	Meningococcal infection (A39)	2	0.8	*
11	Diseases of appendix (K35-K38)	2	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	54	21.3	2.4
Hispanic, male, 10-14 years				
...	All causes	295	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	37.6	5.4
2	Malignant neoplasms (C00-C97)	46	15.6	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	8.8	1.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	4.1	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	3.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.4	*
6	Cerebrovascular diseases (I60-I69)	7	2.4	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.0	*
9	Diabetes mellitus (E10-E14)	2	0.7	*
10	Salmonella infections (A01-A02)	1	0.3	*
10	Tuberculosis (A16-A19)	1	0.3	*
10	Septicemia (A40-A41)	1	0.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
10	Anemias (D50-D64)	1	0.3	*
10	Influenza and pneumonia (J09-J18)	1	0.3	*
10	Diseases of appendix (K35-K38)	1	0.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	63	21.4	3.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,519	100.0	76.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	608	40.0	30.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	441	29.0	22.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	10.2	7.8
4	Malignant neoplasms (C00-C97)	98	6.5	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	2.0	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	1.4	1.1
7	Cerebrovascular diseases (I60-I69)	9	0.6	*
8	Influenza and pneumonia (J09-J18)	8	0.5	*
9	Chronic lower respiratory diseases (J40-J47)	6	0.4	*
9	Legal intervention (Y35,Y89.0)	6	0.4	*
11	Septicemia (A40-A41)	5	0.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.3	*
13	Anemias (D50-D64)	3	0.2	*
13	Diabetes mellitus (E10-E14)	3	0.2	*
15	Aortic aneurysm and dissection (I71)	2	0.1	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.1	*
...	All other causes (Residual)	117	7.7	5.9
Hispanic, male, 20-24 years				
...	All causes	2,560	100.0	131.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,132	44.2	58.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	646	25.2	33.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	269	10.5	13.8
4	Malignant neoplasms (C00-C97)	119	4.6	6.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	2.5	3.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.7	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.7	*
8	Legal intervention (Y35,Y89.0)	15	0.6	*
9	Diabetes mellitus (E10-E14)	10	0.4	*
10	Cerebrovascular diseases (I60-I69)	8	0.3	*
10	Influenza and pneumonia (J09-J18)	8	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.2	*
14	Anemias (D50-D64)	5	0.2	*
14	Aortic aneurysm and dissection (I71)	5	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
...	All other causes (Residual)	223	8.7	11.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,483	100.0	108.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,027	41.4	44.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	488	19.7	21.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	271	10.9	11.8
4	Malignant neoplasms (C00-C97)	154	6.2	6.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	4.1	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	53	2.1	2.3
7	Cerebrovascular diseases (I60-I69)	30	1.2	1.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	19	0.8	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.7	*
10	Legal intervention (Y35,Y89.0)	16	0.6	*
11	Diabetes mellitus (E10-E14)	12	0.5	*
12	Influenza and pneumonia (J09-J18)	9	0.4	*
13	Septicemia (A40-A41)	8	0.3	*
14	Aortic aneurysm and dissection (I71)	7	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
...	All other causes (Residual)	262	10.6	11.4
Hispanic, male, 30-34 years				
...	All causes	2,356	100.0	106.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	876	37.2	39.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	320	13.6	14.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	197	8.4	8.9
4	Malignant neoplasms (C00-C97)	190	8.1	8.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	161	6.8	7.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	81	3.4	3.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	66	2.8	3.0
8	Cerebrovascular diseases (I60-I69)	51	2.2	2.3
9	Diabetes mellitus (E10-E14)	25	1.1	1.1
10	Influenza and pneumonia (J09-J18)	19	0.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.6	*
13	Septicemia (A40-A41)	10	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	10	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
15	Legal intervention (Y35,Y89.0)	8	0.3	*
...	All other causes (Residual)	307	13.0	13.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,571	100.0	128.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	733	28.5	36.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	264	10.3	13.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	255	9.9	12.8
4	Malignant neoplasms (C00-C97)	222	8.6	11.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	202	7.9	10.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	122	4.7	6.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	117	4.6	5.9
8	Cerebrovascular diseases (I60-I69)	60	2.3	3.0
9	Diabetes mellitus (E10-E14)	56	2.2	2.8
10	Influenza and pneumonia (J09-J18)	32	1.2	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.9	1.1
12	Septicemia (A40-A41)	21	0.8	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.7	*
14	Viral hepatitis (B15-B19)	18	0.7	*
15	Chronic lower respiratory diseases (J40-J47)	15	0.6	*
...	All other causes (Residual)	413	16.1	20.7
Hispanic, male, 40-44 years				
...	All causes	3,353	100.0	193.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	761	22.7	43.9
2	Malignant neoplasms (C00-C97)	428	12.8	24.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	408	12.2	23.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	253	7.5	14.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	217	6.5	12.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	5.6	10.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	185	5.5	10.7
8	Cerebrovascular diseases (I60-I69)	128	3.8	7.4
9	Diabetes mellitus (E10-E14)	96	2.9	5.5
10	Viral hepatitis (B15-B19)	60	1.8	3.5
11	Septicemia (A40-A41)	36	1.1	2.1
12	Influenza and pneumonia (J09-J18)	33	1.0	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.9	1.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	0.6	1.2
15	Chronic lower respiratory diseases (J40-J47)	14	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.4	*
...	All other causes (Residual)	481	14.3	27.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,353	100.0	302.0
1	Malignant neoplasms (C00-C97)	746	17.1	51.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	720	16.5	50.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	688	15.8	47.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	401	9.2	27.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	219	5.0	15.2
6	Cerebrovascular diseases (I60-I69)	172	4.0	11.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	158	3.6	11.0
8	Diabetes mellitus (E10-E14)	148	3.4	10.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	127	2.9	8.8
10	Viral hepatitis (B15-B19)	124	2.8	8.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	1.4	4.2
12	Septicemia (A40-A41)	44	1.0	3.1
13	Influenza and pneumonia (J09-J18)	43	1.0	3.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	0.8	2.3
15	Chronic lower respiratory diseases (J40-J47)	28	0.6	1.9
...	All other causes (Residual)	641	14.7	44.5
Hispanic, male, 50-54 years				
...	All causes	5,116	100.0	464.8
1	Malignant neoplasms (C00-C97)	1,053	20.6	95.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	997	19.5	90.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	529	10.3	48.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	506	9.9	46.0
5	Diabetes mellitus (E10-E14)	228	4.5	20.7
6	Cerebrovascular diseases (I60-I69)	214	4.2	19.4
7	Viral hepatitis (B15-B19)	194	3.8	17.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	2.9	13.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	144	2.8	13.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	1.6	7.4
11	Influenza and pneumonia (J09-J18)	71	1.4	6.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	1.3	5.8
13	Septicemia (A40-A41)	60	1.2	5.5
14	Chronic lower respiratory diseases (J40-J47)	57	1.1	5.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	0.8	3.7
...	All other causes (Residual)	729	14.2	66.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	5,731	100.0	709.9
1	Malignant neoplasms (C00-C97)	1,449	25.3	179.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,299	22.7	160.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	436	7.6	54.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	422	7.4	52.3
5	Diabetes mellitus (E10-E14)	313	5.5	38.8
6	Cerebrovascular diseases (I60-I69)	245	4.3	30.3
7	Viral hepatitis (B15-B19)	185	3.2	22.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	114	2.0	14.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	102	1.8	12.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	1.6	11.4
11	Chronic lower respiratory diseases (J40-J47)	78	1.4	9.7
12	Septicemia (A40-A41)	77	1.3	9.5
13	Influenza and pneumonia (J09-J18)	72	1.3	8.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	1.0	6.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	0.9	6.4
...	All other causes (Residual)	739	12.9	91.5
Hispanic, male, 60-64 years				
...	All causes	5,730	100.0	1,003.0
1	Malignant neoplasms (C00-C97)	1,684	29.4	294.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,393	24.3	243.8
3	Diabetes mellitus (E10-E14)	370	6.5	64.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	346	6.0	60.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	251	4.4	43.9
6	Cerebrovascular diseases (I60-I69)	238	4.2	41.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	2.4	23.6
8	Chronic lower respiratory diseases (J40-J47)	125	2.2	21.9
9	Influenza and pneumonia (J09-J18)	102	1.8	17.9
10	Viral hepatitis (B15-B19)	91	1.6	15.9
11	Septicemia (A40-A41)	69	1.2	12.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	53	0.9	9.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	0.9	9.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	0.8	7.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	0.5	4.6
...	All other causes (Residual)	749	13.1	131.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	5,895	100.0	1,506.5
1	Malignant neoplasms (C00-C97)	1,851	31.4	473.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,432	24.3	365.9
3	Diabetes mellitus (E10-E14)	408	6.9	104.3
4	Cerebrovascular diseases (I60-I69)	264	4.5	67.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	246	4.2	62.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	3.0	44.5
7	Chronic lower respiratory diseases (J40-J47)	172	2.9	44.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	157	2.7	40.1
9	Influenza and pneumonia (J09-J18)	115	2.0	29.4
10	Septicemia (A40-A41)	96	1.6	24.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	71	1.2	18.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	0.7	11.2
13	Alzheimer's disease (G30)	39	0.7	10.0
14	Viral hepatitis (B15-B19)	38	0.6	9.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.6	8.4
...	All other causes (Residual)	755	12.8	192.9
Hispanic, male, 70-74 years				
...	All causes	6,526	100.0	2,261.4
1	Malignant neoplasms (C00-C97)	1,936	29.7	670.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,642	25.2	569.0
3	Diabetes mellitus (E10-E14)	423	6.5	146.6
4	Cerebrovascular diseases (I60-I69)	343	5.3	118.9
5	Chronic lower respiratory diseases (J40-J47)	233	3.6	80.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	197	3.0	68.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	2.5	56.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	155	2.4	53.7
9	Influenza and pneumonia (J09-J18)	138	2.1	47.8
10	Septicemia (A40-A41)	124	1.9	43.0
11	Alzheimer's disease (G30)	79	1.2	27.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	0.9	20.8
13	Parkinson's disease (G20-G21)	59	0.9	20.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	0.8	17.3
15	Pneumonitis due to solids and liquids (J69)	46	0.7	15.9
...	All other causes (Residual)	879	13.5	304.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	7,390	100.0	3,524.8
1	Malignant neoplasms (C00-C97)	1,993	27.0	950.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,912	25.9	912.0
3	Cerebrovascular diseases (I60-I69)	438	5.9	208.9
4	Diabetes mellitus (E10-E14)	435	5.9	207.5
5	Chronic lower respiratory diseases (J40-J47)	361	4.9	172.2
6	Influenza and pneumonia (J09-J18)	212	2.9	101.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	200	2.7	95.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	2.2	79.2
9	Alzheimer's disease (G30)	143	1.9	68.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	125	1.7	59.6
11	Septicemia (A40-A41)	104	1.4	49.6
12	Parkinson's disease (G20-G21)	92	1.2	43.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	86	1.2	41.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.6	21.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	0.6	20.0
...	All other causes (Residual)	1,035	14.0	493.7
Hispanic, male, 80-84 years				
...	All causes	7,461	100.0	5,381.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,058	27.6	1,484.5
2	Malignant neoplasms (C00-C97)	1,709	22.9	1,232.8
3	Cerebrovascular diseases (I60-I69)	513	6.9	370.0
4	Diabetes mellitus (E10-E14)	408	5.5	294.3
5	Chronic lower respiratory diseases (J40-J47)	377	5.1	271.9
6	Influenza and pneumonia (J09-J18)	238	3.2	171.7
7	Alzheimer's disease (G30)	236	3.2	170.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	210	2.8	151.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	1.9	103.2
10	Parkinson's disease (G20-G21)	137	1.8	98.8
11	Septicemia (A40-A41)	117	1.6	84.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	1.2	64.9
13	Pneumonitis due to solids and liquids (J69)	64	0.9	46.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	55	0.7	39.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.7	36.8
...	All other causes (Residual)	1,055	14.1	761.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	5,605	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,652	29.5	---
2	Malignant neoplasms (C00-C97)	1,014	18.1	---
3	Cerebrovascular diseases (I60-I69)	392	7.0	---
4	Chronic lower respiratory diseases (J40-J47)	318	5.7	---
5	Diabetes mellitus (E10-E14)	255	4.5	---
6	Alzheimer's disease (G30)	244	4.4	---
7	Influenza and pneumonia (J09-J18)	211	3.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	158	2.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	1.9	---
10	Septicemia (A40-A41)	98	1.7	---
11	Parkinson's disease (G20-G21)	91	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	1.4	---
13	Pneumonitis due to solids and liquids (J69)	51	0.9	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	43	0.8	---
15	Aortic aneurysm and dissection (I71)	33	0.6	---
...	All other causes (Residual)	856	15.3	---
Hispanic, male, 90-94 years				
...	All causes	2,670	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	888	33.3	---
2	Malignant neoplasms (C00-C97)	363	13.6	---
3	Cerebrovascular diseases (I60-I69)	168	6.3	---
4	Chronic lower respiratory diseases (J40-J47)	153	5.7	---
5	Alzheimer's disease (G30)	145	5.4	---
6	Influenza and pneumonia (J09-J18)	113	4.2	---
7	Diabetes mellitus (E10-E14)	88	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	2.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	47	1.8	---
11	Parkinson's disease (G20-G21)	45	1.7	---
12	Pneumonitis due to solids and liquids (J69)	36	1.3	---
13	Septicemia (A40-A41)	32	1.2	---
14	Atherosclerosis (I70)	11	0.4	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	10	0.4	---
...	All other causes (Residual)	430	16.1	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	938	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	320	34.1	---
2	Malignant neoplasms (C00-C97)	102	10.9	---
3	Alzheimer's disease (G30)	57	6.1	---
3	Cerebrovascular diseases (I60-I69)	57	6.1	---
5	Influenza and pneumonia (J09-J18)	55	5.9	---
6	Chronic lower respiratory diseases (J40-J47)	33	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	3.3	---
8	Diabetes mellitus (E10-E14)	30	3.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.7	---
11	Pneumonitis due to solids and liquids (J69)	10	1.1	---
12	Septicemia (A40-A41)	9	1.0	---
13	Parkinson's disease (G20-G21)	8	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	---
14	Aortic aneurysm and dissection (I71)	5	0.5	---
...	All other causes (Residual)	183	19.5	---
Hispanic, male, 100 years and over				
...	All causes	181	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	37.6	---
2	Influenza and pneumonia (J09-J18)	21	11.6	---
3	Malignant neoplasms (C00-C97)	14	7.7	---
4	Chronic lower respiratory diseases (J40-J47)	12	6.6	---
5	Septicemia (A40-A41)	5	2.8	---
5	Cerebrovascular diseases (I60-I69)	5	2.8	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	2.8	---
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	2.8	---
9	Diabetes mellitus (E10-E14)	4	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	1.7	---
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.1	---
11	Pneumonitis due to solids and liquids (J69)	2	1.1	---
13	Nutritional deficiencies (E40-E64)	1	0.6	---
13	Parkinson's disease (G20-G21)	1	0.6	---
13	Alzheimer's disease (G30)	1	0.6	---
13	Atherosclerosis (I70)	1	0.6	---
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.6	---
...	All other causes (Residual)	30	16.6	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	62,380	100.0	274.9
1	Malignant neoplasms (C00-C97)	13,568	21.8	59.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,453	21.6	59.3
3	Cerebrovascular diseases (I60-I69)	3,751	6.0	16.5
4	Diabetes mellitus (E10-E14)	3,230	5.2	14.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,717	4.4	12.0
6	Alzheimer's disease (G30)	2,039	3.3	9.0
7	Chronic lower respiratory diseases (J40-J47)	1,930	3.1	8.5
8	Influenza and pneumonia (J09-J18)	1,632	2.6	7.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,447	2.3	6.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,241	2.0	5.5
11	Certain conditions originating in the perinatal period (P00-P96)	1,228	2.0	5.4
12	Septicemia (A40-A41)	1,031	1.7	4.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	950	1.5	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	856	1.4	3.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	554	0.9	2.4
...	All other causes (Residual)	12,753	20.4	56.2
Hispanic, female, 1-4 years				
...	All causes	442	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	26.5	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	14.7	3.2
3	Malignant neoplasms (C00-C97)	53	12.0	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	10.0	2.2
5	Influenza and pneumonia (J09-J18)	17	3.8	*
6	Septicemia (A40-A41)	14	3.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.0	*
8	Cerebrovascular diseases (I60-I69)	8	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	7	1.6	*
11	Anemias (D50-D64)	2	0.5	*
11	Pneumonitis due to solids and liquids (J69)	2	0.5	*
13	Tuberculosis (A16-A19)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
13	Peptic ulcer (K25-K28)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	89	20.1	4.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	227	100.0	10.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	23.8	2.5
2	Malignant neoplasms (C00-C97)	45	19.8	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	8.8	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	6.6	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	4.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	4.0	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	5	2.2	*
8	Cerebrovascular diseases (I60-I69)	4	1.8	*
8	Influenza and pneumonia (J09-J18)	4	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.8	*
11	Meningitis (G00,G03)	2	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.9	*
13	Viral hepatitis (B15-B19)	1	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.4	*
13	Peptic ulcer (K25-K28)	1	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	50	22.0	2.3
Hispanic, female, 10-14 years				
...	All causes	245	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	24.5	3.1
2	Malignant neoplasms (C00-C97)	46	18.8	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	7.8	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	5.3	*
5	Cerebrovascular diseases (I60-I69)	11	4.5	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	3.7	*
7	Septicemia (A40-A41)	7	2.9	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.4	*
8	Chronic lower respiratory diseases (J40-J47)	6	2.4	*
10	Influenza and pneumonia (J09-J18)	5	2.0	*
11	Diabetes mellitus (E10-E14)	4	1.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	1.2	*
14	Anemias (D50-D64)	2	0.8	*
14	Meningitis (G00,G03)	2	0.8	*
...	All other causes (Residual)	49	20.0	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	516	100.0	27.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	37.4	10.3
2	Malignant neoplasms (C00-C97)	67	13.0	3.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	12.2	3.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	9.3	2.6
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	2.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.1	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	9	1.7	*
8	Cerebrovascular diseases (I60-I69)	7	1.4	*
9	Influenza and pneumonia (J09-J18)	6	1.2	*
10	Anemias (D50-D64)	5	1.0	*
11	Septicemia (A40-A41)	4	0.8	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.6	*
12	Diabetes mellitus (E10-E14)	3	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	77	14.9	4.1
Hispanic, female, 20-24 years				
...	All causes	608	100.0	35.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	36.0	12.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	11.0	3.9
3	Malignant neoplasms (C00-C97)	66	10.9	3.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	6.4	2.3
5	Pregnancy, childbirth and the puerperium (O00-O99)	30	4.9	1.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.3	1.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	12	2.0	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	2.0	*
9	Cerebrovascular diseases (I60-I69)	9	1.5	*
10	Septicemia (A40-A41)	7	1.2	*
10	Influenza and pneumonia (J09-J18)	7	1.2	*
12	Diabetes mellitus (E10-E14)	5	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
13	Anemias (D50-D64)	4	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
...	All other causes (Residual)	95	15.6	5.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	735	100.0	39.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	27.1	10.8
2	Malignant neoplasms (C00-C97)	108	14.7	5.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	9.9	4.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	6.7	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	5.9	2.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	38	5.2	2.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	2.3	*
8	Cerebrovascular diseases (I60-I69)	14	1.9	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.6	*
10	Influenza and pneumonia (J09-J18)	11	1.5	*
11	Diabetes mellitus (E10-E14)	10	1.4	*
12	Septicemia (A40-A41)	9	1.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.0	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
15	Meningitis (G00,G03)	3	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.4	*
...	All other causes (Residual)	128	17.4	6.9
Hispanic, female, 30-34 years				
...	All causes	822	100.0	45.0
1	Malignant neoplasms (C00-C97)	191	23.2	10.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	19.2	8.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	7.3	3.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	6.7	3.0
5	Pregnancy, childbirth and the puerperium (O00-O99)	36	4.4	2.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	4.1	1.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	29	3.5	1.6
8	Cerebrovascular diseases (I60-I69)	22	2.7	1.2
9	Influenza and pneumonia (J09-J18)	16	1.9	*
10	Diabetes mellitus (E10-E14)	15	1.8	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.7	*
12	Septicemia (A40-A41)	10	1.2	*
13	Chronic lower respiratory diseases (J40-J47)	9	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.7	*
...	All other causes (Residual)	161	19.6	8.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,101	100.0	63.5
1	Malignant neoplasms (C00-C97)	337	30.6	19.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	13.4	8.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	7.8	5.0
4	Cerebrovascular diseases (I60-I69)	50	4.5	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	4.5	2.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	4.0	2.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	33	3.0	1.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	29	2.6	1.7
9	Diabetes mellitus (E10-E14)	23	2.1	1.3
9	Pregnancy, childbirth and the puerperium (O00-O99)	23	2.1	1.3
11	Septicemia (A40-A41)	16	1.5	*
11	Influenza and pneumonia (J09-J18)	16	1.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.9	*
14	Chronic lower respiratory diseases (J40-J47)	10	0.9	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.9	*
...	All other causes (Residual)	201	18.3	11.6
Hispanic, female, 40-44 years				
...	All causes	1,596	100.0	103.2
1	Malignant neoplasms (C00-C97)	513	32.1	33.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	12.0	12.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	8.0	8.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	85	5.3	5.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	70	4.4	4.5
6	Cerebrovascular diseases (I60-I69)	63	3.9	4.1
7	Diabetes mellitus (E10-E14)	50	3.1	3.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	2.7	2.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	2.3	2.3
10	Viral hepatitis (B15-B19)	26	1.6	1.7
11	Influenza and pneumonia (J09-J18)	21	1.3	1.4
12	Septicemia (A40-A41)	20	1.3	1.3
13	Chronic lower respiratory diseases (J40-J47)	17	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.9	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.9	*
...	All other causes (Residual)	306	19.2	19.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,195	100.0	162.1
1	Malignant neoplasms (C00-C97)	785	35.8	58.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	219	10.0	16.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	9.3	15.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	111	5.1	8.2
5	Cerebrovascular diseases (I60-I69)	108	4.9	8.0
6	Diabetes mellitus (E10-E14)	84	3.8	6.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	56	2.6	4.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	2.3	3.8
9	Viral hepatitis (B15-B19)	44	2.0	3.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	1.7	2.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.6	2.6
12	Septicemia (A40-A41)	33	1.5	2.4
13	Influenza and pneumonia (J09-J18)	31	1.4	2.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	16	0.7	*
...	All other causes (Residual)	363	16.5	26.8
Hispanic, female, 50-54 years				
...	All causes	2,706	100.0	249.0
1	Malignant neoplasms (C00-C97)	966	35.7	88.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	375	13.9	34.5
3	Cerebrovascular diseases (I60-I69)	144	5.3	13.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	5.2	13.1
5	Diabetes mellitus (E10-E14)	141	5.2	13.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	141	5.2	13.0
7	Viral hepatitis (B15-B19)	50	1.8	4.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	1.8	4.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	48	1.8	4.4
10	Septicemia (A40-A41)	42	1.6	3.9
11	Influenza and pneumonia (J09-J18)	40	1.5	3.7
12	Chronic lower respiratory diseases (J40-J47)	38	1.4	3.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	1.0	2.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	0.8	2.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	0.8	2.0
...	All other causes (Residual)	457	16.9	42.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	3,281	100.0	389.1
1	Malignant neoplasms (C00-C97)	1,287	39.2	152.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	489	14.9	58.0
3	Diabetes mellitus (E10-E14)	217	6.6	25.7
4	Cerebrovascular diseases (I60-I69)	158	4.8	18.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	116	3.5	13.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	3.1	12.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	3.0	11.6
8	Viral hepatitis (B15-B19)	73	2.2	8.7
9	Septicemia (A40-A41)	57	1.7	6.8
10	Influenza and pneumonia (J09-J18)	45	1.4	5.3
10	Chronic lower respiratory diseases (J40-J47)	45	1.4	5.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	1.3	5.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.7	2.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	0.6	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	0.5	*
...	All other causes (Residual)	493	15.0	58.5
Hispanic, female, 60-64 years				
...	All causes	3,760	100.0	594.7
1	Malignant neoplasms (C00-C97)	1,354	36.0	214.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	686	18.2	108.5
3	Diabetes mellitus (E10-E14)	294	7.8	46.5
4	Cerebrovascular diseases (I60-I69)	154	4.1	24.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	128	3.4	20.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	3.4	20.1
7	Chronic lower respiratory diseases (J40-J47)	104	2.8	16.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	2.3	13.8
9	Influenza and pneumonia (J09-J18)	71	1.9	11.2
10	Septicemia (A40-A41)	61	1.6	9.6
11	Viral hepatitis (B15-B19)	59	1.6	9.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	55	1.5	8.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.6	3.5
14	Alzheimer's disease (G30)	19	0.5	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	14	0.4	*
...	All other causes (Residual)	525	14.0	83.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	4,222	100.0	914.4
1	Malignant neoplasms (C00-C97)	1,384	32.8	299.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	822	19.5	178.0
3	Diabetes mellitus (E10-E14)	330	7.8	71.5
4	Cerebrovascular diseases (I60-I69)	248	5.9	53.7
5	Chronic lower respiratory diseases (J40-J47)	148	3.5	32.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	142	3.4	30.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	134	3.2	29.0
8	Influenza and pneumonia (J09-J18)	80	1.9	17.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	1.9	17.3
10	Septicemia (A40-A41)	76	1.8	16.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	53	1.3	11.5
12	Viral hepatitis (B15-B19)	40	0.9	8.7
13	Alzheimer's disease (G30)	33	0.8	7.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.6	5.8
15	Parkinson's disease (G20-G21)	16	0.4	*
...	All other causes (Residual)	609	14.4	131.9
Hispanic, female, 70-74 years				
...	All causes	5,300	100.0	1,455.9
1	Malignant neoplasms (C00-C97)	1,515	28.6	416.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,207	22.8	331.6
3	Diabetes mellitus (E10-E14)	424	8.0	116.5
4	Cerebrovascular diseases (I60-I69)	304	5.7	83.5
5	Chronic lower respiratory diseases (J40-J47)	184	3.5	50.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	3.2	47.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	154	2.9	42.3
8	Influenza and pneumonia (J09-J18)	115	2.2	31.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	2.0	28.8
10	Septicemia (A40-A41)	96	1.8	26.4
11	Alzheimer's disease (G30)	82	1.5	22.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	1.3	18.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.8	11.3
14	Viral hepatitis (B15-B19)	37	0.7	10.2
14	Parkinson's disease (G20-G21)	37	0.7	10.2
...	All other causes (Residual)	760	14.3	208.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	7,065	100.0	2,464.2
1	Malignant neoplasms (C00-C97)	1,725	24.4	601.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,648	23.3	574.8
3	Cerebrovascular diseases (I60-I69)	512	7.2	178.6
4	Diabetes mellitus (E10-E14)	495	7.0	172.6
5	Chronic lower respiratory diseases (J40-J47)	289	4.1	100.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	2.9	70.5
7	Alzheimer's disease (G30)	199	2.8	69.4
8	Influenza and pneumonia (J09-J18)	181	2.6	63.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	157	2.2	54.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	140	2.0	48.8
11	Septicemia (A40-A41)	127	1.8	44.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	1.4	35.2
13	Parkinson's disease (G20-G21)	64	0.9	22.3
14	Viral hepatitis (B15-B19)	37	0.5	12.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.5	12.6
...	All other causes (Residual)	1,152	16.3	401.8
Hispanic, female, 80-84 years				
...	All causes	8,346	100.0	4,023.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,151	25.8	1,037.0
2	Malignant neoplasms (C00-C97)	1,526	18.3	735.7
3	Cerebrovascular diseases (I60-I69)	670	8.0	323.0
4	Diabetes mellitus (E10-E14)	476	5.7	229.5
5	Alzheimer's disease (G30)	414	5.0	199.6
6	Chronic lower respiratory diseases (J40-J47)	383	4.6	184.6
7	Influenza and pneumonia (J09-J18)	277	3.3	133.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	223	2.7	107.5
9	Septicemia (A40-A41)	175	2.1	84.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	1.8	71.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	135	1.6	65.1
12	Parkinson's disease (G20-G21)	101	1.2	48.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	95	1.1	45.8
14	Pneumonitis due to solids and liquids (J69)	59	0.7	28.4
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	54	0.6	26.0
...	All other causes (Residual)	1,458	17.5	702.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	8,095	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,416	29.8	---
2	Malignant neoplasms (C00-C97)	1,023	12.6	---
3	Cerebrovascular diseases (I60-I69)	631	7.8	---
4	Alzheimer's disease (G30)	585	7.2	---
5	Diabetes mellitus (E10-E14)	377	4.7	---
6	Chronic lower respiratory diseases (J40-J47)	366	4.5	---
7	Influenza and pneumonia (J09-J18)	277	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	186	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	156	1.9	---
11	Septicemia (A40-A41)	122	1.5	---
12	Parkinson's disease (G20-G21)	85	1.1	---
13	Pneumonitis due to solids and liquids (J69)	59	0.7	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	50	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.5	---
...	All other causes (Residual)	1,561	19.3	---
Hispanic, female, 90-94 years				
...	All causes	5,291	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,806	34.1	---
2	Alzheimer's disease (G30)	440	8.3	---
3	Malignant neoplasms (C00-C97)	413	7.8	---
4	Cerebrovascular diseases (I60-I69)	399	7.5	---
5	Influenza and pneumonia (J09-J18)	223	4.2	---
6	Chronic lower respiratory diseases (J40-J47)	200	3.8	---
7	Diabetes mellitus (E10-E14)	196	3.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	111	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	1.5	---
11	Septicemia (A40-A41)	76	1.4	---
12	Pneumonitis due to solids and liquids (J69)	54	1.0	---
13	Parkinson's disease (G20-G21)	38	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.5	---
15	Atherosclerosis (I70)	26	0.5	---
...	All other causes (Residual)	1,098	20.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	2,634	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	992	37.7	---
2	Alzheimer's disease (G30)	214	8.1	---
3	Cerebrovascular diseases (I60-I69)	186	7.1	---
4	Malignant neoplasms (C00-C97)	138	5.2	---
5	Influenza and pneumonia (J09-J18)	124	4.7	---
6	Chronic lower respiratory diseases (J40-J47)	80	3.0	---
7	Diabetes mellitus (E10-E14)	74	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	61	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	1.7	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	1.7	---
11	Septicemia (A40-A41)	37	1.4	---
12	Pneumonitis due to solids and liquids (J69)	29	1.1	---
13	Atherosclerosis (I70)	28	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.5	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	12	0.5	---
...	All other causes (Residual)	557	21.1	---
Hispanic, female, 100 years and over				
...	All causes	582	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	228	39.2	---
2	Alzheimer's disease (G30)	47	8.1	---
3	Influenza and pneumonia (J09-J18)	42	7.2	---
4	Cerebrovascular diseases (I60-I69)	39	6.7	---
5	Malignant neoplasms (C00-C97)	19	3.3	---
6	Chronic lower respiratory diseases (J40-J47)	15	2.6	---
7	Diabetes mellitus (E10-E14)	11	1.9	---
8	Septicemia (A40-A41)	10	1.7	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	1.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	1.4	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.2	---
12	Atherosclerosis (I70)	6	1.0	---
13	Parkinson's disease (G20-G21)	5	0.9	---
13	Pneumonitis due to solids and liquids (J69)	5	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	---
15	Nutritional deficiencies (E40-E64)	3	0.5	---
...	All other causes (Residual)	125	21.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	1,981,034	100.0	982.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	503,096	25.4	249.4
2	Malignant neoplasms (C00-C97)	457,084	23.1	226.6
3	Chronic lower respiratory diseases (J40-J47)	126,146	6.4	62.5
4	Cerebrovascular diseases (I60-I69)	106,134	5.4	52.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94,722	4.8	47.0
6	Alzheimer's disease (G30)	73,219	3.7	36.3
7	Diabetes mellitus (E10-E14)	49,388	2.5	24.5
8	Influenza and pneumonia (J09-J18)	45,780	2.3	22.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,439	1.8	17.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,281	1.5	15.0
11	Septicemia (A40-A41)	26,692	1.3	13.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	22,210	1.1	11.0
13	Parkinson's disease (G20-G21)	18,474	0.9	9.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,351	0.9	9.1
15	Pneumonitis due to solids and liquids (J69)	14,135	0.7	7.0
...	All other causes (Residual)	359,883	18.2	178.4
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,373	100.0	26.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	788	33.2	8.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	256	10.8	2.8
3	Malignant neoplasms (C00-C97)	225	9.5	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	7.1	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.4	0.9
6	Influenza and pneumonia (J09-J18)	60	2.5	0.7
7	Septicemia (A40-A41)	41	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	30	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	22	0.9	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	16	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	12	0.5	*
13	Anemias (D50-D64)	10	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
15	Meningococcal infection (A39)	8	0.3	*
15	Meningitis (G00,G03)	8	0.3	*
...	All other causes (Residual)	612	25.8	6.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-9 years				
...	All causes	1,325	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	471	35.5	4.1
2	Malignant neoplasms (C00-C97)	260	19.6	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	6.7	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	3.5	0.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	3.1	0.4
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	2.3	0.3
7	Cerebrovascular diseases (I60-I69)	21	1.6	0.2
8	Influenza and pneumonia (J09-J18)	15	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	15	1.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	13	1.0	*
11	Septicemia (A40-A41)	11	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.6	*
13	Diabetes mellitus (E10-E14)	7	0.5	*
14	Anemias (D50-D64)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	286	21.6	2.5
Non-Hispanic, white, both sexes, 10-14 years				
...	All causes	1,745	100.0	14.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	620	35.5	5.2
2	Malignant neoplasms (C00-C97)	249	14.3	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	8.3	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	4.6	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	4.0	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	3.6	0.5
7	Influenza and pneumonia (J09-J18)	34	1.9	0.3
8	Cerebrovascular diseases (I60-I69)	26	1.5	0.2
9	Septicemia (A40-A41)	21	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.8	*
12	Diabetes mellitus (E10-E14)	11	0.6	*
13	Anemias (D50-D64)	8	0.5	*
14	Meningitis (G00,G03)	5	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	371	21.3	3.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-19 years				
...	All causes	7,046	100.0	53.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,884	55.1	29.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,105	15.7	8.4
3	Malignant neoplasms (C00-C97)	380	5.4	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	324	4.6	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	172	2.4	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	129	1.8	1.0
7	Cerebrovascular diseases (I60-I69)	43	0.6	0.3
8	Influenza and pneumonia (J09-J18)	38	0.5	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.4	0.2
10	Diabetes mellitus (E10-E14)	29	0.4	0.2
11	Septicemia (A40-A41)	27	0.4	0.2
12	Chronic lower respiratory diseases (J40-J47)	22	0.3	0.2
13	Pregnancy, childbirth and the puerperium (O00-O99)	14	0.2	*
14	Aortic aneurysm and dissection (I71)	10	0.1	*
14	Pneumonitis due to solids and liquids (J69)	10	0.1	*
...	All other causes (Residual)	828	11.8	6.3
Non-Hispanic, white, both sexes, 20-24 years				
...	All causes	11,514	100.0	87.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,027	52.3	45.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,920	16.7	14.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	590	5.1	4.5
4	Malignant neoplasms (C00-C97)	569	4.9	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	339	2.9	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	1.2	1.0
7	Influenza and pneumonia (J09-J18)	87	0.8	0.7
8	Diabetes mellitus (E10-E14)	81	0.7	0.6
9	Cerebrovascular diseases (I60-I69)	64	0.6	0.5
10	Pregnancy, childbirth and the puerperium (O00-O99)	49	0.4	0.4
11	Septicemia (A40-A41)	46	0.4	0.4
12	Chronic lower respiratory diseases (J40-J47)	40	0.3	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.3	0.2
14	Legal intervention (Y35,Y89.0)	23	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.2	0.2
...	All other causes (Residual)	1,490	12.9	11.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 25-29 years				
...	All causes	12,154	100.0	94.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,598	46.1	43.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,972	16.2	15.3
3	Malignant neoplasms (C00-C97)	837	6.9	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	627	5.2	4.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	566	4.7	4.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	1.0	1.0
7	Diabetes mellitus (E10-E14)	107	0.9	0.8
8	Cerebrovascular diseases (I60-I69)	93	0.8	0.7
9	Influenza and pneumonia (J09-J18)	89	0.7	0.7
10	Septicemia (A40-A41)	82	0.7	0.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	66	0.5	0.5
11	Pregnancy, childbirth and the puerperium (O00-O99)	66	0.5	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.5	0.5
14	Chronic lower respiratory diseases (J40-J47)	54	0.4	0.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	51	0.4	0.4
...	All other causes (Residual)	1,759	14.5	13.6
Non-Hispanic, white, both sexes, 30-34 years				
...	All causes	12,698	100.0	109.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,608	36.3	39.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,005	15.8	17.3
3	Malignant neoplasms (C00-C97)	1,247	9.8	10.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,096	8.6	9.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	518	4.1	4.5
6	Diabetes mellitus (E10-E14)	206	1.6	1.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	167	1.3	1.4
8	Cerebrovascular diseases (I60-I69)	155	1.2	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	1.0	1.1
10	Influenza and pneumonia (J09-J18)	122	1.0	1.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	116	0.9	1.0
12	Septicemia (A40-A41)	110	0.9	1.0
13	Pregnancy, childbirth and the puerperium (O00-O99)	81	0.6	0.7
14	Chronic lower respiratory diseases (J40-J47)	61	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.4	0.5
...	All other causes (Residual)	2,019	15.9	17.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-39 years				
...	All causes	18,699	100.0	142.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,351	28.6	40.9
2	Malignant neoplasms (C00-C97)	2,650	14.2	20.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,546	13.6	19.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,256	12.1	17.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	512	2.7	3.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	448	2.4	3.4
7	Diabetes mellitus (E10-E14)	348	1.9	2.7
8	Cerebrovascular diseases (I60-I69)	338	1.8	2.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	288	1.5	2.2
10	Influenza and pneumonia (J09-J18)	197	1.1	1.5
11	Septicemia (A40-A41)	170	0.9	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	0.8	1.1
13	Chronic lower respiratory diseases (J40-J47)	135	0.7	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	0.7	1.0
15	Aortic aneurysm and dissection (I71)	84	0.4	0.6
...	All other causes (Residual)	3,108	16.6	23.7
Non-Hispanic, white, both sexes, 40-44 years				
...	All causes	30,392	100.0	214.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,514	21.4	45.9
2	Malignant neoplasms (C00-C97)	5,596	18.4	39.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,934	16.2	34.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,025	10.0	21.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,239	4.1	8.7
6	Diabetes mellitus (E10-E14)	688	2.3	4.8
7	Cerebrovascular diseases (I60-I69)	667	2.2	4.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	546	1.8	3.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	535	1.8	3.8
10	Chronic lower respiratory diseases (J40-J47)	391	1.3	2.8
11	Influenza and pneumonia (J09-J18)	348	1.1	2.5
12	Septicemia (A40-A41)	344	1.1	2.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	247	0.8	1.7
14	Viral hepatitis (B15-B19)	199	0.7	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	169	0.6	1.2
...	All other causes (Residual)	4,950	16.3	34.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-49 years				
...	All causes	53,459	100.0	332.6
1	Malignant neoplasms (C00-C97)	13,021	24.4	81.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,107	18.9	62.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,228	15.4	51.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,763	7.0	23.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,467	4.6	15.4
6	Diabetes mellitus (E10-E14)	1,334	2.5	8.3
6	Cerebrovascular diseases (I60-I69)	1,334	2.5	8.3
8	Chronic lower respiratory diseases (J40-J47)	1,101	2.1	6.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	633	1.2	3.9
10	Viral hepatitis (B15-B19)	628	1.2	3.9
11	Influenza and pneumonia (J09-J18)	624	1.2	3.9
12	Septicemia (A40-A41)	593	1.1	3.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	592	1.1	3.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	415	0.8	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	250	0.5	1.6
...	All other causes (Residual)	8,369	15.7	52.1
Non-Hispanic, white, both sexes, 50-54 years				
...	All causes	76,964	100.0	489.8
1	Malignant neoplasms (C00-C97)	23,279	30.2	148.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,136	21.0	102.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,469	9.7	47.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,579	4.7	22.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,426	4.5	21.8
6	Chronic lower respiratory diseases (J40-J47)	2,388	3.1	15.2
7	Diabetes mellitus (E10-E14)	2,144	2.8	13.6
8	Cerebrovascular diseases (I60-I69)	1,915	2.5	12.2
9	Viral hepatitis (B15-B19)	1,234	1.6	7.9
10	Septicemia (A40-A41)	955	1.2	6.1
11	Influenza and pneumonia (J09-J18)	902	1.2	5.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	706	0.9	4.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	542	0.7	3.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	442	0.6	2.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	363	0.5	2.3
...	All other causes (Residual)	11,484	14.9	73.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-59 years				
...	All causes	97,100	100.0	694.1
1	Malignant neoplasms (C00-C97)	33,923	34.9	242.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,025	21.7	150.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,634	5.8	40.3
4	Chronic lower respiratory diseases (J40-J47)	4,228	4.4	30.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,435	3.5	24.6
6	Diabetes mellitus (E10-E14)	3,167	3.3	22.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,922	3.0	20.9
8	Cerebrovascular diseases (I60-I69)	2,778	2.9	19.9
9	Septicemia (A40-A41)	1,272	1.3	9.1
10	Viral hepatitis (B15-B19)	1,196	1.2	8.5
11	Influenza and pneumonia (J09-J18)	1,083	1.1	7.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,069	1.1	7.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	586	0.6	4.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	474	0.5	3.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	380	0.4	2.7
...	All other causes (Residual)	13,928	14.3	99.6
Non-Hispanic, white, both sexes, 60-64 years				
...	All causes	123,155	100.0	1,046.1
1	Malignant neoplasms (C00-C97)	46,309	37.6	393.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,510	22.3	233.7
3	Chronic lower respiratory diseases (J40-J47)	7,869	6.4	66.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,302	3.5	36.5
5	Diabetes mellitus (E10-E14)	4,255	3.5	36.1
6	Cerebrovascular diseases (I60-I69)	3,828	3.1	32.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,916	2.4	24.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,071	1.7	17.6
9	Septicemia (A40-A41)	1,808	1.5	15.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,579	1.3	13.4
11	Influenza and pneumonia (J09-J18)	1,527	1.2	13.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	765	0.6	6.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	628	0.5	5.3
14	Viral hepatitis (B15-B19)	573	0.5	4.9
15	Aortic aneurysm and dissection (I71)	555	0.5	4.7
...	All other causes (Residual)	16,660	13.5	141.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65-69 years				
...	All causes	143,045	100.0	1,599.0
1	Malignant neoplasms (C00-C97)	54,725	38.3	611.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,934	21.6	345.8
3	Chronic lower respiratory diseases (J40-J47)	12,094	8.5	135.2
4	Cerebrovascular diseases (I60-I69)	5,274	3.7	59.0
5	Diabetes mellitus (E10-E14)	4,741	3.3	53.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,429	2.4	38.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,184	1.5	24.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,176	1.5	24.3
9	Septicemia (A40-A41)	2,116	1.5	23.7
10	Influenza and pneumonia (J09-J18)	1,863	1.3	20.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,399	1.0	15.6
12	Alzheimer's disease (G30)	978	0.7	10.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	905	0.6	10.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	855	0.6	9.6
15	Aortic aneurysm and dissection (I71)	830	0.6	9.3
...	All other causes (Residual)	18,542	13.0	207.3
Non-Hispanic, white, both sexes, 70-74 years				
...	All causes	173,637	100.0	2,504.8
1	Malignant neoplasms (C00-C97)	59,692	34.4	861.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,481	22.2	555.1
3	Chronic lower respiratory diseases (J40-J47)	16,593	9.6	239.4
4	Cerebrovascular diseases (I60-I69)	7,636	4.4	110.2
5	Diabetes mellitus (E10-E14)	5,482	3.2	79.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,828	2.2	55.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,179	1.8	45.9
8	Influenza and pneumonia (J09-J18)	2,818	1.6	40.7
9	Alzheimer's disease (G30)	2,643	1.5	38.1
10	Septicemia (A40-A41)	2,517	1.4	36.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,934	1.1	27.9
12	Parkinson's disease (G20-G21)	1,549	0.9	22.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,230	0.7	17.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,198	0.7	17.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,102	0.6	15.9
...	All other causes (Residual)	23,755	13.7	342.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-79 years				
...	All causes	236,819	100.0	4,016.9
1	Malignant neoplasms (C00-C97)	67,658	28.6	1,147.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,367	23.8	956.1
3	Chronic lower respiratory diseases (J40-J47)	22,037	9.3	373.8
4	Cerebrovascular diseases (I60-I69)	12,756	5.4	216.4
5	Alzheimer's disease (G30)	7,171	3.0	121.6
6	Diabetes mellitus (E10-E14)	6,671	2.8	113.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,155	2.2	87.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,775	2.0	81.0
9	Influenza and pneumonia (J09-J18)	4,772	2.0	80.9
10	Septicemia (A40-A41)	3,683	1.6	62.5
11	Parkinson's disease (G20-G21)	3,280	1.4	55.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,858	0.8	31.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,772	0.7	30.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,669	0.7	28.3
15	Pneumonitis due to solids and liquids (J69)	1,578	0.7	26.8
...	All other causes (Residual)	35,617	15.0	604.1
Non-Hispanic, white, both sexes, 80-84 years				
...	All causes	312,267	100.0	6,531.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,038	26.3	1,716.0
2	Malignant neoplasms (C00-C97)	68,544	22.0	1,433.7
3	Chronic lower respiratory diseases (J40-J47)	24,947	8.0	521.8
4	Cerebrovascular diseases (I60-I69)	20,542	6.6	429.7
5	Alzheimer's disease (G30)	14,969	4.8	313.1
6	Diabetes mellitus (E10-E14)	7,907	2.5	165.4
7	Influenza and pneumonia (J09-J18)	7,858	2.5	164.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,998	2.2	146.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,706	2.1	140.3
10	Parkinson's disease (G20-G21)	5,085	1.6	106.4
11	Septicemia (A40-A41)	4,728	1.5	98.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,105	1.0	64.9
13	Pneumonitis due to solids and liquids (J69)	2,704	0.9	56.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,399	0.8	50.2
15	Aortic aneurysm and dissection (I71)	1,788	0.6	37.4
...	All other causes (Residual)	51,949	16.6	1,086.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 85-89 years				
...	All causes	325,961	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	97,589	29.9	---
2	Malignant neoplasms (C00-C97)	49,425	15.2	---
3	Cerebrovascular diseases (I60-I69)	24,003	7.4	---
4	Alzheimer's disease (G30)	21,226	6.5	---
5	Chronic lower respiratory diseases (J40-J47)	20,403	6.3	---
6	Influenza and pneumonia (J09-J18)	10,039	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,580	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,364	2.3	---
9	Diabetes mellitus (E10-E14)	7,202	2.2	---
10	Parkinson's disease (G20-G21)	4,757	1.5	---
11	Septicemia (A40-A41)	4,447	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,930	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,422	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,258	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,632	0.5	---
...	All other causes (Residual)	60,684	18.6	---
Non-Hispanic, white, both sexes, 90-94 years				
...	All causes	223,256	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,936	33.6	---
2	Malignant neoplasms (C00-C97)	22,177	9.9	---
3	Alzheimer's disease (G30)	17,080	7.7	---
4	Cerebrovascular diseases (I60-I69)	17,034	7.6	---
5	Chronic lower respiratory diseases (J40-J47)	10,335	4.6	---
6	Influenza and pneumonia (J09-J18)	8,370	3.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,255	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,863	2.2	---
9	Diabetes mellitus (E10-E14)	3,824	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,243	1.5	---
11	Septicemia (A40-A41)	2,633	1.2	---
12	Pneumonitis due to solids and liquids (J69)	2,587	1.2	---
13	Parkinson's disease (G20-G21)	2,212	1.0	---
14	Atherosclerosis (I70)	1,483	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,307	0.6	---
...	All other causes (Residual)	45,917	20.6	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 95-99 years				
...	All causes	87,347	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,506	36.1	---
2	Alzheimer's disease (G30)	7,185	8.2	---
3	Cerebrovascular diseases (I60-I69)	6,354	7.3	---
4	Malignant neoplasms (C00-C97)	5,590	6.4	---
5	Influenza and pneumonia (J09-J18)	3,867	4.4	---
6	Chronic lower respiratory diseases (J40-J47)	2,969	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,947	2.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,720	2.0	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,527	1.7	---
10	Pneumonitis due to solids and liquids (J69)	1,068	1.2	---
11	Diabetes mellitus (E10-E14)	1,043	1.2	---
12	Septicemia (A40-A41)	846	1.0	---
13	Atherosclerosis (I70)	727	0.8	---
14	Parkinson's disease (G20-G21)	504	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	341	0.4	---
...	All other causes (Residual)	20,153	23.1	---
Non-Hispanic, white, both sexes, 100 years and over				
...	All causes	17,480	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,662	38.1	---
2	Alzheimer's disease (G30)	1,234	7.1	---
3	Cerebrovascular diseases (I60-I69)	1,172	6.7	---
4	Influenza and pneumonia (J09-J18)	960	5.5	---
5	Malignant neoplasms (C00-C97)	688	3.9	---
6	Chronic lower respiratory diseases (J40-J47)	417	2.4	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	410	2.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	351	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	285	1.6	---
10	Atherosclerosis (I70)	209	1.2	---
11	Pneumonitis due to solids and liquids (J69)	194	1.1	---
12	Septicemia (A40-A41)	140	0.8	---
13	Diabetes mellitus (E10-E14)	136	0.8	---
14	Anemias (D50-D64)	67	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.3	---
...	All other causes (Residual)	4,500	25.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	969,288	100.0	978.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	252,572	26.1	254.9
2	Malignant neoplasms (C00-C97)	238,846	24.6	241.1
3	Chronic lower respiratory diseases (J40-J47)	59,284	6.1	59.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59,173	6.1	59.7
5	Cerebrovascular diseases (I60-I69)	41,092	4.2	41.5
6	Diabetes mellitus (E10-E14)	25,303	2.6	25.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23,835	2.5	24.1
8	Alzheimer's disease (G30)	21,780	2.2	22.0
9	Influenza and pneumonia (J09-J18)	20,516	2.1	20.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,535	1.8	17.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	14,452	1.5	14.6
12	Septicemia (A40-A41)	12,092	1.2	12.2
13	Parkinson's disease (G20-G21)	10,845	1.1	10.9
14	Pneumonitis due to solids and liquids (J69)	7,369	0.8	7.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,088	0.7	7.2
...	All other causes (Residual)	157,506	16.2	159.0
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,365	100.0	29.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	479	35.1	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	10.0	2.9
3	Malignant neoplasms (C00-C97)	127	9.3	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	7.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	2.9	0.8
6	Influenza and pneumonia (J09-J18)	27	2.0	0.6
7	Septicemia (A40-A41)	17	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	14	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.0	*
9	Cerebrovascular diseases (I60-I69)	13	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	7	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
14	Anemias (D50-D64)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	367	26.9	7.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-9 years				
...	All causes	737	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	39.1	4.9
2	Malignant neoplasms (C00-C97)	127	17.2	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	6.0	0.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	4.3	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	3.3	0.4
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	2.3	*
7	Cerebrovascular diseases (I60-I69)	12	1.6	*
8	Influenza and pneumonia (J09-J18)	10	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	9	1.2	*
10	Septicemia (A40-A41)	6	0.8	*
10	Diabetes mellitus (E10-E14)	6	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
15	Anemias (D50-D64)	2	0.3	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.3	*
...	All other causes (Residual)	145	19.7	2.5
Non-Hispanic, white, male, 10-14 years				
...	All causes	1,069	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	417	39.0	6.8
2	Malignant neoplasms (C00-C97)	137	12.8	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	9.9	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	3.9	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.5	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	3.5	0.6
7	Influenza and pneumonia (J09-J18)	17	1.6	*
8	Septicemia (A40-A41)	11	1.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
10	Cerebrovascular diseases (I60-I69)	9	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
12	Anemias (D50-D64)	5	0.5	*
12	Diabetes mellitus (E10-E14)	5	0.5	*
14	Aortic aneurysm and dissection (I71)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
...	All other causes (Residual)	217	20.3	3.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 15-19 years				
...	All causes	4,940	100.0	73.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,705	54.8	40.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	905	18.3	13.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	242	4.9	3.6
4	Malignant neoplasms (C00-C97)	223	4.5	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	115	2.3	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	1.8	1.3
7	Cerebrovascular diseases (I60-I69)	27	0.5	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.5	0.4
9	Influenza and pneumonia (J09-J18)	20	0.4	0.3
10	Diabetes mellitus (E10-E14)	17	0.3	*
11	Septicemia (A40-A41)	13	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	11	0.2	*
13	Aortic aneurysm and dissection (I71)	8	0.2	*
13	Pneumonitis due to solids and liquids (J69)	8	0.2	*
15	Anemias (D50-D64)	6	0.1	*
...	All other causes (Residual)	528	10.7	7.8
Non-Hispanic, white, male, 20-24 years				
...	All causes	8,635	100.0	128.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,677	54.2	69.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,625	18.8	24.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	447	5.2	6.6
4	Malignant neoplasms (C00-C97)	361	4.2	5.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	236	2.7	3.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	0.9	1.1
7	Influenza and pneumonia (J09-J18)	64	0.7	0.9
8	Diabetes mellitus (E10-E14)	52	0.6	0.8
9	Cerebrovascular diseases (I60-I69)	38	0.4	0.6
10	Chronic lower respiratory diseases (J40-J47)	23	0.3	0.3
10	Legal intervention (Y35,Y89.0)	23	0.3	0.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.2	*
13	Septicemia (A40-A41)	16	0.2	*
14	Pneumonitis due to solids and liquids (J69)	13	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.1	*
...	All other causes (Residual)	954	11.0	14.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-29 years				
...	All causes	8,718	100.0	133.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,315	49.5	66.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,610	18.5	24.7
3	Malignant neoplasms (C00-C97)	456	5.2	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	429	4.9	6.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	405	4.6	6.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	0.7	1.0
7	Diabetes mellitus (E10-E14)	60	0.7	0.9
8	Influenza and pneumonia (J09-J18)	51	0.6	0.8
9	Septicemia (A40-A41)	50	0.6	0.8
10	Cerebrovascular diseases (I60-I69)	48	0.6	0.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	42	0.5	0.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	39	0.4	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.4	0.6
14	Chronic lower respiratory diseases (J40-J47)	28	0.3	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.3	0.4
...	All other causes (Residual)	1,061	12.2	16.3
Non-Hispanic, white, male, 30-34 years				
...	All causes	8,612	100.0	147.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,417	39.7	58.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,592	18.5	27.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	759	8.8	13.0
4	Malignant neoplasms (C00-C97)	607	7.0	10.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	378	4.4	6.5
6	Diabetes mellitus (E10-E14)	122	1.4	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	108	1.3	1.9
8	Cerebrovascular diseases (I60-I69)	86	1.0	1.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	77	0.9	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	0.9	1.3
11	Influenza and pneumonia (J09-J18)	66	0.8	1.1
12	Septicemia (A40-A41)	55	0.6	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	0.4	0.6
14	Aortic aneurysm and dissection (I71)	28	0.3	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.3	0.5
15	Chronic lower respiratory diseases (J40-J47)	27	0.3	0.5
...	All other causes (Residual)	1,151	13.4	19.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 35-39 years				
...	All causes	11,998	100.0	182.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,746	31.2	57.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,977	16.5	30.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,574	13.1	23.9
4	Malignant neoplasms (C00-C97)	1,198	10.0	18.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	348	2.9	5.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	284	2.4	4.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	225	1.9	3.4
8	Diabetes mellitus (E10-E14)	211	1.8	3.2
9	Cerebrovascular diseases (I60-I69)	186	1.6	2.8
10	Influenza and pneumonia (J09-J18)	101	0.8	1.5
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	0.7	1.2
12	Septicemia (A40-A41)	76	0.6	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	0.6	1.1
14	Aortic aneurysm and dissection (I71)	71	0.6	1.1
15	Chronic lower respiratory diseases (J40-J47)	63	0.5	1.0
...	All other causes (Residual)	1,783	14.9	27.1
Non-Hispanic, white, male, 40-44 years				
...	All causes	18,812	100.0	265.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,412	23.5	62.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,534	18.8	49.8
3	Malignant neoplasms (C00-C97)	2,465	13.1	34.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,238	11.9	31.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	805	4.3	11.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	437	2.3	6.2
7	Diabetes mellitus (E10-E14)	434	2.3	6.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	374	2.0	5.3
9	Cerebrovascular diseases (I60-I69)	339	1.8	4.8
10	Influenza and pneumonia (J09-J18)	187	1.0	2.6
11	Septicemia (A40-A41)	176	0.9	2.5
11	Chronic lower respiratory diseases (J40-J47)	176	0.9	2.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	0.7	2.0
14	Viral hepatitis (B15-B19)	128	0.7	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	0.5	1.4
...	All other causes (Residual)	2,868	15.2	40.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-49 years				
...	All causes	33,044	100.0	413.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,355	22.3	92.0
2	Malignant neoplasms (C00-C97)	6,378	19.3	79.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,475	16.6	68.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,818	8.5	35.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,637	5.0	20.5
6	Diabetes mellitus (E10-E14)	864	2.6	10.8
7	Cerebrovascular diseases (I60-I69)	694	2.1	8.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	521	1.6	6.5
9	Chronic lower respiratory diseases (J40-J47)	500	1.5	6.3
10	Viral hepatitis (B15-B19)	413	1.2	5.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	411	1.2	5.1
12	Influenza and pneumonia (J09-J18)	325	1.0	4.1
13	Septicemia (A40-A41)	318	1.0	4.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	253	0.8	3.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	157	0.5	2.0
...	All other causes (Residual)	4,925	14.9	61.6
Non-Hispanic, white, male, 50-54 years				
...	All causes	48,287	100.0	620.9
1	Malignant neoplasms (C00-C97)	12,339	25.6	158.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,953	24.8	153.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,040	10.4	64.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,694	5.6	34.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,447	5.1	31.5
6	Diabetes mellitus (E10-E14)	1,373	2.8	17.7
7	Chronic lower respiratory diseases (J40-J47)	1,236	2.6	15.9
8	Cerebrovascular diseases (I60-I69)	1,074	2.2	13.8
9	Viral hepatitis (B15-B19)	910	1.9	11.7
10	Septicemia (A40-A41)	511	1.1	6.6
11	Influenza and pneumonia (J09-J18)	509	1.1	6.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	440	0.9	5.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	440	0.9	5.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	302	0.6	3.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	243	0.5	3.1
...	All other causes (Residual)	6,776	14.0	87.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 55-59 years				
...	All causes	60,557	100.0	881.4
1	Malignant neoplasms (C00-C97)	18,754	31.0	273.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,425	25.5	224.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,859	6.4	56.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,478	4.1	36.1
5	Chronic lower respiratory diseases (J40-J47)	2,262	3.7	32.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,233	3.7	32.5
7	Diabetes mellitus (E10-E14)	1,969	3.3	28.7
8	Cerebrovascular diseases (I60-I69)	1,556	2.6	22.6
9	Viral hepatitis (B15-B19)	896	1.5	13.0
10	Septicemia (A40-A41)	674	1.1	9.8
11	Influenza and pneumonia (J09-J18)	636	1.1	9.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	629	1.0	9.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	383	0.6	5.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	331	0.5	4.8
15	Aortic aneurysm and dissection (I71)	282	0.5	4.1
...	All other causes (Residual)	8,190	13.5	119.2
Non-Hispanic, white, male, 60-64 years				
...	All causes	73,733	100.0	1,290.7
1	Malignant neoplasms (C00-C97)	25,663	34.8	449.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,099	25.9	334.3
3	Chronic lower respiratory diseases (J40-J47)	4,022	5.5	70.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,930	4.0	51.3
5	Diabetes mellitus (E10-E14)	2,545	3.5	44.6
6	Cerebrovascular diseases (I60-I69)	2,144	2.9	37.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,051	2.8	35.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,626	2.2	28.5
9	Septicemia (A40-A41)	956	1.3	16.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	891	1.2	15.6
11	Influenza and pneumonia (J09-J18)	877	1.2	15.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	443	0.6	7.8
13	Aortic aneurysm and dissection (I71)	402	0.5	7.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	376	0.5	6.6
15	Viral hepatitis (B15-B19)	374	0.5	6.5
...	All other causes (Residual)	9,334	12.7	163.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-69 years				
...	All causes	82,493	100.0	1,946.7
1	Malignant neoplasms (C00-C97)	30,196	36.6	712.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,403	24.7	481.5
3	Chronic lower respiratory diseases (J40-J47)	6,077	7.4	143.4
4	Cerebrovascular diseases (I60-I69)	2,844	3.4	67.1
5	Diabetes mellitus (E10-E14)	2,764	3.4	65.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,245	2.7	53.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,412	1.7	33.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,202	1.5	28.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,144	1.4	27.0
10	Septicemia (A40-A41)	1,063	1.3	25.1
11	Influenza and pneumonia (J09-J18)	1,051	1.3	24.8
12	Aortic aneurysm and dissection (I71)	585	0.7	13.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	508	0.6	12.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	496	0.6	11.7
15	Parkinson's disease (G20-G21)	427	0.5	10.1
...	All other causes (Residual)	10,076	12.2	237.8
Non-Hispanic, white, male, 70-74 years				
...	All causes	96,237	100.0	3,036.5
1	Malignant neoplasms (C00-C97)	32,728	34.0	1,032.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,678	24.6	747.1
3	Chronic lower respiratory diseases (J40-J47)	8,236	8.6	259.9
4	Cerebrovascular diseases (I60-I69)	3,804	4.0	120.0
5	Diabetes mellitus (E10-E14)	3,070	3.2	96.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,348	2.4	74.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,719	1.8	54.2
8	Influenza and pneumonia (J09-J18)	1,559	1.6	49.2
9	Septicemia (A40-A41)	1,264	1.3	39.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,183	1.2	37.3
11	Alzheimer's disease (G30)	1,117	1.2	35.2
12	Parkinson's disease (G20-G21)	1,016	1.1	32.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	923	1.0	29.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	689	0.7	21.7
15	Aortic aneurysm and dissection (I71)	682	0.7	21.5
...	All other causes (Residual)	12,221	12.7	385.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 75-79 years				
...	All causes	124,175	100.0	4,883.7
1	Malignant neoplasms (C00-C97)	36,430	29.3	1,432.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,737	25.6	1,248.2
3	Chronic lower respiratory diseases (J40-J47)	10,801	8.7	424.8
4	Cerebrovascular diseases (I60-I69)	5,878	4.7	231.2
5	Diabetes mellitus (E10-E14)	3,437	2.8	135.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,888	2.3	113.6
7	Alzheimer's disease (G30)	2,882	2.3	113.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,555	2.1	100.5
9	Influenza and pneumonia (J09-J18)	2,506	2.0	98.6
10	Parkinson's disease (G20-G21)	2,156	1.7	84.8
11	Septicemia (A40-A41)	1,783	1.4	70.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,038	0.8	40.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	929	0.7	36.5
14	Pneumonitis due to solids and liquids (J69)	898	0.7	35.3
15	Aortic aneurysm and dissection (I71)	895	0.7	35.2
...	All other causes (Residual)	17,362	14.0	682.8
Non-Hispanic, white, male, 80-84 years				
...	All causes	148,307	100.0	7,905.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,778	27.5	2,173.6
2	Malignant neoplasms (C00-C97)	35,378	23.9	1,885.7
3	Chronic lower respiratory diseases (J40-J47)	11,795	8.0	628.7
4	Cerebrovascular diseases (I60-I69)	8,340	5.6	444.5
5	Alzheimer's disease (G30)	5,261	3.5	280.4
6	Influenza and pneumonia (J09-J18)	3,791	2.6	202.1
7	Diabetes mellitus (E10-E14)	3,744	2.5	199.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,542	2.4	188.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,447	2.3	183.7
10	Parkinson's disease (G20-G21)	3,087	2.1	164.5
11	Septicemia (A40-A41)	2,130	1.4	113.5
12	Pneumonitis due to solids and liquids (J69)	1,513	1.0	80.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,295	0.9	69.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,196	0.8	63.8
15	Aortic aneurysm and dissection (I71)	989	0.7	52.7
...	All other causes (Residual)	22,021	14.8	1,173.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85-89 years				
...	All causes	134,152	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,457	30.9	---
2	Malignant neoplasms (C00-C97)	23,730	17.7	---
3	Chronic lower respiratory diseases (J40-J47)	9,042	6.7	---
4	Cerebrovascular diseases (I60-I69)	8,179	6.1	---
5	Alzheimer's disease (G30)	6,439	4.8	---
6	Influenza and pneumonia (J09-J18)	4,315	3.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,438	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,424	2.6	---
9	Diabetes mellitus (E10-E14)	3,051	2.3	---
10	Parkinson's disease (G20-G21)	2,646	2.0	---
11	Septicemia (A40-A41)	1,760	1.3	---
12	Pneumonitis due to solids and liquids (J69)	1,748	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,319	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,050	0.8	---
15	Aortic aneurysm and dissection (I71)	742	0.6	---
...	All other causes (Residual)	21,812	16.3	---
Non-Hispanic, white, male, 90-94 years				
...	All causes	73,443	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,216	34.3	---
2	Malignant neoplasms (C00-C97)	9,437	12.8	---
3	Cerebrovascular diseases (I60-I69)	4,472	6.1	---
4	Alzheimer's disease (G30)	4,157	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	3,986	5.4	---
6	Influenza and pneumonia (J09-J18)	3,059	4.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,978	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,966	2.7	---
9	Diabetes mellitus (E10-E14)	1,287	1.8	---
10	Pneumonitis due to solids and liquids (J69)	1,195	1.6	---
11	Parkinson's disease (G20-G21)	1,066	1.5	---
12	Septicemia (A40-A41)	924	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	880	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	538	0.7	---
15	Atherosclerosis (I70)	418	0.6	---
...	All other causes (Residual)	12,864	17.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 95-99 years				
...	All causes	20,226	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,544	37.3	---
2	Malignant neoplasms (C00-C97)	1,930	9.5	---
3	Cerebrovascular diseases (I60-I69)	1,180	5.8	---
4	Influenza and pneumonia (J09-J18)	1,103	5.5	---
5	Alzheimer's disease (G30)	1,064	5.3	---
6	Chronic lower respiratory diseases (J40-J47)	872	4.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	568	2.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	543	2.7	---
9	Pneumonitis due to solids and liquids (J69)	383	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	285	1.4	---
11	Diabetes mellitus (E10-E14)	268	1.3	---
12	Septicemia (A40-A41)	215	1.1	---
13	Parkinson's disease (G20-G21)	174	0.9	---
14	Atherosclerosis (I70)	133	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	104	0.5	---
...	All other causes (Residual)	3,860	19.1	---
Non-Hispanic, white, male, 100 years and over				
...	All causes	2,674	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,074	40.2	---
2	Influenza and pneumonia (J09-J18)	181	6.8	---
3	Malignant neoplasms (C00-C97)	155	5.8	---
4	Cerebrovascular diseases (I60-I69)	139	5.2	---
5	Alzheimer's disease (G30)	107	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	92	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	2.4	---
9	Pneumonitis due to solids and liquids (J69)	53	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	1.5	---
11	Atherosclerosis (I70)	35	1.3	---
12	Septicemia (A40-A41)	26	1.0	---
13	Diabetes mellitus (E10-E14)	22	0.8	---
14	Parkinson's disease (G20-G21)	17	0.6	---
15	Anemias (D50-D64)	15	0.6	---
...	All other causes (Residual)	582	21.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, all ages				
...	All causes	1,011,746	100.0	985.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250,524	24.8	244.0
2	Malignant neoplasms (C00-C97)	218,238	21.6	212.6
3	Chronic lower respiratory diseases (J40-J47)	66,862	6.6	65.1
4	Cerebrovascular diseases (I60-I69)	65,042	6.4	63.4
5	Alzheimer's disease (G30)	51,439	5.1	50.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,549	3.5	34.6
7	Influenza and pneumonia (J09-J18)	25,264	2.5	24.6
8	Diabetes mellitus (E10-E14)	24,085	2.4	23.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,904	1.8	17.4
10	Septicemia (A40-A41)	14,600	1.4	14.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,263	1.1	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	7,758	0.8	7.6
13	Parkinson's disease (G20-G21)	7,629	0.8	7.4
14	Pneumonitis due to solids and liquids (J69)	6,766	0.7	6.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,446	0.6	6.3
...	All other causes (Residual)	202,377	20.0	197.1
Non-Hispanic, white, female, 1-4 years				
...	All causes	1,008	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	30.7	7.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	11.9	2.7
3	Malignant neoplasms (C00-C97)	98	9.7	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	6.7	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	4.1	0.9
6	Influenza and pneumonia (J09-J18)	33	3.3	0.7
7	Septicemia (A40-A41)	24	2.4	0.5
8	Cerebrovascular diseases (I60-I69)	17	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
12	Meningococcal infection (A39)	5	0.5	*
12	Anemias (D50-D64)	5	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
...	All other causes (Residual)	250	24.8	5.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 5-9 years				
...	All causes	588	100.0	10.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	31.1	3.3
2	Malignant neoplasms (C00-C97)	133	22.6	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	7.7	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	2.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	2.6	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	2.4	*
7	Cerebrovascular diseases (I60-I69)	9	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	8	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.0	*
10	Septicemia (A40-A41)	5	0.9	*
10	Influenza and pneumonia (J09-J18)	5	0.9	*
12	Anemias (D50-D64)	3	0.5	*
12	Meningitis (G00,G03)	3	0.5	*
12	Diseases of appendix (K35-K38)	3	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	136	23.1	2.4
Non-Hispanic, white, female, 10-14 years				
...	All causes	676	100.0	11.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	30.0	3.5
2	Malignant neoplasms (C00-C97)	112	16.6	1.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	5.8	0.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	5.6	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	4.9	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	3.8	0.4
7	Cerebrovascular diseases (I60-I69)	17	2.5	*
7	Influenza and pneumonia (J09-J18)	17	2.5	*
9	Septicemia (A40-A41)	10	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.0	*
11	Diabetes mellitus (E10-E14)	6	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.9	*
13	Anemias (D50-D64)	3	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
15	Meningitis (G00,G03)	2	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	152	22.5	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-19 years				
...	All causes	2,106	100.0	33.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,179	56.0	18.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	200	9.5	3.1
3	Malignant neoplasms (C00-C97)	157	7.5	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	3.9	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	2.7	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	2.0	0.7
7	Influenza and pneumonia (J09-J18)	18	0.9	*
8	Cerebrovascular diseases (I60-I69)	16	0.8	*
9	Septicemia (A40-A41)	14	0.7	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	14	0.7	*
11	Diabetes mellitus (E10-E14)	12	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	11	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.1	*
15	Meningitis (G00,G03)	3	0.1	*
...	All other causes (Residual)	288	13.7	4.5
Non-Hispanic, white, female, 20-24 years				
...	All causes	2,879	100.0	45.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,350	46.9	21.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	295	10.2	4.6
3	Malignant neoplasms (C00-C97)	208	7.2	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	5.0	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	3.6	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	2.1	1.0
7	Pregnancy, childbirth and the puerperium (O00-O99)	49	1.7	0.8
8	Septicemia (A40-A41)	30	1.0	0.5
9	Diabetes mellitus (E10-E14)	29	1.0	0.5
10	Cerebrovascular diseases (I60-I69)	26	0.9	0.4
11	Influenza and pneumonia (J09-J18)	23	0.8	0.4
12	Chronic lower respiratory diseases (J40-J47)	17	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.3	*
...	All other causes (Residual)	512	17.8	8.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 25-29 years				
...	All causes	3,436	100.0	53.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,283	37.3	20.1
2	Malignant neoplasms (C00-C97)	381	11.1	6.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	362	10.5	5.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	198	5.8	3.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	161	4.7	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	66	1.9	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	1.9	1.0
8	Diabetes mellitus (E10-E14)	47	1.4	0.7
9	Cerebrovascular diseases (I60-I69)	45	1.3	0.7
10	Influenza and pneumonia (J09-J18)	38	1.1	0.6
11	Septicemia (A40-A41)	32	0.9	0.5
12	Chronic lower respiratory diseases (J40-J47)	26	0.8	0.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	24	0.7	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.7	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.5	*
...	All other causes (Residual)	669	19.5	10.5
Non-Hispanic, white, female, 30-34 years				
...	All causes	4,086	100.0	71.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,191	29.1	20.7
2	Malignant neoplasms (C00-C97)	640	15.7	11.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	413	10.1	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	337	8.2	5.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	140	3.4	2.4
6	Diabetes mellitus (E10-E14)	84	2.1	1.5
7	Pregnancy, childbirth and the puerperium (O00-O99)	81	2.0	1.4
8	Cerebrovascular diseases (I60-I69)	69	1.7	1.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	59	1.4	1.0
10	Influenza and pneumonia (J09-J18)	56	1.4	1.0
11	Septicemia (A40-A41)	55	1.3	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	1.3	0.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	39	1.0	0.7
14	Chronic lower respiratory diseases (J40-J47)	34	0.8	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.6	0.4
...	All other causes (Residual)	811	19.8	14.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-39 years				
...	All causes	6,701	100.0	102.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,605	24.0	24.6
2	Malignant neoplasms (C00-C97)	1,452	21.7	22.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	682	10.2	10.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	569	8.5	8.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	164	2.4	2.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	2.4	2.5
7	Cerebrovascular diseases (I60-I69)	152	2.3	2.3
8	Diabetes mellitus (E10-E14)	137	2.0	2.1
9	Influenza and pneumonia (J09-J18)	96	1.4	1.5
10	Septicemia (A40-A41)	94	1.4	1.4
11	Chronic lower respiratory diseases (J40-J47)	72	1.1	1.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	63	0.9	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	63	0.9	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	0.9	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	0.8	0.8
...	All other causes (Residual)	1,275	19.0	19.6
Non-Hispanic, white, female, 40-44 years				
...	All causes	11,580	100.0	163.3
1	Malignant neoplasms (C00-C97)	3,131	27.0	44.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,102	18.2	29.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,400	12.1	19.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	787	6.8	11.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	434	3.7	6.1
6	Cerebrovascular diseases (I60-I69)	328	2.8	4.6
7	Diabetes mellitus (E10-E14)	254	2.2	3.6
8	Chronic lower respiratory diseases (J40-J47)	215	1.9	3.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	172	1.5	2.4
10	Septicemia (A40-A41)	168	1.5	2.4
11	Influenza and pneumonia (J09-J18)	161	1.4	2.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	108	0.9	1.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	98	0.8	1.4
14	Viral hepatitis (B15-B19)	71	0.6	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	0.6	1.0
...	All other causes (Residual)	2,082	18.0	29.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-49 years				
...	All causes	20,415	100.0	252.7
1	Malignant neoplasms (C00-C97)	6,643	32.5	82.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,753	13.5	34.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,752	13.5	34.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	945	4.6	11.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	830	4.1	10.3
6	Cerebrovascular diseases (I60-I69)	640	3.1	7.9
7	Chronic lower respiratory diseases (J40-J47)	601	2.9	7.4
8	Diabetes mellitus (E10-E14)	470	2.3	5.8
9	Influenza and pneumonia (J09-J18)	299	1.5	3.7
10	Septicemia (A40-A41)	275	1.3	3.4
11	Viral hepatitis (B15-B19)	215	1.1	2.7
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	181	0.9	2.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	162	0.8	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	0.6	1.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	112	0.5	1.4
...	All other causes (Residual)	3,424	16.8	42.4
Non-Hispanic, white, female, 50-54 years				
...	All causes	28,677	100.0	361.3
1	Malignant neoplasms (C00-C97)	10,940	38.1	137.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,183	14.6	52.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,429	8.5	30.6
4	Chronic lower respiratory diseases (J40-J47)	1,152	4.0	14.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	979	3.4	12.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	885	3.1	11.2
7	Cerebrovascular diseases (I60-I69)	841	2.9	10.6
8	Diabetes mellitus (E10-E14)	771	2.7	9.7
9	Septicemia (A40-A41)	444	1.5	5.6
10	Influenza and pneumonia (J09-J18)	393	1.4	5.0
11	Viral hepatitis (B15-B19)	324	1.1	4.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	266	0.9	3.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	0.6	2.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	140	0.5	1.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	136	0.5	1.7
...	All other causes (Residual)	4,634	16.2	58.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-59 years				
...	All causes	36,543	100.0	513.3
1	Malignant neoplasms (C00-C97)	15,169	41.5	213.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,600	15.3	78.7
3	Chronic lower respiratory diseases (J40-J47)	1,966	5.4	27.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,775	4.9	24.9
5	Cerebrovascular diseases (I60-I69)	1,222	3.3	17.2
6	Diabetes mellitus (E10-E14)	1,198	3.3	16.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	957	2.6	13.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	689	1.9	9.7
9	Septicemia (A40-A41)	598	1.6	8.4
10	Influenza and pneumonia (J09-J18)	447	1.2	6.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	440	1.2	6.2
12	Viral hepatitis (B15-B19)	300	0.8	4.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	203	0.6	2.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	202	0.6	2.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	176	0.5	2.5
...	All other causes (Residual)	5,601	15.3	78.7
Non-Hispanic, white, female, 60-64 years				
...	All causes	49,422	100.0	815.5
1	Malignant neoplasms (C00-C97)	20,646	41.8	340.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,411	17.0	138.8
3	Chronic lower respiratory diseases (J40-J47)	3,847	7.8	63.5
4	Diabetes mellitus (E10-E14)	1,710	3.5	28.2
5	Cerebrovascular diseases (I60-I69)	1,684	3.4	27.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,372	2.8	22.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	865	1.8	14.3
8	Septicemia (A40-A41)	852	1.7	14.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	688	1.4	11.4
10	Influenza and pneumonia (J09-J18)	650	1.3	10.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	445	0.9	7.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	322	0.7	5.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	252	0.5	4.2
14	Alzheimer's disease (G30)	237	0.5	3.9
15	Viral hepatitis (B15-B19)	199	0.4	3.3
...	All other causes (Residual)	7,242	14.7	119.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 65-69 years				
...	All causes	60,552	100.0	1,286.1
1	Malignant neoplasms (C00-C97)	24,529	40.5	521.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,531	17.4	223.7
3	Chronic lower respiratory diseases (J40-J47)	6,017	9.9	127.8
4	Cerebrovascular diseases (I60-I69)	2,430	4.0	51.6
5	Diabetes mellitus (E10-E14)	1,977	3.3	42.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,184	2.0	25.1
7	Septicemia (A40-A41)	1,053	1.7	22.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	974	1.6	20.7
9	Influenza and pneumonia (J09-J18)	812	1.3	17.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	772	1.3	16.4
11	Alzheimer's disease (G30)	553	0.9	11.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	409	0.7	8.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	347	0.6	7.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	255	0.4	5.4
15	Aortic aneurysm and dissection (I71)	245	0.4	5.2
...	All other causes (Residual)	8,464	14.0	179.8
Non-Hispanic, white, female, 70-74 years				
...	All causes	77,400	100.0	2,057.0
1	Malignant neoplasms (C00-C97)	26,964	34.8	716.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,803	19.1	393.4
3	Chronic lower respiratory diseases (J40-J47)	8,357	10.8	222.1
4	Cerebrovascular diseases (I60-I69)	3,832	5.0	101.8
5	Diabetes mellitus (E10-E14)	2,412	3.1	64.1
6	Alzheimer's disease (G30)	1,526	2.0	40.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,480	1.9	39.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,460	1.9	38.8
9	Influenza and pneumonia (J09-J18)	1,259	1.6	33.5
10	Septicemia (A40-A41)	1,253	1.6	33.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	751	1.0	20.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	560	0.7	14.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	541	0.7	14.4
14	Parkinson's disease (G20-G21)	533	0.7	14.2
15	Aortic aneurysm and dissection (I71)	393	0.5	10.4
...	All other causes (Residual)	11,276	14.6	299.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-79 years				
...	All causes	112,644	100.0	3,359.6
1	Malignant neoplasms (C00-C97)	31,228	27.7	931.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,630	21.9	734.6
3	Chronic lower respiratory diseases (J40-J47)	11,236	10.0	335.1
4	Cerebrovascular diseases (I60-I69)	6,878	6.1	205.1
5	Alzheimer's disease (G30)	4,289	3.8	127.9
6	Diabetes mellitus (E10-E14)	3,234	2.9	96.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,267	2.0	67.6
8	Influenza and pneumonia (J09-J18)	2,266	2.0	67.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,220	2.0	66.2
10	Septicemia (A40-A41)	1,900	1.7	56.7
11	Parkinson's disease (G20-G21)	1,124	1.0	33.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,012	0.9	30.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	740	0.7	22.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	734	0.7	21.9
15	Pneumonitis due to solids and liquids (J69)	680	0.6	20.3
...	All other causes (Residual)	18,206	16.2	543.0
Non-Hispanic, white, female, 80-84 years				
...	All causes	163,960	100.0	5,644.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,260	25.2	1,420.4
2	Malignant neoplasms (C00-C97)	33,166	20.2	1,141.8
3	Chronic lower respiratory diseases (J40-J47)	13,152	8.0	452.8
4	Cerebrovascular diseases (I60-I69)	12,202	7.4	420.1
5	Alzheimer's disease (G30)	9,708	5.9	334.2
6	Diabetes mellitus (E10-E14)	4,163	2.5	143.3
7	Influenza and pneumonia (J09-J18)	4,067	2.5	140.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,456	2.1	119.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,259	2.0	112.2
10	Septicemia (A40-A41)	2,598	1.6	89.4
11	Parkinson's disease (G20-G21)	1,998	1.2	68.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,909	1.2	65.7
13	Pneumonitis due to solids and liquids (J69)	1,191	0.7	41.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,104	0.7	38.0
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	854	0.5	29.4
...	All other causes (Residual)	29,873	18.2	1,028.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 85-89 years				
...	All causes	191,809	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,132	29.3	---
2	Malignant neoplasms (C00-C97)	25,695	13.4	---
3	Cerebrovascular diseases (I60-I69)	15,824	8.2	---
4	Alzheimer's disease (G30)	14,787	7.7	---
5	Chronic lower respiratory diseases (J40-J47)	11,361	5.9	---
6	Influenza and pneumonia (J09-J18)	5,724	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,156	2.2	---
8	Diabetes mellitus (E10-E14)	4,151	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,926	2.0	---
10	Septicemia (A40-A41)	2,687	1.4	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,611	1.4	---
12	Parkinson's disease (G20-G21)	2,111	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,674	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,208	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,042	0.5	---
...	All other causes (Residual)	38,720	20.2	---
Non-Hispanic, white, female, 90-94 years				
...	All causes	149,813	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,720	33.2	---
2	Alzheimer's disease (G30)	12,923	8.6	---
3	Malignant neoplasms (C00-C97)	12,740	8.5	---
4	Cerebrovascular diseases (I60-I69)	12,562	8.4	---
5	Chronic lower respiratory diseases (J40-J47)	6,349	4.2	---
6	Influenza and pneumonia (J09-J18)	5,311	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,289	2.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,885	1.9	---
9	Diabetes mellitus (E10-E14)	2,537	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,363	1.6	---
11	Septicemia (A40-A41)	1,709	1.1	---
12	Pneumonitis due to solids and liquids (J69)	1,392	0.9	---
13	Parkinson's disease (G20-G21)	1,146	0.8	---
14	Atherosclerosis (I70)	1,065	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	769	0.5	---
...	All other causes (Residual)	33,053	22.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 95-99 years				
...	All causes	67,121	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,962	35.7	---
2	Alzheimer's disease (G30)	6,121	9.1	---
3	Cerebrovascular diseases (I60-I69)	5,174	7.7	---
4	Malignant neoplasms (C00-C97)	3,660	5.5	---
5	Influenza and pneumonia (J09-J18)	2,764	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	2,097	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,404	2.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,242	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,152	1.7	---
10	Diabetes mellitus (E10-E14)	775	1.2	---
11	Pneumonitis due to solids and liquids (J69)	685	1.0	---
12	Septicemia (A40-A41)	631	0.9	---
13	Atherosclerosis (I70)	594	0.9	---
14	Parkinson's disease (G20-G21)	330	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	237	0.4	---
...	All other causes (Residual)	16,293	24.3	---
Non-Hispanic, white, female, 100 years and over				
...	All causes	14,806	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,588	37.7	---
2	Alzheimer's disease (G30)	1,127	7.6	---
3	Cerebrovascular diseases (I60-I69)	1,033	7.0	---
4	Influenza and pneumonia (J09-J18)	779	5.3	---
5	Malignant neoplasms (C00-C97)	533	3.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	369	2.5	---
7	Chronic lower respiratory diseases (J40-J47)	325	2.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	281	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	220	1.5	---
10	Atherosclerosis (I70)	174	1.2	---
11	Pneumonitis due to solids and liquids (J69)	141	1.0	---
12	Septicemia (A40-A41)	114	0.8	---
12	Diabetes mellitus (E10-E14)	114	0.8	---
14	Anemias (D50-D64)	52	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.3	---
...	All other causes (Residual)	3,911	26.4	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	285,522	100.0	745.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,918	24.5	182.5
2	Malignant neoplasms (C00-C97)	63,279	22.2	165.2
3	Cerebrovascular diseases (I60-I69)	16,527	5.8	43.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,215	4.3	31.9
5	Diabetes mellitus (E10-E14)	11,934	4.2	31.1
6	Chronic lower respiratory diseases (J40-J47)	8,679	3.0	22.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,564	3.0	22.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,206	2.9	21.4
9	Septicemia (A40-A41)	6,359	2.2	16.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5,686	2.0	14.8
11	Influenza and pneumonia (J09-J18)	5,396	1.9	14.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,028	1.8	13.1
13	Alzheimer's disease (G30)	5,024	1.8	13.1
14	Certain conditions originating in the perinatal period (P00-P96)	4,627	1.6	12.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,464	0.9	6.4
...	All other causes (Residual)	51,616	18.1	134.7
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,076	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	26.2	11.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	14.0	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	10.4	4.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	5.9	2.6
5	Malignant neoplasms (C00-C97)	51	4.7	2.1
6	Influenza and pneumonia (J09-J18)	44	4.1	1.8
7	Chronic lower respiratory diseases (J40-J47)	22	2.0	0.9
8	Septicemia (A40-A41)	21	2.0	0.9
9	Anemias (D50-D64)	18	1.7	*
10	Certain conditions originating in the perinatal period (P00-P96)	15	1.4	*
11	Cerebrovascular diseases (I60-I69)	13	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	258	24.0	10.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-9 years				
...	All causes	560	100.0	19.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	35.0	6.6
2	Malignant neoplasms (C00-C97)	77	13.8	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	7.0	1.3
4	Chronic lower respiratory diseases (J40-J47)	31	5.5	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	5.5	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.9	0.7
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	2.7	*
8	Cerebrovascular diseases (I60-I69)	11	2.0	*
9	Influenza and pneumonia (J09-J18)	9	1.6	*
10	Anemias (D50-D64)	8	1.4	*
11	Septicemia (A40-A41)	7	1.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.4	*
...	All other causes (Residual)	102	18.2	3.5
Non-Hispanic, black, both sexes, 10-14 years				
...	All causes	698	100.0	22.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	26.4	6.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	13.5	3.1
3	Malignant neoplasms (C00-C97)	71	10.2	2.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	6.0	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	6.0	1.4
6	Chronic lower respiratory diseases (J40-J47)	41	5.9	1.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	4.0	0.9
8	Diabetes mellitus (E10-E14)	10	1.4	*
9	Cerebrovascular diseases (I60-I69)	8	1.1	*
10	Influenza and pneumonia (J09-J18)	7	1.0	*
11	Septicemia (A40-A41)	6	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
11	Anemias (D50-D64)	6	0.9	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
15	Meningococcal infection (A39)	1	0.1	*
15	Meningitis (G00,G03)	1	0.1	*
15	Atherosclerosis (I70)	1	0.1	*
15	Pneumonitis due to solids and liquids (J69)	1	0.1	*
15	Diseases of appendix (K35-K38)	1	0.1	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.1	*
...	All other causes (Residual)	144	20.6	4.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 15-19 years				
...	All causes	2,806	100.0	82.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,217	43.4	35.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	632	22.5	18.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	189	6.7	5.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	4.6	3.8
5	Malignant neoplasms (C00-C97)	111	4.0	3.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	1.9	1.6
7	Chronic lower respiratory diseases (J40-J47)	37	1.3	1.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.7	0.6
8	Anemias (D50-D64)	21	0.7	0.6
10	Septicemia (A40-A41)	17	0.6	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	17	0.6	*
12	Diabetes mellitus (E10-E14)	16	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
14	Legal intervention (Y35,Y89.0)	11	0.4	*
15	Cerebrovascular diseases (I60-I69)	10	0.4	*
15	Influenza and pneumonia (J09-J18)	10	0.4	*
...	All other causes (Residual)	301	10.7	8.8
Non-Hispanic, black, both sexes, 20-24 years				
...	All causes	4,286	100.0	138.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,775	41.4	57.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	848	19.8	27.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	312	7.3	10.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	243	5.7	7.8
5	Malignant neoplasms (C00-C97)	169	3.9	5.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	92	2.1	3.0
7	Diabetes mellitus (E10-E14)	51	1.2	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	1.2	1.6
9	Anemias (D50-D64)	43	1.0	1.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	39	0.9	1.3
11	Chronic lower respiratory diseases (J40-J47)	38	0.9	1.2
12	Cerebrovascular diseases (I60-I69)	31	0.7	1.0
12	Influenza and pneumonia (J09-J18)	31	0.7	1.0
14	Septicemia (A40-A41)	28	0.7	0.9
15	Legal intervention (Y35,Y89.0)	26	0.6	0.8
...	All other causes (Residual)	510	11.9	16.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-29 years				
...	All causes	4,490	100.0	151.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,437	32.0	48.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	841	18.7	28.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	425	9.5	14.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	265	5.9	9.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	233	5.2	7.9
6	Malignant neoplasms (C00-C97)	228	5.1	7.7
7	Diabetes mellitus (E10-E14)	64	1.4	2.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	56	1.2	1.9
9	Septicemia (A40-A41)	55	1.2	1.9
9	Anemias (D50-D64)	55	1.2	1.9
11	Chronic lower respiratory diseases (J40-J47)	47	1.0	1.6
12	Cerebrovascular diseases (I60-I69)	46	1.0	1.6
13	Influenza and pneumonia (J09-J18)	39	0.9	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	0.8	1.2
15	Legal intervention (Y35,Y89.0)	27	0.6	0.9
...	All other causes (Residual)	637	14.2	21.5
Non-Hispanic, black, both sexes, 30-34 years				
...	All causes	4,626	100.0	182.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,002	21.7	39.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	702	15.2	27.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	618	13.4	24.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	370	8.0	14.6
5	Malignant neoplasms (C00-C97)	339	7.3	13.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	226	4.9	8.9
7	Diabetes mellitus (E10-E14)	109	2.4	4.3
8	Cerebrovascular diseases (I60-I69)	96	2.1	3.8
9	Anemias (D50-D64)	70	1.5	2.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	1.3	2.4
11	Septicemia (A40-A41)	58	1.3	2.3
12	Chronic lower respiratory diseases (J40-J47)	52	1.1	2.1
13	Pregnancy, childbirth and the puerperium (O00-O99)	49	1.1	1.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	1.0	1.8
15	Influenza and pneumonia (J09-J18)	43	0.9	1.7
...	All other causes (Residual)	785	17.0	31.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 35-39 years				
...	All causes	6,176	100.0	233.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,015	16.4	38.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	838	13.6	31.6
3	Malignant neoplasms (C00-C97)	753	12.2	28.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	706	11.4	26.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	614	9.9	23.2
6	Cerebrovascular diseases (I60-I69)	199	3.2	7.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	194	3.1	7.3
8	Diabetes mellitus (E10-E14)	190	3.1	7.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	109	1.8	4.1
10	Septicemia (A40-A41)	89	1.4	3.4
11	Chronic lower respiratory diseases (J40-J47)	79	1.3	3.0
12	Anemias (D50-D64)	75	1.2	2.8
13	Influenza and pneumonia (J09-J18)	64	1.0	2.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	58	0.9	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	55	0.9	2.1
...	All other causes (Residual)	1,138	18.4	43.0
Non-Hispanic, black, both sexes, 40-44 years				
...	All causes	9,403	100.0	349.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,857	19.7	68.9
2	Malignant neoplasms (C00-C97)	1,560	16.6	57.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,000	10.6	37.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	928	9.9	34.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	497	5.3	18.4
6	Cerebrovascular diseases (I60-I69)	414	4.4	15.4
7	Diabetes mellitus (E10-E14)	332	3.5	12.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	198	2.1	7.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	2.0	7.1
10	Septicemia (A40-A41)	158	1.7	5.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	138	1.5	5.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	134	1.4	5.0
13	Chronic lower respiratory diseases (J40-J47)	126	1.3	4.7
14	Influenza and pneumonia (J09-J18)	113	1.2	4.2
15	Anemias (D50-D64)	76	0.8	2.8
...	All other causes (Residual)	1,680	17.9	62.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-49 years				
...	All causes	14,751	100.0	536.1
1	Malignant neoplasms (C00-C97)	3,313	22.5	120.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,233	21.9	117.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,255	8.5	45.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,063	7.2	38.6
5	Cerebrovascular diseases (I60-I69)	740	5.0	26.9
6	Diabetes mellitus (E10-E14)	508	3.4	18.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	448	3.0	16.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	334	2.3	12.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	288	2.0	10.5
10	Septicemia (A40-A41)	264	1.8	9.6
11	Chronic lower respiratory diseases (J40-J47)	239	1.6	8.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	230	1.6	8.4
13	Influenza and pneumonia (J09-J18)	186	1.3	6.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	179	1.2	6.5
15	Viral hepatitis (B15-B19)	112	0.8	4.1
...	All other causes (Residual)	2,359	16.0	85.7
Non-Hispanic, black, both sexes, 50-54 years				
...	All causes	20,776	100.0	845.3
1	Malignant neoplasms (C00-C97)	5,341	25.7	217.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,967	23.9	202.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,249	6.0	50.8
4	Cerebrovascular diseases (I60-I69)	1,158	5.6	47.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	955	4.6	38.9
6	Diabetes mellitus (E10-E14)	836	4.0	34.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	513	2.5	20.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	486	2.3	19.8
9	Septicemia (A40-A41)	441	2.1	17.9
10	Chronic lower respiratory diseases (J40-J47)	423	2.0	17.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	358	1.7	14.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	312	1.5	12.7
13	Influenza and pneumonia (J09-J18)	277	1.3	11.3
14	Viral hepatitis (B15-B19)	256	1.2	10.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	146	0.7	5.9
...	All other causes (Residual)	3,058	14.7	124.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 55-59 years				
...	All causes	24,278	100.0	1,221.3
1	Malignant neoplasms (C00-C97)	7,198	29.6	362.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,040	24.9	303.8
3	Cerebrovascular diseases (I60-I69)	1,307	5.4	65.7
4	Diabetes mellitus (E10-E14)	1,113	4.6	56.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,009	4.2	50.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	703	2.9	35.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	702	2.9	35.3
8	Chronic lower respiratory diseases (J40-J47)	597	2.5	30.0
9	Septicemia (A40-A41)	513	2.1	25.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	492	2.0	24.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	413	1.7	20.8
12	Influenza and pneumonia (J09-J18)	339	1.4	17.1
13	Viral hepatitis (B15-B19)	293	1.2	14.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	184	0.8	9.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	0.4	5.0
...	All other causes (Residual)	3,275	13.5	164.7
Non-Hispanic, black, both sexes, 60-64 years				
...	All causes	23,950	100.0	1,691.2
1	Malignant neoplasms (C00-C97)	7,643	31.9	539.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,255	26.1	441.7
3	Cerebrovascular diseases (I60-I69)	1,235	5.2	87.2
4	Diabetes mellitus (E10-E14)	1,167	4.9	82.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	798	3.3	56.4
6	Chronic lower respiratory diseases (J40-J47)	746	3.1	52.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	582	2.4	41.1
8	Septicemia (A40-A41)	555	2.3	39.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	424	1.8	29.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	364	1.5	25.7
11	Influenza and pneumonia (J09-J18)	357	1.5	25.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	357	1.5	25.2
13	Viral hepatitis (B15-B19)	168	0.7	11.9
14	Pneumonitis due to solids and liquids (J69)	90	0.4	6.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	0.3	5.4
...	All other causes (Residual)	3,132	13.1	221.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-69 years				
...	All causes	24,971	100.0	2,408.1
1	Malignant neoplasms (C00-C97)	8,139	32.6	784.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,449	25.8	621.9
3	Cerebrovascular diseases (I60-I69)	1,439	5.8	138.8
4	Diabetes mellitus (E10-E14)	1,366	5.5	131.7
5	Chronic lower respiratory diseases (J40-J47)	975	3.9	94.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	939	3.8	90.6
7	Septicemia (A40-A41)	586	2.3	56.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	464	1.9	44.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	423	1.7	40.8
10	Influenza and pneumonia (J09-J18)	381	1.5	36.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	245	1.0	23.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	177	0.7	17.1
13	Alzheimer's disease (G30)	115	0.5	11.1
14	Pneumonitis due to solids and liquids (J69)	110	0.4	10.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.4	8.8
...	All other causes (Residual)	3,072	12.3	296.3
Non-Hispanic, black, both sexes, 70-74 years				
...	All causes	26,249	100.0	3,277.9
1	Malignant neoplasms (C00-C97)	7,826	29.8	977.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,946	26.5	867.4
3	Cerebrovascular diseases (I60-I69)	1,593	6.1	198.9
4	Diabetes mellitus (E10-E14)	1,485	5.7	185.4
5	Chronic lower respiratory diseases (J40-J47)	1,125	4.3	140.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	999	3.8	124.8
7	Septicemia (A40-A41)	678	2.6	84.7
8	Influenza and pneumonia (J09-J18)	504	1.9	62.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	486	1.9	60.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	1.6	51.4
11	Alzheimer's disease (G30)	281	1.1	35.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	136	0.5	17.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	126	0.5	15.7
13	Pneumonitis due to solids and liquids (J69)	126	0.5	15.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	98	0.4	12.2
...	All other causes (Residual)	3,428	13.1	428.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 75-79 years				
...	All causes	28,665	100.0	4,829.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,780	27.1	1,310.7
2	Malignant neoplasms (C00-C97)	7,378	25.7	1,242.9
3	Cerebrovascular diseases (I60-I69)	2,050	7.2	345.4
4	Diabetes mellitus (E10-E14)	1,485	5.2	250.2
5	Chronic lower respiratory diseases (J40-J47)	1,231	4.3	207.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,087	3.8	183.1
7	Septicemia (A40-A41)	744	2.6	125.3
8	Alzheimer's disease (G30)	633	2.2	106.6
9	Influenza and pneumonia (J09-J18)	631	2.2	106.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	546	1.9	92.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	386	1.3	65.0
12	Pneumonitis due to solids and liquids (J69)	195	0.7	32.9
13	Parkinson's disease (G20-G21)	142	0.5	23.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	131	0.5	22.1
15	Aortic aneurysm and dissection (I71)	97	0.3	16.3
...	All other causes (Residual)	4,149	14.5	699.0
Non-Hispanic, black, both sexes, 80-84 years				
...	All causes	29,816	100.0	7,030.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,375	28.1	1,974.7
2	Malignant neoplasms (C00-C97)	6,394	21.4	1,507.6
3	Cerebrovascular diseases (I60-I69)	2,252	7.6	531.0
4	Diabetes mellitus (E10-E14)	1,438	4.8	339.1
5	Chronic lower respiratory diseases (J40-J47)	1,254	4.2	295.7
6	Alzheimer's disease (G30)	1,052	3.5	248.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,025	3.4	241.7
8	Septicemia (A40-A41)	805	2.7	189.8
9	Influenza and pneumonia (J09-J18)	770	2.6	181.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	626	2.1	147.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	349	1.2	82.3
12	Pneumonitis due to solids and liquids (J69)	250	0.8	58.9
13	Parkinson's disease (G20-G21)	195	0.7	46.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	148	0.5	34.9
15	Aortic aneurysm and dissection (I71)	108	0.4	25.5
...	All other causes (Residual)	4,775	16.0	1,125.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85-89 years				
...	All causes	25,208	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,392	29.3	---
2	Malignant neoplasms (C00-C97)	4,241	16.8	---
3	Cerebrovascular diseases (I60-I69)	1,959	7.8	---
4	Alzheimer's disease (G30)	1,250	5.0	---
5	Diabetes mellitus (E10-E14)	1,039	4.1	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	948	3.8	---
7	Chronic lower respiratory diseases (J40-J47)	919	3.6	---
8	Influenza and pneumonia (J09-J18)	716	2.8	---
9	Septicemia (A40-A41)	690	2.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	571	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	294	1.2	---
12	Pneumonitis due to solids and liquids (J69)	242	1.0	---
13	Parkinson's disease (G20-G21)	142	0.6	---
14	Atherosclerosis (I70)	117	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	104	0.4	---
...	All other causes (Residual)	4,584	18.2	---
Non-Hispanic, black, both sexes, 90-94 years				
...	All causes	15,306	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,867	31.8	---
2	Malignant neoplasms (C00-C97)	1,774	11.6	---
3	Cerebrovascular diseases (I60-I69)	1,244	8.1	---
4	Alzheimer's disease (G30)	960	6.3	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	493	3.2	---
6	Diabetes mellitus (E10-E14)	491	3.2	---
7	Influenza and pneumonia (J09-J18)	473	3.1	---
8	Chronic lower respiratory diseases (J40-J47)	455	3.0	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	400	2.6	---
10	Septicemia (A40-A41)	351	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	1.2	---
12	Pneumonitis due to solids and liquids (J69)	149	1.0	---
13	Atherosclerosis (I70)	85	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.5	---
15	Parkinson's disease (G20-G21)	77	0.5	---
...	All other causes (Residual)	3,228	21.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 95-99 years				
...	All causes	7,006	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,348	33.5	---
2	Malignant neoplasms (C00-C97)	533	7.6	---
3	Cerebrovascular diseases (I60-I69)	524	7.5	---
4	Alzheimer's disease (G30)	521	7.4	---
5	Influenza and pneumonia (J09-J18)	253	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	224	3.2	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	186	2.7	---
7	Chronic lower respiratory diseases (J40-J47)	186	2.7	---
9	Diabetes mellitus (E10-E14)	179	2.6	---
10	Septicemia (A40-A41)	152	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	1.2	---
12	Pneumonitis due to solids and liquids (J69)	78	1.1	---
13	Atherosclerosis (I70)	45	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.4	---
15	Anemias (D50-D64)	26	0.4	---
...	All other causes (Residual)	1,639	23.4	---
Non-Hispanic, black, both sexes, 100 years and over				
...	All causes	2,218	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	736	33.2	---
2	Cerebrovascular diseases (I60-I69)	158	7.1	---
3	Alzheimer's disease (G30)	144	6.5	---
4	Malignant neoplasms (C00-C97)	124	5.6	---
5	Influenza and pneumonia (J09-J18)	88	4.0	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	84	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	2.9	---
8	Diabetes mellitus (E10-E14)	49	2.2	---
9	Chronic lower respiratory diseases (J40-J47)	44	2.0	---
10	Septicemia (A40-A41)	37	1.7	---
11	Pneumonitis due to solids and liquids (J69)	26	1.2	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	1.1	---
13	Atherosclerosis (I70)	17	0.8	---
14	Anemias (D50-D64)	14	0.6	---
15	Nutritional deficiencies (E40-E64)	7	0.3	---
...	All other causes (Residual)	600	27.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, all ages				
...	All causes	145,168	100.0	794.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,965	24.1	191.3
2	Malignant neoplasms (C00-C97)	32,654	22.5	178.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,282	5.7	45.3
4	Cerebrovascular diseases (I60-I69)	7,135	4.9	39.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,047	4.9	38.6
6	Diabetes mellitus (E10-E14)	5,378	3.7	29.4
7	Chronic lower respiratory diseases (J40-J47)	4,497	3.1	24.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,893	2.7	21.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,724	2.6	20.4
10	Septicemia (A40-A41)	2,841	2.0	15.5
11	Certain conditions originating in the perinatal period (P00-P96)	2,608	1.8	14.3
12	Influenza and pneumonia (J09-J18)	2,542	1.8	13.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,194	1.5	12.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,721	1.2	9.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,653	1.1	9.0
...	All other causes (Residual)	24,034	16.6	131.5
Non-Hispanic, black, male, 1-4 years				
...	All causes	619	100.0	50.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	27.5	13.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	14.5	7.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	10.3	5.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	5.7	2.8
5	Malignant neoplasms (C00-C97)	32	5.2	2.6
6	Influenza and pneumonia (J09-J18)	26	4.2	2.1
7	Septicemia (A40-A41)	11	1.8	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.8	*
9	Anemias (D50-D64)	9	1.5	*
10	Cerebrovascular diseases (I60-I69)	8	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	144	23.3	11.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 5-9 years				
...	All causes	336	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	36.0	8.1
2	Malignant neoplasms (C00-C97)	37	11.0	2.5
3	Chronic lower respiratory diseases (J40-J47)	25	7.4	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	6.8	1.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	6.0	1.3
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	3.6	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	3.0	*
8	Cerebrovascular diseases (I60-I69)	7	2.1	*
8	Influenza and pneumonia (J09-J18)	7	2.1	*
10	Anemias (D50-D64)	4	1.2	*
11	Septicemia (A40-A41)	3	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.9	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.6	*
14	Aortic aneurysm and dissection (I71)	1	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
14	Hernia (K40-K46)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	57	17.0	3.8
Non-Hispanic, black, male, 10-14 years				
...	All causes	414	100.0	26.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	29.2	7.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	15.2	4.0
3	Malignant neoplasms (C00-C97)	31	7.5	2.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	6.0	1.6
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	5.6	1.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	4.8	1.3
7	Chronic lower respiratory diseases (J40-J47)	19	4.6	*
8	Diabetes mellitus (E10-E14)	7	1.7	*
8	Cerebrovascular diseases (I60-I69)	7	1.7	*
10	Septicemia (A40-A41)	3	0.7	*
10	Influenza and pneumonia (J09-J18)	3	0.7	*
12	Anemias (D50-D64)	2	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Atherosclerosis (I70)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (Residual)	83	20.0	5.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-19 years				
...	All causes	2,144	100.0	124.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,084	50.6	62.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	469	21.9	27.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	7.1	8.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	4.1	5.1
5	Malignant neoplasms (C00-C97)	69	3.2	4.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.4	1.7
7	Chronic lower respiratory diseases (J40-J47)	20	0.9	1.2
8	Anemias (D50-D64)	13	0.6	*
9	Septicemia (A40-A41)	10	0.5	*
9	Legal intervention (Y35,Y89.0)	10	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.4	*
11	Diabetes mellitus (E10-E14)	8	0.4	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.3	*
14	Cerebrovascular diseases (I60-I69)	6	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
...	All other causes (Residual)	164	7.6	9.5
Non-Hispanic, black, male, 20-24 years				
...	All causes	3,279	100.0	209.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,588	48.4	101.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	632	19.3	40.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	265	8.1	16.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	167	5.1	10.7
5	Malignant neoplasms (C00-C97)	94	2.9	6.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	52	1.6	3.3
7	Diabetes mellitus (E10-E14)	33	1.0	2.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.0	2.1
9	Anemias (D50-D64)	30	0.9	1.9
10	Legal intervention (Y35,Y89.0)	26	0.8	1.7
11	Chronic lower respiratory diseases (J40-J47)	21	0.6	1.3
12	Cerebrovascular diseases (I60-I69)	17	0.5	*
13	Influenza and pneumonia (J09-J18)	14	0.4	*
14	Septicemia (A40-A41)	12	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.3	*
...	All other causes (Residual)	286	8.7	18.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 25-29 years				
...	All causes	3,211	100.0	220.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,299	40.5	89.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	641	20.0	43.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	249	7.8	17.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	219	6.8	15.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	140	4.4	9.6
6	Malignant neoplasms (C00-C97)	124	3.9	8.5
7	Diabetes mellitus (E10-E14)	30	0.9	2.1
7	Chronic lower respiratory diseases (J40-J47)	30	0.9	2.1
9	Anemias (D50-D64)	27	0.8	1.8
10	Influenza and pneumonia (J09-J18)	26	0.8	1.8
10	Legal intervention (Y35,Y89.0)	26	0.8	1.8
12	Septicemia (A40-A41)	23	0.7	1.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.5	*
14	Cerebrovascular diseases (I60-I69)	15	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.5	*
...	All other causes (Residual)	331	10.3	22.7
Non-Hispanic, black, male, 30-34 years				
...	All causes	3,037	100.0	251.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	891	29.3	73.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	530	17.5	43.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	385	12.7	31.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	194	6.4	16.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	194	6.4	16.0
6	Malignant neoplasms (C00-C97)	136	4.5	11.2
7	Diabetes mellitus (E10-E14)	54	1.8	4.5
8	Cerebrovascular diseases (I60-I69)	51	1.7	4.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	1.1	2.6
10	Septicemia (A40-A41)	31	1.0	2.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.0	2.6
12	Anemias (D50-D64)	29	1.0	2.4
12	Chronic lower respiratory diseases (J40-J47)	29	1.0	2.4
14	Influenza and pneumonia (J09-J18)	21	0.7	1.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	19	0.6	*
...	All other causes (Residual)	410	13.5	33.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-39 years				
...	All causes	3,727	100.0	298.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	655	17.6	52.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	612	16.4	49.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	601	16.1	48.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	355	9.5	28.4
5	Malignant neoplasms (C00-C97)	263	7.1	21.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	163	4.4	13.0
7	Diabetes mellitus (E10-E14)	106	2.8	8.5
8	Cerebrovascular diseases (I60-I69)	99	2.7	7.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	1.6	4.6
10	Chronic lower respiratory diseases (J40-J47)	50	1.3	4.0
11	Septicemia (A40-A41)	40	1.1	3.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.0	3.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.0	3.0
14	Influenza and pneumonia (J09-J18)	36	1.0	2.9
15	Anemias (D50-D64)	29	0.8	2.3
15	Aortic aneurysm and dissection (I71)	29	0.8	2.3
...	All other causes (Residual)	556	14.9	44.5
Non-Hispanic, black, male, 40-44 years				
...	All causes	5,343	100.0	423.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,125	21.1	89.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	685	12.8	54.3
3	Malignant neoplasms (C00-C97)	628	11.8	49.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	579	10.8	45.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	401	7.5	31.8
6	Cerebrovascular diseases (I60-I69)	203	3.8	16.1
7	Diabetes mellitus (E10-E14)	187	3.5	14.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	160	3.0	12.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	2.1	8.7
10	Septicemia (A40-A41)	97	1.8	7.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	1.4	6.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	74	1.4	5.9
13	Influenza and pneumonia (J09-J18)	66	1.2	5.2
14	Chronic lower respiratory diseases (J40-J47)	56	1.0	4.4
15	Anemias (D50-D64)	41	0.8	3.2
...	All other causes (Residual)	855	16.0	67.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-49 years				
...	All causes	8,279	100.0	648.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,940	23.4	151.9
2	Malignant neoplasms (C00-C97)	1,511	18.3	118.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	857	10.4	67.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	714	8.6	55.9
5	Cerebrovascular diseases (I60-I69)	374	4.5	29.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	360	4.3	28.2
7	Diabetes mellitus (E10-E14)	284	3.4	22.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	204	2.5	16.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	151	1.8	11.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	1.8	11.6
11	Septicemia (A40-A41)	136	1.6	10.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	130	1.6	10.2
13	Chronic lower respiratory diseases (J40-J47)	105	1.3	8.2
14	Influenza and pneumonia (J09-J18)	103	1.2	8.1
15	Viral hepatitis (B15-B19)	65	0.8	5.1
...	All other causes (Residual)	1,197	14.5	93.7
Non-Hispanic, black, male, 50-54 years				
...	All causes	12,125	100.0	1,074.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,031	25.0	268.7
2	Malignant neoplasms (C00-C97)	2,761	22.8	244.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	882	7.3	78.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	663	5.5	58.8
5	Cerebrovascular diseases (I60-I69)	641	5.3	56.8
6	Diabetes mellitus (E10-E14)	469	3.9	41.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	343	2.8	30.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	285	2.4	25.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	271	2.2	24.0
10	Septicemia (A40-A41)	222	1.8	19.7
11	Chronic lower respiratory diseases (J40-J47)	221	1.8	19.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	195	1.6	17.3
13	Viral hepatitis (B15-B19)	190	1.6	16.8
14	Influenza and pneumonia (J09-J18)	151	1.2	13.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	121	1.0	10.7
...	All other causes (Residual)	1,679	13.8	148.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-59 years				
...	All causes	14,563	100.0	1,629.3
1	Malignant neoplasms (C00-C97)	4,028	27.7	450.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,834	26.3	428.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	737	5.1	82.5
4	Cerebrovascular diseases (I60-I69)	731	5.0	81.8
5	Diabetes mellitus (E10-E14)	589	4.0	65.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	508	3.5	56.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	404	2.8	45.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	363	2.5	40.6
9	Chronic lower respiratory diseases (J40-J47)	331	2.3	37.0
10	Septicemia (A40-A41)	269	1.8	30.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	247	1.7	27.6
12	Viral hepatitis (B15-B19)	210	1.4	23.5
13	Influenza and pneumonia (J09-J18)	199	1.4	22.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	1.0	16.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	0.6	9.3
...	All other causes (Residual)	1,879	12.9	210.2
Non-Hispanic, black, male, 60-64 years				
...	All causes	13,781	100.0	2,213.3
1	Malignant neoplasms (C00-C97)	4,241	30.8	681.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,813	27.7	612.4
3	Cerebrovascular diseases (I60-I69)	669	4.9	107.4
4	Diabetes mellitus (E10-E14)	611	4.4	98.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	425	3.1	68.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	409	3.0	65.7
7	Chronic lower respiratory diseases (J40-J47)	406	2.9	65.2
8	Septicemia (A40-A41)	282	2.0	45.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	277	2.0	44.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	252	1.8	40.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	234	1.7	37.6
12	Influenza and pneumonia (J09-J18)	214	1.6	34.4
13	Viral hepatitis (B15-B19)	121	0.9	19.4
14	Pneumonitis due to solids and liquids (J69)	59	0.4	9.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	0.4	9.5
...	All other causes (Residual)	1,709	12.4	274.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 65-69 years				
...	All causes	13,935	100.0	3,174.3
1	Malignant neoplasms (C00-C97)	4,401	31.6	1,002.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,873	27.8	882.2
3	Cerebrovascular diseases (I60-I69)	797	5.7	181.6
4	Diabetes mellitus (E10-E14)	682	4.9	155.4
5	Chronic lower respiratory diseases (J40-J47)	543	3.9	123.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	481	3.5	109.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	316	2.3	72.0
8	Septicemia (A40-A41)	301	2.2	68.6
9	Influenza and pneumonia (J09-J18)	215	1.5	49.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	213	1.5	48.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	163	1.2	37.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	123	0.9	28.0
13	Pneumonitis due to solids and liquids (J69)	70	0.5	15.9
14	Alzheimer's disease (G30)	65	0.5	14.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	0.4	11.6
...	All other causes (Residual)	1,641	11.8	373.8
Non-Hispanic, black, male, 70-74 years				
...	All causes	13,718	100.0	4,197.5
1	Malignant neoplasms (C00-C97)	4,249	31.0	1,300.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,778	27.5	1,156.0
3	Cerebrovascular diseases (I60-I69)	812	5.9	248.5
4	Diabetes mellitus (E10-E14)	668	4.9	204.4
5	Chronic lower respiratory diseases (J40-J47)	609	4.4	186.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	453	3.3	138.6
7	Septicemia (A40-A41)	299	2.2	91.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	263	1.9	80.5
9	Influenza and pneumonia (J09-J18)	243	1.8	74.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	229	1.7	70.1
11	Alzheimer's disease (G30)	110	0.8	33.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	87	0.6	26.6
13	Pneumonitis due to solids and liquids (J69)	75	0.5	22.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	66	0.5	20.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.5	19.3
...	All other causes (Residual)	1,714	12.5	524.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-79 years				
...	All causes	13,885	100.0	6,192.9
1	Malignant neoplasms (C00-C97)	3,901	28.1	1,739.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,837	27.6	1,711.4
3	Cerebrovascular diseases (I60-I69)	840	6.0	374.7
4	Chronic lower respiratory diseases (J40-J47)	683	4.9	304.6
5	Diabetes mellitus (E10-E14)	630	4.5	281.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	462	3.3	206.1
7	Influenza and pneumonia (J09-J18)	328	2.4	146.3
8	Septicemia (A40-A41)	323	2.3	144.1
9	Alzheimer's disease (G30)	235	1.7	104.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	229	1.6	102.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	218	1.6	97.2
12	Pneumonitis due to solids and liquids (J69)	109	0.8	48.6
13	Parkinson's disease (G20-G21)	80	0.6	35.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.4	27.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	56	0.4	25.0
...	All other causes (Residual)	1,893	13.6	844.3
Non-Hispanic, black, male, 80-84 years				
...	All causes	12,876	100.0	8,835.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,549	27.6	2,435.4
2	Malignant neoplasms (C00-C97)	3,187	24.8	2,187.0
3	Cerebrovascular diseases (I60-I69)	836	6.5	573.7
4	Chronic lower respiratory diseases (J40-J47)	645	5.0	442.6
5	Diabetes mellitus (E10-E14)	546	4.2	374.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	436	3.4	299.2
7	Influenza and pneumonia (J09-J18)	350	2.7	240.2
8	Alzheimer's disease (G30)	339	2.6	232.6
9	Septicemia (A40-A41)	330	2.6	226.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	231	1.8	158.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	1.4	125.6
12	Pneumonitis due to solids and liquids (J69)	122	0.9	83.7
13	Parkinson's disease (G20-G21)	97	0.8	66.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.5	47.3
15	Atherosclerosis (I70)	48	0.4	32.9
...	All other causes (Residual)	1,908	14.8	1,309.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 85-89 years				
...	All causes	9,191	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,572	28.0	---
2	Malignant neoplasms (C00-C97)	1,995	21.7	---
3	Cerebrovascular diseases (I60-I69)	618	6.7	---
4	Chronic lower respiratory diseases (J40-J47)	427	4.6	---
5	Alzheimer's disease (G30)	333	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	331	3.6	---
7	Diabetes mellitus (E10-E14)	307	3.3	---
8	Influenza and pneumonia (J09-J18)	269	2.9	---
9	Septicemia (A40-A41)	253	2.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	173	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	1.3	---
12	Pneumonitis due to solids and liquids (J69)	115	1.3	---
13	Parkinson's disease (G20-G21)	74	0.8	---
14	Atherosclerosis (I70)	45	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.5	---
...	All other causes (Residual)	1,516	16.5	---
Non-Hispanic, black, male, 90-94 years				
...	All causes	4,297	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,334	31.0	---
2	Malignant neoplasms (C00-C97)	718	16.7	---
3	Cerebrovascular diseases (I60-I69)	283	6.6	---
4	Alzheimer's disease (G30)	186	4.3	---
5	Chronic lower respiratory diseases (J40-J47)	179	4.2	---
6	Influenza and pneumonia (J09-J18)	148	3.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	147	3.4	---
8	Diabetes mellitus (E10-E14)	118	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	104	2.4	---
10	Septicemia (A40-A41)	103	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	1.3	---
12	Pneumonitis due to solids and liquids (J69)	52	1.2	---
13	Parkinson's disease (G20-G21)	33	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	---
15	Atherosclerosis (I70)	20	0.5	---
...	All other causes (Residual)	794	18.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 95-99 years				
...	All causes	1,494	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	495	33.1	---
2	Malignant neoplasms (C00-C97)	200	13.4	---
3	Alzheimer's disease (G30)	80	5.4	---
4	Cerebrovascular diseases (I60-I69)	74	5.0	---
5	Influenza and pneumonia (J09-J18)	64	4.3	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	4.1	---
7	Chronic lower respiratory diseases (J40-J47)	61	4.1	---
8	Diabetes mellitus (E10-E14)	38	2.5	---
9	Septicemia (A40-A41)	34	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	1.6	---
12	Pneumonitis due to solids and liquids (J69)	22	1.5	---
13	Atherosclerosis (I70)	9	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	---
15	Anemias (D50-D64)	5	0.3	---
...	All other causes (Residual)	286	19.1	---
Non-Hispanic, black, male, 100 years and over				
...	All causes	354	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	29.7	---
2	Malignant neoplasms (C00-C97)	36	10.2	---
3	Cerebrovascular diseases (I60-I69)	26	7.3	---
4	Influenza and pneumonia (J09-J18)	22	6.2	---
5	Alzheimer's disease (G30)	15	4.2	---
5	Chronic lower respiratory diseases (J40-J47)	15	4.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	4.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	2.5	---
8	Pneumonitis due to solids and liquids (J69)	9	2.5	---
10	Diabetes mellitus (E10-E14)	8	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	2.0	---
12	Septicemia (A40-A41)	5	1.4	---
13	Atherosclerosis (I70)	3	0.8	---
14	Anemias (D50-D64)	2	0.6	---
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	---
...	All other causes (Residual)	76	21.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	140,354	100.0	700.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,953	24.9	174.4
2	Malignant neoplasms (C00-C97)	30,625	21.8	152.8
3	Cerebrovascular diseases (I60-I69)	9,392	6.7	46.9
4	Diabetes mellitus (E10-E14)	6,556	4.7	32.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,671	3.3	23.3
6	Chronic lower respiratory diseases (J40-J47)	4,182	3.0	20.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,933	2.8	19.6
8	Alzheimer's disease (G30)	3,630	2.6	18.1
9	Septicemia (A40-A41)	3,518	2.5	17.6
10	Influenza and pneumonia (J09-J18)	2,854	2.0	14.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,834	2.0	14.1
12	Certain conditions originating in the perinatal period (P00-P96)	2,019	1.4	10.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,962	1.4	9.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,159	0.8	5.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	811	0.6	4.0
...	All other causes (Residual)	27,255	19.4	136.0
Non-Hispanic, black, female, 1-4 years				
...	All causes	457	100.0	38.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	24.5	9.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	13.3	5.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	10.5	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	6.3	2.4
5	Malignant neoplasms (C00-C97)	19	4.2	*
6	Influenza and pneumonia (J09-J18)	18	3.9	*
7	Chronic lower respiratory diseases (J40-J47)	11	2.4	*
8	Septicemia (A40-A41)	10	2.2	*
9	Anemias (D50-D64)	9	2.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	8	1.8	*
11	Cerebrovascular diseases (I60-I69)	5	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	5	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
15	Salmonella infections (A01-A02)	1	0.2	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	111	24.3	9.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-9 years				
...	All causes	224	100.0	15.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75	33.5	5.2
2	Malignant neoplasms (C00-C97)	40	17.9	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	8.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	5.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	3.6	*
6	Chronic lower respiratory diseases (J40-J47)	6	2.7	*
7	Septicemia (A40-A41)	4	1.8	*
7	Anemias (D50-D64)	4	1.8	*
7	Cerebrovascular diseases (I60-I69)	4	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	3	1.3	*
12	Diabetes mellitus (E10-E14)	2	0.9	*
12	Influenza and pneumonia (J09-J18)	2	0.9	*
14	Viral hepatitis (B15-B19)	1	0.4	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
14	Diseases of appendix (K35-K38)	1	0.4	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	*
...	All other causes (Residual)	36	16.1	2.5
Non-Hispanic, black, female, 10-14 years				
...	All causes	284	100.0	18.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	22.2	4.2
2	Malignant neoplasms (C00-C97)	40	14.1	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	10.9	2.1
4	Chronic lower respiratory diseases (J40-J47)	22	7.7	1.5
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	6.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	6.0	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	2.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.8	*
9	Anemias (D50-D64)	4	1.4	*
9	Influenza and pneumonia (J09-J18)	4	1.4	*
11	Septicemia (A40-A41)	3	1.1	*
11	Diabetes mellitus (E10-E14)	3	1.1	*
13	Meningococcal infection (A39)	1	0.4	*
13	Meningitis (G00,G03)	1	0.4	*
13	Cerebrovascular diseases (I60-I69)	1	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	61	21.5	4.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-19 years				
...	All causes	662	100.0	39.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	163	24.6	9.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	20.1	7.9
3	Malignant neoplasms (C00-C97)	42	6.3	2.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	6.2	2.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	5.6	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	3.5	1.4
7	Chronic lower respiratory diseases (J40-J47)	17	2.6	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	17	2.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	14	2.1	*
10	Anemias (D50-D64)	8	1.2	*
10	Diabetes mellitus (E10-E14)	8	1.2	*
12	Septicemia (A40-A41)	7	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
13	Influenza and pneumonia (J09-J18)	6	0.9	*
15	Cerebrovascular diseases (I60-I69)	4	0.6	*
...	All other causes (Residual)	136	20.5	8.1
Non-Hispanic, black, female, 20-24 years				
...	All causes	1,007	100.0	65.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	21.4	14.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	18.6	12.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	7.5	5.0
4	Malignant neoplasms (C00-C97)	75	7.4	4.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	4.7	3.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	40	4.0	2.6
7	Pregnancy, childbirth and the puerperium (O00-O99)	39	3.9	2.5
8	Diabetes mellitus (E10-E14)	18	1.8	*
9	Influenza and pneumonia (J09-J18)	17	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	17	1.7	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.7	*
12	Septicemia (A40-A41)	16	1.6	*
13	Cerebrovascular diseases (I60-I69)	14	1.4	*
14	Anemias (D50-D64)	13	1.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.0	*
...	All other causes (Residual)	205	20.4	13.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-29 years				
...	All causes	1,279	100.0	85.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	15.6	13.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	176	13.8	11.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	10.8	9.2
4	Malignant neoplasms (C00-C97)	104	8.1	6.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	93	7.3	6.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	56	4.4	3.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	3.6	3.1
8	Diabetes mellitus (E10-E14)	34	2.7	2.3
9	Septicemia (A40-A41)	32	2.5	2.1
10	Cerebrovascular diseases (I60-I69)	31	2.4	2.1
11	Anemias (D50-D64)	28	2.2	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.6	1.3
13	Chronic lower respiratory diseases (J40-J47)	17	1.3	*
14	Influenza and pneumonia (J09-J18)	13	1.0	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.8	*
...	All other causes (Residual)	281	22.0	18.8
Non-Hispanic, black, female, 30-34 years				
...	All causes	1,589	100.0	119.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	233	14.7	17.6
2	Malignant neoplasms (C00-C97)	203	12.8	15.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	176	11.1	13.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	10.8	13.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	111	7.0	8.4
6	Diabetes mellitus (E10-E14)	55	3.5	4.1
7	Pregnancy, childbirth and the puerperium (O00-O99)	49	3.1	3.7
8	Cerebrovascular diseases (I60-I69)	45	2.8	3.4
9	Anemias (D50-D64)	41	2.6	3.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	2.0	2.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	1.9	2.3
12	Septicemia (A40-A41)	27	1.7	2.0
13	Chronic lower respiratory diseases (J40-J47)	23	1.4	1.7
14	Influenza and pneumonia (J09-J18)	22	1.4	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.9	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.9	*
...	All other causes (Residual)	342	21.5	25.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 35-39 years				
...	All causes	2,449	100.0	175.0
1	Malignant neoplasms (C00-C97)	490	20.0	35.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	360	14.7	25.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	259	10.6	18.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	226	9.2	16.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	4.3	7.5
6	Cerebrovascular diseases (I60-I69)	100	4.1	7.1
7	Diabetes mellitus (E10-E14)	84	3.4	6.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	2.1	3.6
9	Septicemia (A40-A41)	49	2.0	3.5
10	Anemias (D50-D64)	46	1.9	3.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	44	1.8	3.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	1.3	2.2
13	Chronic lower respiratory diseases (J40-J47)	29	1.2	2.1
14	Influenza and pneumonia (J09-J18)	28	1.1	2.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	0.9	1.5
...	All other causes (Residual)	526	21.5	37.6
Non-Hispanic, black, female, 40-44 years				
...	All causes	4,060	100.0	283.4
1	Malignant neoplasms (C00-C97)	932	23.0	65.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	732	18.0	51.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	349	8.6	24.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	315	7.8	22.0
5	Cerebrovascular diseases (I60-I69)	211	5.2	14.7
6	Diabetes mellitus (E10-E14)	145	3.6	10.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	2.4	6.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	2.0	5.7
9	Chronic lower respiratory diseases (J40-J47)	70	1.7	4.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	62	1.5	4.3
11	Septicemia (A40-A41)	61	1.5	4.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	60	1.5	4.2
13	Influenza and pneumonia (J09-J18)	47	1.2	3.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	0.9	2.7
15	Anemias (D50-D64)	35	0.9	2.4
...	All other causes (Residual)	825	20.3	57.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-49 years				
...	All causes	6,472	100.0	438.9
1	Malignant neoplasms (C00-C97)	1,802	27.8	122.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,293	20.0	87.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	398	6.1	27.0
4	Cerebrovascular diseases (I60-I69)	366	5.7	24.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	349	5.4	23.7
6	Diabetes mellitus (E10-E14)	224	3.5	15.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	137	2.1	9.3
8	Chronic lower respiratory diseases (J40-J47)	134	2.1	9.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	130	2.0	8.8
10	Septicemia (A40-A41)	128	2.0	8.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	100	1.5	6.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	1.4	6.0
13	Influenza and pneumonia (J09-J18)	83	1.3	5.6
14	Viral hepatitis (B15-B19)	47	0.7	3.2
15	Anemias (D50-D64)	32	0.5	2.2
...	All other causes (Residual)	1,161	17.9	78.7
Non-Hispanic, black, female, 50-54 years				
...	All causes	8,651	100.0	650.5
1	Malignant neoplasms (C00-C97)	2,580	29.8	194.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,936	22.4	145.6
3	Cerebrovascular diseases (I60-I69)	517	6.0	38.9
4	Diabetes mellitus (E10-E14)	367	4.2	27.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	367	4.2	27.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	292	3.4	22.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	228	2.6	17.1
8	Septicemia (A40-A41)	219	2.5	16.5
9	Chronic lower respiratory diseases (J40-J47)	202	2.3	15.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	163	1.9	12.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	143	1.7	10.8
12	Influenza and pneumonia (J09-J18)	126	1.5	9.5
13	Viral hepatitis (B15-B19)	66	0.8	5.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	0.5	3.1
15	Anemias (D50-D64)	29	0.3	2.2
...	All other causes (Residual)	1,375	15.9	103.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 55-59 years				
...	All causes	9,715	100.0	887.9
1	Malignant neoplasms (C00-C97)	3,170	32.6	289.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,206	22.7	201.6
3	Cerebrovascular diseases (I60-I69)	576	5.9	52.6
4	Diabetes mellitus (E10-E14)	524	5.4	47.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	299	3.1	27.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	2.8	24.9
7	Chronic lower respiratory diseases (J40-J47)	266	2.7	24.3
8	Septicemia (A40-A41)	244	2.5	22.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	194	2.0	17.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	166	1.7	15.2
11	Influenza and pneumonia (J09-J18)	140	1.4	12.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	129	1.3	11.8
13	Viral hepatitis (B15-B19)	83	0.9	7.6
14	Anemias (D50-D64)	37	0.4	3.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	0.3	3.0
...	All other causes (Residual)	1,376	14.2	125.8
Non-Hispanic, black, female, 60-64 years				
...	All causes	10,169	100.0	1,281.6
1	Malignant neoplasms (C00-C97)	3,402	33.5	428.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,442	24.0	307.8
3	Cerebrovascular diseases (I60-I69)	566	5.6	71.3
4	Diabetes mellitus (E10-E14)	556	5.5	70.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	389	3.8	49.0
6	Chronic lower respiratory diseases (J40-J47)	340	3.3	42.8
7	Septicemia (A40-A41)	273	2.7	34.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	190	1.9	23.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	1.5	19.8
10	Influenza and pneumonia (J09-J18)	143	1.4	18.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	105	1.0	13.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	87	0.9	11.0
13	Viral hepatitis (B15-B19)	47	0.5	5.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.3	4.0
15	Anemias (D50-D64)	31	0.3	3.9
15	Pneumonitis due to solids and liquids (J69)	31	0.3	3.9
...	All other causes (Residual)	1,378	13.6	173.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-69 years				
...	All causes	11,036	100.0	1,845.6
1	Malignant neoplasms (C00-C97)	3,738	33.9	625.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,576	23.3	430.8
3	Diabetes mellitus (E10-E14)	684	6.2	114.4
4	Cerebrovascular diseases (I60-I69)	642	5.8	107.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	458	4.2	76.6
6	Chronic lower respiratory diseases (J40-J47)	432	3.9	72.2
7	Septicemia (A40-A41)	285	2.6	47.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	210	1.9	35.1
9	Influenza and pneumonia (J09-J18)	166	1.5	27.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	1.3	24.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	82	0.7	13.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	54	0.5	9.0
13	Alzheimer's disease (G30)	50	0.5	8.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.4	7.7
15	Pneumonitis due to solids and liquids (J69)	40	0.4	6.7
...	All other causes (Residual)	1,425	12.9	238.3
Non-Hispanic, black, female, 70-74 years				
...	All causes	12,531	100.0	2,643.8
1	Malignant neoplasms (C00-C97)	3,577	28.5	754.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,168	25.3	668.4
3	Diabetes mellitus (E10-E14)	817	6.5	172.4
4	Cerebrovascular diseases (I60-I69)	781	6.2	164.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	546	4.4	115.2
6	Chronic lower respiratory diseases (J40-J47)	516	4.1	108.9
7	Septicemia (A40-A41)	379	3.0	80.0
8	Influenza and pneumonia (J09-J18)	261	2.1	55.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	257	2.1	54.2
10	Alzheimer's disease (G30)	171	1.4	36.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	1.2	31.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.5	13.3
13	Pneumonitis due to solids and liquids (J69)	51	0.4	10.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	49	0.4	10.3
15	Aortic aneurysm and dissection (I71)	40	0.3	8.4
...	All other causes (Residual)	1,706	13.6	359.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 75-79 years				
...	All causes	14,780	100.0	4,001.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,943	26.7	1,067.5
2	Malignant neoplasms (C00-C97)	3,477	23.5	941.3
3	Cerebrovascular diseases (I60-I69)	1,210	8.2	327.6
4	Diabetes mellitus (E10-E14)	855	5.8	231.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	625	4.2	169.2
6	Chronic lower respiratory diseases (J40-J47)	548	3.7	148.4
7	Septicemia (A40-A41)	421	2.8	114.0
8	Alzheimer's disease (G30)	398	2.7	107.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	317	2.1	85.8
10	Influenza and pneumonia (J09-J18)	303	2.1	82.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	168	1.1	45.5
12	Pneumonitis due to solids and liquids (J69)	86	0.6	23.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.5	19.0
14	Parkinson's disease (G20-G21)	62	0.4	16.8
15	Aortic aneurysm and dissection (I71)	51	0.3	13.8
...	All other causes (Residual)	2,246	15.2	608.0
Non-Hispanic, black, female, 80-84 years				
...	All causes	16,940	100.0	6,085.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,826	28.5	1,733.6
2	Malignant neoplasms (C00-C97)	3,207	18.9	1,152.0
3	Cerebrovascular diseases (I60-I69)	1,416	8.4	508.7
4	Diabetes mellitus (E10-E14)	892	5.3	320.4
5	Alzheimer's disease (G30)	713	4.2	256.1
6	Chronic lower respiratory diseases (J40-J47)	609	3.6	218.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	589	3.5	211.6
8	Septicemia (A40-A41)	475	2.8	170.6
9	Influenza and pneumonia (J09-J18)	420	2.5	150.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	395	2.3	141.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	1.0	59.6
12	Pneumonitis due to solids and liquids (J69)	128	0.8	46.0
13	Parkinson's disease (G20-G21)	98	0.6	35.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	79	0.5	28.4
15	Aortic aneurysm and dissection (I71)	67	0.4	24.1
...	All other causes (Residual)	2,860	16.9	1,027.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85-89 years				
...	All causes	16,017	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,820	30.1	---
2	Malignant neoplasms (C00-C97)	2,246	14.0	---
3	Cerebrovascular diseases (I60-I69)	1,341	8.4	---
4	Alzheimer's disease (G30)	917	5.7	---
5	Diabetes mellitus (E10-E14)	732	4.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	617	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	492	3.1	---
8	Influenza and pneumonia (J09-J18)	447	2.8	---
9	Septicemia (A40-A41)	437	2.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	398	2.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	1.1	---
12	Pneumonitis due to solids and liquids (J69)	127	0.8	---
13	Atherosclerosis (I70)	72	0.4	---
14	Parkinson's disease (G20-G21)	68	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.4	---
...	All other causes (Residual)	3,068	19.2	---
Non-Hispanic, black, female, 90-94 years				
...	All causes	11,009	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,533	32.1	---
2	Malignant neoplasms (C00-C97)	1,056	9.6	---
3	Cerebrovascular diseases (I60-I69)	961	8.7	---
4	Alzheimer's disease (G30)	774	7.0	---
5	Diabetes mellitus (E10-E14)	373	3.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	346	3.1	---
7	Influenza and pneumonia (J09-J18)	325	3.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	296	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	276	2.5	---
10	Septicemia (A40-A41)	248	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	1.1	---
12	Pneumonitis due to solids and liquids (J69)	97	0.9	---
13	Atherosclerosis (I70)	65	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.5	---
15	Parkinson's disease (G20-G21)	44	0.4	---
...	All other causes (Residual)	2,434	22.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 95-99 years				
...	All causes	5,512	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,853	33.6	---
2	Cerebrovascular diseases (I60-I69)	450	8.2	---
3	Alzheimer's disease (G30)	441	8.0	---
4	Malignant neoplasms (C00-C97)	333	6.0	---
5	Influenza and pneumonia (J09-J18)	189	3.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	162	2.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	153	2.8	---
8	Diabetes mellitus (E10-E14)	141	2.6	---
9	Chronic lower respiratory diseases (J40-J47)	125	2.3	---
10	Septicemia (A40-A41)	118	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	1.1	---
12	Pneumonitis due to solids and liquids (J69)	56	1.0	---
13	Atherosclerosis (I70)	36	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.4	---
14	Anemias (D50-D64)	21	0.4	---
...	All other causes (Residual)	1,353	24.5	---
Non-Hispanic, black, female, 100 years and over				
...	All causes	1,864	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	631	33.9	---
2	Cerebrovascular diseases (I60-I69)	132	7.1	---
3	Alzheimer's disease (G30)	129	6.9	---
4	Malignant neoplasms (C00-C97)	88	4.7	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	75	4.0	---
6	Influenza and pneumonia (J09-J18)	66	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	2.7	---
8	Diabetes mellitus (E10-E14)	41	2.2	---
9	Septicemia (A40-A41)	32	1.7	---
10	Chronic lower respiratory diseases (J40-J47)	29	1.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	1.0	---
12	Pneumonitis due to solids and liquids (J69)	17	0.9	---
13	Atherosclerosis (I70)	14	0.8	---
14	Anemias (D50-D64)	12	0.6	---
15	Nutritional deficiencies (E40-E64)	6	0.3	---
...	All other causes (Residual)	523	28.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available