

A revised mortality data file for 2014 was released on April 3, 2017 to include corrections from Tennessee and Massachusetts affecting 125 records previously coded to Accidental discharge of firearms (W32-W34). After the revision, 50% of the changed records are now classified as Assault (homicide) by discharge of firearms (\*U01.4 and X93-X95); 42% as Intentional self-harm (suicide) by discharge of firearms (X72-X74); 4% as Discharge of firearms, undetermined intent (Y22-Y24); and the remaining 4% to various other causes. For more information, please see "[Deaths: Final Data for 2014](#)." This table was updated on May 3, 2017 to correct errors. The shaded areas in this table indicate the changes that were made.

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	169,387	100.0	305.8
1	Malignant neoplasms (C00-C97)	36,447	21.5	65.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,021	20.1	61.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,413	7.3	22.4
4	Cerebrovascular diseases (I60-I69)	8,713	5.1	15.7
5	Diabetes mellitus (E10-E14)	7,795	4.6	14.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	5,658	3.3	10.2
7	Alzheimer's disease (G30)	4,934	2.9	8.9
8	Chronic lower respiratory diseases (J40-J47)	4,795	2.8	8.7
9	Influenza and pneumonia (J09-J18)	3,875	2.3	7.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,273	1.9	5.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,246	1.9	5.9
12	Septicemia (A40-A41)	2,613	1.5	4.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,610	1.5	4.7
14	Certain conditions originating in the perinatal period (P00-P96)	2,422	1.4	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,139	1.3	3.9
...	All other causes (residual)	34,433	20.3	62.2
Hispanic, both sexes, 1-4 years				
...	All causes	769	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	28.7	5.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	12.6	2.4
3	Malignant neoplasms (C00-C97)	81	10.5	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	9.1	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.8	0.7
6	Influenza and pneumonia (J09-J18)	20	2.6	0.5
7	Chronic lower respiratory diseases (J40-J47)	10	1.3	*
7	Certain conditions originating in the perinatal period (P00-P96)	10	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.0	*
10	Cerebrovascular diseases (I60-I69)	7	0.9	*
11	Septicemia (A40-A41)	5	0.7	*
12	Nutritional deficiencies (E40-E64)	4	0.5	*
12	Meningitis (G00,G03)	4	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (residual)	196	25.5	4.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	531	100.0	10.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	31.5	3.2
2	Malignant neoplasms (C00-C97)	118	22.2	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	7.9	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	4.0	0.4
5	Influenza and pneumonia (J09-J18)	16	3.0	*
6	Cerebrovascular diseases (I60-I69)	15	2.8	*
6	Chronic lower respiratory diseases (J40-J47)	15	2.8	*
8	Septicemia (A40-A41)	11	2.1	*
9	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
11	Nutritional deficiencies (E40-E64)	2	0.4	*
11	Meningitis (G00,G03)	2	0.4	*
11	Hernia (K40-K46)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Anemias (D50-D64)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
...	All other causes (residual)	102	19.2	2.0
Hispanic, both sexes, 10-14 years				
...	All causes	585	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	25.0	3.0
2	Malignant neoplasms (C00-C97)	101	17.3	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	11.3	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.5	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	5.8	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.2	*
7	Cerebrovascular diseases (I60-I69)	12	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.2	*
9	Influenza and pneumonia (J09-J18)	7	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	7	1.2	*
12	Septicemia (A40-A41)	5	0.9	*
13	Anemias (D50-D64)	4	0.7	*
14	Diabetes mellitus (E10-E14)	3	0.5	*
14	Diseases of appendix (K35-K38)	3	0.5	*
...	All other causes (residual)	124	21.2	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	1,767	100.0	38.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	685	38.8	14.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	304	17.2	6.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	279	15.8	6.0
4	Malignant neoplasms (C00-C97)	136	7.7	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	3.1	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	2.0	0.8
7	Cerebrovascular diseases (I60-I69)	9	0.5	*
8	Influenza and pneumonia (J09-J18)	8	0.5	*
9	Septicemia (A40-A41)	7	0.4	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	7	0.4	*
11	Diabetes mellitus (E10-E14)	6	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.3	*
12	Legal intervention (Y35,Y89.0)	5	0.3	*
14	Anemias (D50-D64)	4	0.2	*
14	Pneumonitis due to solids and liquids (J69)	4	0.2	*
...	All other causes (residual)	218	12.3	4.7
Hispanic, both sexes, 20-24 years				
...	All causes	3,138	100.0	65.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,359	43.3	28.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	508	16.2	10.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	441	14.1	9.2
4	Malignant neoplasms (C00-C97)	228	7.3	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	2.4	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	0.9	0.6
7	Cerebrovascular diseases (I60-I69)	25	0.8	0.5
8	Influenza and pneumonia (J09-J18)	23	0.7	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	22	0.7	0.5
10	Septicemia (A40-A41)	17	0.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	16	0.5	*
13	Legal intervention (Y35,Y89.0)	15	0.5	*
14	Diabetes mellitus (E10-E14)	12	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (residual)	336	10.7	7.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	3,245	100.0	73.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,242	38.3	27.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	416	12.8	9.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	385	11.9	8.7
4	Malignant neoplasms (C00-C97)	271	8.4	6.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	4.7	3.5
6	Cerebrovascular diseases (I60-I69)	50	1.5	1.1
7	Influenza and pneumonia (J09-J18)	46	1.4	1.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	38	1.2	0.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.2	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.0	0.7
11	Diabetes mellitus (E10-E14)	32	1.0	0.7
12	Pregnancy, childbirth and the puerperium (O00-O99)	31	1.0	0.7
13	Septicemia (A40-A41)	23	0.7	0.5
14	Legal intervention (Y35,Y89.0)	22	0.7	0.5
15	Chronic lower respiratory diseases (J40-J47)	18	0.6	*
...	All other causes (residual)	446	13.7	10.0
Hispanic, both sexes, 30-34 years				
...	All causes	3,330	100.0	75.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,101	33.1	24.8
2	Malignant neoplasms (C00-C97)	360	10.8	8.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	327	9.8	7.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	326	9.8	7.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	241	7.2	5.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	113	3.4	2.5
7	Diabetes mellitus (E10-E14)	69	2.1	1.6
8	Cerebrovascular diseases (I60-I69)	62	1.9	1.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.8	1.4
10	Influenza and pneumonia (J09-J18)	56	1.7	1.3
11	Septicemia (A40-A41)	30	0.9	0.7
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.8	0.6
13	Pregnancy, childbirth and the puerperium (O00-O99)	25	0.8	0.6
14	Chronic lower respiratory diseases (J40-J47)	24	0.7	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.5	*
...	All other causes (residual)	490	14.7	11.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,972	100.0	96.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	968	24.4	23.4
2	Malignant neoplasms (C00-C97)	670	16.9	16.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	364	9.2	8.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	316	8.0	7.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	283	7.1	6.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	186	4.7	4.5
7	Cerebrovascular diseases (I60-I69)	111	2.8	2.7
8	Diabetes mellitus (E10-E14)	101	2.5	2.4
9	Influenza and pneumonia (J09-J18)	85	2.1	2.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	76	1.9	1.8
11	Septicemia (A40-A41)	50	1.3	1.2
12	Pregnancy, childbirth and the puerperium (O00-O99)	32	0.8	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	0.8	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.7	0.7
15	Legal intervention (Y35,Y89.0)	21	0.5	0.5
...	All other causes (residual)	651	16.4	15.7
Hispanic, both sexes, 40-44 years				
...	All causes	5,196	100.0	134.7
1	Malignant neoplasms (C00-C97)	1,028	19.8	26.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,012	19.5	26.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	670	12.9	17.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	347	6.7	9.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	289	5.6	7.5
6	Cerebrovascular diseases (I60-I69)	207	4.0	5.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	207	4.0	5.4
8	Diabetes mellitus (E10-E14)	158	3.0	4.1
9	Influenza and pneumonia (J09-J18)	108	2.1	2.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	97	1.9	2.5
11	Septicemia (A40-A41)	73	1.4	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	1.2	1.6
13	Viral hepatitis (B15-B19)	40	0.8	1.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	0.7	0.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	29	0.6	0.8
...	All other causes (residual)	834	16.1	21.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	6,951	100.0	205.0
1	Malignant neoplasms (C00-C97)	1,618	23.3	47.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,059	15.2	31.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	975	14.0	28.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	557	8.0	16.4
5	Diabetes mellitus (E10-E14)	297	4.3	8.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	281	4.0	8.3
7	Cerebrovascular diseases (I60-I69)	278	4.0	8.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	139	2.0	4.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	1.9	3.8
10	Influenza and pneumonia (J09-J18)	121	1.7	3.6
11	Viral hepatitis (B15-B19)	93	1.3	2.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	1.3	2.7
13	Septicemia (A40-A41)	78	1.1	2.3
14	Chronic lower respiratory diseases (J40-J47)	63	0.9	1.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	0.7	1.5
...	All other causes (residual)	1,118	16.1	33.0
Hispanic, both sexes, 50-54 years				
...	All causes	9,754	100.0	331.7
1	Malignant neoplasms (C00-C97)	2,615	26.8	88.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,655	17.0	56.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	897	9.2	30.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	865	8.9	29.4
5	Diabetes mellitus (E10-E14)	483	5.0	16.4
6	Cerebrovascular diseases (I60-I69)	388	4.0	13.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	230	2.4	7.8
8	Viral hepatitis (B15-B19)	206	2.1	7.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	192	2.0	6.5
10	Septicemia (A40-A41)	169	1.7	5.7
11	Influenza and pneumonia (J09-J18)	159	1.6	5.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	1.6	5.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	113	1.2	3.8
14	Chronic lower respiratory diseases (J40-J47)	96	1.0	3.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	94	1.0	3.2
...	All other causes (residual)	1,436	14.7	48.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	11,637	100.0	500.7
1	Malignant neoplasms (C00-C97)	3,424	29.4	147.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,100	18.0	90.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	945	8.1	40.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	707	6.1	30.4
5	Diabetes mellitus (E10-E14)	670	5.8	28.8
6	Cerebrovascular diseases (I60-I69)	464	4.0	20.0
7	Viral hepatitis (B15-B19)	257	2.2	11.1
8	Influenza and pneumonia (J09-J18)	240	2.1	10.3
9	Septicemia (A40-A41)	220	1.9	9.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	189	1.6	8.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	188	1.6	8.1
12	Chronic lower respiratory diseases (J40-J47)	184	1.6	7.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	123	1.1	5.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	114	1.0	4.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	0.7	3.3
...	All other causes (residual)	1,735	14.9	74.7
Hispanic, both sexes, 60-64 years				
...	All causes	13,064	100.0	755.3
1	Malignant neoplasms (C00-C97)	4,064	31.1	235.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,688	20.6	155.4
3	Diabetes mellitus (E10-E14)	826	6.3	47.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	799	6.1	46.2
5	Cerebrovascular diseases (I60-I69)	546	4.2	31.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	543	4.2	31.4
7	Influenza and pneumonia (J09-J18)	296	2.3	17.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	283	2.2	16.4
9	Chronic lower respiratory diseases (J40-J47)	253	1.9	14.6
10	Septicemia (A40-A41)	242	1.9	14.0
11	Viral hepatitis (B15-B19)	223	1.7	12.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	140	1.1	8.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	127	1.0	7.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	76	0.6	4.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	64	0.5	3.7
...	All other causes (residual)	1,894	14.5	109.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	13,647	100.0	1,076.5
1	Malignant neoplasms (C00-C97)	4,388	32.2	346.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,838	20.8	223.9
3	Diabetes mellitus (E10-E14)	923	6.8	72.8
4	Cerebrovascular diseases (I60-I69)	610	4.5	48.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	599	4.4	47.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	415	3.0	32.7
7	Chronic lower respiratory diseases (J40-J47)	390	2.9	30.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	382	2.8	30.1
9	Influenza and pneumonia (J09-J18)	281	2.1	22.2
10	Septicemia (A40-A41)	236	1.7	18.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	184	1.3	14.5
12	Viral hepatitis (B15-B19)	114	0.8	9.0
13	Alzheimer's disease (G30)	104	0.8	8.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	0.7	7.2
15	Parkinson's disease (G20-G21)	60	0.4	4.7
...	All other causes (residual)	2,032	14.9	160.3
Hispanic, both sexes, 70-74 years				
...	All causes	14,389	100.0	1,669.6
1	Malignant neoplasms (C00-C97)	4,327	30.1	502.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,150	21.9	365.5
3	Diabetes mellitus (E10-E14)	901	6.3	104.5
4	Cerebrovascular diseases (I60-I69)	770	5.4	89.3
5	Chronic lower respiratory diseases (J40-J47)	505	3.5	58.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	472	3.3	54.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	350	2.4	40.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	344	2.4	39.9
9	Influenza and pneumonia (J09-J18)	304	2.1	35.3
10	Septicemia (A40-A41)	267	1.9	31.0
11	Alzheimer's disease (G30)	232	1.6	26.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	181	1.3	21.0
13	Parkinson's disease (G20-G21)	141	1.0	16.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.7	10.9
15	Viral hepatitis (B15-B19)	85	0.6	9.9
...	All other causes (residual)	2,266	15.7	262.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	16,528	100.0	2,679.8
1	Malignant neoplasms (C00-C97)	4,335	26.2	702.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,808	23.0	617.4
3	Cerebrovascular diseases (I60-I69)	1,036	6.3	168.0
4	Diabetes mellitus (E10-E14)	1,001	6.1	162.3
5	Chronic lower respiratory diseases (J40-J47)	683	4.1	110.7
6	Alzheimer's disease (G30)	511	3.1	82.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	420	2.5	68.1
8	Influenza and pneumonia (J09-J18)	417	2.5	67.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	358	2.2	58.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	322	1.9	52.2
11	Septicemia (A40-A41)	306	1.9	49.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	243	1.5	39.4
13	Parkinson's disease (G20-G21)	236	1.4	38.3
14	Pneumonitis due to solids and liquids (J69)	102	0.6	16.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	101	0.6	16.4
...	All other causes (residual)	2,649	16.0	429.5
Hispanic, both sexes, 80-84 years				
...	All causes	18,814	100.0	4,501.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,544	24.2	1,087.2
2	Malignant neoplasms (C00-C97)	3,919	20.8	937.6
3	Cerebrovascular diseases (I60-I69)	1,349	7.2	322.8
4	Diabetes mellitus (E10-E14)	976	5.2	233.5
5	Alzheimer's disease (G30)	927	4.9	221.8
6	Chronic lower respiratory diseases (J40-J47)	869	4.6	207.9
7	Influenza and pneumonia (J09-J18)	498	2.6	119.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	487	2.6	116.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	2.0	91.9
10	Parkinson's disease (G20-G21)	328	1.7	78.5
11	Septicemia (A40-A41)	327	1.7	78.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	315	1.7	75.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	232	1.2	55.5
14	Pneumonitis due to solids and liquids (J69)	139	0.7	33.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	128	0.7	30.6
...	All other causes (residual)	3,392	18.0	811.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	18,877	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,056	26.8	---
2	Malignant neoplasms (C00-C97)	2,999	15.9	---
3	Cerebrovascular diseases (I60-I69)	1,405	7.4	---
4	Alzheimer's disease (G30)	1,381	7.3	---
5	Chronic lower respiratory diseases (J40-J47)	839	4.4	---
6	Diabetes mellitus (E10-E14)	767	4.1	---
7	Influenza and pneumonia (J09-J18)	545	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	441	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	354	1.9	---
11	Parkinson's disease (G20-G21)	314	1.7	---
12	Septicemia (A40-A41)	278	1.5	---
13	Pneumonitis due to solids and liquids (J69)	153	0.8	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	123	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	112	0.6	---
...	All other causes (residual)	3,739	19.8	---
Hispanic, both sexes, 90-94 years				
...	All causes	12,741	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,676	28.9	---
2	Malignant neoplasms (C00-C97)	1,353	10.6	---
3	Alzheimer's disease (G30)	1,147	9.0	---
4	Cerebrovascular diseases (I60-I69)	946	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	544	4.3	---
6	Diabetes mellitus (E10-E14)	438	3.4	---
7	Influenza and pneumonia (J09-J18)	374	2.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	265	2.1	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	255	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	244	1.9	---
11	Parkinson's disease (G20-G21)	190	1.5	---
12	Septicemia (A40-A41)	179	1.4	---
13	Pneumonitis due to solids and liquids (J69)	114	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.7	---
15	Atherosclerosis (I70)	54	0.4	---
...	All other causes (residual)	2,875	22.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	4,498	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,360	30.2	---
2	Alzheimer's disease (G30)	458	10.2	---
3	Malignant neoplasms (C00-C97)	349	7.8	---
4	Cerebrovascular diseases (I60-I69)	340	7.6	---
5	Chronic lower respiratory diseases (J40-J47)	189	4.2	---
6	Influenza and pneumonia (J09-J18)	180	4.0	---
7	Diabetes mellitus (E10-E14)	105	2.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	95	2.1	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	92	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	1.6	---
11	Septicemia (A40-A41)	43	1.0	---
12	Pneumonitis due to solids and liquids (J69)	41	0.9	---
13	Parkinson's disease (G20-G21)	38	0.8	---
14	Atherosclerosis (I70)	22	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	---
...	All other causes (residual)	1,099	24.4	---
Hispanic, both sexes, 100 years and over				
...	All causes	1,176	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	34.9	---
2	Alzheimer's disease (G30)	104	8.8	---
3	Cerebrovascular diseases (I60-I69)	68	5.8	---
4	Influenza and pneumonia (J09-J18)	56	4.8	---
5	Malignant neoplasms (C00-C97)	52	4.4	---
6	Chronic lower respiratory diseases (J40-J47)	40	3.4	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	2.8	---
8	Diabetes mellitus (E10-E14)	24	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	1.5	---
10	Pneumonitis due to solids and liquids (J69)	14	1.2	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.8	---
12	Septicemia (A40-A41)	8	0.7	---
13	Nutritional deficiencies (E40-E64)	5	0.4	---
13	Parkinson's disease (G20-G21)	5	0.4	---
13	Atherosclerosis (I70)	5	0.4	---
...	All other causes (residual)	325	27.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	92,474	100.0	330.1
1	Malignant neoplasms (C00-C97)	19,040	20.6	68.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,884	20.4	67.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,969	9.7	32.0
4	Diabetes mellitus (E10-E14)	4,149	4.5	14.8
5	Cerebrovascular diseases (I60-I69)	4,092	4.4	14.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,911	4.2	14.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,584	2.8	9.2
8	Chronic lower respiratory diseases (J40-J47)	2,433	2.6	8.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,122	2.3	7.6
10	Influenza and pneumonia (J09-J18)	1,975	2.1	7.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,700	1.8	6.1
12	Alzheimer's disease (G30)	1,600	1.7	5.7
13	Certain conditions originating in the perinatal period (P00-P96)	1,369	1.5	4.9
14	Septicemia (A40-A41)	1,319	1.4	4.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,016	1.1	3.6
...	All other causes (residual)	17,311	18.7	61.8
Hispanic, male, 1-4 years				
...	All causes	421	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	32.8	6.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	11.2	2.2
3	Malignant neoplasms (C00-C97)	37	8.8	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	7.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.3	*
6	Influenza and pneumonia (J09-J18)	9	2.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	7	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
9	Cerebrovascular diseases (I60-I69)	5	1.2	*
10	Nutritional deficiencies (E40-E64)	3	0.7	*
11	Septicemia (A40-A41)	2	0.5	*
11	Pneumonitis due to solids and liquids (J69)	2	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
13	Aortic aneurysm and dissection (I71)	1	0.2	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	111	26.4	5.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	301	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	33.9	3.9
2	Malignant neoplasms (C00-C97)	68	22.6	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	7.3	0.8
4	Chronic lower respiratory diseases (J40-J47)	9	3.0	*
5	Septicemia (A40-A41)	8	2.7	*
6	Influenza and pneumonia (J09-J18)	6	2.0	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	2.0	*
8	Cerebrovascular diseases (I60-I69)	5	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.3	*
10	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	1.0	*
11	Nutritional deficiencies (E40-E64)	2	0.7	*
11	Meningitis (G00,G03)	2	0.7	*
13	Anemias (D50-D64)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (residual)	62	20.6	2.3
Hispanic, male, 10-14 years				
...	All causes	337	100.0	13.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	27.0	3.7
2	Malignant neoplasms (C00-C97)	64	19.0	2.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	10.4	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	6.2	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	5.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	3.3	*
7	Cerebrovascular diseases (I60-I69)	7	2.1	*
8	Influenza and pneumonia (J09-J18)	5	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	5	1.5	*
10	Septicemia (A40-A41)	3	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.9	*
10	Anemias (D50-D64)	3	0.9	*
13	Diseases of appendix (K35-K38)	2	0.6	*
14	Salmonella infections (A01-A02)	1	0.3	*
14	Meningococcal infection (A39)	1	0.3	*
14	Nutritional deficiencies (E40-E64)	1	0.3	*
14	Meningitis (G00,G03)	1	0.3	*
14	Atherosclerosis (I70)	1	0.3	*
14	Pneumonitis due to solids and liquids (J69)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (residual)	63	18.7	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,278	100.0	53.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	507	39.7	21.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	261	20.4	11.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	211	16.5	8.9
4	Malignant neoplasms (C00-C97)	85	6.7	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	2.7	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.3	*
7	Influenza and pneumonia (J09-J18)	7	0.5	*
8	Cerebrovascular diseases (I60-I69)	6	0.5	*
9	Septicemia (A40-A41)	5	0.4	*
10	Diabetes mellitus (E10-E14)	4	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.3	*
10	Legal intervention (Y35,Y89.0)	4	0.3	*
13	Aortic aneurysm and dissection (I71)	3	0.2	*
14	Tuberculosis (A16-A19)	1	0.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.1	*
14	Anemias (D50-D64)	1	0.1	*
14	Meningitis (G00,G03)	1	0.1	*
14	Pneumonitis due to solids and liquids (J69)	1	0.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.1	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.1	*
...	All other causes (residual)	122	9.5	5.1
Hispanic, male, 20-24 years				
...	All causes	2,391	100.0	96.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,065	44.5	42.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	430	18.0	17.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	352	14.7	14.1
4	Malignant neoplasms (C00-C97)	152	6.4	6.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	2.4	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.8	0.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.7	*
8	Cerebrovascular diseases (I60-I69)	15	0.6	*
8	Legal intervention (Y35,Y89.0)	15	0.6	*
10	Influenza and pneumonia (J09-J18)	12	0.5	*
11	Septicemia (A40-A41)	8	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.3	*
13	Diabetes mellitus (E10-E14)	6	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
15	Anemias (D50-D64)	4	0.2	*
15	Pneumonitis due to solids and liquids (J69)	4	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.2	*
...	All other causes (residual)	218	9.1	8.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,396	100.0	102.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	976	40.7	41.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	333	13.9	14.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	328	13.7	14.0
4	Malignant neoplasms (C00-C97)	150	6.3	6.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	5.1	5.2
6	Cerebrovascular diseases (I60-I69)	32	1.3	1.4
7	Influenza and pneumonia (J09-J18)	31	1.3	1.3
8	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.1	1.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.0	1.1
10	Legal intervention (Y35,Y89.0)	22	0.9	0.9
11	Diabetes mellitus (E10-E14)	20	0.8	0.9
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.8	*
13	Septicemia (A40-A41)	11	0.5	*
13	Chronic lower respiratory diseases (J40-J47)	11	0.5	*
15	Aortic aneurysm and dissection (I71)	8	0.3	*
...	All other causes (residual)	282	11.8	12.0
Hispanic, male, 30-34 years				
...	All causes	2,440	100.0	105.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	891	36.5	38.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	278	11.4	12.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	276	11.3	11.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	190	7.8	8.2
5	Malignant neoplasms (C00-C97)	179	7.3	7.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	89	3.6	3.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	45	1.8	1.9
8	Diabetes mellitus (E10-E14)	41	1.7	1.8
9	Cerebrovascular diseases (I60-I69)	38	1.6	1.6
10	Influenza and pneumonia (J09-J18)	33	1.4	1.4
11	Septicemia (A40-A41)	17	0.7	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	13	0.5	*
14	Aortic aneurysm and dissection (I71)	12	0.5	*
15	Legal intervention (Y35,Y89.0)	10	0.4	*
...	All other causes (residual)	311	12.7	13.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,706	100.0	127.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	765	28.3	36.0
2	Malignant neoplasms (C00-C97)	284	10.5	13.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	266	9.8	12.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	266	9.8	12.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	243	9.0	11.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	149	5.5	7.0
7	Cerebrovascular diseases (I60-I69)	70	2.6	3.3
8	Diabetes mellitus (E10-E14)	61	2.3	2.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	59	2.2	2.8
10	Influenza and pneumonia (J09-J18)	51	1.9	2.4
11	Septicemia (A40-A41)	27	1.0	1.3
12	Legal intervention (Y35,Y89.0)	21	0.8	1.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.6	*
15	Aortic aneurysm and dissection (I71)	13	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	13	0.5	*
...	All other causes (residual)	386	14.3	18.2
Hispanic, male, 40-44 years				
...	All causes	3,393	100.0	173.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	790	23.3	40.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	490	14.4	25.0
3	Malignant neoplasms (C00-C97)	420	12.4	21.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	260	7.7	13.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	231	6.8	11.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	162	4.8	8.3
7	Cerebrovascular diseases (I60-I69)	123	3.6	6.3
8	Diabetes mellitus (E10-E14)	100	2.9	5.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	78	2.3	4.0
10	Influenza and pneumonia (J09-J18)	71	2.1	3.6
11	Septicemia (A40-A41)	52	1.5	2.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	1.2	2.0
13	Viral hepatitis (B15-B19)	28	0.8	1.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	0.8	1.4
15	Aortic aneurysm and dissection (I71)	17	0.5	*
...	All other causes (residual)	503	14.8	25.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,514	100.0	261.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	801	17.7	46.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	745	16.5	43.2
3	Malignant neoplasms (C00-C97)	727	16.1	42.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	428	9.5	24.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	225	5.0	13.0
6	Diabetes mellitus (E10-E14)	195	4.3	11.3
7	Cerebrovascular diseases (I60-I69)	172	3.8	10.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	113	2.5	6.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	2.2	5.8
10	Viral hepatitis (B15-B19)	71	1.6	4.1
10	Influenza and pneumonia (J09-J18)	71	1.6	4.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	1.3	3.3
13	Septicemia (A40-A41)	51	1.1	3.0
14	Chronic lower respiratory diseases (J40-J47)	37	0.8	2.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.7	1.9
...	All other causes (residual)	689	15.3	40.0
Hispanic, male, 50-54 years				
...	All causes	6,380	100.0	434.0
1	Malignant neoplasms (C00-C97)	1,353	21.2	92.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,244	19.5	84.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	667	10.5	45.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	658	10.3	44.8
5	Diabetes mellitus (E10-E14)	320	5.0	21.8
6	Cerebrovascular diseases (I60-I69)	253	4.0	17.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	2.7	11.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	158	2.5	10.7
9	Viral hepatitis (B15-B19)	144	2.3	9.8
10	Influenza and pneumonia (J09-J18)	105	1.6	7.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	102	1.6	6.9
12	Septicemia (A40-A41)	101	1.6	6.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	1.6	6.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	63	1.0	4.3
15	Chronic lower respiratory diseases (J40-J47)	59	0.9	4.0
...	All other causes (residual)	879	13.8	59.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	7,392	100.0	651.4
1	Malignant neoplasms (C00-C97)	1,819	24.6	160.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,516	20.5	133.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	716	9.7	63.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	536	7.3	47.2
5	Diabetes mellitus (E10-E14)	423	5.7	37.3
6	Cerebrovascular diseases (I60-I69)	306	4.1	27.0
7	Viral hepatitis (B15-B19)	185	2.5	16.3
8	Influenza and pneumonia (J09-J18)	139	1.9	12.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	139	1.9	12.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	1.6	10.3
11	Septicemia (A40-A41)	116	1.6	10.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	104	1.4	9.2
13	Chronic lower respiratory diseases (J40-J47)	103	1.4	9.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	1.0	6.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	0.7	4.7
...	All other causes (residual)	1,047	14.2	92.3
Hispanic, male, 60-64 years				
...	All causes	8,006	100.0	976.2
1	Malignant neoplasms (C00-C97)	2,243	28.0	273.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,798	22.5	219.2
3	Chronic liver disease and cirrhosis (K70,K73-K74)	605	7.6	73.8
4	Diabetes mellitus (E10-E14)	487	6.1	59.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	388	4.8	47.3
6	Cerebrovascular diseases (I60-I69)	343	4.3	41.8
7	Influenza and pneumonia (J09-J18)	162	2.0	19.8
8	Viral hepatitis (B15-B19)	160	2.0	19.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	1.9	18.8
10	Septicemia (A40-A41)	142	1.8	17.3
11	Chronic lower respiratory diseases (J40-J47)	137	1.7	16.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	102	1.3	12.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	96	1.2	11.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	62	0.8	7.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	0.4	4.3
...	All other causes (residual)	1,092	13.6	133.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	7,926	100.0	1,355.6
1	Malignant neoplasms (C00-C97)	2,394	30.2	409.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,864	23.5	318.8
3	Diabetes mellitus (E10-E14)	514	6.5	87.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	376	4.7	64.3
5	Cerebrovascular diseases (I60-I69)	326	4.1	55.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	283	3.6	48.4
7	Chronic lower respiratory diseases (J40-J47)	226	2.9	38.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	2.6	35.6
9	Influenza and pneumonia (J09-J18)	157	2.0	26.9
10	Septicemia (A40-A41)	124	1.6	21.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	108	1.4	18.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	0.9	12.5
13	Viral hepatitis (B15-B19)	72	0.9	12.3
14	Alzheimer's disease (G30)	41	0.5	7.0
15	Parkinson's disease (G20-G21)	38	0.5	6.5
...	All other causes (residual)	1,122	14.2	191.9
Hispanic, male, 70-74 years				
...	All causes	8,022	100.0	2,117.2
1	Malignant neoplasms (C00-C97)	2,404	30.0	634.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,909	23.8	503.8
3	Diabetes mellitus (E10-E14)	472	5.9	124.6
4	Cerebrovascular diseases (I60-I69)	416	5.2	109.8
5	Chronic lower respiratory diseases (J40-J47)	297	3.7	78.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	270	3.4	71.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	2.7	57.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	171	2.1	45.1
9	Influenza and pneumonia (J09-J18)	166	2.1	43.8
10	Septicemia (A40-A41)	150	1.9	39.6
11	Alzheimer's disease (G30)	103	1.3	27.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	95	1.2	25.1
13	Parkinson's disease (G20-G21)	85	1.1	22.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.7	15.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	0.7	14.3
...	All other causes (residual)	1,154	14.4	304.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	8,594	100.0	3,300.2
1	Malignant neoplasms (C00-C97)	2,336	27.2	897.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,075	24.1	796.8
3	Diabetes mellitus (E10-E14)	511	5.9	196.2
4	Cerebrovascular diseases (I60-I69)	490	5.7	188.2
5	Chronic lower respiratory diseases (J40-J47)	362	4.2	139.0
6	Alzheimer's disease (G30)	220	2.6	84.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	2.5	83.7
8	Influenza and pneumonia (J09-J18)	211	2.5	81.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	2.4	78.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	165	1.9	63.4
11	Septicemia (A40-A41)	143	1.7	54.9
12	Parkinson's disease (G20-G21)	139	1.6	53.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	105	1.2	40.3
14	Pneumonitis due to solids and liquids (J69)	61	0.7	23.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.7	23.0
...	All other causes (residual)	1,294	15.1	496.9
Hispanic, male, 80-84 years				
...	All causes	8,950	100.0	5,354.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,285	25.5	1,367.0
2	Malignant neoplasms (C00-C97)	2,053	22.9	1,228.2
3	Cerebrovascular diseases (I60-I69)	561	6.3	335.6
4	Diabetes mellitus (E10-E14)	456	5.1	272.8
5	Chronic lower respiratory diseases (J40-J47)	424	4.7	253.7
6	Alzheimer's disease (G30)	324	3.6	193.8
7	Influenza and pneumonia (J09-J18)	241	2.7	144.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	231	2.6	138.2
9	Parkinson's disease (G20-G21)	195	2.2	116.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	2.1	113.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	155	1.7	92.7
12	Septicemia (A40-A41)	141	1.6	84.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	94	1.1	56.2
14	Pneumonitis due to solids and liquids (J69)	83	0.9	49.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.7	40.1
...	All other causes (residual)	1,450	16.2	867.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	8,060	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,260	28.0	---
2	Malignant neoplasms (C00-C97)	1,459	18.1	---
3	Cerebrovascular diseases (I60-I69)	515	6.4	---
4	Alzheimer's disease (G30)	434	5.4	---
5	Chronic lower respiratory diseases (J40-J47)	395	4.9	---
6	Diabetes mellitus (E10-E14)	334	4.1	---
7	Influenza and pneumonia (J09-J18)	237	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	2.4	---
10	Parkinson's disease (G20-G21)	159	2.0	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	120	1.5	---
12	Septicemia (A40-A41)	117	1.5	---
13	Pneumonitis due to solids and liquids (J69)	87	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.7	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	48	0.6	---
...	All other causes (residual)	1,443	17.9	---
Hispanic, male, 90-94 years				
...	All causes	4,758	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,418	29.8	---
2	Malignant neoplasms (C00-C97)	652	13.7	---
3	Alzheimer's disease (G30)	331	7.0	---
4	Cerebrovascular diseases (I60-I69)	306	6.4	---
5	Chronic lower respiratory diseases (J40-J47)	219	4.6	---
6	Diabetes mellitus (E10-E14)	173	3.6	---
7	Influenza and pneumonia (J09-J18)	154	3.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	93	2.0	---
11	Parkinson's disease (G20-G21)	83	1.7	---
12	Septicemia (A40-A41)	63	1.3	---
13	Pneumonitis due to solids and liquids (J69)	62	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.7	---
15	Atherosclerosis (I70)	22	0.5	---
...	All other causes (residual)	935	19.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	1,319	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	409	31.0	---
2	Malignant neoplasms (C00-C97)	134	10.2	---
3	Alzheimer's disease (G30)	98	7.4	---
4	Chronic lower respiratory diseases (J40-J47)	83	6.3	---
5	Cerebrovascular diseases (I60-I69)	82	6.2	---
6	Influenza and pneumonia (J09-J18)	68	5.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	2.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	2.5	---
9	Diabetes mellitus (E10-E14)	26	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.5	---
11	Parkinson's disease (G20-G21)	18	1.4	---
12	Septicemia (A40-A41)	14	1.1	---
12	Pneumonitis due to solids and liquids (J69)	14	1.1	---
14	Atherosclerosis (I70)	7	0.5	---
15	Anemias (D50-D64)	4	0.3	---
15	Hyperplasia of prostate (N40)	4	0.3	---
...	All other causes (residual)	272	20.6	---
Hispanic, male, 100 years and over				
...	All causes	258	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	31.4	---
2	Malignant neoplasms (C00-C97)	22	8.5	---
3	Influenza and pneumonia (J09-J18)	19	7.4	---
4	Alzheimer's disease (G30)	14	5.4	---
5	Cerebrovascular diseases (I60-I69)	13	5.0	---
6	Chronic lower respiratory diseases (J40-J47)	10	3.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	2.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	2.3	---
9	Diabetes mellitus (E10-E14)	5	1.9	---
10	Pneumonitis due to solids and liquids (J69)	4	1.6	---
11	Parkinson's disease (G20-G21)	3	1.2	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.2	---
13	Septicemia (A40-A41)	1	0.4	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	---
13	Anemias (D50-D64)	1	0.4	---
13	Nutritional deficiencies (E40-E64)	1	0.4	---
13	Atherosclerosis (I70)	1	0.4	---
13	Aortic aneurysm and dissection (I71)	1	0.4	---
13	Hernia (K40-K46)	1	0.4	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	---
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.4	---
...	All other causes (residual)	63	24.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	76,913	100.0	281.0
1	Malignant neoplasms (C00-C97)	17,407	22.6	63.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,137	19.7	55.3
3	Cerebrovascular diseases (I60-I69)	4,621	6.0	16.9
4	Diabetes mellitus (E10-E14)	3,646	4.7	13.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,444	4.5	12.6
6	Alzheimer's disease (G30)	3,334	4.3	12.2
7	Chronic lower respiratory diseases (J40-J47)	2,362	3.1	8.6
8	Influenza and pneumonia (J09-J18)	1,900	2.5	6.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,747	2.3	6.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,573	2.0	5.7
11	Septicemia (A40-A41)	1,294	1.7	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,123	1.5	4.1
13	Certain conditions originating in the perinatal period (P00-P96)	1,053	1.4	3.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	873	1.1	3.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	662	0.9	2.4
...	All other causes (residual)	16,737	21.8	61.2
Hispanic, female, 1-4 years				
...	All causes	348	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	23.9	4.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	14.4	2.5
3	Malignant neoplasms (C00-C97)	44	12.6	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	11.2	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	4.3	*
6	Influenza and pneumonia (J09-J18)	11	3.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	2.0	*
8	Meningitis (G00,G03)	4	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.1	*
10	Septicemia (A40-A41)	3	0.9	*
10	Acute bronchitis and bronchiolitis (J20-J21)	3	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	3	0.9	*
13	Cerebrovascular diseases (I60-I69)	2	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
15	Anemias (D50-D64)	1	0.3	*
15	Diabetes mellitus (E10-E14)	1	0.3	*
15	Nutritional deficiencies (E40-E64)	1	0.3	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.3	*
15	Hernia (K40-K46)	1	0.3	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	71	20.4	3.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	230	100.0	9.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	28.3	2.6
2	Malignant neoplasms (C00-C97)	50	21.7	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	8.7	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	6.5	*
5	Cerebrovascular diseases (I60-I69)	10	4.3	*
5	Influenza and pneumonia (J09-J18)	10	4.3	*
7	Chronic lower respiratory diseases (J40-J47)	6	2.6	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.2	*
9	Septicemia (A40-A41)	3	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	*
10	Hernia (K40-K46)	2	0.9	*
12	Diabetes mellitus (E10-E14)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (residual)	40	17.4	1.6
Hispanic, female, 10-14 years				
...	All causes	248	100.0	10.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	22.2	2.3
2	Malignant neoplasms (C00-C97)	37	14.9	1.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	12.5	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	6.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	6.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.2	*
7	Certain conditions originating in the perinatal period (P00-P96)	7	2.8	*
8	Cerebrovascular diseases (I60-I69)	5	2.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	4	1.6	*
11	Diabetes mellitus (E10-E14)	3	1.2	*
12	Septicemia (A40-A41)	2	0.8	*
12	Influenza and pneumonia (J09-J18)	2	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.8	*
15	Meningococcal infection (A39)	1	0.4	*
15	Anemias (D50-D64)	1	0.4	*
15	Meningitis (G00,G03)	1	0.4	*
15	Diseases of appendix (K35-K38)	1	0.4	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
...	All other causes (residual)	49	19.8	2.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	489	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	36.4	7.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	13.9	3.0
3	Malignant neoplasms (C00-C97)	51	10.4	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	8.8	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.9	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	3.9	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	7	1.4	*
8	Anemias (D50-D64)	3	0.6	*
8	Cerebrovascular diseases (I60-I69)	3	0.6	*
8	Pneumonitis due to solids and liquids (J69)	3	0.6	*
11	Septicemia (A40-A41)	2	0.4	*
11	Diabetes mellitus (E10-E14)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Meningococcal infection (A39)	1	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
15	Influenza and pneumonia (J09-J18)	1	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
15	Legal intervention (Y35,Y89.0)	1	0.2	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	79	16.2	3.5
Hispanic, female, 20-24 years				
...	All causes	747	100.0	32.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	294	39.4	12.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	89	11.9	3.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	10.4	3.4
4	Malignant neoplasms (C00-C97)	76	10.2	3.3
5	Pregnancy, childbirth and the puerperium (O00-O99)	22	2.9	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.3	*
7	Influenza and pneumonia (J09-J18)	11	1.5	*
8	Cerebrovascular diseases (I60-I69)	10	1.3	*
9	Septicemia (A40-A41)	9	1.2	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	8	1.1	*
12	Diabetes mellitus (E10-E14)	6	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (residual)	106	14.2	4.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	849	100.0	40.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	266	31.3	12.7
2	Malignant neoplasms (C00-C97)	121	14.3	5.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	10.4	4.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	6.1	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	3.7	1.5
5	Pregnancy, childbirth and the puerperium (O00-O99)	31	3.7	1.5
7	Cerebrovascular diseases (I60-I69)	18	2.1	*
8	Influenza and pneumonia (J09-J18)	15	1.8	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	1.8	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	13	1.5	*
11	Septicemia (A40-A41)	12	1.4	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.4	*
11	Diabetes mellitus (E10-E14)	12	1.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
14	Chronic lower respiratory diseases (J40-J47)	7	0.8	*
...	All other causes (residual)	149	17.6	7.1
Hispanic, female, 30-34 years				
...	All causes	890	100.0	42.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	23.6	9.9
2	Malignant neoplasms (C00-C97)	181	20.3	8.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	5.7	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	5.7	2.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	5.4	2.3
6	Diabetes mellitus (E10-E14)	28	3.1	1.3
7	Pregnancy, childbirth and the puerperium (O00-O99)	25	2.8	1.2
8	Cerebrovascular diseases (I60-I69)	24	2.7	1.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	24	2.7	1.1
10	Influenza and pneumonia (J09-J18)	23	2.6	1.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	16	1.8	*
12	Septicemia (A40-A41)	13	1.5	*
13	Chronic lower respiratory diseases (J40-J47)	11	1.2	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.0	*
...	All other causes (residual)	166	18.7	7.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,266	100.0	63.0
1	Malignant neoplasms (C00-C97)	386	30.5	19.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	16.0	10.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	7.7	4.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	3.9	2.5
5	Cerebrovascular diseases (I60-I69)	41	3.2	2.0
6	Diabetes mellitus (E10-E14)	40	3.2	2.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	3.2	2.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	37	2.9	1.8
9	Influenza and pneumonia (J09-J18)	34	2.7	1.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	32	2.5	1.6
11	Septicemia (A40-A41)	23	1.8	1.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.3	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
...	All other causes (residual)	230	18.2	11.4
Hispanic, female, 40-44 years				
...	All causes	1,803	100.0	94.8
1	Malignant neoplasms (C00-C97)	608	33.7	32.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	222	12.3	11.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	10.0	9.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	87	4.8	4.6
5	Cerebrovascular diseases (I60-I69)	84	4.7	4.4
6	Diabetes mellitus (E10-E14)	58	3.2	3.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	3.2	3.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	2.5	2.4
9	Influenza and pneumonia (J09-J18)	37	2.1	1.9
10	Pregnancy, childbirth and the puerperium (O00-O99)	29	1.6	1.5
11	Septicemia (A40-A41)	21	1.2	1.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.2	1.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	19	1.1	*
14	Viral hepatitis (B15-B19)	12	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	0.7	*
...	All other causes (residual)	310	17.2	16.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,437	100.0	146.3
1	Malignant neoplasms (C00-C97)	891	36.6	53.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	258	10.6	15.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	9.4	13.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	129	5.3	7.7
5	Cerebrovascular diseases (I60-I69)	106	4.3	6.4
6	Diabetes mellitus (E10-E14)	102	4.2	6.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	2.3	3.4
8	Influenza and pneumonia (J09-J18)	50	2.1	3.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	1.5	2.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	1.2	1.7
11	Septicemia (A40-A41)	27	1.1	1.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.1	1.6
12	Chronic lower respiratory diseases (J40-J47)	26	1.1	1.6
14	Viral hepatitis (B15-B19)	22	0.9	1.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.8	1.2
...	All other causes (residual)	429	17.6	25.8
Hispanic, female, 50-54 years				
...	All causes	3,374	100.0	229.4
1	Malignant neoplasms (C00-C97)	1,262	37.4	85.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	411	12.2	27.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	7.1	16.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	198	5.9	13.5
5	Diabetes mellitus (E10-E14)	163	4.8	11.1
6	Cerebrovascular diseases (I60-I69)	135	4.0	9.2
7	Septicemia (A40-A41)	68	2.0	4.6
8	Viral hepatitis (B15-B19)	62	1.8	4.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	1.6	3.7
10	Influenza and pneumonia (J09-J18)	54	1.6	3.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	1.6	3.7
12	Chronic lower respiratory diseases (J40-J47)	37	1.1	2.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	34	1.0	2.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	0.9	2.1
15	Pregnancy, childbirth and the puerperium (O00-O99)	17	0.5	*
...	All other causes (residual)	554	16.4	37.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	4,245	100.0	357.0
1	Malignant neoplasms (C00-C97)	1,605	37.8	135.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	584	13.8	49.1
3	Diabetes mellitus (E10-E14)	247	5.8	20.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	229	5.4	19.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	4.0	14.4
6	Cerebrovascular diseases (I60-I69)	158	3.7	13.3
7	Septicemia (A40-A41)	104	2.4	8.7
8	Influenza and pneumonia (J09-J18)	101	2.4	8.5
9	Chronic lower respiratory diseases (J40-J47)	81	1.9	6.8
10	Viral hepatitis (B15-B19)	72	1.7	6.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	1.7	6.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	1.2	4.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	1.0	3.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	0.6	2.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.4	*
...	All other causes (residual)	669	15.8	56.3
Hispanic, female, 60-64 years				
...	All causes	5,058	100.0	556.2
1	Malignant neoplasms (C00-C97)	1,821	36.0	200.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	890	17.6	97.9
3	Diabetes mellitus (E10-E14)	339	6.7	37.3
4	Cerebrovascular diseases (I60-I69)	203	4.0	22.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	194	3.8	21.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	3.1	17.0
7	Influenza and pneumonia (J09-J18)	134	2.6	14.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	2.6	14.2
9	Chronic lower respiratory diseases (J40-J47)	116	2.3	12.8
10	Septicemia (A40-A41)	100	2.0	11.0
11	Viral hepatitis (B15-B19)	63	1.2	6.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	0.9	4.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.6	3.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	0.5	2.7
15	Pneumonitis due to solids and liquids (J69)	22	0.4	2.4
...	All other causes (residual)	792	15.7	87.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	5,721	100.0	837.5
1	Malignant neoplasms (C00-C97)	1,994	34.9	291.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	974	17.0	142.6
3	Diabetes mellitus (E10-E14)	409	7.1	59.9
4	Cerebrovascular diseases (I60-I69)	284	5.0	41.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	223	3.9	32.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	174	3.0	25.5
7	Chronic lower respiratory diseases (J40-J47)	164	2.9	24.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	2.3	19.3
9	Influenza and pneumonia (J09-J18)	124	2.2	18.2
10	Septicemia (A40-A41)	112	2.0	16.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	1.3	11.1
12	Alzheimer's disease (G30)	63	1.1	9.2
13	Viral hepatitis (B15-B19)	42	0.7	6.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.4	3.4
15	Parkinson's disease (G20-G21)	22	0.4	3.2
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	22	0.4	3.2
...	All other causes (residual)	883	15.4	129.3
Hispanic, female, 70-74 years				
...	All causes	6,367	100.0	1,318.4
1	Malignant neoplasms (C00-C97)	1,923	30.2	398.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,241	19.5	257.0
3	Diabetes mellitus (E10-E14)	429	6.7	88.8
4	Cerebrovascular diseases (I60-I69)	354	5.6	73.3
5	Chronic lower respiratory diseases (J40-J47)	208	3.3	43.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	202	3.2	41.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	179	2.8	37.1
8	Influenza and pneumonia (J09-J18)	138	2.2	28.6
9	Alzheimer's disease (G30)	129	2.0	26.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	2.0	25.9
11	Septicemia (A40-A41)	117	1.8	24.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	86	1.4	17.8
13	Parkinson's disease (G20-G21)	56	0.9	11.6
14	Viral hepatitis (B15-B19)	45	0.7	9.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.6	7.7
...	All other causes (residual)	1,098	17.2	227.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	7,934	100.0	2,226.5
1	Malignant neoplasms (C00-C97)	1,999	25.2	561.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,733	21.8	486.3
3	Cerebrovascular diseases (I60-I69)	546	6.9	153.2
4	Diabetes mellitus (E10-E14)	490	6.2	137.5
5	Chronic lower respiratory diseases (J40-J47)	321	4.0	90.1
6	Alzheimer's disease (G30)	291	3.7	81.7
7	Influenza and pneumonia (J09-J18)	206	2.6	57.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	2.5	56.7
9	Septicemia (A40-A41)	163	2.1	45.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	157	2.0	44.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	1.9	43.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	138	1.7	38.7
13	Parkinson's disease (G20-G21)	97	1.2	27.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.5	11.5
14	Pneumonitis due to solids and liquids (J69)	41	0.5	11.5
...	All other causes (residual)	1,355	17.1	380.2
Hispanic, female, 80-84 years				
...	All causes	9,864	100.0	3,932.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,259	22.9	900.7
2	Malignant neoplasms (C00-C97)	1,866	18.9	744.0
3	Cerebrovascular diseases (I60-I69)	788	8.0	314.2
4	Alzheimer's disease (G30)	603	6.1	240.4
5	Diabetes mellitus (E10-E14)	520	5.3	207.3
6	Chronic lower respiratory diseases (J40-J47)	445	4.5	177.4
7	Influenza and pneumonia (J09-J18)	257	2.6	102.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	256	2.6	102.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	2.0	77.3
10	Septicemia (A40-A41)	186	1.9	74.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	160	1.6	63.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	138	1.4	55.0
13	Parkinson's disease (G20-G21)	133	1.3	53.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.6	24.3
15	Pneumonitis due to solids and liquids (J69)	56	0.6	22.3
...	All other causes (residual)	1,942	19.7	774.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	10,817	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,796	25.8	---
2	Malignant neoplasms (C00-C97)	1,540	14.2	---
3	Alzheimer's disease (G30)	947	8.8	---
4	Cerebrovascular diseases (I60-I69)	890	8.2	---
5	Chronic lower respiratory diseases (J40-J47)	444	4.1	---
6	Diabetes mellitus (E10-E14)	433	4.0	---
7	Influenza and pneumonia (J09-J18)	308	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	238	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	234	2.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	1.7	---
11	Septicemia (A40-A41)	161	1.5	---
12	Parkinson's disease (G20-G21)	155	1.4	---
13	Chronic liver disease and cirrhosis (K70,K73-K74)	75	0.7	---
14	Pneumonitis due to solids and liquids (J69)	66	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.5	---
...	All other causes (residual)	2,296	21.2	---
Hispanic, female, 90-94 years				
...	All causes	7,983	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,258	28.3	---
2	Alzheimer's disease (G30)	816	10.2	---
3	Malignant neoplasms (C00-C97)	701	8.8	---
4	Cerebrovascular diseases (I60-I69)	640	8.0	---
5	Chronic lower respiratory diseases (J40-J47)	325	4.1	---
6	Diabetes mellitus (E10-E14)	265	3.3	---
7	Influenza and pneumonia (J09-J18)	220	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	172	2.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	1.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	1.6	---
11	Septicemia (A40-A41)	116	1.5	---
12	Parkinson's disease (G20-G21)	107	1.3	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.7	---
14	Pneumonitis due to solids and liquids (J69)	52	0.7	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	34	0.4	---
...	All other causes (residual)	1,938	24.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	3,179	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	951	29.9	---
2	Alzheimer's disease (G30)	360	11.3	---
3	Cerebrovascular diseases (I60-I69)	258	8.1	---
4	Malignant neoplasms (C00-C97)	215	6.8	---
5	Influenza and pneumonia (J09-J18)	112	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	106	3.3	---
7	Diabetes mellitus (E10-E14)	79	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	75	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	1.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	1.3	---
11	Septicemia (A40-A41)	29	0.9	---
12	Pneumonitis due to solids and liquids (J69)	27	0.8	---
13	Parkinson's disease (G20-G21)	20	0.6	---
14	Atherosclerosis (I70)	15	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.4	---
...	All other causes (residual)	821	25.8	---
Hispanic, female, 100 years and over				
...	All causes	918	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	329	35.8	---
2	Alzheimer's disease (G30)	90	9.8	---
3	Cerebrovascular diseases (I60-I69)	55	6.0	---
4	Influenza and pneumonia (J09-J18)	37	4.0	---
5	Malignant neoplasms (C00-C97)	30	3.3	---
5	Chronic lower respiratory diseases (J40-J47)	30	3.3	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	2.9	---
8	Diabetes mellitus (E10-E14)	19	2.1	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	1.3	---
10	Pneumonitis due to solids and liquids (J69)	10	1.1	---
11	Septicemia (A40-A41)	7	0.8	---
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.7	---
13	Nutritional deficiencies (E40-E64)	4	0.4	---
13	Atherosclerosis (I70)	4	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.2	---
15	Parkinson's disease (G20-G21)	2	0.2	---
...	All other causes (residual)	254	27.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,066,949	100.0	1,028.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	489,926	23.7	243.7
2	Malignant neoplasms (C00-C97)	466,271	22.6	231.9
3	Chronic lower respiratory diseases (J40-J47)	129,519	6.3	64.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104,682	5.1	52.1
5	Cerebrovascular diseases (I60-I69)	102,326	5.0	50.9
6	Alzheimer's disease (G30)	80,014	3.9	39.8
7	Diabetes mellitus (E10-E14)	51,944	2.5	25.8
8	Influenza and pneumonia (J09-J18)	43,377	2.1	21.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35,444	1.7	17.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,691	1.7	17.3
11	Septicemia (A40-A41)	28,868	1.4	14.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27,830	1.3	13.8
13	Parkinson's disease (G20-G21)	22,917	1.1	11.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21,511	1.0	10.7
15	Pneumonitis due to solids and liquids (J69)	15,644	0.8	7.8
...	All other causes (residual)	411,985	19.9	204.9
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,876	100.0	22.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	655	34.9	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	9.6	2.2
3	Malignant neoplasms (C00-C97)	176	9.4	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	7.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	3.0	0.7
6	Influenza and pneumonia (J09-J18)	53	2.8	0.6
7	Septicemia (A40-A41)	32	1.7	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.1	0.2
9	Chronic lower respiratory diseases (J40-J47)	18	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	14	0.7	*
11	Cerebrovascular diseases (I60-I69)	13	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
13	Meningitis (G00,G03)	8	0.4	*
14	Anemias (D50-D64)	6	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.3	*
...	All other causes (residual)	493	26.3	5.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-9 years				
...	All causes	1,176	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	376	32.0	3.5
2	Malignant neoplasms (C00-C97)	229	19.5	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	8.6	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	4.4	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.2	0.3
6	Influenza and pneumonia (J09-J18)	27	2.3	0.2
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.6	*
8	Cerebrovascular diseases (I60-I69)	17	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	13	1.1	*
10	Septicemia (A40-A41)	11	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	11	0.9	*
12	Diabetes mellitus (E10-E14)	4	0.3	*
13	Anemias (D50-D64)	3	0.3	*
13	Pneumonitis due to solids and liquids (J69)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (residual)	269	22.9	2.5
Non-Hispanic, white, both sexes, 10-14 years				
...	All causes	1,508	100.0	13.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	393	26.1	3.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	284	18.8	2.5
3	Malignant neoplasms (C00-C97)	222	14.7	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	5.0	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	3.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	3.4	0.5
7	Cerebrovascular diseases (I60-I69)	25	1.7	0.2
7	Influenza and pneumonia (J09-J18)	25	1.7	0.2
9	Chronic lower respiratory diseases (J40-J47)	24	1.6	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.3	*
11	Septicemia (A40-A41)	15	1.0	*
12	Diabetes mellitus (E10-E14)	8	0.5	*
13	Anemias (D50-D64)	5	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.3	*
...	All other causes (residual)	299	19.8	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-19 years				
...	All causes	5,249	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,335	44.5	19.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,290	24.6	10.9
3	Malignant neoplasms (C00-C97)	344	6.6	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	181	3.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	2.8	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	2.1	0.9
7	Influenza and pneumonia (J09-J18)	46	0.9	0.4
8	Cerebrovascular diseases (I60-I69)	33	0.6	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.5	0.2
10	Chronic lower respiratory diseases (J40-J47)	25	0.5	0.2
11	Diabetes mellitus (E10-E14)	21	0.4	0.2
12	Septicemia (A40-A41)	20	0.4	0.2
13	Pregnancy, childbirth and the puerperium (O00-O99)	13	0.2	*
14	Legal intervention (Y35,Y89.0)	9	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.1	*
...	All other causes (residual)	643	12.2	5.4
Non-Hispanic, white, both sexes, 20-24 years				
...	All causes	10,791	100.0	83.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,408	50.1	41.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,213	20.5	17.1
3	Malignant neoplasms (C00-C97)	515	4.8	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	426	3.9	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	348	3.2	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	1.0	0.9
7	Diabetes mellitus (E10-E14)	80	0.7	0.6
8	Influenza and pneumonia (J09-J18)	77	0.7	0.6
9	Pregnancy, childbirth and the puerperium (O00-O99)	58	0.5	0.4
10	Cerebrovascular diseases (I60-I69)	55	0.5	0.4
11	Chronic lower respiratory diseases (J40-J47)	50	0.5	0.4
12	Septicemia (A40-A41)	45	0.4	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.2	0.2
14	Pneumonitis due to solids and liquids (J69)	21	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.2	*
...	All other causes (residual)	1,339	12.4	10.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 25-29 years				
...	All causes	13,178	100.0	103.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,145	46.6	48.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,380	18.1	18.7
3	Malignant neoplasms (C00-C97)	793	6.0	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	603	4.6	4.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	489	3.7	3.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	1.1	1.1
7	Influenza and pneumonia (J09-J18)	130	1.0	1.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	123	0.9	1.0
9	Diabetes mellitus (E10-E14)	118	0.9	0.9
10	Cerebrovascular diseases (I60-I69)	104	0.8	0.8
11	Chronic lower respiratory diseases (J40-J47)	88	0.7	0.7
12	Pregnancy, childbirth and the puerperium (O00-O99)	80	0.6	0.6
13	Septicemia (A40-A41)	77	0.6	0.6
14	Legal intervention (Y35,Y89.0)	46	0.3	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	0.3	0.3
...	All other causes (residual)	1,818	13.8	14.3
Non-Hispanic, white, both sexes, 30-34 years				
...	All causes	15,791	100.0	126.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,203	39.3	49.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,528	16.0	20.2
3	Malignant neoplasms (C00-C97)	1,351	8.6	10.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,191	7.5	9.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	491	3.1	3.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	318	2.0	2.5
7	Diabetes mellitus (E10-E14)	219	1.4	1.8
8	Influenza and pneumonia (J09-J18)	192	1.2	1.5
9	Cerebrovascular diseases (I60-I69)	177	1.1	1.4
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	0.9	1.2
11	Septicemia (A40-A41)	128	0.8	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	108	0.7	0.9
13	Chronic lower respiratory diseases (J40-J47)	104	0.7	0.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	76	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	0.5	0.6
...	All other causes (residual)	2,486	15.7	19.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-39 years				
...	All causes	18,060	100.0	156.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,704	31.6	49.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,503	13.9	21.7
3	Malignant neoplasms (C00-C97)	2,269	12.6	19.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,995	11.0	17.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	506	2.8	4.4
6	Diabetes mellitus (E10-E14)	390	2.2	3.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	379	2.1	3.3
8	Cerebrovascular diseases (I60-I69)	294	1.6	2.5
9	Influenza and pneumonia (J09-J18)	240	1.3	2.1
10	Septicemia (A40-A41)	179	1.0	1.6
11	Chronic lower respiratory diseases (J40-J47)	143	0.8	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	0.6	1.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	0.6	1.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	108	0.6	0.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	101	0.6	0.9
...	All other causes (residual)	3,022	16.7	26.2
Non-Hispanic, white, both sexes, 40-44 years				
...	All causes	26,505	100.0	213.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,887	22.2	47.3
2	Malignant neoplasms (C00-C97)	4,665	17.6	37.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,114	15.5	33.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,869	10.8	23.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,138	4.3	9.1
6	Diabetes mellitus (E10-E14)	703	2.7	5.7
7	Cerebrovascular diseases (I60-I69)	524	2.0	4.2
8	Influenza and pneumonia (J09-J18)	434	1.6	3.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	420	1.6	3.4
10	Chronic lower respiratory diseases (J40-J47)	347	1.3	2.8
11	Septicemia (A40-A41)	303	1.1	2.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	226	0.9	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	0.7	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	0.6	1.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	119	0.4	1.0
...	All other causes (residual)	4,421	16.7	35.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-49 years				
...	All causes	43,261	100.0	322.8
1	Malignant neoplasms (C00-C97)	9,723	22.5	72.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,775	18.0	58.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,854	15.8	51.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,515	8.1	26.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,146	5.0	16.0
6	Diabetes mellitus (E10-E14)	1,260	2.9	9.4
7	Cerebrovascular diseases (I60-I69)	1,008	2.3	7.5
8	Chronic lower respiratory diseases (J40-J47)	851	2.0	6.4
9	Influenza and pneumonia (J09-J18)	700	1.6	5.2
10	Septicemia (A40-A41)	554	1.3	4.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	452	1.0	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	354	0.8	2.6
13	Viral hepatitis (B15-B19)	340	0.8	2.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	316	0.7	2.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	252	0.6	1.9
...	All other causes (residual)	7,161	16.6	53.4
Non-Hispanic, white, both sexes, 50-54 years				
...	All causes	77,418	100.0	500.1
1	Malignant neoplasms (C00-C97)	21,409	27.7	138.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,387	19.9	99.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,668	11.2	56.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,079	5.3	26.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,964	5.1	25.6
6	Chronic lower respiratory diseases (J40-J47)	2,662	3.4	17.2
7	Diabetes mellitus (E10-E14)	2,310	3.0	14.9
8	Cerebrovascular diseases (I60-I69)	1,901	2.5	12.3
9	Influenza and pneumonia (J09-J18)	1,137	1.5	7.3
10	Septicemia (A40-A41)	1,043	1.3	6.7
11	Viral hepatitis (B15-B19)	873	1.1	5.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	690	0.9	4.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	513	0.7	3.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	444	0.6	2.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	430	0.6	2.8
...	All other causes (residual)	11,908	15.4	76.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-59 years				
...	All causes	112,289	100.0	728.5
1	Malignant neoplasms (C00-C97)	36,530	32.5	237.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,150	20.6	150.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,762	6.9	50.4
4	Chronic lower respiratory diseases (J40-J47)	5,465	4.9	35.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,846	4.3	31.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,847	3.4	25.0
7	Diabetes mellitus (E10-E14)	3,495	3.1	22.7
8	Cerebrovascular diseases (I60-I69)	3,007	2.7	19.5
9	Septicemia (A40-A41)	1,662	1.5	10.8
10	Influenza and pneumonia (J09-J18)	1,657	1.5	10.8
11	Viral hepatitis (B15-B19)	1,321	1.2	8.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,150	1.0	7.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	812	0.7	5.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	488	0.4	3.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	467	0.4	3.0
...	All other causes (residual)	16,630	14.8	107.9
Non-Hispanic, white, both sexes, 60-64 years				
...	All causes	140,009	100.0	1,016.2
1	Malignant neoplasms (C00-C97)	49,046	35.0	356.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,561	21.1	214.6
3	Chronic lower respiratory diseases (J40-J47)	8,440	6.0	61.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,908	4.2	42.9
5	Diabetes mellitus (E10-E14)	4,861	3.5	35.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,459	3.2	32.4
7	Cerebrovascular diseases (I60-I69)	4,163	3.0	30.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,908	2.1	21.1
9	Septicemia (A40-A41)	2,115	1.5	15.4
10	Influenza and pneumonia (J09-J18)	2,053	1.5	14.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,683	1.2	12.2
12	Viral hepatitis (B15-B19)	1,317	0.9	9.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,156	0.8	8.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	744	0.5	5.4
15	Aortic aneurysm and dissection (I71)	605	0.4	4.4
...	All other causes (residual)	20,990	15.0	152.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65-69 years				
...	All causes	171,100	100.0	1,452.1
1	Malignant neoplasms (C00-C97)	61,118	35.7	518.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,127	21.1	306.6
3	Chronic lower respiratory diseases (J40-J47)	13,205	7.7	112.1
4	Diabetes mellitus (E10-E14)	6,159	3.6	52.3
5	Cerebrovascular diseases (I60-I69)	5,903	3.5	50.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,792	2.8	40.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,373	2.0	28.6
8	Septicemia (A40-A41)	2,793	1.6	23.7
9	Influenza and pneumonia (J09-J18)	2,669	1.6	22.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,562	1.5	21.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,093	1.2	17.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,333	0.8	11.3
13	Alzheimer's disease (G30)	1,177	0.7	10.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,059	0.6	9.0
15	Parkinson's disease (G20-G21)	858	0.5	7.3
...	All other causes (residual)	25,879	15.1	219.6
Non-Hispanic, white, both sexes, 70-74 years				
...	All causes	196,679	100.0	2,270.6
1	Malignant neoplasms (C00-C97)	65,592	33.3	757.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,249	21.0	476.2
3	Chronic lower respiratory diseases (J40-J47)	18,356	9.3	211.9
4	Cerebrovascular diseases (I60-I69)	8,014	4.1	92.5
5	Diabetes mellitus (E10-E14)	6,237	3.2	72.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,728	2.4	54.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,354	1.7	38.7
8	Influenza and pneumonia (J09-J18)	3,270	1.7	37.8
9	Septicemia (A40-A41)	3,204	1.6	37.0
10	Alzheimer's disease (G30)	3,086	1.6	35.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,517	1.3	29.1
12	Parkinson's disease (G20-G21)	2,067	1.1	23.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,640	0.8	18.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,565	0.8	18.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,381	0.7	15.9
...	All other causes (residual)	30,419	15.5	351.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-79 years				
...	All causes	226,557	100.0	3,638.3
1	Malignant neoplasms (C00-C97)	64,498	28.5	1,035.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,605	21.9	796.6
3	Chronic lower respiratory diseases (J40-J47)	20,438	9.0	328.2
4	Cerebrovascular diseases (I60-I69)	11,610	5.1	186.4
5	Alzheimer's disease (G30)	7,069	3.1	113.5
6	Diabetes mellitus (E10-E14)	6,467	2.9	103.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,553	2.5	89.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,313	1.9	69.3
9	Influenza and pneumonia (J09-J18)	4,293	1.9	68.9
10	Parkinson's disease (G20-G21)	3,773	1.7	60.6
11	Septicemia (A40-A41)	3,540	1.6	56.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,961	0.9	31.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,924	0.8	30.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,803	0.8	29.0
15	Pneumonitis due to solids and liquids (J69)	1,623	0.7	26.1
...	All other causes (residual)	38,087	16.8	611.6
Non-Hispanic, white, both sexes, 80-84 years				
...	All causes	283,724	100.0	6,129.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,696	24.2	1,484.2
2	Malignant neoplasms (C00-C97)	62,209	21.9	1,344.0
3	Chronic lower respiratory diseases (J40-J47)	22,309	7.9	482.0
4	Cerebrovascular diseases (I60-I69)	16,834	5.9	363.7
5	Alzheimer's disease (G30)	14,304	5.0	309.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,232	2.5	156.2
7	Diabetes mellitus (E10-E14)	7,089	2.5	153.2
8	Influenza and pneumonia (J09-J18)	6,151	2.2	132.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,736	2.0	123.9
10	Parkinson's disease (G20-G21)	5,724	2.0	123.7
11	Septicemia (A40-A41)	4,169	1.5	90.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,002	1.1	64.9
13	Pneumonitis due to solids and liquids (J69)	2,509	0.9	54.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,301	0.8	49.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,327	0.5	28.7
...	All other causes (residual)	54,132	19.1	1,169.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 85-89 years				
...	All causes	325,374	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	88,684	27.3	---
2	Malignant neoplasms (C00-C97)	50,859	15.6	---
3	Alzheimer's disease (G30)	21,855	6.7	---
4	Cerebrovascular diseases (I60-I69)	21,757	6.7	---
5	Chronic lower respiratory diseases (J40-J47)	20,397	6.3	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,847	2.7	---
7	Influenza and pneumonia (J09-J18)	8,219	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,811	2.1	---
9	Diabetes mellitus (E10-E14)	6,707	2.1	---
10	Parkinson's disease (G20-G21)	5,943	1.8	---
11	Septicemia (A40-A41)	4,528	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,241	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,390	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,485	0.8	---
15	Aortic aneurysm and dissection (I71)	1,256	0.4	---
...	All other causes (residual)	69,395	21.3	---
Non-Hispanic, white, both sexes, 90-94 years				
...	All causes	261,264	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	79,619	30.5	---
2	Malignant neoplasms (C00-C97)	27,021	10.3	---
3	Alzheimer's disease (G30)	20,872	8.0	---
4	Cerebrovascular diseases (I60-I69)	18,510	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	12,320	4.7	---
6	Influenza and pneumonia (J09-J18)	7,658	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,423	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,246	2.0	---
9	Diabetes mellitus (E10-E14)	4,277	1.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,010	1.5	---
11	Septicemia (A40-A41)	3,179	1.2	---
12	Parkinson's disease (G20-G21)	3,168	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,092	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,660	0.6	---
15	Atherosclerosis (I70)	1,206	0.5	---
...	All other causes (residual)	62,003	23.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 95-99 years				
...	All causes	103,206	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,809	32.8	---
2	Alzheimer's disease (G30)	9,103	8.8	---
3	Cerebrovascular diseases (I60-I69)	7,026	6.8	---
4	Malignant neoplasms (C00-C97)	6,819	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	3,703	3.6	---
6	Influenza and pneumonia (J09-J18)	3,422	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,714	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,954	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,953	1.9	---
10	Diabetes mellitus (E10-E14)	1,346	1.3	---
11	Pneumonitis due to solids and liquids (J69)	1,262	1.2	---
12	Septicemia (A40-A41)	1,035	1.0	---
13	Parkinson's disease (G20-G21)	756	0.7	---
14	Atherosclerosis (I70)	633	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	549	0.5	---
...	All other causes (residual)	27,122	26.3	---
Non-Hispanic, white, both sexes, 100 years and over				
...	All causes	21,496	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,572	35.2	---
2	Alzheimer's disease (G30)	1,804	8.4	---
3	Cerebrovascular diseases (I60-I69)	1,298	6.0	---
4	Influenza and pneumonia (J09-J18)	853	4.0	---
5	Malignant neoplasms (C00-C97)	840	3.9	---
6	Chronic lower respiratory diseases (J40-J47)	550	2.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	501	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	348	1.6	---
10	Pneumonitis due to solids and liquids (J69)	248	1.2	---
11	Diabetes mellitus (E10-E14)	190	0.9	---
12	Atherosclerosis (I70)	183	0.9	---
13	Septicemia (A40-A41)	178	0.8	---
14	Parkinson's disease (G20-G21)	94	0.4	---
15	Anemias (D50-D64)	69	0.3	---
15	Nutritional deficiencies (E40-E64)	69	0.3	---
...	All other causes (residual)	6,178	28.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	1,035,345	100.0	1,045.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	258,473	25.0	261.0
2	Malignant neoplasms (C00-C97)	246,872	23.8	249.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63,807	6.2	64.4
4	Chronic lower respiratory diseases (J40-J47)	60,428	5.8	61.0
5	Cerebrovascular diseases (I60-I69)	41,410	4.0	41.8
6	Diabetes mellitus (E10-E14)	28,743	2.8	29.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,410	2.6	27.7
8	Alzheimer's disease (G30)	24,321	2.3	24.6
9	Influenza and pneumonia (J09-J18)	20,633	2.0	20.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,878	1.7	18.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17,844	1.7	18.0
12	Parkinson's disease (G20-G21)	13,830	1.3	14.0
13	Septicemia (A40-A41)	13,514	1.3	13.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,984	0.9	9.1
15	Pneumonitis due to solids and liquids (J69)	8,623	0.8	8.7
...	All other causes (residual)	182,575	17.6	184.3
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,060	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	36.9	9.2
2	Malignant neoplasms (C00-C97)	110	10.4	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	9.5	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	7.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	2.7	0.7
5	Influenza and pneumonia (J09-J18)	29	2.7	0.7
7	Septicemia (A40-A41)	13	1.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	10	0.9	*
10	Cerebrovascular diseases (I60-I69)	7	0.7	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.7	*
12	Meningitis (G00,G03)	4	0.4	*
13	Anemias (D50-D64)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
15	Diseases of appendix (K35-K38)	2	0.2	*
...	All other causes (residual)	259	24.4	6.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-9 years				
...	All causes	680	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	35.4	4.3
2	Malignant neoplasms (C00-C97)	129	19.0	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	6.2	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	3.8	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	2.9	0.4
6	Influenza and pneumonia (J09-J18)	14	2.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
7	Cerebrovascular diseases (I60-I69)	9	1.3	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	1.0	*
11	Diabetes mellitus (E10-E14)	3	0.4	*
12	Septicemia (A40-A41)	2	0.3	*
12	Pneumonitis due to solids and liquids (J69)	2	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
15	Meningococcal infection (A39)	1	0.1	*
15	Anemias (D50-D64)	1	0.1	*
15	Meningitis (G00,G03)	1	0.1	*
15	Diseases of appendix (K35-K38)	1	0.1	*
15	Hernia (K40-K46)	1	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.1	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.1	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.1	*
...	All other causes (residual)	157	23.1	2.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 10-14 years				
...	All causes	940	100.0	16.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	270	28.7	4.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	190	20.2	3.3
3	Malignant neoplasms (C00-C97)	127	13.5	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	5.3	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	4.3	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	2.8	0.4
7	Chronic lower respiratory diseases (J40-J47)	16	1.7	*
8	Cerebrovascular diseases (I60-I69)	15	1.6	*
9	Influenza and pneumonia (J09-J18)	12	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.1	*
11	Septicemia (A40-A41)	8	0.9	*
12	Diabetes mellitus (E10-E14)	7	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
14	Anemias (D50-D64)	3	0.3	*
15	Whooping cough (A37)	2	0.2	*
15	Hernia (K40-K46)	2	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (residual)	154	16.4	2.6
Non-Hispanic, white, male, 15-19 years				
...	All causes	3,632	100.0	59.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,666	45.9	27.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,011	27.8	16.6
3	Malignant neoplasms (C00-C97)	195	5.4	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	3.4	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	2.5	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	1.7	1.0
7	Influenza and pneumonia (J09-J18)	23	0.6	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.4	*
9	Chronic lower respiratory diseases (J40-J47)	15	0.4	*
10	Diabetes mellitus (E10-E14)	13	0.4	*
10	Cerebrovascular diseases (I60-I69)	13	0.4	*
12	Septicemia (A40-A41)	11	0.3	*
13	Legal intervention (Y35,Y89.0)	6	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.2	*
15	Aortic aneurysm and dissection (I71)	4	0.1	*
...	All other causes (residual)	379	10.4	6.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 20-24 years				
...	All causes	7,876	100.0	119.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,041	51.3	61.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,839	23.3	27.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	322	4.1	4.9
4	Malignant neoplasms (C00-C97)	314	4.0	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	229	2.9	3.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	0.8	1.0
7	Diabetes mellitus (E10-E14)	56	0.7	0.8
8	Influenza and pneumonia (J09-J18)	43	0.5	0.7
9	Cerebrovascular diseases (I60-I69)	35	0.4	0.5
10	Chronic lower respiratory diseases (J40-J47)	34	0.4	0.5
11	Septicemia (A40-A41)	27	0.3	0.4
12	Legal intervention (Y35,Y89.0)	18	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.2	*
14	Pneumonitis due to solids and liquids (J69)	12	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	11	0.1	*
...	All other causes (residual)	816	10.4	12.3
Non-Hispanic, white, male, 25-29 years				
...	All causes	9,244	100.0	143.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,489	48.6	69.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,901	20.6	29.5
3	Malignant neoplasms (C00-C97)	427	4.6	6.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	401	4.3	6.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	352	3.8	5.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	1.1	1.6
7	Diabetes mellitus (E10-E14)	77	0.8	1.2
8	Influenza and pneumonia (J09-J18)	74	0.8	1.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	68	0.7	1.1
10	Cerebrovascular diseases (I60-I69)	64	0.7	1.0
11	Chronic lower respiratory diseases (J40-J47)	53	0.6	0.8
12	Septicemia (A40-A41)	45	0.5	0.7
13	Legal intervention (Y35,Y89.0)	41	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.3	0.4
15	Pneumonitis due to solids and liquids (J69)	21	0.2	0.3
...	All other causes (residual)	1,107	12.0	17.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 30-34 years				
...	All causes	10,601	100.0	168.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,502	42.5	71.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,982	18.7	31.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	812	7.7	12.9
4	Malignant neoplasms (C00-C97)	670	6.3	10.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	320	3.0	5.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	185	1.7	2.9
7	Diabetes mellitus (E10-E14)	140	1.3	2.2
8	Influenza and pneumonia (J09-J18)	90	0.8	1.4
9	Cerebrovascular diseases (I60-I69)	88	0.8	1.4
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	0.7	1.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	64	0.6	1.0
12	Chronic lower respiratory diseases (J40-J47)	61	0.6	1.0
13	Septicemia (A40-A41)	59	0.6	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.4	0.7
15	Legal intervention (Y35,Y89.0)	35	0.3	0.6
...	All other causes (residual)	1,473	13.9	23.4
Non-Hispanic, white, male, 35-39 years				
...	All causes	11,379	100.0	195.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,918	34.4	67.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,898	16.7	32.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,376	12.1	23.7
4	Malignant neoplasms (C00-C97)	990	8.7	17.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	283	2.5	4.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	260	2.3	4.5
7	Diabetes mellitus (E10-E14)	247	2.2	4.2
8	Cerebrovascular diseases (I60-I69)	156	1.4	2.7
9	Influenza and pneumonia (J09-J18)	130	1.1	2.2
10	Septicemia (A40-A41)	82	0.7	1.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	82	0.7	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	0.6	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	0.6	1.2
14	Chronic lower respiratory diseases (J40-J47)	63	0.6	1.1
15	Aortic aneurysm and dissection (I71)	47	0.4	0.8
...	All other causes (residual)	1,708	15.0	29.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 40-44 years				
...	All causes	16,034	100.0	256.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,874	24.2	62.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,831	17.7	45.3
3	Malignant neoplasms (C00-C97)	2,097	13.1	33.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,097	13.1	33.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	712	4.4	11.4
6	Diabetes mellitus (E10-E14)	439	2.7	7.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	289	1.8	4.6
8	Cerebrovascular diseases (I60-I69)	286	1.8	4.6
9	Influenza and pneumonia (J09-J18)	226	1.4	3.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	168	1.0	2.7
11	Chronic lower respiratory diseases (J40-J47)	154	1.0	2.5
12	Septicemia (A40-A41)	148	0.9	2.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	0.7	1.7
14	Aortic aneurysm and dissection (I71)	83	0.5	1.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	0.5	1.3
...	All other causes (residual)	2,445	15.2	39.2
Non-Hispanic, white, male, 45-49 years				
...	All causes	26,234	100.0	391.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,532	21.1	82.6
2	Malignant neoplasms (C00-C97)	4,625	17.6	69.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,494	17.1	67.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,589	9.9	38.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,366	5.2	20.4
6	Diabetes mellitus (E10-E14)	818	3.1	12.2
7	Cerebrovascular diseases (I60-I69)	562	2.1	8.4
8	Chronic lower respiratory diseases (J40-J47)	385	1.5	5.8
9	Influenza and pneumonia (J09-J18)	380	1.4	5.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	319	1.2	4.8
11	Septicemia (A40-A41)	285	1.1	4.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	251	1.0	3.7
13	Viral hepatitis (B15-B19)	235	0.9	3.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	0.8	3.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	156	0.6	2.3
...	All other causes (residual)	4,035	15.4	60.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 50-54 years				
...	All causes	47,177	100.0	615.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,979	23.3	143.2
2	Malignant neoplasms (C00-C97)	10,953	23.2	142.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,736	12.2	74.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,946	6.2	38.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,626	5.6	34.3
6	Diabetes mellitus (E10-E14)	1,467	3.1	19.1
7	Chronic lower respiratory diseases (J40-J47)	1,199	2.5	15.6
8	Cerebrovascular diseases (I60-I69)	1,059	2.2	13.8
9	Influenza and pneumonia (J09-J18)	604	1.3	7.9
10	Viral hepatitis (B15-B19)	588	1.2	7.7
11	Septicemia (A40-A41)	563	1.2	7.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	451	1.0	5.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	385	0.8	5.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	341	0.7	4.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	296	0.6	3.9
...	All other causes (residual)	6,984	14.8	91.1
Non-Hispanic, white, male, 55-59 years				
...	All causes	69,125	100.0	913.4
1	Malignant neoplasms (C00-C97)	20,117	29.1	265.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,667	24.1	220.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,100	7.4	67.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,288	4.8	43.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,855	4.1	37.7
6	Chronic lower respiratory diseases (J40-J47)	2,729	3.9	36.1
7	Diabetes mellitus (E10-E14)	2,240	3.2	29.6
8	Cerebrovascular diseases (I60-I69)	1,705	2.5	22.5
9	Influenza and pneumonia (J09-J18)	944	1.4	12.5
10	Viral hepatitis (B15-B19)	922	1.3	12.2
11	Septicemia (A40-A41)	842	1.2	11.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	673	1.0	8.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	528	0.8	7.0
14	Aortic aneurysm and dissection (I71)	317	0.5	4.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	304	0.4	4.0
...	All other causes (residual)	9,894	14.3	130.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 60-64 years				
...	All causes	85,555	100.0	1,280.0
1	Malignant neoplasms (C00-C97)	27,627	32.3	413.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,897	24.4	312.6
3	Chronic lower respiratory diseases (J40-J47)	4,460	5.2	66.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,984	4.7	59.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,124	3.7	46.7
6	Diabetes mellitus (E10-E14)	3,073	3.6	46.0
7	Cerebrovascular diseases (I60-I69)	2,387	2.8	35.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,242	2.6	33.5
9	Influenza and pneumonia (J09-J18)	1,167	1.4	17.5
10	Septicemia (A40-A41)	1,148	1.3	17.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	984	1.2	14.7
12	Viral hepatitis (B15-B19)	946	1.1	14.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	721	0.8	10.8
14	Aortic aneurysm and dissection (I71)	448	0.5	6.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	439	0.5	6.6
...	All other causes (residual)	11,908	13.9	178.2
Non-Hispanic, white, male, 65-69 years				
...	All causes	100,288	100.0	1,771.9
1	Malignant neoplasms (C00-C97)	34,005	33.9	600.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,283	24.2	429.0
3	Chronic lower respiratory diseases (J40-J47)	6,767	6.7	119.6
4	Diabetes mellitus (E10-E14)	3,764	3.8	66.5
5	Cerebrovascular diseases (I60-I69)	3,245	3.2	57.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,118	3.1	55.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,254	2.2	39.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,624	1.6	28.7
9	Influenza and pneumonia (J09-J18)	1,484	1.5	26.2
10	Septicemia (A40-A41)	1,459	1.5	25.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,435	1.4	25.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	799	0.8	14.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	672	0.7	11.9
14	Parkinson's disease (G20-G21)	571	0.6	10.1
15	Aortic aneurysm and dissection (I71)	553	0.6	9.8
...	All other causes (residual)	14,255	14.2	251.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 70-74 years				
...	All causes	110,520	100.0	2,725.7
1	Malignant neoplasms (C00-C97)	36,150	32.7	891.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,128	23.6	644.4
3	Chronic lower respiratory diseases (J40-J47)	9,017	8.2	222.4
4	Cerebrovascular diseases (I60-I69)	4,202	3.8	103.6
5	Diabetes mellitus (E10-E14)	3,761	3.4	92.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,851	2.6	70.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,832	1.7	45.2
8	Influenza and pneumonia (J09-J18)	1,776	1.6	43.8
9	Septicemia (A40-A41)	1,641	1.5	40.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,545	1.4	38.1
11	Parkinson's disease (G20-G21)	1,382	1.3	34.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,363	1.2	33.6
13	Alzheimer's disease (G30)	1,320	1.2	32.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	830	0.8	20.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	811	0.7	20.0
...	All other causes (residual)	15,911	14.4	392.4
Non-Hispanic, white, male, 75-79 years				
...	All causes	121,086	100.0	4,314.2
1	Malignant neoplasms (C00-C97)	34,906	28.8	1,243.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,259	24.2	1,042.5
3	Chronic lower respiratory diseases (J40-J47)	9,900	8.2	352.7
4	Cerebrovascular diseases (I60-I69)	5,580	4.6	198.8
5	Diabetes mellitus (E10-E14)	3,613	3.0	128.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,153	2.6	112.3
7	Alzheimer's disease (G30)	2,812	2.3	100.2
8	Parkinson's disease (G20-G21)	2,543	2.1	90.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,352	1.9	83.8
10	Influenza and pneumonia (J09-J18)	2,293	1.9	81.7
11	Septicemia (A40-A41)	1,695	1.4	60.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,092	0.9	38.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,068	0.9	38.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	982	0.8	35.0
15	Pneumonitis due to solids and liquids (J69)	976	0.8	34.8
...	All other causes (residual)	18,862	15.6	672.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 80-84 years				
...	All causes	140,288	100.0	7,207.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,436	26.0	1,871.9
2	Malignant neoplasms (C00-C97)	32,783	23.4	1,684.2
3	Chronic lower respiratory diseases (J40-J47)	10,497	7.5	539.3
4	Cerebrovascular diseases (I60-I69)	7,056	5.0	362.5
5	Alzheimer's disease (G30)	5,169	3.7	265.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,840	2.7	197.3
7	Diabetes mellitus (E10-E14)	3,709	2.6	190.6
8	Parkinson's disease (G20-G21)	3,576	2.5	183.7
9	Influenza and pneumonia (J09-J18)	3,104	2.2	159.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,040	2.2	156.2
11	Septicemia (A40-A41)	1,947	1.4	100.0
12	Pneumonitis due to solids and liquids (J69)	1,447	1.0	74.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,303	0.9	66.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,246	0.9	64.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	897	0.6	46.1
...	All other causes (residual)	24,238	17.3	1,245.2
Non-Hispanic, white, male, 85-89 years				
...	All causes	141,893	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,841	28.8	---
2	Malignant neoplasms (C00-C97)	25,622	18.1	---
3	Chronic lower respiratory diseases (J40-J47)	8,943	6.3	---
4	Cerebrovascular diseases (I60-I69)	7,793	5.5	---
5	Alzheimer's disease (G30)	6,932	4.9	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,049	2.9	---
7	Influenza and pneumonia (J09-J18)	3,759	2.6	---
8	Parkinson's disease (G20-G21)	3,447	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,409	2.4	---
10	Diabetes mellitus (E10-E14)	3,197	2.3	---
11	Septicemia (A40-A41)	1,967	1.4	---
12	Pneumonitis due to solids and liquids (J69)	1,893	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,572	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,246	0.9	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	647	0.5	---
...	All other causes (residual)	26,576	18.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 90-94 years				
...	All causes	93,996	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,300	32.2	---
2	Malignant neoplasms (C00-C97)	12,155	12.9	---
3	Alzheimer's disease (G30)	5,410	5.8	---
4	Cerebrovascular diseases (I60-I69)	5,389	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	4,777	5.1	---
6	Influenza and pneumonia (J09-J18)	3,124	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,864	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,386	2.5	---
9	Diabetes mellitus (E10-E14)	1,650	1.8	---
10	Parkinson's disease (G20-G21)	1,623	1.7	---
11	Pneumonitis due to solids and liquids (J69)	1,582	1.7	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,170	1.2	---
13	Septicemia (A40-A41)	1,159	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	738	0.8	---
15	Atherosclerosis (I70)	397	0.4	---
...	All other causes (residual)	19,272	20.5	---
Non-Hispanic, white, male, 95-99 years				
...	All causes	28,133	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,822	34.9	---
2	Malignant neoplasms (C00-C97)	2,602	9.2	---
3	Alzheimer's disease (G30)	1,680	6.0	---
4	Cerebrovascular diseases (I60-I69)	1,538	5.5	---
5	Chronic lower respiratory diseases (J40-J47)	1,217	4.3	---
6	Influenza and pneumonia (J09-J18)	1,116	4.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	776	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	728	2.6	---
9	Pneumonitis due to solids and liquids (J69)	562	2.0	---
10	Diabetes mellitus (E10-E14)	439	1.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	438	1.6	---
12	Septicemia (A40-A41)	328	1.2	---
13	Parkinson's disease (G20-G21)	313	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	185	0.7	---
15	Atherosclerosis (I70)	165	0.6	---
...	All other causes (residual)	6,224	22.1	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 100 years and over				
...	All causes	3,743	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,432	38.3	---
2	Malignant neoplasms (C00-C97)	243	6.5	---
3	Influenza and pneumonia (J09-J18)	203	5.4	---
4	Cerebrovascular diseases (I60-I69)	190	5.1	---
5	Alzheimer's disease (G30)	167	4.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	3.1	---
7	Chronic lower respiratory diseases (J40-J47)	112	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	2.6	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	91	2.4	---
10	Pneumonitis due to solids and liquids (J69)	80	2.1	---
11	Septicemia (A40-A41)	52	1.4	---
12	Atherosclerosis (I70)	33	0.9	---
13	Diabetes mellitus (E10-E14)	28	0.7	---
14	Parkinson's disease (G20-G21)	26	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.3	---
...	All other causes (residual)	861	23.0	---
Non-Hispanic, white, female, all ages				
...	All causes	1,031,604	100.0	1,011.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	231,453	22.4	226.9
2	Malignant neoplasms (C00-C97)	219,399	21.3	215.1
3	Chronic lower respiratory diseases (J40-J47)	69,091	6.7	67.7
4	Cerebrovascular diseases (I60-I69)	60,916	5.9	59.7
5	Alzheimer's disease (G30)	55,693	5.4	54.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40,875	4.0	40.1
7	Diabetes mellitus (E10-E14)	23,201	2.2	22.7
8	Influenza and pneumonia (J09-J18)	22,744	2.2	22.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,813	1.6	16.5
10	Septicemia (A40-A41)	15,354	1.5	15.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,527	1.2	12.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,986	1.0	9.8
13	Parkinson's disease (G20-G21)	9,087	0.9	8.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,034	0.8	7.9
15	Pneumonitis due to solids and liquids (J69)	7,021	0.7	6.9
...	All other causes (residual)	229,410	22.2	224.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	816	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	264	32.4	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	9.7	2.0
3	Malignant neoplasms (C00-C97)	66	8.1	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	7.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	3.3	0.7
6	Influenza and pneumonia (J09-J18)	24	2.9	0.6
7	Septicemia (A40-A41)	19	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	8	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.9	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.7	*
13	Meningitis (G00,G03)	4	0.5	*
14	Anemias (D50-D64)	3	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.4	*
14	Pneumonitis due to solids and liquids (J69)	3	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	3	0.4	*
...	All other causes (residual)	221	27.1	5.5
Non-Hispanic, white, female, 5-9 years				
...	All causes	496	100.0	9.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	27.2	2.5
2	Malignant neoplasms (C00-C97)	100	20.2	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	11.9	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	5.2	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.6	*
6	Influenza and pneumonia (J09-J18)	13	2.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	2.0	*
8	Septicemia (A40-A41)	9	1.8	*
9	Cerebrovascular diseases (I60-I69)	8	1.6	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.8	*
10	Certain conditions originating in the perinatal period (P00-P96)	4	0.8	*
12	Tuberculosis (A16-A19)	2	0.4	*
12	Anemias (D50-D64)	2	0.4	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	103	20.8	1.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 10-14 years				
...	All causes	568	100.0	10.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	21.7	2.2
2	Malignant neoplasms (C00-C97)	95	16.7	1.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	16.5	1.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.6	0.5
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	4.4	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	2.5	*
7	Influenza and pneumonia (J09-J18)	13	2.3	*
8	Cerebrovascular diseases (I60-I69)	10	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.6	*
10	Chronic lower respiratory diseases (J40-J47)	8	1.4	*
11	Septicemia (A40-A41)	7	1.2	*
12	Anemias (D50-D64)	2	0.4	*
12	Pneumonitis due to solids and liquids (J69)	2	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
15	Shigellosis and amebiasis (A03,A06)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.2	*
...	All other causes (residual)	132	23.2	2.4
Non-Hispanic, white, female, 15-19 years				
...	All causes	1,617	100.0	28.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	669	41.4	11.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	279	17.3	4.8
3	Malignant neoplasms (C00-C97)	149	9.2	2.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	3.5	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	3.5	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	3.0	0.8
7	Influenza and pneumonia (J09-J18)	23	1.4	0.4
8	Cerebrovascular diseases (I60-I69)	20	1.2	0.3
9	Pregnancy, childbirth and the puerperium (O00-O99)	13	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	10	0.6	*
12	Septicemia (A40-A41)	9	0.6	*
13	Diabetes mellitus (E10-E14)	8	0.5	*
14	Anemias (D50-D64)	4	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.2	*
...	All other causes (residual)	252	15.6	4.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 20-24 years				
...	All causes	2,915	100.0	46.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,367	46.9	21.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	374	12.8	5.9
3	Malignant neoplasms (C00-C97)	201	6.9	3.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	119	4.1	1.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	3.6	1.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	58	2.0	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	1.6	0.7
8	Influenza and pneumonia (J09-J18)	34	1.2	0.5
9	Diabetes mellitus (E10-E14)	24	0.8	0.4
10	Cerebrovascular diseases (I60-I69)	20	0.7	0.3
11	Septicemia (A40-A41)	18	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	16	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
15	Pneumonitis due to solids and liquids (J69)	9	0.3	*
...	All other causes (residual)	502	17.2	8.0
Non-Hispanic, white, female, 25-29 years				
...	All causes	3,934	100.0	62.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,656	42.1	26.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	479	12.2	7.6
3	Malignant neoplasms (C00-C97)	366	9.3	5.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	202	5.1	3.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	137	3.5	2.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	80	2.0	1.3
7	Influenza and pneumonia (J09-J18)	56	1.4	0.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	55	1.4	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	1.1	0.7
10	Diabetes mellitus (E10-E14)	41	1.0	0.7
11	Cerebrovascular diseases (I60-I69)	40	1.0	0.6
12	Chronic lower respiratory diseases (J40-J47)	35	0.9	0.6
13	Septicemia (A40-A41)	32	0.8	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.3	*
...	All other causes (residual)	682	17.3	10.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 30-34 years				
...	All causes	5,190	100.0	83.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,701	32.8	27.5
2	Malignant neoplasms (C00-C97)	681	13.1	11.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	546	10.5	8.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	379	7.3	6.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	171	3.3	2.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	133	2.6	2.1
7	Pregnancy, childbirth and the puerperium (O00-O99)	108	2.1	1.7
8	Influenza and pneumonia (J09-J18)	102	2.0	1.6
9	Cerebrovascular diseases (I60-I69)	89	1.7	1.4
10	Diabetes mellitus (E10-E14)	79	1.5	1.3
11	Septicemia (A40-A41)	69	1.3	1.1
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	1.3	1.1
13	Chronic lower respiratory diseases (J40-J47)	43	0.8	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.6	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.4	0.4
...	All other causes (residual)	968	18.7	15.6
Non-Hispanic, white, female, 35-39 years				
...	All causes	6,681	100.0	116.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,786	26.7	31.2
2	Malignant neoplasms (C00-C97)	1,279	19.1	22.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	619	9.3	10.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	605	9.1	10.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	223	3.3	3.9
6	Diabetes mellitus (E10-E14)	143	2.1	2.5
7	Cerebrovascular diseases (I60-I69)	138	2.1	2.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	1.8	2.1
9	Influenza and pneumonia (J09-J18)	110	1.6	1.9
10	Pregnancy, childbirth and the puerperium (O00-O99)	101	1.5	1.8
11	Septicemia (A40-A41)	97	1.5	1.7
12	Chronic lower respiratory diseases (J40-J47)	80	1.2	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	0.7	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	0.6	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.5	0.6
...	All other causes (residual)	1,260	18.9	22.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 40-44 years				
...	All causes	10,471	100.0	169.0
1	Malignant neoplasms (C00-C97)	2,568	24.5	41.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,013	19.2	32.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,283	12.3	20.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	772	7.4	12.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	426	4.1	6.9
6	Diabetes mellitus (E10-E14)	264	2.5	4.3
7	Cerebrovascular diseases (I60-I69)	238	2.3	3.8
8	Influenza and pneumonia (J09-J18)	208	2.0	3.4
9	Chronic lower respiratory diseases (J40-J47)	193	1.8	3.1
10	Septicemia (A40-A41)	155	1.5	2.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	131	1.3	2.1
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	0.8	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	0.7	1.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	58	0.6	0.9
14	Pregnancy, childbirth and the puerperium (O00-O99)	58	0.6	0.9
...	All other causes (residual)	1,949	18.6	31.5
Non-Hispanic, white, female, 45-49 years				
...	All causes	17,027	100.0	253.9
1	Malignant neoplasms (C00-C97)	5,098	29.9	76.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,360	13.9	35.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,243	13.2	33.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	926	5.4	13.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	780	4.6	11.6
6	Chronic lower respiratory diseases (J40-J47)	466	2.7	6.9
7	Cerebrovascular diseases (I60-I69)	446	2.6	6.6
8	Diabetes mellitus (E10-E14)	442	2.6	6.6
9	Influenza and pneumonia (J09-J18)	320	1.9	4.8
10	Septicemia (A40-A41)	269	1.6	4.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	0.9	2.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	0.8	2.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	0.8	1.9
14	Viral hepatitis (B15-B19)	105	0.6	1.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	0.5	1.2
...	All other causes (residual)	3,077	18.1	45.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 50-54 years				
...	All causes	30,241	100.0	387.0
1	Malignant neoplasms (C00-C97)	10,456	34.6	133.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,408	14.6	56.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,932	9.7	37.5
4	Chronic lower respiratory diseases (J40-J47)	1,463	4.8	18.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,338	4.4	17.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,133	3.7	14.5
7	Diabetes mellitus (E10-E14)	843	2.8	10.8
8	Cerebrovascular diseases (I60-I69)	842	2.8	10.8
9	Influenza and pneumonia (J09-J18)	533	1.8	6.8
10	Septicemia (A40-A41)	480	1.6	6.1
11	Viral hepatitis (B15-B19)	285	0.9	3.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	239	0.8	3.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	172	0.6	2.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	155	0.5	2.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	134	0.4	1.7
...	All other causes (residual)	4,828	16.0	61.8
Non-Hispanic, white, female, 55-59 years				
...	All causes	43,164	100.0	550.2
1	Malignant neoplasms (C00-C97)	16,413	38.0	209.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,483	15.0	82.6
3	Chronic lower respiratory diseases (J40-J47)	2,736	6.3	34.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,662	6.2	33.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,558	3.6	19.9
6	Cerebrovascular diseases (I60-I69)	1,302	3.0	16.6
7	Diabetes mellitus (E10-E14)	1,255	2.9	16.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	992	2.3	12.6
9	Septicemia (A40-A41)	820	1.9	10.5
10	Influenza and pneumonia (J09-J18)	713	1.7	9.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	477	1.1	6.1
12	Viral hepatitis (B15-B19)	399	0.9	5.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	284	0.7	3.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	215	0.5	2.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	184	0.4	2.3
...	All other causes (residual)	6,671	15.5	85.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 60-64 years				
...	All causes	54,454	100.0	767.7
1	Malignant neoplasms (C00-C97)	21,419	39.3	302.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,664	15.9	122.1
3	Chronic lower respiratory diseases (J40-J47)	3,980	7.3	56.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,924	3.5	27.1
5	Diabetes mellitus (E10-E14)	1,788	3.3	25.2
6	Cerebrovascular diseases (I60-I69)	1,776	3.3	25.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,335	2.5	18.8
8	Septicemia (A40-A41)	967	1.8	13.6
9	Influenza and pneumonia (J09-J18)	886	1.6	12.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	699	1.3	9.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	666	1.2	9.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	435	0.8	6.1
13	Viral hepatitis (B15-B19)	371	0.7	5.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	305	0.6	4.3
15	Alzheimer's disease (G30)	291	0.5	4.1
...	All other causes (residual)	8,948	16.4	126.1
Non-Hispanic, white, female, 65-69 years				
...	All causes	70,812	100.0	1,156.5
1	Malignant neoplasms (C00-C97)	27,113	38.3	442.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,844	16.7	193.4
3	Chronic lower respiratory diseases (J40-J47)	6,438	9.1	105.1
4	Cerebrovascular diseases (I60-I69)	2,658	3.8	43.4
5	Diabetes mellitus (E10-E14)	2,395	3.4	39.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,674	2.4	27.3
7	Septicemia (A40-A41)	1,334	1.9	21.8
8	Influenza and pneumonia (J09-J18)	1,185	1.7	19.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,127	1.6	18.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,119	1.6	18.3
11	Alzheimer's disease (G30)	660	0.9	10.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	534	0.8	8.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	469	0.7	7.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	387	0.5	6.3
15	Pneumonitis due to solids and liquids (J69)	353	0.5	5.8
...	All other causes (residual)	11,522	16.3	188.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 70-74 years				
...	All causes	86,159	100.0	1,870.1
1	Malignant neoplasms (C00-C97)	29,442	34.2	639.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,121	17.6	328.2
3	Chronic lower respiratory diseases (J40-J47)	9,339	10.8	202.7
4	Cerebrovascular diseases (I60-I69)	3,812	4.4	82.7
5	Diabetes mellitus (E10-E14)	2,476	2.9	53.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,877	2.2	40.7
7	Alzheimer's disease (G30)	1,766	2.0	38.3
8	Septicemia (A40-A41)	1,563	1.8	33.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,522	1.8	33.0
10	Influenza and pneumonia (J09-J18)	1,494	1.7	32.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	972	1.1	21.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	735	0.9	16.0
13	Parkinson's disease (G20-G21)	685	0.8	14.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	570	0.7	12.4
15	Pneumonitis due to solids and liquids (J69)	442	0.5	9.6
...	All other causes (residual)	14,343	16.6	311.3
Non-Hispanic, white, female, 75-79 years				
...	All causes	105,471	100.0	3,083.6
1	Malignant neoplasms (C00-C97)	29,592	28.1	865.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,346	19.3	594.9
3	Chronic lower respiratory diseases (J40-J47)	10,538	10.0	308.1
4	Cerebrovascular diseases (I60-I69)	6,030	5.7	176.3
5	Alzheimer's disease (G30)	4,257	4.0	124.5
6	Diabetes mellitus (E10-E14)	2,854	2.7	83.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,400	2.3	70.2
8	Influenza and pneumonia (J09-J18)	2,000	1.9	58.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,961	1.9	57.3
10	Septicemia (A40-A41)	1,845	1.7	53.9
11	Parkinson's disease (G20-G21)	1,230	1.2	36.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,028	1.0	30.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	832	0.8	24.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	735	0.7	21.5
15	Pneumonitis due to solids and liquids (J69)	647	0.6	18.9
...	All other causes (residual)	19,176	18.2	560.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 80-84 years				
...	All causes	143,436	100.0	5,348.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,260	22.5	1,202.8
2	Malignant neoplasms (C00-C97)	29,426	20.5	1,097.2
3	Chronic lower respiratory diseases (J40-J47)	11,812	8.2	440.4
4	Cerebrovascular diseases (I60-I69)	9,778	6.8	364.6
5	Alzheimer's disease (G30)	9,135	6.4	340.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,392	2.4	126.5
7	Diabetes mellitus (E10-E14)	3,380	2.4	126.0
8	Influenza and pneumonia (J09-J18)	3,047	2.1	113.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,696	1.9	100.5
10	Septicemia (A40-A41)	2,222	1.5	82.8
11	Parkinson's disease (G20-G21)	2,148	1.5	80.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,756	1.2	65.5
13	Pneumonitis due to solids and liquids (J69)	1,062	0.7	39.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	998	0.7	37.2
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	658	0.5	24.5
...	All other causes (residual)	29,666	20.7	1,106.1
Non-Hispanic, white, female, 85-89 years				
...	All causes	183,481	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,843	26.1	---
2	Malignant neoplasms (C00-C97)	25,237	13.8	---
3	Alzheimer's disease (G30)	14,923	8.1	---
4	Cerebrovascular diseases (I60-I69)	13,964	7.6	---
5	Chronic lower respiratory diseases (J40-J47)	11,454	6.2	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,798	2.6	---
7	Influenza and pneumonia (J09-J18)	4,460	2.4	---
8	Diabetes mellitus (E10-E14)	3,510	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,402	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,669	1.5	---
11	Septicemia (A40-A41)	2,561	1.4	---
12	Parkinson's disease (G20-G21)	2,496	1.4	---
13	Pneumonitis due to solids and liquids (J69)	1,497	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,239	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	768	0.4	---
...	All other causes (residual)	42,660	23.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 90-94 years				
...	All causes	167,268	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,319	29.5	---
2	Alzheimer's disease (G30)	15,462	9.2	---
3	Malignant neoplasms (C00-C97)	14,866	8.9	---
4	Cerebrovascular diseases (I60-I69)	13,121	7.8	---
5	Chronic lower respiratory diseases (J40-J47)	7,543	4.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,559	2.7	---
7	Influenza and pneumonia (J09-J18)	4,534	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,860	1.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,840	1.7	---
10	Diabetes mellitus (E10-E14)	2,627	1.6	---
11	Septicemia (A40-A41)	2,020	1.2	---
12	Parkinson's disease (G20-G21)	1,545	0.9	---
13	Pneumonitis due to solids and liquids (J69)	1,510	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	922	0.6	---
15	Atherosclerosis (I70)	809	0.5	---
...	All other causes (residual)	42,731	25.5	---
Non-Hispanic, white, female, 95-99 years				
...	All causes	75,073	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,987	32.0	---
2	Alzheimer's disease (G30)	7,423	9.9	---
3	Cerebrovascular diseases (I60-I69)	5,488	7.3	---
4	Malignant neoplasms (C00-C97)	4,217	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	2,486	3.3	---
6	Influenza and pneumonia (J09-J18)	2,306	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,938	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,516	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,225	1.6	---
10	Diabetes mellitus (E10-E14)	907	1.2	---
11	Septicemia (A40-A41)	707	0.9	---
12	Pneumonitis due to solids and liquids (J69)	700	0.9	---
13	Atherosclerosis (I70)	468	0.6	---
14	Parkinson's disease (G20-G21)	443	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	364	0.5	---
...	All other causes (residual)	20,898	27.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 100 years and over				
...	All causes	17,753	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,140	34.6	---
2	Alzheimer's disease (G30)	1,637	9.2	---
3	Cerebrovascular diseases (I60-I69)	1,108	6.2	---
4	Influenza and pneumonia (J09-J18)	650	3.7	---
5	Malignant neoplasms (C00-C97)	597	3.4	---
6	Chronic lower respiratory diseases (J40-J47)	438	2.5	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	410	2.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	406	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	250	1.4	---
10	Pneumonitis due to solids and liquids (J69)	168	0.9	---
11	Diabetes mellitus (E10-E14)	162	0.9	---
12	Atherosclerosis (I70)	150	0.8	---
13	Septicemia (A40-A41)	126	0.7	---
14	Parkinson's disease (G20-G21)	68	0.4	---
15	Anemias (D50-D64)	59	0.3	---
15	Nutritional deficiencies (E40-E64)	59	0.3	---
...	All other causes (residual)	5,325	30.0	---
Non-Hispanic, black, both sexes, all ages				
...	All causes	303,844	100.0	735.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,894	23.7	174.0
2	Malignant neoplasms (C00-C97)	68,104	22.4	164.8
3	Cerebrovascular diseases (I60-I69)	16,883	5.6	40.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,820	4.5	33.4
5	Diabetes mellitus (E10-E14)	13,264	4.4	32.1
6	Chronic lower respiratory diseases (J40-J47)	9,762	3.2	23.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,496	2.8	20.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,767	2.6	18.8
9	Alzheimer's disease (G30)	6,493	2.1	15.7
10	Septicemia (A40-A41)	6,321	2.1	15.3
11	Influenza and pneumonia (J09-J18)	5,462	1.8	13.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,322	1.8	12.9
13	Certain conditions originating in the perinatal period (P00-P96)	3,810	1.3	9.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,527	1.2	8.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,011	1.0	7.3
...	All other causes (residual)	59,908	19.7	145.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	967	100.0	39.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	28.9	11.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	14.8	5.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	10.0	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	5.6	2.2
5	Malignant neoplasms (C00-C97)	50	5.2	2.0
6	Influenza and pneumonia (J09-J18)	28	2.9	1.1
7	Chronic lower respiratory diseases (J40-J47)	20	2.1	0.8
8	Septicemia (A40-A41)	12	1.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
10	Cerebrovascular diseases (I60-I69)	9	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
11	Acute bronchitis and bronchiolitis (J20-J21)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
14	Anemias (D50-D64)	6	0.6	*
15	Meningitis (G00,G03)	4	0.4	*
...	All other causes (residual)	232	24.0	9.5
Non-Hispanic, black, both sexes, 5-9 years				
...	All causes	521	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	28.6	4.8
2	Malignant neoplasms (C00-C97)	75	14.4	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	8.3	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	7.1	1.2
5	Chronic lower respiratory diseases (J40-J47)	35	6.7	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.6	*
7	Influenza and pneumonia (J09-J18)	11	2.1	*
8	Cerebrovascular diseases (I60-I69)	10	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.7	*
10	Septicemia (A40-A41)	8	1.5	*
10	Anemias (D50-D64)	8	1.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	1.0	*
13	Meningitis (G00,G03)	3	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	99	19.0	3.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 10-14 years				
...	All causes	641	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	27.8	5.8
2	Malignant neoplasms (C00-C97)	67	10.5	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	9.4	1.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	8.3	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	6.6	1.4
6	Chronic lower respiratory diseases (J40-J47)	36	5.6	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	5.1	1.1
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.7	*
9	Anemias (D50-D64)	9	1.4	*
10	Diabetes mellitus (E10-E14)	8	1.2	*
11	Septicemia (A40-A41)	7	1.1	*
12	Cerebrovascular diseases (I60-I69)	5	0.8	*
12	Influenza and pneumonia (J09-J18)	5	0.8	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (residual)	117	18.3	3.8
Non-Hispanic, black, both sexes, 15-19 years				
...	All causes	2,104	100.0	65.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	887	42.2	27.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	530	25.2	16.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	146	6.9	4.5
4	Malignant neoplasms (C00-C97)	96	4.6	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	4.1	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	1.2	0.8
7	Chronic lower respiratory diseases (J40-J47)	23	1.1	0.7
8	Anemias (D50-D64)	17	0.8	*
9	Septicemia (A40-A41)	12	0.6	*
9	Cerebrovascular diseases (I60-I69)	12	0.6	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.5	*
12	Influenza and pneumonia (J09-J18)	8	0.4	*
12	Legal intervention (Y35,Y89.0)	8	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.3	*
14	Diabetes mellitus (E10-E14)	7	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
...	All other causes (residual)	222	10.6	6.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 20-24 years				
...	All causes	4,321	100.0	119.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,745	40.4	48.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	936	21.7	25.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	373	8.6	10.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	4.5	5.4
5	Malignant neoplasms (C00-C97)	165	3.8	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	76	1.8	2.1
7	Diabetes mellitus (E10-E14)	52	1.2	1.4
8	Chronic lower respiratory diseases (J40-J47)	48	1.1	1.3
9	Anemias (D50-D64)	47	1.1	1.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	45	1.0	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.9	1.1
12	Cerebrovascular diseases (I60-I69)	36	0.8	1.0
13	Influenza and pneumonia (J09-J18)	28	0.6	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.5	0.6
15	Legal intervention (Y35,Y89.0)	21	0.5	0.6
...	All other causes (residual)	493	11.4	13.6
Non-Hispanic, black, both sexes, 25-29 years				
...	All causes	4,420	100.0	143.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,371	31.0	44.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	932	21.1	30.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	7.6	10.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	296	6.7	9.6
5	Malignant neoplasms (C00-C97)	219	5.0	7.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	160	3.6	5.2
7	Diabetes mellitus (E10-E14)	88	2.0	2.8
8	Cerebrovascular diseases (I60-I69)	60	1.4	1.9
9	Chronic lower respiratory diseases (J40-J47)	57	1.3	1.8
9	Pregnancy, childbirth and the puerperium (O00-O99)	57	1.3	1.8
11	Anemias (D50-D64)	50	1.1	1.6
12	Influenza and pneumonia (J09-J18)	43	1.0	1.4
13	Septicemia (A40-A41)	36	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.7	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	0.7	1.1
...	All other causes (residual)	649	14.7	21.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 30-34 years				
...	All causes	4,953	100.0	174.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	977	19.7	34.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	972	19.6	34.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	662	13.4	23.3
4	Malignant neoplasms (C00-C97)	404	8.2	14.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	257	5.2	9.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	209	4.2	7.3
7	Diabetes mellitus (E10-E14)	145	2.9	5.1
8	Cerebrovascular diseases (I60-I69)	91	1.8	3.2
9	Pregnancy, childbirth and the puerperium (O00-O99)	79	1.6	2.8
10	Anemias (D50-D64)	75	1.5	2.6
11	Chronic lower respiratory diseases (J40-J47)	64	1.3	2.2
12	Influenza and pneumonia (J09-J18)	55	1.1	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.0	1.8
14	Septicemia (A40-A41)	45	0.9	1.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	35	0.7	1.2
...	All other causes (residual)	833	16.8	29.3
Non-Hispanic, black, both sexes, 35-39 years				
...	All causes	5,868	100.0	225.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,018	17.3	39.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	885	15.1	34.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	710	12.1	27.3
4	Malignant neoplasms (C00-C97)	683	11.6	26.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	272	4.6	10.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	227	3.9	8.7
7	Diabetes mellitus (E10-E14)	204	3.5	7.8
8	Cerebrovascular diseases (I60-I69)	175	3.0	6.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	1.7	3.9
10	Influenza and pneumonia (J09-J18)	84	1.4	3.2
11	Chronic lower respiratory diseases (J40-J47)	77	1.3	3.0
12	Septicemia (A40-A41)	73	1.2	2.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.2	2.8
14	Anemias (D50-D64)	66	1.1	2.5
15	Pregnancy, childbirth and the puerperium (O00-O99)	61	1.0	2.3
...	All other causes (residual)	1,160	19.8	44.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 40-44 years				
...	All causes	7,905	100.0	297.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,728	21.9	65.1
2	Malignant neoplasms (C00-C97)	1,260	15.9	47.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	928	11.7	35.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	491	6.2	18.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	361	4.6	13.6
6	Diabetes mellitus (E10-E14)	340	4.3	12.8
7	Cerebrovascular diseases (I60-I69)	332	4.2	12.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	202	2.6	7.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	2.1	6.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	123	1.6	4.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	122	1.5	4.6
12	Septicemia (A40-A41)	114	1.4	4.3
13	Influenza and pneumonia (J09-J18)	111	1.4	4.2
14	Chronic lower respiratory diseases (J40-J47)	108	1.4	4.1
15	Anemias (D50-D64)	75	0.9	2.8
...	All other causes (residual)	1,446	18.3	54.5
Non-Hispanic, black, both sexes, 45-49 years				
...	All causes	11,889	100.0	448.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,830	23.8	106.8
2	Malignant neoplasms (C00-C97)	2,410	20.3	91.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,087	9.1	41.0
4	Diabetes mellitus (E10-E14)	566	4.8	21.4
5	Cerebrovascular diseases (I60-I69)	513	4.3	19.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	502	4.2	18.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	396	3.3	14.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	230	1.9	8.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	223	1.9	8.4
10	Chronic lower respiratory diseases (J40-J47)	203	1.7	7.7
11	Septicemia (A40-A41)	202	1.7	7.6
12	Influenza and pneumonia (J09-J18)	200	1.7	7.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	182	1.5	6.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	172	1.4	6.5
15	Aortic aneurysm and dissection (I71)	71	0.6	2.7
...	All other causes (residual)	2,102	17.7	79.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 50-54 years				
...	All causes	19,408	100.0	700.6
1	Malignant neoplasms (C00-C97)	4,956	25.5	178.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,742	24.4	171.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,386	7.1	50.0
4	Cerebrovascular diseases (I60-I69)	942	4.9	34.0
5	Diabetes mellitus (E10-E14)	860	4.4	31.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	582	3.0	21.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	479	2.5	17.3
8	Chronic lower respiratory diseases (J40-J47)	440	2.3	15.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	421	2.2	15.2
10	Septicemia (A40-A41)	382	2.0	13.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	321	1.7	11.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	294	1.5	10.6
13	Influenza and pneumonia (J09-J18)	288	1.5	10.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	1.0	6.8
15	Viral hepatitis (B15-B19)	140	0.7	5.1
...	All other causes (residual)	2,988	15.4	107.9
Non-Hispanic, black, both sexes, 55-59 years				
...	All causes	27,460	100.0	1,082.8
1	Malignant neoplasms (C00-C97)	7,989	29.1	315.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,648	24.2	262.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,413	5.1	55.7
4	Cerebrovascular diseases (I60-I69)	1,316	4.8	51.9
5	Diabetes mellitus (E10-E14)	1,310	4.8	51.7
6	Chronic lower respiratory diseases (J40-J47)	799	2.9	31.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	665	2.4	26.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	644	2.3	25.4
9	Septicemia (A40-A41)	604	2.2	23.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	576	2.1	22.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	444	1.6	17.5
12	Influenza and pneumonia (J09-J18)	406	1.5	16.0
13	Viral hepatitis (B15-B19)	293	1.1	11.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	215	0.8	8.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	0.5	5.4
...	All other causes (residual)	4,000	14.6	157.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 60-64 years				
...	All causes	31,619	100.0	1,565.7
1	Malignant neoplasms (C00-C97)	9,828	31.1	486.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,744	24.5	383.5
3	Cerebrovascular diseases (I60-I69)	1,618	5.1	80.1
4	Diabetes mellitus (E10-E14)	1,555	4.9	77.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,008	3.2	49.9
6	Chronic lower respiratory diseases (J40-J47)	1,006	3.2	49.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	916	2.9	45.4
8	Septicemia (A40-A41)	707	2.2	35.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	612	1.9	30.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	562	1.8	27.8
11	Influenza and pneumonia (J09-J18)	506	1.6	25.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	400	1.3	19.8
13	Viral hepatitis (B15-B19)	343	1.1	17.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	134	0.4	6.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	125	0.4	6.2
...	All other causes (residual)	4,555	14.4	225.6
Non-Hispanic, black, both sexes, 65-69 years				
...	All causes	30,534	100.0	2,076.2
1	Malignant neoplasms (C00-C97)	9,710	31.8	660.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,548	24.7	513.2
3	Cerebrovascular diseases (I60-I69)	1,660	5.4	112.9
4	Diabetes mellitus (E10-E14)	1,563	5.1	106.3
5	Chronic lower respiratory diseases (J40-J47)	1,222	4.0	83.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,000	3.3	68.0
7	Septicemia (A40-A41)	725	2.4	49.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	660	2.2	44.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	497	1.6	33.8
10	Influenza and pneumonia (J09-J18)	463	1.5	31.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	395	1.3	26.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	208	0.7	14.1
13	Viral hepatitis (B15-B19)	165	0.5	11.2
14	Pneumonitis due to solids and liquids (J69)	152	0.5	10.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	149	0.5	10.1
...	All other causes (residual)	4,417	14.5	300.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 70-74 years				
...	All causes	29,534	100.0	2,957.4
1	Malignant neoplasms (C00-C97)	8,727	29.5	873.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,333	24.8	734.3
3	Cerebrovascular diseases (I60-I69)	1,816	6.1	181.8
4	Diabetes mellitus (E10-E14)	1,615	5.5	161.7
5	Chronic lower respiratory diseases (J40-J47)	1,329	4.5	133.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,065	3.6	106.6
7	Septicemia (A40-A41)	746	2.5	74.7
8	Influenza and pneumonia (J09-J18)	523	1.8	52.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	507	1.7	50.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	500	1.7	50.1
11	Alzheimer's disease (G30)	367	1.2	36.7
12	Pneumonitis due to solids and liquids (J69)	207	0.7	20.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	182	0.6	18.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	157	0.5	15.7
15	Parkinson's disease (G20-G21)	119	0.4	11.9
...	All other causes (residual)	4,341	14.7	434.7
Non-Hispanic, black, both sexes, 75-79 years				
...	All causes	29,864	100.0	4,296.2
1	Malignant neoplasms (C00-C97)	7,704	25.8	1,108.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,540	25.2	1,084.7
3	Cerebrovascular diseases (I60-I69)	1,993	6.7	286.7
4	Diabetes mellitus (E10-E14)	1,597	5.3	229.7
5	Chronic lower respiratory diseases (J40-J47)	1,306	4.4	187.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,032	3.5	148.5
7	Alzheimer's disease (G30)	796	2.7	114.5
8	Septicemia (A40-A41)	717	2.4	103.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	571	1.9	82.1
10	Influenza and pneumonia (J09-J18)	564	1.9	81.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	466	1.6	67.0
12	Pneumonitis due to solids and liquids (J69)	236	0.8	34.0
13	Parkinson's disease (G20-G21)	229	0.8	32.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	156	0.5	22.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	120	0.4	17.3
...	All other causes (residual)	4,837	16.2	695.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 80-84 years				
...	All causes	29,619	100.0	6,422.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,755	26.2	1,681.5
2	Malignant neoplasms (C00-C97)	6,368	21.5	1,380.7
3	Cerebrovascular diseases (I60-I69)	2,131	7.2	462.1
4	Diabetes mellitus (E10-E14)	1,372	4.6	297.5
5	Alzheimer's disease (G30)	1,262	4.3	273.6
6	Chronic lower respiratory diseases (J40-J47)	1,202	4.1	260.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,032	3.5	223.8
8	Septicemia (A40-A41)	654	2.2	141.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	633	2.1	137.3
10	Influenza and pneumonia (J09-J18)	610	2.1	132.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	383	1.3	83.0
12	Parkinson's disease (G20-G21)	254	0.9	55.1
13	Pneumonitis due to solids and liquids (J69)	245	0.8	53.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	152	0.5	33.0
15	Aortic aneurysm and dissection (I71)	91	0.3	19.7
...	All other causes (residual)	5,475	18.5	1,187.1
Non-Hispanic, black, both sexes, 85-89 years				
...	All causes	26,857	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,233	26.9	---
2	Malignant neoplasms (C00-C97)	4,450	16.6	---
3	Cerebrovascular diseases (I60-I69)	1,982	7.4	---
4	Alzheimer's disease (G30)	1,554	5.8	---
5	Diabetes mellitus (E10-E14)	1,107	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	993	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	945	3.5	---
8	Influenza and pneumonia (J09-J18)	653	2.4	---
9	Septicemia (A40-A41)	604	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	604	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	343	1.3	---
12	Pneumonitis due to solids and liquids (J69)	234	0.9	---
13	Parkinson's disease (G20-G21)	232	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	149	0.6	---
15	Nutritional deficiencies (E40-E64)	86	0.3	---
...	All other causes (residual)	5,688	21.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 90-94 years				
...	All causes	18,522	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,253	28.4	---
2	Malignant neoplasms (C00-C97)	2,180	11.8	---
3	Alzheimer's disease (G30)	1,414	7.6	---
4	Cerebrovascular diseases (I60-I69)	1,397	7.5	---
5	Diabetes mellitus (E10-E14)	633	3.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	545	2.9	---
7	Chronic lower respiratory diseases (J40-J47)	538	2.9	---
8	Influenza and pneumonia (J09-J18)	510	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	466	2.5	---
10	Septicemia (A40-A41)	416	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	245	1.3	---
12	Pneumonitis due to solids and liquids (J69)	173	0.9	---
13	Parkinson's disease (G20-G21)	139	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.4	---
15	Atherosclerosis (I70)	68	0.4	---
...	All other causes (residual)	4,463	24.1	---
Non-Hispanic, black, both sexes, 95-99 years				
...	All causes	7,622	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,262	29.7	---
2	Alzheimer's disease (G30)	649	8.5	---
3	Malignant neoplasms (C00-C97)	634	8.3	---
4	Cerebrovascular diseases (I60-I69)	584	7.7	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	224	2.9	---
6	Influenza and pneumonia (J09-J18)	219	2.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	203	2.7	---
8	Diabetes mellitus (E10-E14)	188	2.5	---
9	Chronic lower respiratory diseases (J40-J47)	184	2.4	---
10	Septicemia (A40-A41)	146	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	1.5	---
12	Pneumonitis due to solids and liquids (J69)	77	1.0	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.5	---
14	Parkinson's disease (G20-G21)	34	0.4	---
15	Nutritional deficiencies (E40-E64)	33	0.4	---
...	All other causes (residual)	2,032	26.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 100 years and over				
...	All causes	2,498	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	750	30.0	---
2	Alzheimer's disease (G30)	227	9.1	---
3	Cerebrovascular diseases (I60-I69)	178	7.1	---
4	Malignant neoplasms (C00-C97)	119	4.8	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	3.6	---
6	Influenza and pneumonia (J09-J18)	76	3.0	---
7	Chronic lower respiratory diseases (J40-J47)	64	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	2.4	---
9	Diabetes mellitus (E10-E14)	50	2.0	---
10	Septicemia (A40-A41)	39	1.6	---
11	Pneumonitis due to solids and liquids (J69)	30	1.2	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	0.9	---
13	Atherosclerosis (I70)	19	0.8	---
14	Nutritional deficiencies (E40-E64)	12	0.5	---
15	Anemias (D50-D64)	9	0.4	---
...	All other causes (residual)	753	30.1	---
Non-Hispanic, black, male, all ages				
...	All causes	154,836	100.0	783.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,265	24.1	188.5
2	Malignant neoplasms (C00-C97)	34,539	22.3	174.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,316	6.0	47.1
4	Cerebrovascular diseases (I60-I69)	7,650	4.9	38.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,718	4.3	34.0
6	Diabetes mellitus (E10-E14)	6,369	4.1	32.2
7	Chronic lower respiratory diseases (J40-J47)	4,940	3.2	25.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,983	2.6	20.2
9	Septicemia (A40-A41)	2,938	1.9	14.9
10	Influenza and pneumonia (J09-J18)	2,656	1.7	13.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,440	1.6	12.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,321	1.5	11.7
13	Certain conditions originating in the perinatal period (P00-P96)	2,134	1.4	10.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,876	1.2	9.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,873	1.2	9.5
...	All other causes (residual)	27,818	18.0	140.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	560	100.0	45.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	30.9	13.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	13.9	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	10.0	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.6	2.1
5	Malignant neoplasms (C00-C97)	25	4.5	2.0
6	Influenza and pneumonia (J09-J18)	18	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.6	*
9	Septicemia (A40-A41)	7	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
11	Anemias (D50-D64)	5	0.9	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
14	Meningitis (G00,G03)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (residual)	133	23.8	10.7
Non-Hispanic, black, male, 5-9 years				
...	All causes	302	100.0	19.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	32.5	6.2
2	Malignant neoplasms (C00-C97)	36	11.9	2.3
3	Chronic lower respiratory diseases (J40-J47)	25	8.3	1.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	8.3	1.6
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	6.6	1.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	4.6	*
7	Influenza and pneumonia (J09-J18)	7	2.3	*
8	Septicemia (A40-A41)	5	1.7	*
9	Anemias (D50-D64)	4	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	4	1.3	*
11	Cerebrovascular diseases (I60-I69)	3	1.0	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.7	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Diseases of appendix (K35-K38)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	52	17.2	3.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 10-14 years				
...	All causes	399	100.0	25.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	30.3	7.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	11.0	2.8
3	Malignant neoplasms (C00-C97)	39	9.8	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	9.8	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	6.5	1.7
6	Chronic lower respiratory diseases (J40-J47)	22	5.5	1.4
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	5.0	1.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.3	*
9	Anemias (D50-D64)	4	1.0	*
9	Diabetes mellitus (E10-E14)	4	1.0	*
9	Cerebrovascular diseases (I60-I69)	4	1.0	*
12	Septicemia (A40-A41)	3	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.8	*
14	Influenza and pneumonia (J09-J18)	2	0.5	*
15	Salmonella infections (A01-A02)	1	0.3	*
15	Aortic aneurysm and dissection (I71)	1	0.3	*
15	Pneumonitis due to solids and liquids (J69)	1	0.3	*
15	Peptic ulcer (K25-K28)	1	0.3	*
15	Diseases of appendix (K35-K38)	1	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
15	Legal intervention (Y35,Y89.0)	1	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	55	13.8	3.5
Non-Hispanic, black, male, 15-19 years				
...	All causes	1,627	100.0	99.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	797	49.0	48.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	24.3	24.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	111	6.8	6.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	3.6	3.6
5	Malignant neoplasms (C00-C97)	58	3.6	3.6
6	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.9	*
8	Cerebrovascular diseases (I60-I69)	9	0.6	*
9	Legal intervention (Y35,Y89.0)	8	0.5	*
10	Influenza and pneumonia (J09-J18)	7	0.4	*
11	Diabetes mellitus (E10-E14)	6	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.3	*
13	Septicemia (A40-A41)	4	0.2	*
13	Anemias (D50-D64)	4	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (residual)	129	7.9	7.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 20-24 years				
...	All causes	3,308	100.0	181.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,591	48.1	87.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	697	21.1	38.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	317	9.6	17.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	3.4	6.1
5	Malignant neoplasms (C00-C97)	95	2.9	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	44	1.3	2.4
7	Chronic lower respiratory diseases (J40-J47)	30	0.9	1.6
8	Anemias (D50-D64)	28	0.8	1.5
9	Diabetes mellitus (E10-E14)	27	0.8	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.7	1.3
11	Legal intervention (Y35,Y89.0)	21	0.6	1.1
12	Cerebrovascular diseases (I60-I69)	20	0.6	1.1
13	Influenza and pneumonia (J09-J18)	19	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
15	Septicemia (A40-A41)	8	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
...	All other causes (residual)	256	7.7	14.0
Non-Hispanic, black, male, 25-29 years				
...	All causes	3,227	100.0	212.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,220	37.8	80.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	703	21.8	46.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	256	7.9	16.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	216	6.7	14.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	125	3.9	8.2
6	Malignant neoplasms (C00-C97)	99	3.1	6.5
7	Diabetes mellitus (E10-E14)	46	1.4	3.0
8	Chronic lower respiratory diseases (J40-J47)	34	1.1	2.2
9	Cerebrovascular diseases (I60-I69)	30	0.9	2.0
10	Anemias (D50-D64)	29	0.9	1.9
10	Legal intervention (Y35,Y89.0)	29	0.9	1.9
12	Septicemia (A40-A41)	21	0.7	1.4
13	Influenza and pneumonia (J09-J18)	20	0.6	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.6	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.5	*
...	All other causes (residual)	364	11.3	24.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 30-34 years				
...	All causes	3,271	100.0	240.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	879	26.9	64.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	717	21.9	52.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	406	12.4	29.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	210	6.4	15.4
5	Malignant neoplasms (C00-C97)	157	4.8	11.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	122	3.7	9.0
7	Diabetes mellitus (E10-E14)	82	2.5	6.0
8	Cerebrovascular diseases (I60-I69)	59	1.8	4.3
9	Anemias (D50-D64)	49	1.5	3.6
10	Chronic lower respiratory diseases (J40-J47)	37	1.1	2.7
11	Influenza and pneumonia (J09-J18)	27	0.8	2.0
12	Septicemia (A40-A41)	22	0.7	1.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.7	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.7	1.6
15	Aortic aneurysm and dissection (I71)	18	0.6	*
...	All other causes (residual)	442	13.5	32.4
Non-Hispanic, black, male, 35-39 years				
...	All causes	3,568	100.0	289.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	628	17.6	51.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	626	17.5	50.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	623	17.5	50.6
4	Malignant neoplasms (C00-C97)	260	7.3	21.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	5.2	15.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	152	4.3	12.3
7	Diabetes mellitus (E10-E14)	112	3.1	9.1
8	Cerebrovascular diseases (I60-I69)	108	3.0	8.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	1.7	4.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	50	1.4	4.1
11	Chronic lower respiratory diseases (J40-J47)	45	1.3	3.7
12	Influenza and pneumonia (J09-J18)	41	1.1	3.3
13	Anemias (D50-D64)	33	0.9	2.7
14	Septicemia (A40-A41)	29	0.8	2.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	27	0.8	2.2
...	All other causes (residual)	587	16.5	47.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 40-44 years				
...	All causes	4,498	100.0	360.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,076	23.9	86.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	630	14.0	50.5
3	Malignant neoplasms (C00-C97)	495	11.0	39.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	411	9.1	33.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	208	4.6	16.7
6	Cerebrovascular diseases (I60-I69)	192	4.3	15.4
7	Diabetes mellitus (E10-E14)	186	4.1	14.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	3.4	12.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	2.0	7.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	78	1.7	6.3
11	Septicemia (A40-A41)	62	1.4	5.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	60	1.3	4.8
13	Influenza and pneumonia (J09-J18)	59	1.3	4.7
14	Chronic lower respiratory diseases (J40-J47)	54	1.2	4.3
15	Aortic aneurysm and dissection (I71)	48	1.1	3.8
...	All other causes (residual)	694	15.4	55.7
Non-Hispanic, black, male, 45-49 years				
...	All causes	6,755	100.0	541.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,794	26.6	143.8
2	Malignant neoplasms (C00-C97)	1,019	15.1	81.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	751	11.1	60.2
4	Diabetes mellitus (E10-E14)	332	4.9	26.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	323	4.8	25.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	322	4.8	25.8
7	Cerebrovascular diseases (I60-I69)	281	4.2	22.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	142	2.1	11.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	1.9	10.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	115	1.7	9.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	114	1.7	9.1
12	Influenza and pneumonia (J09-J18)	111	1.6	8.9
13	Septicemia (A40-A41)	102	1.5	8.2
14	Chronic lower respiratory diseases (J40-J47)	85	1.3	6.8
15	Aortic aneurysm and dissection (I71)	54	0.8	4.3
...	All other causes (residual)	1,080	16.0	86.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 50-54 years				
...	All causes	10,913	100.0	841.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,934	26.9	226.3
2	Malignant neoplasms (C00-C97)	2,373	21.7	183.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	938	8.6	72.3
4	Cerebrovascular diseases (I60-I69)	502	4.6	38.7
5	Diabetes mellitus (E10-E14)	483	4.4	37.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	385	3.5	29.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	301	2.8	23.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	261	2.4	20.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	217	2.0	16.7
10	Septicemia (A40-A41)	195	1.8	15.0
11	Chronic lower respiratory diseases (J40-J47)	189	1.7	14.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	173	1.6	13.3
13	Influenza and pneumonia (J09-J18)	143	1.3	11.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	1.3	11.0
15	Viral hepatitis (B15-B19)	89	0.8	6.9
...	All other causes (residual)	1,588	14.6	122.5
Non-Hispanic, black, male, 55-59 years				
...	All causes	15,913	100.0	1,358.1
1	Malignant neoplasms (C00-C97)	4,200	26.4	358.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,193	26.3	357.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	991	6.2	84.6
4	Cerebrovascular diseases (I60-I69)	780	4.9	66.6
5	Diabetes mellitus (E10-E14)	728	4.6	62.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	427	2.7	36.4
7	Chronic lower respiratory diseases (J40-J47)	404	2.5	34.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	397	2.5	33.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	373	2.3	31.8
10	Septicemia (A40-A41)	312	2.0	26.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	269	1.7	23.0
12	Influenza and pneumonia (J09-J18)	217	1.4	18.5
13	Viral hepatitis (B15-B19)	194	1.2	16.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	174	1.1	14.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	103	0.6	8.8
...	All other causes (residual)	2,151	13.5	183.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 60-64 years				
...	All causes	18,355	100.0	2,029.0
1	Malignant neoplasms (C00-C97)	5,417	29.5	598.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,827	26.3	533.6
3	Cerebrovascular diseases (I60-I69)	919	5.0	101.6
4	Diabetes mellitus (E10-E14)	853	4.6	94.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	703	3.8	77.7
6	Chronic lower respiratory diseases (J40-J47)	519	2.8	57.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	502	2.7	55.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	408	2.2	45.1
9	Septicemia (A40-A41)	366	2.0	40.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	312	1.7	34.5
11	Influenza and pneumonia (J09-J18)	298	1.6	32.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	291	1.6	32.2
13	Viral hepatitis (B15-B19)	239	1.3	26.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	106	0.6	11.7
15	Pneumonitis due to solids and liquids (J69)	82	0.4	9.1
...	All other causes (residual)	2,513	13.7	277.8
Non-Hispanic, black, male, 65-69 years				
...	All causes	17,182	100.0	2,675.8
1	Malignant neoplasms (C00-C97)	5,295	30.8	824.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,507	26.2	701.9
3	Cerebrovascular diseases (I60-I69)	908	5.3	141.4
4	Diabetes mellitus (E10-E14)	851	5.0	132.5
5	Chronic lower respiratory diseases (J40-J47)	676	3.9	105.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	495	2.9	77.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	453	2.6	70.5
8	Septicemia (A40-A41)	372	2.2	57.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	278	1.6	43.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	275	1.6	42.8
10	Influenza and pneumonia (J09-J18)	275	1.6	42.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	151	0.9	23.5
13	Viral hepatitis (B15-B19)	115	0.7	17.9
14	Pneumonitis due to solids and liquids (J69)	100	0.6	15.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	0.5	12.3
...	All other causes (residual)	2,352	13.7	366.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 70-74 years				
...	All causes	15,780	100.0	3,778.9
1	Malignant neoplasms (C00-C97)	4,733	30.0	1,133.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,061	25.7	972.5
3	Cerebrovascular diseases (I60-I69)	934	5.9	223.7
4	Diabetes mellitus (E10-E14)	786	5.0	188.2
5	Chronic lower respiratory diseases (J40-J47)	764	4.8	183.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	498	3.2	119.3
7	Septicemia (A40-A41)	378	2.4	90.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	317	2.0	75.9
9	Influenza and pneumonia (J09-J18)	308	2.0	73.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	265	1.7	63.5
11	Alzheimer's disease (G30)	136	0.9	32.6
12	Pneumonitis due to solids and liquids (J69)	129	0.8	30.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	125	0.8	29.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.5	19.2
15	Parkinson's disease (G20-G21)	77	0.5	18.4
...	All other causes (residual)	2,189	13.9	524.2
Non-Hispanic, black, male, 75-79 years				
...	All causes	14,687	100.0	5,342.8
1	Malignant neoplasms (C00-C97)	3,904	26.6	1,420.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,878	26.4	1,410.7
3	Cerebrovascular diseases (I60-I69)	929	6.3	338.0
4	Diabetes mellitus (E10-E14)	763	5.2	277.6
5	Chronic lower respiratory diseases (J40-J47)	676	4.6	245.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	473	3.2	172.1
7	Septicemia (A40-A41)	338	2.3	123.0
8	Alzheimer's disease (G30)	310	2.1	112.8
9	Influenza and pneumonia (J09-J18)	302	2.1	109.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	1.9	98.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	262	1.8	95.3
12	Parkinson's disease (G20-G21)	139	0.9	50.6
13	Pneumonitis due to solids and liquids (J69)	131	0.9	47.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.5	27.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	62	0.4	22.6
...	All other causes (residual)	2,172	14.8	790.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 80-84 years				
...	All causes	12,985	100.0	7,860.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,506	27.0	2,122.4
2	Malignant neoplasms (C00-C97)	3,147	24.2	1,905.0
3	Cerebrovascular diseases (I60-I69)	839	6.5	507.9
4	Chronic lower respiratory diseases (J40-J47)	579	4.5	350.5
5	Diabetes mellitus (E10-E14)	508	3.9	307.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	430	3.3	260.3
7	Alzheimer's disease (G30)	407	3.1	246.4
8	Septicemia (A40-A41)	296	2.3	179.2
9	Influenza and pneumonia (J09-J18)	261	2.0	158.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	241	1.9	145.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	206	1.6	124.7
12	Parkinson's disease (G20-G21)	150	1.2	90.8
13	Pneumonitis due to solids and liquids (J69)	121	0.9	73.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.4	34.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.3	20.0
...	All other causes (residual)	2,204	17.0	1,334.2
Non-Hispanic, black, male, 85-89 years				
...	All causes	10,069	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,651	26.3	---
2	Malignant neoplasms (C00-C97)	2,026	20.1	---
3	Cerebrovascular diseases (I60-I69)	653	6.5	---
4	Chronic lower respiratory diseases (J40-J47)	470	4.7	---
5	Alzheimer's disease (G30)	432	4.3	---
6	Diabetes mellitus (E10-E14)	365	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	356	3.5	---
8	Influenza and pneumonia (J09-J18)	283	2.8	---
9	Septicemia (A40-A41)	222	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	186	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	1.6	---
12	Parkinson's disease (G20-G21)	127	1.3	---
13	Pneumonitis due to solids and liquids (J69)	107	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	---
15	Nutritional deficiencies (E40-E64)	34	0.3	---
...	All other causes (residual)	1,943	19.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 90-94 years				
...	All causes	5,592	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,627	29.1	---
2	Malignant neoplasms (C00-C97)	910	16.3	---
3	Cerebrovascular diseases (I60-I69)	331	5.9	---
4	Alzheimer's disease (G30)	322	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	212	3.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	190	3.4	---
7	Diabetes mellitus (E10-E14)	179	3.2	---
8	Influenza and pneumonia (J09-J18)	162	2.9	---
9	Septicemia (A40-A41)	123	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	1.7	---
12	Parkinson's disease (G20-G21)	69	1.2	---
13	Pneumonitis due to solids and liquids (J69)	67	1.2	---
14	Nutritional deficiencies (E40-E64)	26	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.4	---
...	All other causes (residual)	1,132	20.2	---
Non-Hispanic, black, male, 95-99 years				
...	All causes	1,778	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	551	31.0	---
2	Malignant neoplasms (C00-C97)	218	12.3	---
3	Cerebrovascular diseases (I60-I69)	115	6.5	---
4	Alzheimer's disease (G30)	96	5.4	---
5	Chronic lower respiratory diseases (J40-J47)	65	3.7	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	3.7	---
7	Influenza and pneumonia (J09-J18)	52	2.9	---
8	Diabetes mellitus (E10-E14)	50	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	2.4	---
10	Septicemia (A40-A41)	42	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	2.0	---
12	Pneumonitis due to solids and liquids (J69)	23	1.3	---
13	Parkinson's disease (G20-G21)	12	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.5	---
15	Nutritional deficiencies (E40-E64)	8	0.4	---
...	All other causes (residual)	394	22.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 100 years and over				
...	All causes	371	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	118	31.8	---
2	Malignant neoplasms (C00-C97)	31	8.4	---
3	Chronic lower respiratory diseases (J40-J47)	22	5.9	---
4	Influenza and pneumonia (J09-J18)	17	4.6	---
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	4.6	---
6	Alzheimer's disease (G30)	16	4.3	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	4.3	---
6	Cerebrovascular diseases (I60-I69)	16	4.3	---
9	Pneumonitis due to solids and liquids (J69)	8	2.2	---
10	Diabetes mellitus (E10-E14)	5	1.3	---
11	Septicemia (A40-A41)	4	1.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	1.1	---
13	Atherosclerosis (I70)	3	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	---
14	Nutritional deficiencies (E40-E64)	2	0.5	---
...	All other causes (residual)	90	24.3	---
Non-Hispanic, black, female, all ages				
...	All causes	149,008	100.0	691.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,629	23.2	160.7
2	Malignant neoplasms (C00-C97)	33,565	22.5	155.8
3	Cerebrovascular diseases (I60-I69)	9,233	6.2	42.8
4	Diabetes mellitus (E10-E14)	6,895	4.6	32.0
5	Chronic lower respiratory diseases (J40-J47)	4,822	3.2	22.4
6	Alzheimer's disease (G30)	4,675	3.1	21.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,513	3.0	20.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,504	3.0	20.9
9	Septicemia (A40-A41)	3,383	2.3	15.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,882	1.9	13.4
11	Influenza and pneumonia (J09-J18)	2,806	1.9	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,676	1.1	7.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,206	0.8	5.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,138	0.8	5.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,049	0.7	4.9
...	All other causes (residual)	32,032	21.5	148.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 1-4 years				
...	All causes	407	100.0	34.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	26.0	8.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	16.0	5.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	10.1	3.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	6.9	2.3
5	Malignant neoplasms (C00-C97)	25	6.1	2.1
6	Chronic lower respiratory diseases (J40-J47)	11	2.7	*
7	Influenza and pneumonia (J09-J18)	10	2.5	*
8	Septicemia (A40-A41)	5	1.2	*
9	Cerebrovascular diseases (I60-I69)	4	1.0	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	4	1.0	*
11	Acute bronchitis and bronchiolitis (J20-J21)	3	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
13	Salmonella infections (A01-A02)	1	0.2	*
13	Meningococcal infection (A39)	1	0.2	*
13	Malaria (B50-B54)	1	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Nutritional deficiencies (E40-E64)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (residual)	94	23.1	7.8
Non-Hispanic, black, female, 5-9 years				
...	All causes	219	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	23.3	3.3
2	Malignant neoplasms (C00-C97)	39	17.8	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	8.2	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	7.8	*
5	Chronic lower respiratory diseases (J40-J47)	10	4.6	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	4.1	*
7	Cerebrovascular diseases (I60-I69)	7	3.2	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.3	*
9	Anemias (D50-D64)	4	1.8	*
9	Influenza and pneumonia (J09-J18)	4	1.8	*
11	Septicemia (A40-A41)	3	1.4	*
12	Meningitis (G00,G03)	2	0.9	*
13	Diabetes mellitus (E10-E14)	1	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (residual)	45	20.5	2.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 10-14 years				
...	All causes	242	100.0	15.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	23.6	3.7
2	Malignant neoplasms (C00-C97)	28	11.6	1.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	6.6	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	6.6	*
5	Chronic lower respiratory diseases (J40-J47)	14	5.8	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	5.8	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	5.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	2.5	*
9	Anemias (D50-D64)	5	2.1	*
10	Septicemia (A40-A41)	4	1.7	*
10	Diabetes mellitus (E10-E14)	4	1.7	*
12	Influenza and pneumonia (J09-J18)	3	1.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.8	*
15	Cerebrovascular diseases (I60-I69)	1	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (residual)	56	23.1	3.7
Non-Hispanic, black, female, 15-19 years				
...	All causes	477	100.0	30.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	28.3	8.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	18.9	5.7
3	Malignant neoplasms (C00-C97)	38	8.0	2.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	7.3	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	5.7	1.7
6	Anemias (D50-D64)	13	2.7	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	11	2.3	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	2.3	*
9	Septicemia (A40-A41)	8	1.7	*
10	Chronic lower respiratory diseases (J40-J47)	6	1.3	*
11	Cerebrovascular diseases (I60-I69)	3	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Influenza and pneumonia (J09-J18)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (residual)	88	18.4	5.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 20-24 years				
...	All causes	1,013	100.0	56.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	23.6	13.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	15.2	8.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	8.2	4.6
4	Malignant neoplasms (C00-C97)	70	6.9	3.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	5.5	3.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	45	4.4	2.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	32	3.2	1.8
8	Diabetes mellitus (E10-E14)	25	2.5	1.4
9	Anemias (D50-D64)	19	1.9	*
10	Chronic lower respiratory diseases (J40-J47)	18	1.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.7	*
12	Cerebrovascular diseases (I60-I69)	16	1.6	*
13	Septicemia (A40-A41)	12	1.2	*
14	Influenza and pneumonia (J09-J18)	9	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.9	*
...	All other causes (residual)	209	20.6	11.7
Non-Hispanic, black, female, 25-29 years				
...	All causes	1,193	100.0	75.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	229	19.2	14.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	12.7	9.6
3	Malignant neoplasms (C00-C97)	120	10.1	7.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	10.1	7.6
5	Pregnancy, childbirth and the puerperium (O00-O99)	57	4.8	3.6
6	Diabetes mellitus (E10-E14)	42	3.5	2.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	3.4	2.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	35	2.9	2.2
9	Cerebrovascular diseases (I60-I69)	30	2.5	1.9
10	Influenza and pneumonia (J09-J18)	23	1.9	1.5
10	Chronic lower respiratory diseases (J40-J47)	23	1.9	1.5
12	Anemias (D50-D64)	21	1.8	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.5	*
14	Septicemia (A40-A41)	15	1.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.1	*
...	All other causes (residual)	256	21.5	16.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 30-34 years				
...	All causes	1,682	100.0	113.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	256	15.2	17.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	255	15.2	17.2
3	Malignant neoplasms (C00-C97)	247	14.7	16.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	5.8	6.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	87	5.2	5.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	79	4.7	5.3
7	Diabetes mellitus (E10-E14)	63	3.7	4.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	2.8	3.2
9	Cerebrovascular diseases (I60-I69)	32	1.9	2.2
10	Influenza and pneumonia (J09-J18)	28	1.7	1.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	1.7	1.9
12	Chronic lower respiratory diseases (J40-J47)	27	1.6	1.8
13	Anemias (D50-D64)	26	1.5	1.8
14	Septicemia (A40-A41)	23	1.4	1.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	23	1.4	1.5
...	All other causes (residual)	363	21.6	24.5
Non-Hispanic, black, female, 35-39 years				
...	All causes	2,300	100.0	167.4
1	Malignant neoplasms (C00-C97)	423	18.4	30.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	395	17.2	28.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	257	11.2	18.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	120	5.2	8.7
5	Diabetes mellitus (E10-E14)	92	4.0	6.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	3.7	6.1
7	Cerebrovascular diseases (I60-I69)	67	2.9	4.9
8	Pregnancy, childbirth and the puerperium (O00-O99)	61	2.7	4.4
9	Septicemia (A40-A41)	44	1.9	3.2
10	Influenza and pneumonia (J09-J18)	43	1.9	3.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.8	3.0
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	1.7	2.9
13	Anemias (D50-D64)	33	1.4	2.4
14	Chronic lower respiratory diseases (J40-J47)	32	1.4	2.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	29	1.3	2.1
...	All other causes (residual)	539	23.4	39.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 40-44 years				
...	All causes	3,407	100.0	242.2
1	Malignant neoplasms (C00-C97)	765	22.5	54.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	652	19.1	46.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	298	8.7	21.2
4	Diabetes mellitus (E10-E14)	154	4.5	10.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	153	4.5	10.9
6	Cerebrovascular diseases (I60-I69)	140	4.1	10.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	2.3	5.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	2.2	5.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	62	1.8	4.4
10	Chronic lower respiratory diseases (J40-J47)	54	1.6	3.8
11	Septicemia (A40-A41)	52	1.5	3.7
11	Influenza and pneumonia (J09-J18)	52	1.5	3.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	1.4	3.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	1.3	3.2
15	Pregnancy, childbirth and the puerperium (O00-O99)	42	1.2	3.0
...	All other causes (residual)	737	21.6	52.4
Non-Hispanic, black, female, 45-49 years				
...	All causes	5,134	100.0	366.1
1	Malignant neoplasms (C00-C97)	1,391	27.1	99.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,036	20.2	73.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	336	6.5	24.0
4	Diabetes mellitus (E10-E14)	234	4.6	16.7
5	Cerebrovascular diseases (I60-I69)	232	4.5	16.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	179	3.5	12.8
7	Chronic lower respiratory diseases (J40-J47)	118	2.3	8.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	108	2.1	7.7
9	Septicemia (A40-A41)	100	1.9	7.1
10	Influenza and pneumonia (J09-J18)	89	1.7	6.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	1.7	6.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	1.4	5.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	68	1.3	4.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	0.8	3.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	36	0.7	2.6
...	All other causes (residual)	1,003	19.5	71.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 50-54 years				
...	All causes	8,495	100.0	576.5
1	Malignant neoplasms (C00-C97)	2,583	30.4	175.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,808	21.3	122.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	448	5.3	30.4
4	Cerebrovascular diseases (I60-I69)	440	5.2	29.9
5	Diabetes mellitus (E10-E14)	377	4.4	25.6
6	Chronic lower respiratory diseases (J40-J47)	251	3.0	17.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	204	2.4	13.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	197	2.3	13.4
9	Septicemia (A40-A41)	187	2.2	12.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	178	2.1	12.1
11	Influenza and pneumonia (J09-J18)	145	1.7	9.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.4	8.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	0.7	4.1
14	Viral hepatitis (B15-B19)	51	0.6	3.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	0.5	3.1
...	All other causes (residual)	1,400	16.5	95.0
Non-Hispanic, black, female, 55-59 years				
...	All causes	11,547	100.0	846.3
1	Malignant neoplasms (C00-C97)	3,789	32.8	277.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,455	21.3	179.9
3	Diabetes mellitus (E10-E14)	582	5.0	42.7
4	Cerebrovascular diseases (I60-I69)	536	4.6	39.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	422	3.7	30.9
6	Chronic lower respiratory diseases (J40-J47)	395	3.4	29.0
7	Septicemia (A40-A41)	292	2.5	21.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	271	2.3	19.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	238	2.1	17.4
10	Influenza and pneumonia (J09-J18)	189	1.6	13.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	179	1.6	13.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	175	1.5	12.8
13	Viral hepatitis (B15-B19)	99	0.9	7.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.4	3.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	0.4	3.0
...	All other causes (residual)	1,842	16.0	135.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 60-64 years				
...	All causes	13,264	100.0	1,189.8
1	Malignant neoplasms (C00-C97)	4,411	33.3	395.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,917	22.0	261.7
3	Diabetes mellitus (E10-E14)	702	5.3	63.0
4	Cerebrovascular diseases (I60-I69)	699	5.3	62.7
5	Chronic lower respiratory diseases (J40-J47)	487	3.7	43.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	414	3.1	37.1
7	Septicemia (A40-A41)	341	2.6	30.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	2.3	27.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	250	1.9	22.4
10	Influenza and pneumonia (J09-J18)	208	1.6	18.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	204	1.5	18.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	109	0.8	9.8
13	Viral hepatitis (B15-B19)	104	0.8	9.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.4	4.8
15	Aortic aneurysm and dissection (I71)	42	0.3	3.8
15	Pneumonitis due to solids and liquids (J69)	42	0.3	3.8
...	All other causes (residual)	1,975	14.9	177.2
Non-Hispanic, black, female, 65-69 years				
...	All causes	13,352	100.0	1,611.5
1	Malignant neoplasms (C00-C97)	4,415	33.1	532.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,041	22.8	367.0
3	Cerebrovascular diseases (I60-I69)	752	5.6	90.8
4	Diabetes mellitus (E10-E14)	712	5.3	85.9
5	Chronic lower respiratory diseases (J40-J47)	546	4.1	65.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	505	3.8	61.0
7	Septicemia (A40-A41)	353	2.6	42.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	222	1.7	26.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	1.6	25.0
10	Influenza and pneumonia (J09-J18)	188	1.4	22.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	117	0.9	14.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.6	9.8
13	Alzheimer's disease (G30)	64	0.5	7.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	57	0.4	6.9
15	Pneumonitis due to solids and liquids (J69)	52	0.4	6.3
...	All other causes (residual)	2,040	15.3	246.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 70-74 years				
...	All causes	13,754	100.0	2,367.0
1	Malignant neoplasms (C00-C97)	3,994	29.0	687.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,272	23.8	563.1
3	Cerebrovascular diseases (I60-I69)	882	6.4	151.8
4	Diabetes mellitus (E10-E14)	829	6.0	142.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	567	4.1	97.6
6	Chronic lower respiratory diseases (J40-J47)	565	4.1	97.2
7	Septicemia (A40-A41)	368	2.7	63.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	242	1.8	41.6
9	Alzheimer's disease (G30)	231	1.7	39.8
10	Influenza and pneumonia (J09-J18)	215	1.6	37.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	1.3	31.5
12	Pneumonitis due to solids and liquids (J69)	78	0.6	13.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.6	13.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	57	0.4	9.8
15	Aortic aneurysm and dissection (I71)	51	0.4	8.8
...	All other causes (residual)	2,143	15.6	368.8
Non-Hispanic, black, female, 75-79 years				
...	All causes	15,177	100.0	3,611.6
1	Malignant neoplasms (C00-C97)	3,800	25.0	904.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,662	24.1	871.4
3	Cerebrovascular diseases (I60-I69)	1,064	7.0	253.2
4	Diabetes mellitus (E10-E14)	834	5.5	198.5
5	Chronic lower respiratory diseases (J40-J47)	630	4.2	149.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	559	3.7	133.0
7	Alzheimer's disease (G30)	486	3.2	115.7
8	Septicemia (A40-A41)	379	2.5	90.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	309	2.0	73.5
10	Influenza and pneumonia (J09-J18)	262	1.7	62.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	1.3	46.2
12	Pneumonitis due to solids and liquids (J69)	105	0.7	25.0
13	Parkinson's disease (G20-G21)	90	0.6	21.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.5	19.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	58	0.4	13.8
...	All other causes (residual)	2,665	17.6	634.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 80-84 years				
...	All causes	16,634	100.0	5,619.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,249	25.5	1,435.4
2	Malignant neoplasms (C00-C97)	3,221	19.4	1,088.2
3	Cerebrovascular diseases (I60-I69)	1,292	7.8	436.5
4	Diabetes mellitus (E10-E14)	864	5.2	291.9
5	Alzheimer's disease (G30)	855	5.1	288.8
6	Chronic lower respiratory diseases (J40-J47)	623	3.7	210.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	602	3.6	203.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	392	2.4	132.4
9	Septicemia (A40-A41)	358	2.2	120.9
10	Influenza and pneumonia (J09-J18)	349	2.1	117.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	1.1	59.8
12	Pneumonitis due to solids and liquids (J69)	124	0.7	41.9
13	Parkinson's disease (G20-G21)	104	0.6	35.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	95	0.6	32.1
15	Aortic aneurysm and dissection (I71)	59	0.4	19.9
...	All other causes (residual)	3,270	19.7	1,104.7
Non-Hispanic, black, female, 85-89 years				
...	All causes	16,788	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,582	27.3	---
2	Malignant neoplasms (C00-C97)	2,424	14.4	---
3	Cerebrovascular diseases (I60-I69)	1,329	7.9	---
4	Alzheimer's disease (G30)	1,122	6.7	---
5	Diabetes mellitus (E10-E14)	742	4.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	589	3.5	---
7	Chronic lower respiratory diseases (J40-J47)	523	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	418	2.5	---
9	Septicemia (A40-A41)	382	2.3	---
10	Influenza and pneumonia (J09-J18)	370	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	1.1	---
12	Pneumonitis due to solids and liquids (J69)	127	0.8	---
13	Parkinson's disease (G20-G21)	105	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	97	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	61	0.4	---
...	All other causes (residual)	3,736	22.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 90-94 years				
...	All causes	12,930	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,626	28.0	---
2	Malignant neoplasms (C00-C97)	1,270	9.8	---
3	Alzheimer's disease (G30)	1,092	8.4	---
4	Cerebrovascular diseases (I60-I69)	1,066	8.2	---
5	Diabetes mellitus (E10-E14)	454	3.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	355	2.7	---
7	Influenza and pneumonia (J09-J18)	348	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	345	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	326	2.5	---
10	Septicemia (A40-A41)	293	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	1.2	---
12	Pneumonitis due to solids and liquids (J69)	106	0.8	---
13	Parkinson's disease (G20-G21)	70	0.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.4	---
15	Atherosclerosis (I70)	49	0.4	---
...	All other causes (residual)	3,324	25.7	---
Non-Hispanic, black, female, 95-99 years				
...	All causes	5,844	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,711	29.3	---
2	Alzheimer's disease (G30)	553	9.5	---
3	Cerebrovascular diseases (I60-I69)	469	8.0	---
4	Malignant neoplasms (C00-C97)	416	7.1	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	181	3.1	---
6	Influenza and pneumonia (J09-J18)	167	2.9	---
7	Diabetes mellitus (E10-E14)	138	2.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	138	2.4	---
9	Chronic lower respiratory diseases (J40-J47)	119	2.0	---
10	Septicemia (A40-A41)	104	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	1.3	---
12	Pneumonitis due to solids and liquids (J69)	54	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.5	---
14	Nutritional deficiencies (E40-E64)	25	0.4	---
15	Atherosclerosis (I70)	24	0.4	---
...	All other causes (residual)	1,636	28.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 100 years and over				
...	All causes	2,127	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	632	29.7	---
2	Alzheimer's disease (G30)	211	9.9	---
3	Cerebrovascular diseases (I60-I69)	162	7.6	---
4	Malignant neoplasms (C00-C97)	88	4.1	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	74	3.5	---
6	Influenza and pneumonia (J09-J18)	59	2.8	---
7	Diabetes mellitus (E10-E14)	45	2.1	---
8	Chronic lower respiratory diseases (J40-J47)	42	2.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.0	---
10	Septicemia (A40-A41)	35	1.6	---
11	Pneumonitis due to solids and liquids (J69)	22	1.0	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	0.9	---
13	Atherosclerosis (I70)	16	0.8	---
14	Nutritional deficiencies (E40-E64)	10	0.5	---
15	Anemias (D50-D64)	8	0.4	---
...	All other causes (residual)	662	31.1	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014