

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	133,004	100.0	300.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,921	21.7	65.3
2	Malignant neoplasms (C00-C97)	26,633	20.0	60.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,052	9.1	27.2
4	Cerebrovascular diseases (I60-I69)	7,005	5.3	15.8
5	Diabetes mellitus (E10-E14)	6,287	4.7	14.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,592	2.7	8.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,524	2.6	8.0
8	Chronic lower respiratory diseases (J40-J47)	3,310	2.5	7.5
9	Influenza and pneumonia (J10-J18)	2,966	2.2	6.7
10	Certain conditions originating in the perinatal period (P00-P96)	2,804	2.1	6.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,592	1.9	5.8
12	Alzheimer's disease (G30)	2,399	1.8	5.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,177	1.6	4.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,952	1.5	4.4
15	Septicemia (A40-A41)	1,829	1.4	4.1
...	All other causes (Residual)	24,961	18.8	56.3
Hispanic, both sexes, 1-4 years				
...	All causes	987	100.0	26.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	361	36.6	9.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	12.7	3.3
3	Malignant neoplasms (C00-C97)	82	8.3	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	6.3	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	3.1	0.8
6	Influenza and pneumonia (J10-J18)	28	2.8	0.7
7	Septicemia (A40-A41)	18	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
10	Cerebrovascular diseases (I60-I69)	9	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
12	Meningitis (G00,G03)	6	0.6	*
13	Anemias (D50-D64)	4	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	223	22.6	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,137	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	36.2	5.1
2	Malignant neoplasms (C00-C97)	183	16.1	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	8.4	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	5.4	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.0	0.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	2.2	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.6	*
8	Cerebrovascular diseases (I60-I69)	17	1.5	*
9	Septicemia (A40-A41)	16	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	16	1.4	*
11	Influenza and pneumonia (J10-J18)	15	1.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	10	0.9	*
13	Diabetes mellitus (E10-E14)	6	0.5	*
14	Anemias (D50-D64)	3	0.3	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
14	Diseases of appendix (K35-K38)	3	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	206	18.1	2.6
Hispanic, both sexes, 15-24 years				
...	All causes	5,932	100.0	80.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,662	44.9	36.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,343	22.6	18.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	543	9.2	7.4
4	Malignant neoplasms (C00-C97)	321	5.4	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	130	2.2	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	1.2	1.0
7	Cerebrovascular diseases (I60-I69)	52	0.9	0.7
8	Pregnancy, childbirth and the puerperium (O00-099)	39	0.7	0.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.5	0.4
10	Legal intervention (Y35,Y89.0)	25	0.4	0.3
11	Influenza and pneumonia (J10-J18)	24	0.4	0.3
12	Septicemia (A40-A41)	22	0.4	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.4	0.3
14	Diabetes mellitus (E10-E14)	18	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	15	0.3	*
...	All other causes (Residual)	614	10.4	8.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	6,592	100.0	81.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,400	36.4	29.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	995	15.1	12.3
3	Malignant neoplasms (C00-C97)	579	8.8	7.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	574	8.7	7.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	374	5.7	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	195	3.0	2.4
7	Cerebrovascular diseases (I60-I69)	94	1.4	1.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	85	1.3	1.1
9	Diabetes mellitus (E10-E14)	74	1.1	0.9
10	Pregnancy, childbirth and the puerperium (O00-099)	70	1.1	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	0.9	0.7
12	Influenza and pneumonia (J10-J18)	49	0.7	0.6
13	Septicemia (A40-A41)	41	0.6	0.5
14	Legal intervention (Y35,Y89.0)	39	0.6	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.6	0.5
...	All other causes (Residual)	929	14.1	11.5
Hispanic, both sexes, 35-44 years				
...	All causes	8,951	100.0	134.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,102	23.5	31.6
2	Malignant neoplasms (C00-C97)	1,440	16.1	21.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	930	10.4	14.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	584	6.5	8.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	537	6.0	8.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	444	5.0	6.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	441	4.9	6.6
8	Cerebrovascular diseases (I60-I69)	308	3.4	4.6
9	Diabetes mellitus (E10-E14)	185	2.1	2.8
10	Viral hepatitis (B15-B19)	104	1.2	1.6
11	Septicemia (A40-A41)	79	0.9	1.2
12	Influenza and pneumonia (J10-J18)	77	0.9	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	0.7	1.0
14	Chronic lower respiratory diseases (J40-J47)	49	0.5	0.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	0.5	0.6
...	All other causes (Residual)	1,563	17.5	23.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	14,035	100.0	310.7
1	Malignant neoplasms (C00-C97)	3,374	24.0	74.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,337	16.7	51.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,609	11.5	35.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,072	7.6	23.7
5	Cerebrovascular diseases (I60-I69)	652	4.6	14.4
6	Diabetes mellitus (E10-E14)	595	4.2	13.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	527	3.8	11.7
8	Viral hepatitis (B15-B19)	423	3.0	9.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	294	2.1	6.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	263	1.9	5.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	222	1.6	4.9
12	Septicemia (A40-A41)	179	1.3	4.0
13	Influenza and pneumonia (J10-J18)	165	1.2	3.7
14	Chronic lower respiratory diseases (J40-J47)	118	0.8	2.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	106	0.8	2.3
...	All other causes (Residual)	2,099	15.0	46.5
Hispanic, both sexes, 55-64 years				
...	All causes	16,705	100.0	657.6
1	Malignant neoplasms (C00-C97)	5,044	30.2	198.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,612	21.6	142.2
3	Diabetes mellitus (E10-E14)	1,081	6.5	42.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	870	5.2	34.2
5	Cerebrovascular diseases (I60-I69)	869	5.2	34.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	844	5.1	33.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	366	2.2	14.4
8	Viral hepatitis (B15-B19)	329	2.0	13.0
9	Chronic lower respiratory diseases (J40-J47)	305	1.8	12.0
10	Septicemia (A40-A41)	251	1.5	9.9
11	Influenza and pneumonia (J10-J18)	249	1.5	9.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	194	1.2	7.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	181	1.1	7.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	153	0.9	6.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	0.6	3.7
...	All other causes (Residual)	2,263	13.5	89.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	21,061	100.0	1,531.0
1	Malignant neoplasms (C00-C97)	6,354	30.2	461.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,189	24.6	377.2
3	Diabetes mellitus (E10-E14)	1,623	7.7	118.0
4	Cerebrovascular diseases (I60-I69)	1,201	5.7	87.3
5	Chronic lower respiratory diseases (J40-J47)	626	3.0	45.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	591	2.8	43.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	554	2.6	40.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	517	2.5	37.6
9	Influenza and pneumonia (J10-J18)	423	2.0	30.8
10	Septicemia (A40-A41)	342	1.6	24.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	233	1.1	16.9
12	Alzheimer's disease (G30)	172	0.8	12.5
13	Viral hepatitis (B15-B19)	148	0.7	10.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	122	0.6	8.9
14	Parkinson's disease (G20-G21)	122	0.6	8.9
...	All other causes (Residual)	2,844	13.5	206.7
Hispanic, both sexes, 75-84 years				
...	All causes	28,772	100.0	3,745.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,180	28.4	1,064.9
2	Malignant neoplasms (C00-C97)	6,483	22.5	844.0
3	Cerebrovascular diseases (I60-I69)	2,013	7.0	262.1
4	Diabetes mellitus (E10-E14)	1,758	6.1	228.9
5	Chronic lower respiratory diseases (J40-J47)	1,179	4.1	153.5
6	Influenza and pneumonia (J10-J18)	877	3.0	114.2
7	Alzheimer's disease (G30)	848	2.9	110.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	764	2.7	99.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	575	2.0	74.9
10	Septicemia (A40-A41)	500	1.7	65.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	418	1.5	54.4
12	Parkinson's disease (G20-G21)	319	1.1	41.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	308	1.1	40.1
14	Pneumonitis due to solids and liquids (J69)	199	0.7	25.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	192	0.7	25.0
...	All other causes (Residual)	4,159	14.5	541.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	23,061	100.0	9,022.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,027	34.8	3,140.6
2	Malignant neoplasms (C00-C97)	2,754	11.9	1,077.5
3	Cerebrovascular diseases (I60-I69)	1,756	7.6	687.0
4	Alzheimer's disease (G30)	1,315	5.7	514.5
5	Influenza and pneumonia (J10-J18)	999	4.3	390.9
6	Chronic lower respiratory diseases (J40-J47)	957	4.1	374.4
7	Diabetes mellitus (E10-E14)	945	4.1	369.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	527	2.3	206.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	376	1.6	147.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	365	1.6	142.8
11	Septicemia (A40-A41)	314	1.4	122.9
12	Parkinson's disease (G20-G21)	240	1.0	93.9
13	Pneumonitis due to solids and liquids (J69)	198	0.9	77.5
14	Atherosclerosis (I70)	129	0.6	50.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	124	0.5	48.5
...	All other causes (Residual)	4,035	17.5	1,578.7
Hispanic, male, all ages				
...	All causes	74,250	100.0	323.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,518	20.9	67.7
2	Malignant neoplasms (C00-C97)	13,856	18.7	60.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,102	12.3	39.7
4	Cerebrovascular diseases (I60-I69)	3,269	4.4	14.3
5	Diabetes mellitus (E10-E14)	3,140	4.2	13.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,004	4.0	13.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,527	3.4	11.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,813	2.4	7.9
9	Chronic lower respiratory diseases (J40-J47)	1,708	2.3	7.5
10	Certain conditions originating in the perinatal period (P00-P96)	1,565	2.1	6.8
11	Influenza and pneumonia (J10-J18)	1,439	1.9	6.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,295	1.7	5.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,275	1.7	5.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,046	1.4	4.6
15	Septicemia (A40-A41)	932	1.3	4.1
...	All other causes (Residual)	12,761	17.2	55.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	549	100.0	28.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	39.2	11.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	12.6	3.6
3	Malignant neoplasms (C00-C97)	49	8.9	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	6.4	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.5	*
6	Influenza and pneumonia (J10-J18)	13	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
8	Septicemia (A40-A41)	6	1.1	*
8	Cerebrovascular diseases (I60-I69)	6	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
12	Meningococcal infection (A39)	2	0.4	*
12	Anemias (D50-D64)	2	0.4	*
12	Meningitis (G00,G03)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	111	20.2	5.8
Hispanic, male, 5-14 years				
...	All causes	673	100.0	16.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	259	38.5	6.3
2	Malignant neoplasms (C00-C97)	95	14.1	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	9.4	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	5.1	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.0	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	2.7	*
7	Chronic lower respiratory diseases (J40-J47)	13	1.9	*
8	Cerebrovascular diseases (I60-I69)	12	1.8	*
9	Septicemia (A40-A41)	10	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	1.0	*
12	Influenza and pneumonia (J10-J18)	6	0.9	*
13	Diabetes mellitus (E10-E14)	3	0.4	*
13	Pneumonitis due to solids and liquids (J69)	3	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	112	16.6	2.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,712	100.0	120.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,169	46.0	55.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,210	25.7	31.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	452	9.6	11.6
4	Malignant neoplasms (C00-C97)	201	4.3	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	2.1	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	1.0	1.2
7	Cerebrovascular diseases (I60-I69)	31	0.7	0.8
8	Legal intervention (Y35,Y89.0)	25	0.5	0.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.4	*
10	Diabetes mellitus (E10-E14)	14	0.3	*
11	Influenza and pneumonia (J10-J18)	13	0.3	*
12	Septicemia (A40-A41)	11	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.2	*
13	Chronic lower respiratory diseases (J40-J47)	10	0.2	*
15	Meningitis (G00,G03)	6	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.1	*
...	All other causes (Residual)	388	8.2	9.9
Hispanic, male, 25-34 years				
...	All causes	5,024	100.0	112.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,000	39.8	44.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	880	17.5	19.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	486	9.7	10.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	291	5.8	6.5
5	Malignant neoplasms (C00-C97)	270	5.4	6.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	152	3.0	3.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	64	1.3	1.4
8	Cerebrovascular diseases (I60-I69)	61	1.2	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	0.8	0.9
10	Legal intervention (Y35,Y89.0)	38	0.8	0.9
11	Diabetes mellitus (E10-E14)	36	0.7	0.8
12	Influenza and pneumonia (J10-J18)	33	0.7	0.7
13	Septicemia (A40-A41)	22	0.4	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.4	*
15	Aortic aneurysm and dissection (I71)	16	0.3	*
...	All other causes (Residual)	617	12.3	13.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	6,223	100.0	176.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,691	27.2	48.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	709	11.4	20.1
3	Malignant neoplasms (C00-C97)	565	9.1	16.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	456	7.3	12.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	440	7.1	12.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	373	6.0	10.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	350	5.6	9.9
8	Cerebrovascular diseases (I60-I69)	193	3.1	5.5
9	Diabetes mellitus (E10-E14)	118	1.9	3.3
10	Viral hepatitis (B15-B19)	88	1.4	2.5
11	Influenza and pneumonia (J10-J18)	50	0.8	1.4
12	Septicemia (A40-A41)	43	0.7	1.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.7	1.2
14	Aortic aneurysm and dissection (I71)	34	0.5	1.0
15	Chronic lower respiratory diseases (J40-J47)	26	0.4	0.7
...	All other causes (Residual)	1,044	16.8	29.6
Hispanic, male, 45-54 years				
...	All causes	9,234	100.0	403.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,729	18.7	75.6
2	Malignant neoplasms (C00-C97)	1,633	17.7	71.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,236	13.4	54.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	855	9.3	37.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	426	4.6	18.6
6	Cerebrovascular diseases (I60-I69)	389	4.2	17.0
7	Diabetes mellitus (E10-E14)	370	4.0	16.2
8	Viral hepatitis (B15-B19)	319	3.5	13.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	232	2.5	10.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	209	2.3	9.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	1.5	6.2
12	Septicemia (A40-A41)	112	1.2	4.9
13	Influenza and pneumonia (J10-J18)	102	1.1	4.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	63	0.7	2.8
14	Chronic lower respiratory diseases (J40-J47)	63	0.7	2.8
...	All other causes (Residual)	1,355	14.7	59.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	10,271	100.0	843.6
1	Malignant neoplasms (C00-C97)	2,737	26.6	224.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,463	24.0	202.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	659	6.4	54.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	623	6.1	51.2
5	Diabetes mellitus (E10-E14)	608	5.9	49.9
6	Cerebrovascular diseases (I60-I69)	501	4.9	41.1
7	Viral hepatitis (B15-B19)	214	2.1	17.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	1.9	15.8
9	Chronic lower respiratory diseases (J40-J47)	171	1.7	14.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	157	1.5	12.9
11	Influenza and pneumonia (J10-J18)	153	1.5	12.6
12	Septicemia (A40-A41)	138	1.3	11.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	128	1.2	10.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	102	1.0	8.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	0.8	6.7
...	All other causes (Residual)	1,343	13.1	110.3
Hispanic, male, 65-74 years				
...	All causes	11,780	100.0	1,910.7
1	Malignant neoplasms (C00-C97)	3,544	30.1	574.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,117	26.5	505.6
3	Diabetes mellitus (E10-E14)	836	7.1	135.6
4	Cerebrovascular diseases (I60-I69)	617	5.2	100.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	351	3.0	56.9
6	Chronic lower respiratory diseases (J40-J47)	349	3.0	56.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	334	2.8	54.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	297	2.5	48.2
9	Influenza and pneumonia (J10-J18)	226	1.9	36.7
10	Septicemia (A40-A41)	188	1.6	30.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	117	1.0	19.0
12	Alzheimer's disease (G30)	81	0.7	13.1
13	Viral hepatitis (B15-B19)	74	0.6	12.0
14	Parkinson's disease (G20-G21)	72	0.6	11.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.6	11.2
...	All other causes (Residual)	1,508	12.8	244.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	14,209	100.0	4,492.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,138	29.1	1,308.4
2	Malignant neoplasms (C00-C97)	3,474	24.4	1,098.4
3	Cerebrovascular diseases (I60-I69)	926	6.5	292.8
4	Diabetes mellitus (E10-E14)	815	5.7	257.7
5	Chronic lower respiratory diseases (J40-J47)	621	4.4	196.3
6	Influenza and pneumonia (J10-J18)	416	2.9	131.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	360	2.5	113.8
8	Alzheimer's disease (G30)	339	2.4	107.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	303	2.1	95.8
10	Septicemia (A40-A41)	242	1.7	76.5
11	Parkinson's disease (G20-G21)	202	1.4	63.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	193	1.4	61.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	123	0.9	38.9
14	Pneumonitis due to solids and liquids (J69)	101	0.7	31.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	96	0.7	30.4
...	All other causes (Residual)	1,860	13.1	588.1
Hispanic, male, 85 years and over				
...	All causes	8,367	100.0	9,435.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,889	34.5	3,257.9
2	Malignant neoplasms (C00-C97)	1,277	15.3	1,440.1
3	Cerebrovascular diseases (I60-I69)	516	6.2	581.9
4	Chronic lower respiratory diseases (J40-J47)	430	5.1	484.9
5	Influenza and pneumonia (J10-J18)	391	4.7	440.9
6	Alzheimer's disease (G30)	359	4.3	404.8
7	Diabetes mellitus (E10-E14)	338	4.0	381.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	214	2.6	241.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	1.9	174.8
10	Parkinson's disease (G20-G21)	134	1.6	151.1
11	Septicemia (A40-A41)	123	1.5	138.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	117	1.4	131.9
13	Pneumonitis due to solids and liquids (J69)	77	0.9	86.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.6	53.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	46	0.5	51.9
...	All other causes (Residual)	1,254	15.0	1,414.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	58,754	100.0	274.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,403	22.8	62.6
2	Malignant neoplasms (C00-C97)	12,777	21.7	59.7
3	Cerebrovascular diseases (I60-I69)	3,736	6.4	17.5
4	Diabetes mellitus (E10-E14)	3,147	5.4	14.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,950	5.0	13.8
6	Chronic lower respiratory diseases (J40-J47)	1,602	2.7	7.5
7	Alzheimer's disease (G30)	1,587	2.7	7.4
8	Influenza and pneumonia (J10-J18)	1,527	2.6	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,297	2.2	6.1
10	Certain conditions originating in the perinatal period (P00-P96)	1,239	2.1	5.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,065	1.8	5.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	906	1.5	4.2
13	Septicemia (A40-A41)	897	1.5	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	697	1.2	3.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	520	0.9	2.4
...	All other causes (Residual)	11,404	19.4	53.3
Hispanic, female, 1-4 years				
...	All causes	438	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	33.3	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	12.8	3.1
3	Malignant neoplasms (C00-C97)	33	7.5	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	6.2	1.5
5	Influenza and pneumonia (J10-J18)	15	3.4	*
6	Septicemia (A40-A41)	12	2.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
9	Meningitis (G00,G03)	4	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	4	0.9	*
12	Cerebrovascular diseases (I60-I69)	3	0.7	*
13	Anemias (D50-D64)	2	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	108	24.7	5.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	464	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	33.0	3.9
2	Malignant neoplasms (C00-C97)	88	19.0	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.9	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	5.8	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.9	*
6	Influenza and pneumonia (J10-J18)	9	1.9	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	1.5	*
9	Septicemia (A40-A41)	6	1.3	*
10	Cerebrovascular diseases (I60-I69)	5	1.1	*
11	Diabetes mellitus (E10-E14)	3	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
14	Meningococcal infection (A39)	2	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
14	Anemias (D50-D64)	2	0.4	*
14	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (Residual)	93	20.0	2.4
Hispanic, female, 15-24 years				
...	All causes	1,220	100.0	35.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	493	40.4	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	10.9	3.8
3	Malignant neoplasms (C00-C97)	120	9.8	3.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	7.5	2.6
5	Pregnancy, childbirth and the puerperium (O00-099)	39	3.2	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.5	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	1.9	0.7
8	Cerebrovascular diseases (I60-I69)	21	1.7	0.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.0	*
11	Septicemia (A40-A41)	11	0.9	*
11	Influenza and pneumonia (J10-J18)	11	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	207	17.0	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,568	100.0	43.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	400	25.5	11.0
2	Malignant neoplasms (C00-C97)	309	19.7	8.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	115	7.3	3.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	5.6	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	5.3	2.3
6	Pregnancy, childbirth and the puerperium (O00-099)	70	4.5	1.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	43	2.7	1.2
8	Diabetes mellitus (E10-E14)	38	2.4	1.0
9	Cerebrovascular diseases (I60-I69)	33	2.1	0.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	21	1.3	0.6
11	Septicemia (A40-A41)	19	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.1	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.1	*
14	Influenza and pneumonia (J10-J18)	16	1.0	*
15	Chronic lower respiratory diseases (J40-J47)	14	0.9	*
...	All other causes (Residual)	283	18.0	7.8
Hispanic, female, 35-44 years				
...	All causes	2,728	100.0	87.1
1	Malignant neoplasms (C00-C97)	875	32.1	27.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	411	15.1	13.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	221	8.1	7.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	128	4.7	4.1
5	Cerebrovascular diseases (I60-I69)	115	4.2	3.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	3.6	3.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	94	3.4	3.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	2.5	2.2
9	Diabetes mellitus (E10-E14)	67	2.5	2.1
10	Septicemia (A40-A41)	36	1.3	1.1
11	Pregnancy, childbirth and the puerperium (O00-099)	34	1.2	1.1
12	Influenza and pneumonia (J10-J18)	27	1.0	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.9	0.8
14	Chronic lower respiratory diseases (J40-J47)	23	0.8	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.7	*
...	All other causes (Residual)	490	18.0	15.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	4,801	100.0	215.3
1	Malignant neoplasms (C00-C97)	1,741	36.3	78.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	608	12.7	27.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	373	7.8	16.7
4	Cerebrovascular diseases (I60-I69)	263	5.5	11.8
5	Diabetes mellitus (E10-E14)	225	4.7	10.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	217	4.5	9.7
7	Viral hepatitis (B15-B19)	104	2.2	4.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	101	2.1	4.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	1.7	3.6
10	Septicemia (A40-A41)	67	1.4	3.0
11	Influenza and pneumonia (J10-J18)	63	1.3	2.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	1.3	2.8
13	Chronic lower respiratory diseases (J40-J47)	55	1.1	2.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	1.1	2.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	0.9	1.9
...	All other causes (Residual)	744	15.5	33.4
Hispanic, female, 55-64 years				
...	All causes	6,434	100.0	486.5
1	Malignant neoplasms (C00-C97)	2,307	35.9	174.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,149	17.9	86.9
3	Diabetes mellitus (E10-E14)	473	7.4	35.8
4	Cerebrovascular diseases (I60-I69)	368	5.7	27.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	3.4	16.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	211	3.3	16.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	174	2.7	13.2
8	Chronic lower respiratory diseases (J40-J47)	134	2.1	10.1
9	Viral hepatitis (B15-B19)	115	1.8	8.7
10	Septicemia (A40-A41)	113	1.8	8.5
11	Influenza and pneumonia (J10-J18)	96	1.5	7.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	1.2	6.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.6	3.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	37	0.6	2.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	0.4	1.9
...	All other causes (Residual)	892	13.9	67.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	9,281	100.0	1,222.7
1	Malignant neoplasms (C00-C97)	2,810	30.3	370.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,072	22.3	273.0
3	Diabetes mellitus (E10-E14)	787	8.5	103.7
4	Cerebrovascular diseases (I60-I69)	584	6.3	76.9
5	Chronic lower respiratory diseases (J40-J47)	277	3.0	36.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	257	2.8	33.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	240	2.6	31.6
8	Influenza and pneumonia (J10-J18)	197	2.1	26.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	2.0	24.1
10	Septicemia (A40-A41)	154	1.7	20.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	116	1.2	15.3
12	Alzheimer's disease (G30)	91	1.0	12.0
13	Viral hepatitis (B15-B19)	74	0.8	9.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.6	7.0
15	Parkinson's disease (G20-G21)	50	0.5	6.6
...	All other causes (Residual)	1,336	14.4	176.0
Hispanic, female, 75-84 years				
...	All causes	14,563	100.0	3,222.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,042	27.8	894.5
2	Malignant neoplasms (C00-C97)	3,009	20.7	665.9
3	Cerebrovascular diseases (I60-I69)	1,087	7.5	240.6
4	Diabetes mellitus (E10-E14)	943	6.5	208.7
5	Chronic lower respiratory diseases (J40-J47)	558	3.8	123.5
6	Alzheimer's disease (G30)	509	3.5	112.6
7	Influenza and pneumonia (J10-J18)	461	3.2	102.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	404	2.8	89.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	1.9	60.2
10	Septicemia (A40-A41)	258	1.8	57.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	225	1.5	49.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	185	1.3	40.9
13	Parkinson's disease (G20-G21)	117	0.8	25.9
14	Pneumonitis due to solids and liquids (J69)	98	0.7	21.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	96	0.7	21.2
...	All other causes (Residual)	2,299	15.8	508.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	14,694	100.0	8,803.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,138	35.0	3,078.3
2	Malignant neoplasms (C00-C97)	1,477	10.1	884.9
3	Cerebrovascular diseases (I60-I69)	1,240	8.4	742.9
4	Alzheimer's disease (G30)	956	6.5	572.8
5	Influenza and pneumonia (J10-J18)	608	4.1	364.3
6	Diabetes mellitus (E10-E14)	607	4.1	363.7
7	Chronic lower respiratory diseases (J40-J47)	527	3.6	315.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	313	2.1	187.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	259	1.8	155.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	1.4	125.8
11	Septicemia (A40-A41)	191	1.3	114.4
12	Pneumonitis due to solids and liquids (J69)	121	0.8	72.5
13	Parkinson's disease (G20-G21)	106	0.7	63.5
14	Atherosclerosis (I70)	89	0.6	53.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.5	46.1
...	All other causes (Residual)	2,775	18.9	1,662.6
White, non-Hispanic, both sexes, all ages				
...	All causes	1,944,617	100.0	968.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	516,883	26.6	257.4
2	Malignant neoplasms (C00-C97)	455,978	23.4	227.1
3	Chronic lower respiratory diseases (J40-J47)	111,559	5.7	55.6
4	Cerebrovascular diseases (I60-I69)	108,886	5.6	54.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91,830	4.7	45.7
6	Alzheimer's disease (G30)	64,660	3.3	32.2
7	Diabetes mellitus (E10-E14)	50,950	2.6	25.4
8	Influenza and pneumonia (J10-J18)	46,419	2.4	23.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33,208	1.7	16.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,952	1.4	13.9
11	Septicemia (A40-A41)	25,556	1.3	12.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,605	1.1	10.3
13	Parkinson's disease (G20-G21)	17,806	0.9	8.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,034	0.9	8.5
15	Pneumonitis due to solids and liquids (J69)	14,563	0.7	7.3
...	All other causes (Residual)	340,728	17.5	169.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,279	100.0	25.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	814	35.7	8.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	253	11.1	2.8
3	Malignant neoplasms (C00-C97)	209	9.2	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	5.8	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	3.5	0.9
6	Influenza and pneumonia (J10-J18)	54	2.4	0.6
7	Septicemia (A40-A41)	44	1.9	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	1.9	0.5
9	Cerebrovascular diseases (I60-I69)	32	1.4	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	29	1.3	0.3
11	Chronic lower respiratory diseases (J40-J47)	15	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.6	*
13	Meningococcal infection (A39)	10	0.4	*
13	Meningitis (G00,G03)	10	0.4	*
15	Anemias (D50-D64)	9	0.4	*
...	All other causes (Residual)	531	23.3	5.8
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	3,353	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,267	37.8	5.3
2	Malignant neoplasms (C00-C97)	540	16.1	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	5.9	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	143	4.3	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	134	4.0	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	128	3.8	0.5
7	Cerebrovascular diseases (I60-I69)	51	1.5	0.2
8	Septicemia (A40-A41)	47	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	1.2	0.2
10	Influenza and pneumonia (J10-J18)	39	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	30	0.9	0.1
12	Diabetes mellitus (E10-E14)	13	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.4	*
14	Anemias (D50-D64)	11	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.3	*
...	All other causes (Residual)	688	20.5	2.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	20,055	100.0	76.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,203	55.9	42.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,944	14.7	11.2
3	Malignant neoplasms (C00-C97)	980	4.9	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	900	4.5	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	608	3.0	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	272	1.4	1.0
7	Influenza and pneumonia (J10-J18)	114	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	105	0.5	0.4
9	Diabetes mellitus (E10-E14)	89	0.4	0.3
10	Pregnancy, childbirth and the puerperium (O00-099)	78	0.4	0.3
11	Septicemia (A40-A41)	69	0.3	0.3
12	Chronic lower respiratory diseases (J40-J47)	54	0.3	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	0.3	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.2	0.2
15	Legal intervention (Y35,Y89.0)	30	0.1	0.1
...	All other causes (Residual)	2,506	12.5	9.5
White, non-Hispanic, both sexes, 25-34 years				
...	All causes	24,606	100.0	102.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,085	41.0	41.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,689	15.0	15.3
3	Malignant neoplasms (C00-C97)	2,226	9.0	9.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,774	7.2	7.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	931	3.8	3.9
6	Diabetes mellitus (E10-E14)	334	1.4	1.4
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	292	1.2	1.2
8	Cerebrovascular diseases (I60-I69)	251	1.0	1.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	209	0.8	0.9
10	Influenza and pneumonia (J10-J18)	189	0.8	0.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	166	0.7	0.7
12	Septicemia (A40-A41)	145	0.6	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	123	0.5	0.5
14	Chronic lower respiratory diseases (J40-J47)	115	0.5	0.5
15	Pregnancy, childbirth and the puerperium (O00-099)	113	0.5	0.5
...	All other causes (Residual)	3,964	16.1	16.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 35-44 years				
...	All causes	53,198	100.0	184.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,559	23.6	43.6
2	Malignant neoplasms (C00-C97)	9,327	17.5	32.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,960	15.0	27.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,508	10.4	19.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,693	3.2	5.9
6	Diabetes mellitus (E10-E14)	1,257	2.4	4.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	1,193	2.2	4.1
8	Cerebrovascular diseases (I60-I69)	1,086	2.0	3.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,039	2.0	3.6
10	Influenza and pneumonia (J10-J18)	549	1.0	1.9
11	Chronic lower respiratory diseases (J40-J47)	539	1.0	1.9
12	Septicemia (A40-A41)	490	0.9	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	347	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	340	0.6	1.2
15	Viral hepatitis (B15-B19)	318	0.6	1.1
...	All other causes (Residual)	8,993	16.9	31.2
White, non-Hispanic, both sexes, 45-54 years				
...	All causes	127,969	100.0	406.6
1	Malignant neoplasms (C00-C97)	36,458	28.5	115.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,138	20.4	83.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,619	11.4	46.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,614	5.2	21.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,541	4.3	17.6
6	Cerebrovascular diseases (I60-I69)	3,495	2.7	11.1
7	Diabetes mellitus (E10-E14)	3,465	2.7	11.0
8	Chronic lower respiratory diseases (J40-J47)	3,113	2.4	9.9
9	Viral hepatitis (B15-B19)	1,991	1.6	6.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,357	1.1	4.3
11	Septicemia (A40-A41)	1,337	1.0	4.2
12	Influenza and pneumonia (J10-J18)	1,327	1.0	4.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,113	0.9	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,021	0.8	3.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	591	0.5	1.9
...	All other causes (Residual)	19,789	15.5	62.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-64 years				
...	All causes	210,318	100.0	861.2
1	Malignant neoplasms (C00-C97)	79,381	37.7	325.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,053	22.8	196.8
3	Chronic lower respiratory diseases (J40-J47)	10,666	5.1	43.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,653	4.1	35.4
5	Diabetes mellitus (E10-E14)	7,398	3.5	30.3
6	Cerebrovascular diseases (I60-I69)	6,555	3.1	26.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,408	2.6	22.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,167	2.0	17.1
9	Septicemia (A40-A41)	2,679	1.3	11.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,512	1.2	10.3
11	Influenza and pneumonia (J10-J18)	2,313	1.1	9.5
12	Viral hepatitis (B15-B19)	1,305	0.6	5.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,185	0.6	4.9
14	Aortic aneurysm and dissection (I71)	1,061	0.5	4.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,020	0.5	4.2
...	All other causes (Residual)	27,962	13.3	114.5
White, non-Hispanic, both sexes, 65-74 years				
...	All causes	307,475	100.0	2,055.0
1	Malignant neoplasms (C00-C97)	111,945	36.4	748.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,146	23.1	475.5
3	Chronic lower respiratory diseases (J40-J47)	25,397	8.3	169.7
4	Cerebrovascular diseases (I60-I69)	13,037	4.2	87.1
5	Diabetes mellitus (E10-E14)	10,305	3.4	68.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,745	2.2	45.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,807	1.6	32.1
8	Influenza and pneumonia (J10-J18)	4,629	1.5	30.9
9	Septicemia (A40-A41)	4,335	1.4	29.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	3,839	1.2	25.7
11	Alzheimer's disease (G30)	3,253	1.1	21.7
12	Aortic aneurysm and dissection (I71)	2,203	0.7	14.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,167	0.7	14.5
14	Parkinson's disease (G20-G21)	1,996	0.6	13.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,969	0.6	13.2
...	All other causes (Residual)	39,702	12.9	265.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 75-84 years				
...	All causes	564,113	100.0	5,205.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	151,129	26.8	1,394.5
2	Malignant neoplasms (C00-C97)	140,646	24.9	1,297.8
3	Chronic lower respiratory diseases (J40-J47)	43,173	7.7	398.4
4	Cerebrovascular diseases (I60-I69)	35,968	6.4	331.9
5	Alzheimer's disease (G30)	20,273	3.6	187.1
6	Diabetes mellitus (E10-E14)	16,147	2.9	149.0
7	Influenza and pneumonia (J10-J18)	13,870	2.5	128.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,943	2.1	110.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,294	2.0	104.2
10	Septicemia (A40-A41)	8,495	1.5	78.4
11	Parkinson's disease (G20-G21)	8,322	1.5	76.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,961	0.9	45.8
13	Pneumonitis due to solids and liquids (J69)	4,552	0.8	42.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,138	0.7	38.2
15	Aortic aneurysm and dissection (I71)	4,087	0.7	37.7
...	All other causes (Residual)	85,115	15.1	785.4
White, non-Hispanic, both sexes, 85 years and over				
...	All causes	618,253	100.0	13,688.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	209,695	33.9	4,642.7
2	Malignant neoplasms (C00-C97)	74,213	12.0	1,643.1
3	Cerebrovascular diseases (I60-I69)	48,250	7.8	1,068.3
4	Alzheimer's disease (G30)	40,494	6.5	896.5
5	Chronic lower respiratory diseases (J40-J47)	28,446	4.6	629.8
6	Influenza and pneumonia (J10-J18)	23,232	3.8	514.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,363	2.2	295.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12,863	2.1	284.8
9	Diabetes mellitus (E10-E14)	11,933	1.9	264.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,165	1.3	180.8
11	Septicemia (A40-A41)	7,824	1.3	173.2
12	Pneumonitis due to solids and liquids (J69)	7,411	1.2	164.1
13	Parkinson's disease (G20-G21)	7,090	1.1	157.0
14	Atherosclerosis (I70)	4,302	0.7	95.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,685	0.6	81.6
...	All other causes (Residual)	117,287	19.0	2,596.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	947,966	100.0	962.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	256,459	27.1	260.3
2	Malignant neoplasms (C00-C97)	236,393	24.9	239.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57,758	6.1	58.6
4	Chronic lower respiratory diseases (J40-J47)	52,270	5.5	53.0
5	Cerebrovascular diseases (I60-I69)	41,941	4.4	42.6
6	Diabetes mellitus (E10-E14)	25,931	2.7	26.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,952	2.3	22.3
8	Influenza and pneumonia (J10-J18)	20,863	2.2	21.2
9	Alzheimer's disease (G30)	18,844	2.0	19.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,415	1.7	16.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13,212	1.4	13.4
12	Septicemia (A40-A41)	11,526	1.2	11.7
13	Parkinson's disease (G20-G21)	10,277	1.1	10.4
14	Pneumonitis due to solids and liquids (J69)	7,694	0.8	7.8
15	Aortic aneurysm and dissection (I71)	6,512	0.7	6.6
...	All other causes (Residual)	149,919	15.8	152.1
White, non-Hispanic, male, 1-4 years				
...	All causes	1,247	100.0	26.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	458	36.7	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	9.8	2.6
3	Malignant neoplasms (C00-C97)	116	9.3	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	6.2	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.3	0.9
6	Influenza and pneumonia (J10-J18)	29	2.3	0.6
7	Septicemia (A40-A41)	24	1.9	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	21	1.7	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.3	*
10	Cerebrovascular diseases (I60-I69)	12	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	10	0.8	*
12	Meningococcal infection (A39)	7	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Anemias (D50-D64)	3	0.2	*
15	Meningitis (G00,G03)	3	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
...	All other causes (Residual)	293	23.5	6.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	1,983	100.0	16.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	766	38.6	6.2
2	Malignant neoplasms (C00-C97)	309	15.6	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	5.7	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	5.0	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	3.7	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	3.4	0.6
7	Cerebrovascular diseases (I60-I69)	34	1.7	0.3
8	Septicemia (A40-A41)	28	1.4	0.2
9	Influenza and pneumonia (J10-J18)	25	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	22	1.1	0.2
12	Diabetes mellitus (E10-E14)	8	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.4	*
14	Aortic aneurysm and dissection (I71)	7	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
...	All other causes (Residual)	392	19.8	3.2
White, non-Hispanic, male, 15-24 years				
...	All causes	14,553	100.0	107.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,290	57.0	61.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,496	17.2	18.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	638	4.4	4.7
4	Malignant neoplasms (C00-C97)	608	4.2	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	2.8	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	158	1.1	1.2
7	Influenza and pneumonia (J10-J18)	62	0.4	0.5
8	Cerebrovascular diseases (I60-I69)	60	0.4	0.4
9	Diabetes mellitus (E10-E14)	47	0.3	0.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.3	0.3
11	Septicemia (A40-A41)	33	0.2	0.2
12	Chronic lower respiratory diseases (J40-J47)	30	0.2	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.2	0.2
14	Legal intervention (Y35,Y89.0)	28	0.2	0.2
15	Aortic aneurysm and dissection (I71)	22	0.2	0.2
...	All other causes (Residual)	1,605	11.0	11.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-34 years				
...	All causes	17,115	100.0	141.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,578	44.3	62.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,960	17.3	24.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,250	7.3	10.3
4	Malignant neoplasms (C00-C97)	1,126	6.6	9.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	664	3.9	5.5
6	Diabetes mellitus (E10-E14)	189	1.1	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	173	1.0	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	152	0.9	1.3
9	Cerebrovascular diseases (I60-I69)	134	0.8	1.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	110	0.6	0.9
11	Influenza and pneumonia (J10-J18)	96	0.6	0.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	0.5	0.6
13	Septicemia (A40-A41)	77	0.4	0.6
14	Chronic lower respiratory diseases (J40-J47)	56	0.3	0.5
15	Legal intervention (Y35,Y89.0)	51	0.3	0.4
...	All other causes (Residual)	2,421	14.1	20.0
White, non-Hispanic, male, 35-44 years				
...	All causes	33,609	100.0	233.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,714	25.9	60.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,738	17.1	39.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,179	12.4	29.0
4	Malignant neoplasms (C00-C97)	4,165	12.4	28.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,072	3.2	7.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	981	2.9	6.8
7	Diabetes mellitus (E10-E14)	777	2.3	5.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	700	2.1	4.9
9	Cerebrovascular diseases (I60-I69)	558	1.7	3.9
10	Influenza and pneumonia (J10-J18)	297	0.9	2.1
11	Septicemia (A40-A41)	262	0.8	1.8
12	Chronic lower respiratory diseases (J40-J47)	254	0.8	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	210	0.6	1.5
14	Viral hepatitis (B15-B19)	203	0.6	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	195	0.6	1.4
...	All other causes (Residual)	5,304	15.8	36.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 45-54 years				
...	All causes	80,427	100.0	515.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,439	24.2	124.5
2	Malignant neoplasms (C00-C97)	18,532	23.0	118.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,124	12.6	64.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,949	6.2	31.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,924	4.9	25.1
6	Diabetes mellitus (E10-E14)	2,211	2.7	14.2
7	Cerebrovascular diseases (I60-I69)	1,893	2.4	12.1
8	Chronic lower respiratory diseases (J40-J47)	1,538	1.9	9.8
9	Viral hepatitis (B15-B19)	1,450	1.8	9.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,155	1.4	7.4
11	Influenza and pneumonia (J10-J18)	771	1.0	4.9
12	Septicemia (A40-A41)	746	0.9	4.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	703	0.9	4.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	679	0.8	4.3
15	Aortic aneurysm and dissection (I71)	409	0.5	2.6
...	All other causes (Residual)	11,904	14.8	76.2
White, non-Hispanic, male, 55-64 years				
...	All causes	126,779	100.0	1,064.0
1	Malignant neoplasms (C00-C97)	43,300	34.2	363.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,901	26.7	284.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,847	4.6	49.1
4	Chronic lower respiratory diseases (J40-J47)	5,407	4.3	45.4
5	Diabetes mellitus (E10-E14)	4,486	3.5	37.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,799	3.0	31.9
7	Cerebrovascular diseases (I60-I69)	3,612	2.8	30.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,129	2.5	26.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,465	1.2	12.3
10	Septicemia (A40-A41)	1,412	1.1	11.9
11	Influenza and pneumonia (J10-J18)	1,373	1.1	11.5
12	Viral hepatitis (B15-B19)	947	0.7	7.9
13	Aortic aneurysm and dissection (I71)	796	0.6	6.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	730	0.6	6.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	579	0.5	4.9
...	All other causes (Residual)	15,996	12.6	134.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65-74 years				
...	All causes	173,056	100.0	2,490.3
1	Malignant neoplasms (C00-C97)	61,359	35.5	883.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,772	25.9	644.3
3	Chronic lower respiratory diseases (J40-J47)	12,546	7.2	180.5
4	Cerebrovascular diseases (I60-I69)	6,704	3.9	96.5
5	Diabetes mellitus (E10-E14)	5,830	3.4	83.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,164	2.4	59.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,622	1.5	37.7
8	Influenza and pneumonia (J10-J18)	2,576	1.5	37.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,359	1.4	33.9
10	Septicemia (A40-A41)	2,191	1.3	31.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,803	1.0	25.9
12	Alzheimer's disease (G30)	1,462	0.8	21.0
13	Aortic aneurysm and dissection (I71)	1,431	0.8	20.6
14	Parkinson's disease (G20-G21)	1,324	0.8	19.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,147	0.7	16.5
...	All other causes (Residual)	20,766	12.0	298.8
White, non-Hispanic, male, 75-84 years				
...	All causes	278,690	100.0	6,278.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,455	28.2	1,767.4
2	Malignant neoplasms (C00-C97)	73,816	26.5	1,662.9
3	Chronic lower respiratory diseases (J40-J47)	20,850	7.5	469.7
4	Cerebrovascular diseases (I60-I69)	15,113	5.4	340.5
5	Diabetes mellitus (E10-E14)	7,971	2.9	179.6
6	Alzheimer's disease (G30)	7,363	2.6	165.9
7	Influenza and pneumonia (J10-J18)	6,966	2.5	156.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,269	2.2	141.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,791	2.1	130.5
10	Parkinson's disease (G20-G21)	5,060	1.8	114.0
11	Septicemia (A40-A41)	3,929	1.4	88.5
12	Pneumonitis due to solids and liquids (J69)	2,626	0.9	59.2
13	Aortic aneurysm and dissection (I71)	2,308	0.8	52.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,222	0.8	50.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,039	0.7	45.9
...	All other causes (Residual)	37,912	13.6	854.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85 years and over				
...	All causes	213,151	100.0	14,841.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,282	33.9	5,032.8
2	Malignant neoplasms (C00-C97)	33,036	15.5	2,300.2
3	Cerebrovascular diseases (I60-I69)	13,790	6.5	960.2
4	Chronic lower respiratory diseases (J40-J47)	11,550	5.4	804.2
5	Alzheimer's disease (G30)	9,715	4.6	676.4
6	Influenza and pneumonia (J10-J18)	8,600	4.0	598.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,467	2.6	380.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,203	2.4	362.3
9	Diabetes mellitus (E10-E14)	4,406	2.1	306.8
10	Parkinson's disease (G20-G21)	3,625	1.7	252.4
11	Pneumonitis due to solids and liquids (J69)	3,445	1.6	239.9
12	Septicemia (A40-A41)	2,776	1.3	193.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,183	1.0	152.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,379	0.6	96.0
15	Aortic aneurysm and dissection (I71)	1,333	0.6	92.8
...	All other causes (Residual)	34,361	16.1	2,392.5
White, non-Hispanic, female, all ages				
...	All causes	996,651	100.0	974.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	260,424	26.1	254.7
2	Malignant neoplasms (C00-C97)	219,585	22.0	214.7
3	Cerebrovascular diseases (I60-I69)	66,945	6.7	65.5
4	Chronic lower respiratory diseases (J40-J47)	59,289	5.9	58.0
5	Alzheimer's disease (G30)	45,816	4.6	44.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,072	3.4	33.3
7	Influenza and pneumonia (J10-J18)	25,556	2.6	25.0
8	Diabetes mellitus (E10-E14)	25,019	2.5	24.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,793	1.7	16.4
10	Septicemia (A40-A41)	14,030	1.4	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,583	1.1	10.3
12	Parkinson's disease (G20-G21)	7,529	0.8	7.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,393	0.7	7.2
14	Pneumonitis due to solids and liquids (J69)	6,869	0.7	6.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,000	0.6	5.9
...	All other causes (Residual)	190,748	19.1	186.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	1,032	100.0	23.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	356	34.5	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	12.7	2.9
3	Malignant neoplasms (C00-C97)	93	9.0	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	5.4	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.7	0.9
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	2.7	0.6
7	Influenza and pneumonia (J10-J18)	25	2.4	0.6
8	Septicemia (A40-A41)	20	1.9	0.4
8	Cerebrovascular diseases (I60-I69)	20	1.9	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	8	0.8	*
11	Meningitis (G00,G03)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
13	Anemias (D50-D64)	6	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
15	Pneumonitis due to solids and liquids (J69)	4	0.4	*
...	All other causes (Residual)	228	22.1	5.1
White, non-Hispanic, female, 5-14 years				
...	All causes	1,370	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	501	36.6	4.3
2	Malignant neoplasms (C00-C97)	231	16.9	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	6.2	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	4.8	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	4.0	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.1	0.4
7	Septicemia (A40-A41)	19	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.2	*
8	Cerebrovascular diseases (I60-I69)	17	1.2	*
10	Influenza and pneumonia (J10-J18)	14	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	8	0.6	*
12	Anemias (D50-D64)	6	0.4	*
13	Diabetes mellitus (E10-E14)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
...	All other causes (Residual)	294	21.5	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,502	100.0	42.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,913	52.9	22.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	448	8.1	3.5
3	Malignant neoplasms (C00-C97)	372	6.8	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	262	4.8	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	198	3.6	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	2.1	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	78	1.4	0.6
8	Influenza and pneumonia (J10-J18)	52	0.9	0.4
9	Cerebrovascular diseases (I60-I69)	45	0.8	0.4
10	Diabetes mellitus (E10-E14)	42	0.8	0.3
11	Septicemia (A40-A41)	36	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	24	0.4	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.4	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.2	*
...	All other causes (Residual)	869	15.8	6.8
White, non-Hispanic, female, 25-34 years				
...	All causes	7,491	100.0	62.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,507	33.5	20.9
2	Malignant neoplasms (C00-C97)	1,100	14.7	9.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	729	9.7	6.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	524	7.0	4.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	267	3.6	2.2
6	Diabetes mellitus (E10-E14)	145	1.9	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	1.6	1.0
8	Cerebrovascular diseases (I60-I69)	117	1.6	1.0
9	Pregnancy, childbirth and the puerperium (O00-099)	113	1.5	0.9
10	Influenza and pneumonia (J10-J18)	93	1.2	0.8
11	Septicemia (A40-A41)	68	0.9	0.6
12	Chronic lower respiratory diseases (J40-J47)	59	0.8	0.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	57	0.8	0.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	56	0.7	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.6	0.4
...	All other causes (Residual)	1,492	19.9	12.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 35-44 years				
...	All causes	19,589	100.0	136.3
1	Malignant neoplasms (C00-C97)	5,162	26.4	35.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,845	19.6	26.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,222	11.3	15.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,329	6.8	9.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	621	3.2	4.3
6	Cerebrovascular diseases (I60-I69)	528	2.7	3.7
7	Diabetes mellitus (E10-E14)	480	2.5	3.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	1.7	2.4
9	Chronic lower respiratory diseases (J40-J47)	285	1.5	2.0
10	Influenza and pneumonia (J10-J18)	252	1.3	1.8
11	Septicemia (A40-A41)	228	1.2	1.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	212	1.1	1.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	0.7	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	137	0.7	1.0
15	Viral hepatitis (B15-B19)	115	0.6	0.8
...	All other causes (Residual)	3,689	18.8	25.7
White, non-Hispanic, female, 45-54 years				
...	All causes	47,542	100.0	299.8
1	Malignant neoplasms (C00-C97)	17,926	37.7	113.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,699	14.1	42.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,495	9.5	28.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,665	3.5	10.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,617	3.4	10.2
6	Cerebrovascular diseases (I60-I69)	1,602	3.4	10.1
7	Chronic lower respiratory diseases (J40-J47)	1,575	3.3	9.9
8	Diabetes mellitus (E10-E14)	1,254	2.6	7.9
9	Septicemia (A40-A41)	591	1.2	3.7
10	Influenza and pneumonia (J10-J18)	556	1.2	3.5
11	Viral hepatitis (B15-B19)	541	1.1	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	434	0.9	2.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	318	0.7	2.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	257	0.5	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	217	0.5	1.4
...	All other causes (Residual)	7,795	16.4	49.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 55-64 years				
...	All causes	83,539	100.0	668.0
1	Malignant neoplasms (C00-C97)	36,081	43.2	288.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,152	16.9	113.2
3	Chronic lower respiratory diseases (J40-J47)	5,259	6.3	42.1
4	Cerebrovascular diseases (I60-I69)	2,943	3.5	23.5
5	Diabetes mellitus (E10-E14)	2,912	3.5	23.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,806	3.4	22.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,609	1.9	12.9
8	Septicemia (A40-A41)	1,267	1.5	10.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,047	1.3	8.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,038	1.2	8.3
11	Influenza and pneumonia (J10-J18)	940	1.1	7.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	455	0.5	3.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	441	0.5	3.5
14	Viral hepatitis (B15-B19)	358	0.4	2.9
15	Alzheimer's disease (G30)	299	0.4	2.4
...	All other causes (Residual)	11,932	14.3	95.4
White, non-Hispanic, female, 65-74 years				
...	All causes	134,419	100.0	1,677.4
1	Malignant neoplasms (C00-C97)	50,586	37.6	631.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,374	19.6	329.1
3	Chronic lower respiratory diseases (J40-J47)	12,851	9.6	160.4
4	Cerebrovascular diseases (I60-I69)	6,333	4.7	79.0
5	Diabetes mellitus (E10-E14)	4,475	3.3	55.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,581	1.9	32.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,185	1.6	27.3
8	Septicemia (A40-A41)	2,144	1.6	26.8
9	Influenza and pneumonia (J10-J18)	2,053	1.5	25.6
10	Alzheimer's disease (G30)	1,791	1.3	22.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,480	1.1	18.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	967	0.7	12.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	822	0.6	10.3
14	Aortic aneurysm and dissection (I71)	772	0.6	9.6
15	Parkinson's disease (G20-G21)	672	0.5	8.4
...	All other causes (Residual)	18,333	13.6	228.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 75-84 years				
...	All causes	285,423	100.0	4,460.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,674	25.5	1,135.8
2	Malignant neoplasms (C00-C97)	66,830	23.4	1,044.4
3	Chronic lower respiratory diseases (J40-J47)	22,323	7.8	348.9
4	Cerebrovascular diseases (I60-I69)	20,855	7.3	325.9
5	Alzheimer's disease (G30)	12,910	4.5	201.8
6	Diabetes mellitus (E10-E14)	8,176	2.9	127.8
7	Influenza and pneumonia (J10-J18)	6,904	2.4	107.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,674	2.0	88.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,503	1.9	86.0
10	Septicemia (A40-A41)	4,566	1.6	71.4
11	Parkinson's disease (G20-G21)	3,262	1.1	51.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,922	1.0	45.7
13	Pneumonitis due to solids and liquids (J69)	1,926	0.7	30.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,916	0.7	29.9
15	Aortic aneurysm and dissection (I71)	1,779	0.6	27.8
...	All other causes (Residual)	47,203	16.5	737.7
White, non-Hispanic, female, 85 years and over				
...	All causes	405,102	100.0	13,150.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	137,413	33.9	4,460.8
2	Malignant neoplasms (C00-C97)	41,177	10.2	1,336.7
3	Cerebrovascular diseases (I60-I69)	34,460	8.5	1,118.7
4	Alzheimer's disease (G30)	30,779	7.6	999.2
5	Chronic lower respiratory diseases (J40-J47)	16,896	4.2	548.5
6	Influenza and pneumonia (J10-J18)	14,632	3.6	475.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,160	2.0	264.9
8	Diabetes mellitus (E10-E14)	7,527	1.9	244.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,396	1.8	240.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,982	1.5	194.2
11	Septicemia (A40-A41)	5,048	1.2	163.9
12	Pneumonitis due to solids and liquids (J69)	3,966	1.0	128.7
13	Parkinson's disease (G20-G21)	3,465	0.9	112.5
14	Atherosclerosis (I70)	3,005	0.7	97.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,306	0.6	74.9
...	All other causes (Residual)	82,890	20.5	2,690.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	286,581	100.0	759.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,461	24.9	189.3
2	Malignant neoplasms (C00-C97)	62,475	21.8	165.5
3	Cerebrovascular diseases (I60-I69)	16,882	5.9	44.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,684	4.8	36.2
5	Diabetes mellitus (E10-E14)	12,671	4.4	33.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,902	3.1	23.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,334	2.9	22.1
8	Chronic lower respiratory diseases (J40-J47)	7,657	2.7	20.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,767	2.4	17.9
10	Septicemia (A40-A41)	6,045	2.1	16.0
11	Influenza and pneumonia (J10-J18)	5,242	1.8	13.9
12	Certain conditions originating in the perinatal period (P00-P96)	4,906	1.7	13.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,851	1.7	12.8
14	Alzheimer's disease (G30)	4,422	1.5	11.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,310	0.8	6.1
...	All other causes (Residual)	49,972	17.4	132.4
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,108	100.0	44.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	355	32.0	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	13.5	6.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	9.6	4.2
4	Malignant neoplasms (C00-C97)	63	5.7	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	4.0	1.8
6	Influenza and pneumonia (J10-J18)	34	3.1	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	22	2.0	0.9
8	Septicemia (A40-A41)	19	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	15	1.4	*
10	Anemias (D50-D64)	12	1.1	*
11	Cerebrovascular diseases (I60-I69)	11	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	256	23.1	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,361	100.0	21.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	460	33.8	7.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	10.8	2.4
3	Malignant neoplasms (C00-C97)	140	10.3	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	4.9	1.1
5	Chronic lower respiratory diseases (J40-J47)	66	4.8	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	4.6	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	2.6	0.6
8	Septicemia (A40-A41)	20	1.5	0.3
8	Cerebrovascular diseases (I60-I69)	20	1.5	0.3
10	Anemias (D50-D64)	19	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.8	*
13	Diabetes mellitus (E10-E14)	9	0.7	*
13	Influenza and pneumonia (J10-J18)	9	0.7	*
15	Meningitis (G00,G03)	8	0.6	*
...	All other causes (Residual)	272	20.0	4.4
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	7,347	100.0	115.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,253	44.3	51.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,703	23.2	26.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	407	5.5	6.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	302	4.1	4.8
5	Malignant neoplasms (C00-C97)	259	3.5	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	143	1.9	2.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	1.2	1.4
8	Chronic lower respiratory diseases (J40-J47)	74	1.0	1.2
9	Anemias (D50-D64)	64	0.9	1.0
10	Pregnancy, childbirth and the puerperium (O00-099)	57	0.8	0.9
11	Diabetes mellitus (E10-E14)	54	0.7	0.8
12	Cerebrovascular diseases (I60-I69)	44	0.6	0.7
13	Septicemia (A40-A41)	41	0.6	0.6
13	Influenza and pneumonia (J10-J18)	41	0.6	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.4	0.5
...	All other causes (Residual)	788	10.7	12.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-34 years				
...	All causes	9,920	100.0	185.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,635	26.6	49.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,903	19.2	35.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,020	10.3	19.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	765	7.7	14.3
5	Malignant neoplasms (C00-C97)	636	6.4	11.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	466	4.7	8.7
7	Diabetes mellitus (E10-E14)	237	2.4	4.4
8	Cerebrovascular diseases (I60-I69)	149	1.5	2.8
9	Pregnancy, childbirth and the puerperium (O00-099)	115	1.2	2.2
10	Anemias (D50-D64)	111	1.1	2.1
11	Chronic lower respiratory diseases (J40-J47)	106	1.1	2.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	106	1.1	2.0
13	Septicemia (A40-A41)	88	0.9	1.6
14	Influenza and pneumonia (J10-J18)	77	0.8	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	0.7	1.3
...	All other causes (Residual)	1,434	14.5	26.8
Black, non-Hispanic, both sexes, 35-44 years				
...	All causes	17,700	100.0	325.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,037	17.2	55.8
2	Malignant neoplasms (C00-C97)	2,495	14.1	45.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,307	13.0	42.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,159	12.2	39.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,331	7.5	24.5
6	Cerebrovascular diseases (I60-I69)	691	3.9	12.7
7	Diabetes mellitus (E10-E14)	578	3.3	10.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	406	2.3	7.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	333	1.9	6.1
10	Septicemia (A40-A41)	270	1.5	5.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	251	1.4	4.6
12	Chronic lower respiratory diseases (J40-J47)	224	1.3	4.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	208	1.2	3.8
14	Influenza and pneumonia (J10-J18)	183	1.0	3.4
15	Anemias (D50-D64)	143	0.8	2.6
...	All other causes (Residual)	3,084	17.4	56.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-54 years				
...	All causes	36,990	100.0	738.5
1	Malignant neoplasms (C00-C97)	8,713	23.6	174.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,520	23.0	170.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,817	7.6	56.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,417	6.5	48.3
5	Cerebrovascular diseases (I60-I69)	1,898	5.1	37.9
6	Diabetes mellitus (E10-E14)	1,402	3.8	28.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	836	2.3	16.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	812	2.2	16.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	792	2.1	15.8
10	Septicemia (A40-A41)	673	1.8	13.4
11	Chronic lower respiratory diseases (J40-J47)	614	1.7	12.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	565	1.5	11.3
13	Influenza and pneumonia (J10-J18)	449	1.2	9.0
14	Viral hepatitis (B15-B19)	391	1.1	7.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	303	0.8	6.0
...	All other causes (Residual)	5,788	15.6	115.6
Black, non-Hispanic, both sexes, 55-64 years				
...	All causes	45,961	100.0	1,472.4
1	Malignant neoplasms (C00-C97)	14,048	30.6	450.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,942	26.0	382.6
3	Cerebrovascular diseases (I60-I69)	2,592	5.6	83.0
4	Diabetes mellitus (E10-E14)	2,475	5.4	79.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,544	3.4	49.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,334	2.9	42.7
7	Chronic lower respiratory diseases (J40-J47)	1,190	2.6	38.1
8	Septicemia (A40-A41)	986	2.1	31.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	962	2.1	30.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	730	1.6	23.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	700	1.5	22.4
12	Influenza and pneumonia (J10-J18)	507	1.1	16.2
13	Viral hepatitis (B15-B19)	391	0.9	12.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	304	0.7	9.7
15	Aortic aneurysm and dissection (I71)	210	0.5	6.7
...	All other causes (Residual)	6,046	13.2	193.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65-74 years				
...	All causes	50,574	100.0	2,887.7
1	Malignant neoplasms (C00-C97)	15,597	30.8	890.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,768	27.2	786.1
3	Cerebrovascular diseases (I60-I69)	3,176	6.3	181.3
4	Diabetes mellitus (E10-E14)	2,977	5.9	170.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,831	3.6	104.5
6	Chronic lower respiratory diseases (J40-J47)	1,802	3.6	102.9
7	Septicemia (A40-A41)	1,230	2.4	70.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	878	1.7	50.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	853	1.7	48.7
10	Influenza and pneumonia (J10-J18)	796	1.6	45.5
11	Alzheimer's disease (G30)	334	0.7	19.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	333	0.7	19.0
13	Aortic aneurysm and dissection (I71)	255	0.5	14.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	250	0.5	14.3
15	Pneumonitis due to solids and liquids (J69)	247	0.5	14.1
...	All other causes (Residual)	6,247	12.4	356.7
Black, non-Hispanic, both sexes, 75-84 years				
...	All causes	59,416	100.0	5,980.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,184	28.9	1,729.7
2	Malignant neoplasms (C00-C97)	14,110	23.7	1,420.3
3	Cerebrovascular diseases (I60-I69)	4,398	7.4	442.7
4	Diabetes mellitus (E10-E14)	3,143	5.3	316.4
5	Chronic lower respiratory diseases (J40-J47)	2,284	3.8	229.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,124	3.6	213.8
7	Septicemia (A40-A41)	1,523	2.6	153.3
8	Alzheimer's disease (G30)	1,483	2.5	149.3
9	Influenza and pneumonia (J10-J18)	1,403	2.4	141.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,181	2.0	118.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	815	1.4	82.0
12	Pneumonitis due to solids and liquids (J69)	426	0.7	42.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	320	0.5	32.2
14	Parkinson's disease (G20-G21)	304	0.5	30.6
15	Aortic aneurysm and dissection (I71)	281	0.5	28.3
...	All other causes (Residual)	8,437	14.2	849.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85 years and over				
...	All causes	47,678	100.0	12,655.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,461	32.4	4,103.9
2	Malignant neoplasms (C00-C97)	6,406	13.4	1,700.4
3	Cerebrovascular diseases (I60-I69)	3,854	8.1	1,023.0
4	Alzheimer's disease (G30)	2,541	5.3	674.5
5	Diabetes mellitus (E10-E14)	1,793	3.8	475.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,693	3.6	449.4
7	Influenza and pneumonia (J10-J18)	1,656	3.5	439.6
8	Chronic lower respiratory diseases (J40-J47)	1,271	2.7	337.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,216	2.6	322.8
10	Septicemia (A40-A41)	1,096	2.3	290.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	577	1.2	153.2
12	Pneumonitis due to solids and liquids (J69)	493	1.0	130.9
13	Atherosclerosis (I70)	274	0.6	72.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	239	0.5	63.4
15	Parkinson's disease (G20-G21)	195	0.4	51.8
...	All other causes (Residual)	8,913	18.7	2,365.8
Black, non-Hispanic, male, all ages				
...	All causes	146,729	100.0	815.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,787	24.4	198.8
2	Malignant neoplasms (C00-C97)	32,253	22.0	179.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,443	6.4	52.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,566	5.2	42.0
5	Cerebrovascular diseases (I60-I69)	7,349	5.0	40.8
6	Diabetes mellitus (E10-E14)	5,701	3.9	31.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,384	3.0	24.4
8	Chronic lower respiratory diseases (J40-J47)	4,091	2.8	22.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,782	2.6	21.0
10	Certain conditions originating in the perinatal period (P00-P96)	2,706	1.8	15.0
11	Septicemia (A40-A41)	2,688	1.8	14.9
12	Influenza and pneumonia (J10-J18)	2,449	1.7	13.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,100	1.4	11.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,630	1.1	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,485	1.0	8.3
...	All other causes (Residual)	23,315	15.9	129.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	612	100.0	48.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	33.8	16.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	13.2	6.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	6.7	3.2
4	Malignant neoplasms (C00-C97)	34	5.6	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.4	2.1
6	Influenza and pneumonia (J10-J18)	16	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	12	2.0	*
8	Septicemia (A40-A41)	11	1.8	*
9	Anemias (D50-D64)	9	1.5	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.5	*
11	Cerebrovascular diseases (I60-I69)	7	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
15	Meningococcal infection (A39)	1	0.2	*
15	Syphilis (A50-A53)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	143	23.4	11.3
Black, non-Hispanic, male, 5-14 years				
...	All causes	814	100.0	25.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	35.4	9.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	12.8	3.3
3	Malignant neoplasms (C00-C97)	80	9.8	2.5
4	Chronic lower respiratory diseases (J40-J47)	41	5.0	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	4.2	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	3.6	0.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	3.4	0.9
8	Septicemia (A40-A41)	13	1.6	*
9	Anemias (D50-D64)	9	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.9	*
10	Cerebrovascular diseases (I60-I69)	7	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
13	Meningitis (G00,G03)	3	0.4	*
13	Influenza and pneumonia (J10-J18)	3	0.4	*
13	Pneumonitis due to solids and liquids (J69)	3	0.4	*
...	All other causes (Residual)	159	19.5	5.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,691	100.0	176.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,946	51.8	91.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,293	22.7	40.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	351	6.2	10.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	3.2	5.6
5	Malignant neoplasms (C00-C97)	151	2.7	4.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	83	1.5	2.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	0.9	1.6
8	Chronic lower respiratory diseases (J40-J47)	46	0.8	1.4
9	Anemias (D50-D64)	37	0.7	1.1
10	Diabetes mellitus (E10-E14)	31	0.5	1.0
11	Legal intervention (Y35,Y89.0)	24	0.4	0.7
12	Influenza and pneumonia (J10-J18)	23	0.4	0.7
13	Septicemia (A40-A41)	21	0.4	0.7
14	Cerebrovascular diseases (I60-I69)	20	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
...	All other causes (Residual)	411	7.2	12.8
Black, non-Hispanic, male, 25-34 years				
...	All causes	6,828	100.0	266.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,315	33.9	90.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,480	21.7	57.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	653	9.6	25.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	416	6.1	16.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	414	6.1	16.1
6	Malignant neoplasms (C00-C97)	271	4.0	10.6
7	Diabetes mellitus (E10-E14)	135	2.0	5.3
8	Cerebrovascular diseases (I60-I69)	75	1.1	2.9
9	Chronic lower respiratory diseases (J40-J47)	61	0.9	2.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.8	2.2
11	Anemias (D50-D64)	55	0.8	2.1
12	Septicemia (A40-A41)	44	0.6	1.7
12	Legal intervention (Y35,Y89.0)	44	0.6	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.6	1.7
15	Influenza and pneumonia (J10-J18)	37	0.5	1.4
...	All other causes (Residual)	729	10.7	28.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-44 years				
...	All causes	10,363	100.0	405.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,864	18.0	72.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,637	15.8	64.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,336	12.9	52.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,052	10.2	41.2
5	Malignant neoplasms (C00-C97)	973	9.4	38.1
6	Diabetes mellitus (E10-E14)	343	3.3	13.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	342	3.3	13.4
8	Cerebrovascular diseases (I60-I69)	339	3.3	13.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	181	1.7	7.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	151	1.5	5.9
11	Septicemia (A40-A41)	132	1.3	5.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	126	1.2	4.9
13	Chronic lower respiratory diseases (J40-J47)	102	1.0	4.0
14	Influenza and pneumonia (J10-J18)	95	0.9	3.7
15	Anemias (D50-D64)	64	0.6	2.5
...	All other causes (Residual)	1,626	15.7	63.6
Black, non-Hispanic, male, 45-54 years				
...	All causes	21,829	100.0	944.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,434	24.9	235.0
2	Malignant neoplasms (C00-C97)	4,323	19.8	187.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,037	9.3	88.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,625	7.4	70.3
5	Cerebrovascular diseases (I60-I69)	1,032	4.7	44.6
6	Diabetes mellitus (E10-E14)	792	3.6	34.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	640	2.9	27.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	530	2.4	22.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	477	2.2	20.6
10	Septicemia (A40-A41)	350	1.6	15.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	328	1.5	14.2
12	Chronic lower respiratory diseases (J40-J47)	314	1.4	13.6
13	Influenza and pneumonia (J10-J18)	270	1.2	11.7
14	Viral hepatitis (B15-B19)	256	1.2	11.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	244	1.1	10.6
...	All other causes (Residual)	3,177	14.6	137.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-64 years				
...	All causes	26,929	100.0	1,935.2
1	Malignant neoplasms (C00-C97)	7,753	28.8	557.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,485	27.8	537.9
3	Cerebrovascular diseases (I60-I69)	1,511	5.6	108.6
4	Diabetes mellitus (E10-E14)	1,289	4.8	92.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,110	4.1	79.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	710	2.6	51.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	700	2.6	50.3
8	Chronic lower respiratory diseases (J40-J47)	679	2.5	48.8
9	Septicemia (A40-A41)	527	2.0	37.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	472	1.8	33.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	399	1.5	28.7
12	Influenza and pneumonia (J10-J18)	304	1.1	21.8
13	Viral hepatitis (B15-B19)	279	1.0	20.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	240	0.9	17.2
15	Aortic aneurysm and dissection (I71)	136	0.5	9.8
...	All other causes (Residual)	3,335	12.4	239.7
Black, non-Hispanic, male, 65-74 years				
...	All causes	27,265	100.0	3,736.8
1	Malignant neoplasms (C00-C97)	8,602	31.5	1,178.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,754	28.4	1,062.7
3	Cerebrovascular diseases (I60-I69)	1,625	6.0	222.7
4	Diabetes mellitus (E10-E14)	1,396	5.1	191.3
5	Chronic lower respiratory diseases (J40-J47)	1,019	3.7	139.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	857	3.1	117.5
7	Septicemia (A40-A41)	563	2.1	77.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	539	2.0	73.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	442	1.6	60.6
10	Influenza and pneumonia (J10-J18)	434	1.6	59.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	214	0.8	29.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	176	0.6	24.1
13	Pneumonitis due to solids and liquids (J69)	155	0.6	21.2
14	Alzheimer's disease (G30)	141	0.5	19.3
14	Aortic aneurysm and dissection (I71)	141	0.5	19.3
...	All other causes (Residual)	3,207	11.8	439.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-84 years				
...	All causes	27,045	100.0	7,520.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,783	28.8	2,164.2
2	Malignant neoplasms (C00-C97)	7,249	26.8	2,015.7
3	Cerebrovascular diseases (I60-I69)	1,724	6.4	479.4
4	Chronic lower respiratory diseases (J40-J47)	1,241	4.6	345.1
5	Diabetes mellitus (E10-E14)	1,202	4.4	334.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	921	3.4	256.1
7	Influenza and pneumonia (J10-J18)	655	2.4	182.1
8	Septicemia (A40-A41)	646	2.4	179.6
9	Alzheimer's disease (G30)	485	1.8	134.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	464	1.7	129.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	422	1.6	117.3
12	Pneumonitis due to solids and liquids (J69)	246	0.9	68.4
13	Parkinson's disease (G20-G21)	185	0.7	51.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	152	0.6	42.3
15	Aortic aneurysm and dissection (I71)	109	0.4	30.3
...	All other causes (Residual)	3,561	13.2	990.2
Black, non-Hispanic, male, 85 years and over				
...	All causes	14,628	100.0	13,403.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,507	30.8	4,129.6
2	Malignant neoplasms (C00-C97)	2,813	19.2	2,577.4
3	Cerebrovascular diseases (I60-I69)	976	6.7	894.3
4	Chronic lower respiratory diseases (J40-J47)	570	3.9	522.3
5	Influenza and pneumonia (J10-J18)	559	3.8	512.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	542	3.7	496.6
7	Alzheimer's disease (G30)	522	3.6	478.3
8	Diabetes mellitus (E10-E14)	511	3.5	468.2
9	Septicemia (A40-A41)	328	2.2	300.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	300	2.1	274.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	1.5	202.5
12	Pneumonitis due to solids and liquids (J69)	194	1.3	177.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	90	0.6	82.5
14	Atherosclerosis (I70)	83	0.6	76.0
15	Parkinson's disease (G20-G21)	78	0.5	71.5
...	All other causes (Residual)	2,334	16.0	2,138.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	139,852	100.0	708.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,674	25.5	180.6
2	Malignant neoplasms (C00-C97)	30,222	21.6	153.0
3	Cerebrovascular diseases (I60-I69)	9,533	6.8	48.3
4	Diabetes mellitus (E10-E14)	6,970	5.0	35.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,552	3.3	23.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,241	3.0	21.5
7	Chronic lower respiratory diseases (J40-J47)	3,566	2.5	18.1
8	Septicemia (A40-A41)	3,357	2.4	17.0
9	Alzheimer's disease (G30)	3,246	2.3	16.4
10	Influenza and pneumonia (J10-J18)	2,793	2.0	14.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,751	2.0	13.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,383	1.7	12.1
13	Certain conditions originating in the perinatal period (P00-P96)	2,200	1.6	11.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,336	1.0	6.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	828	0.6	4.2
...	All other causes (Residual)	26,200	18.7	132.6
Black, non-Hispanic, female, 1-4 years				
...	All causes	496	100.0	40.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	29.8	12.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	13.9	5.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	13.1	5.3
4	Malignant neoplasms (C00-C97)	29	5.8	2.4
5	Influenza and pneumonia (J10-J18)	18	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.6	*
8	Septicemia (A40-A41)	8	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
9	Cerebrovascular diseases (I60-I69)	4	0.8	*
11	Anemias (D50-D64)	3	0.6	*
11	Meningitis (G00,G03)	3	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (Residual)	108	21.8	8.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	547	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	31.4	5.6
2	Malignant neoplasms (C00-C97)	60	11.0	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	7.9	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.9	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.3	0.9
6	Chronic lower respiratory diseases (J40-J47)	25	4.6	0.8
7	Cerebrovascular diseases (I60-I69)	13	2.4	*
8	Anemias (D50-D64)	10	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.5	*
10	Septicemia (A40-A41)	7	1.3	*
10	Diabetes mellitus (E10-E14)	7	1.3	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	1.3	*
13	Influenza and pneumonia (J10-J18)	6	1.1	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.9	*
14	Meningitis (G00,G03)	5	0.9	*
...	All other causes (Residual)	112	20.5	3.7
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,656	100.0	52.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	410	24.8	13.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	307	18.5	9.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	7.4	3.9
4	Malignant neoplasms (C00-C97)	108	6.5	3.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	60	3.6	1.9
6	Pregnancy, childbirth and the puerperium (O00-099)	57	3.4	1.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	3.4	1.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	2.1	1.1
9	Chronic lower respiratory diseases (J40-J47)	28	1.7	0.9
10	Anemias (D50-D64)	27	1.6	0.9
11	Cerebrovascular diseases (I60-I69)	24	1.4	0.8
12	Diabetes mellitus (E10-E14)	23	1.4	0.7
13	Septicemia (A40-A41)	20	1.2	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.1	*
15	Influenza and pneumonia (J10-J18)	18	1.1	*
...	All other causes (Residual)	342	20.7	10.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-34 years				
...	All causes	3,092	100.0	111.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	423	13.7	15.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	367	11.9	13.2
3	Malignant neoplasms (C00-C97)	365	11.8	13.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	349	11.3	12.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	320	10.3	11.5
6	Pregnancy, childbirth and the puerperium (O00-099)	115	3.7	4.1
7	Diabetes mellitus (E10-E14)	102	3.3	3.7
8	Cerebrovascular diseases (I60-I69)	74	2.4	2.7
9	Anemias (D50-D64)	56	1.8	2.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	1.7	1.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.6	1.8
12	Chronic lower respiratory diseases (J40-J47)	45	1.5	1.6
13	Septicemia (A40-A41)	44	1.4	1.6
14	Influenza and pneumonia (J10-J18)	40	1.3	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	0.9	1.0
...	All other causes (Residual)	661	21.4	23.8
Black, non-Hispanic, female, 35-44 years				
...	All causes	7,337	100.0	254.1
1	Malignant neoplasms (C00-C97)	1,522	20.7	52.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,173	16.0	40.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	823	11.2	28.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	670	9.1	23.2
5	Cerebrovascular diseases (I60-I69)	352	4.8	12.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	279	3.8	9.7
7	Diabetes mellitus (E10-E14)	235	3.2	8.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	2.1	5.3
9	Septicemia (A40-A41)	138	1.9	4.8
10	Chronic lower respiratory diseases (J40-J47)	122	1.7	4.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	100	1.4	3.5
12	Influenza and pneumonia (J10-J18)	88	1.2	3.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	1.1	2.8
14	Anemias (D50-D64)	79	1.1	2.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	0.9	2.2
...	All other causes (Residual)	1,458	19.9	50.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 45-54 years				
...	All causes	15,161	100.0	562.2
1	Malignant neoplasms (C00-C97)	4,390	29.0	162.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,086	20.4	114.4
3	Cerebrovascular diseases (I60-I69)	866	5.7	32.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	792	5.2	29.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	780	5.1	28.9
6	Diabetes mellitus (E10-E14)	610	4.0	22.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	359	2.4	13.3
8	Septicemia (A40-A41)	323	2.1	12.0
9	Chronic lower respiratory diseases (J40-J47)	300	2.0	11.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	262	1.7	9.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	237	1.6	8.8
12	Influenza and pneumonia (J10-J18)	179	1.2	6.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	172	1.1	6.4
14	Viral hepatitis (B15-B19)	135	0.9	5.0
15	Anemias (D50-D64)	76	0.5	2.8
...	All other causes (Residual)	2,594	17.1	96.2
Black, non-Hispanic, female, 55-64 years				
...	All causes	19,032	100.0	1,100.2
1	Malignant neoplasms (C00-C97)	6,295	33.1	363.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,457	23.4	257.7
3	Diabetes mellitus (E10-E14)	1,186	6.2	68.6
4	Cerebrovascular diseases (I60-I69)	1,081	5.7	62.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	624	3.3	36.1
6	Chronic lower respiratory diseases (J40-J47)	511	2.7	29.5
7	Septicemia (A40-A41)	459	2.4	26.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	434	2.3	25.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	331	1.7	19.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	262	1.4	15.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	228	1.2	13.2
12	Influenza and pneumonia (J10-J18)	203	1.1	11.7
13	Viral hepatitis (B15-B19)	112	0.6	6.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	84	0.4	4.9
15	Aortic aneurysm and dissection (I71)	74	0.4	4.3
...	All other causes (Residual)	2,691	14.1	155.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 65-74 years				
...	All causes	23,309	100.0	2,281.4
1	Malignant neoplasms (C00-C97)	6,995	30.0	684.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,014	25.8	588.6
3	Diabetes mellitus (E10-E14)	1,581	6.8	154.7
4	Cerebrovascular diseases (I60-I69)	1,551	6.7	151.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	974	4.2	95.3
6	Chronic lower respiratory diseases (J40-J47)	783	3.4	76.6
7	Septicemia (A40-A41)	667	2.9	65.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	436	1.9	42.7
9	Influenza and pneumonia (J10-J18)	362	1.6	35.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	1.3	30.7
11	Alzheimer's disease (G30)	193	0.8	18.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	119	0.5	11.6
13	Aortic aneurysm and dissection (I71)	114	0.5	11.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	98	0.4	9.6
15	Pneumonitis due to solids and liquids (J69)	92	0.4	9.0
...	All other causes (Residual)	3,016	12.9	295.2
Black, non-Hispanic, female, 75-84 years				
...	All causes	32,371	100.0	5,107.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,401	29.0	1,483.2
2	Malignant neoplasms (C00-C97)	6,861	21.2	1,082.5
3	Cerebrovascular diseases (I60-I69)	2,674	8.3	421.9
4	Diabetes mellitus (E10-E14)	1,941	6.0	306.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,203	3.7	189.8
6	Chronic lower respiratory diseases (J40-J47)	1,043	3.2	164.6
7	Alzheimer's disease (G30)	998	3.1	157.5
8	Septicemia (A40-A41)	877	2.7	138.4
9	Influenza and pneumonia (J10-J18)	748	2.3	118.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	717	2.2	113.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	393	1.2	62.0
12	Pneumonitis due to solids and liquids (J69)	180	0.6	28.4
13	Aortic aneurysm and dissection (I71)	172	0.5	27.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	168	0.5	26.5
15	Parkinson's disease (G20-G21)	119	0.4	18.8
...	All other causes (Residual)	4,876	15.1	769.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank <sup>1</sup>	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 85 years and over				
...	All causes	33,050	100.0	12,350.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,954	33.1	4,093.4
2	Malignant neoplasms (C00-C97)	3,593	10.9	1,342.7
3	Cerebrovascular diseases (I60-I69)	2,878	8.7	1,075.5
4	Alzheimer's disease (G30)	2,019	6.1	754.5
5	Diabetes mellitus (E10-E14)	1,282	3.9	479.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,151	3.5	430.1
7	Influenza and pneumonia (J10-J18)	1,097	3.3	409.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	916	2.8	342.3
9	Septicemia (A40-A41)	768	2.3	287.0
10	Chronic lower respiratory diseases (J40-J47)	701	2.1	262.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	356	1.1	133.0
12	Pneumonitis due to solids and liquids (J69)	299	0.9	111.7
13	Atherosclerosis (I70)	191	0.6	71.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	149	0.5	55.7
15	Aortic aneurysm and dissection (I71)	123	0.4	46.0
...	All other causes (Residual)	6,573	19.9	2,456.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable