

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	135,519	100.0	297.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,021	21.4	63.8
2	Malignant neoplasms (C00-C97)	27,660	20.4	60.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,723	8.7	25.8
4	Cerebrovascular diseases (I60-I69)	7,078	5.2	15.6
5	Diabetes mellitus (E10-E14)	6,417	4.7	14.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,913	2.9	8.6
7	Chronic lower respiratory diseases (J40-J47)	3,531	2.6	7.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,466	2.6	7.6
9	Certain conditions originating in the perinatal period (P00-P96)	2,946	2.2	6.5
10	Influenza and pneumonia (J09-J18)	2,735	2.0	6.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,691	2.0	5.9
12	Alzheimer's disease (G30)	2,471	1.8	5.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,465	1.8	5.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,991	1.5	4.4
15	Septicemia (A40-A41)	1,894	1.4	4.2
...	All other causes (Residual)	25,517	18.8	56.1
Hispanic, both sexes, 1-4 years				
...	All causes	1,009	100.0	26.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	348	34.5	9.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	13.1	3.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	76	7.5	2.0
4	Malignant neoplasms (C00-C97)	68	6.7	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	3.9	1.0
6	Influenza and pneumonia (J09-J18)	24	2.4	0.6
7	Septicemia (A40-A41)	18	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	16	1.6	*
10	Cerebrovascular diseases (I60-I69)	8	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
12	Pneumonitis due to solids and liquids (J69)	5	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
15	Anemias (D50-D64)	2	0.2	*
15	Meningitis (G00,G03)	2	0.2	*
15	Diseases of appendix (K35-K38)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	236	23.4	6.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,157	100.0	14.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	387	33.4	4.7
2	Malignant neoplasms (C00-C97)	228	19.7	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	76	6.6	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	6.3	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	3.5	0.5
6	Influenza and pneumonia (J09-J18)	20	1.7	0.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	17	1.5	*
10	Septicemia (A40-A41)	16	1.4	*
10	Cerebrovascular diseases (I60-I69)	16	1.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
13	Anemias (D50-D64)	5	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
15	Meningitis (G00,G03)	3	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	223	19.3	2.7
Hispanic, both sexes, 15-24 years				
...	All causes	5,686	100.0	76.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,511	44.2	33.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,295	22.8	17.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	527	9.3	7.1
4	Malignant neoplasms (C00-C97)	345	6.1	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	2.6	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.0	0.7
7	Cerebrovascular diseases (I60-I69)	37	0.7	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	28	0.5	0.4
10	Legal intervention (Y35,Y89.0)	24	0.4	0.3
11	Chronic lower respiratory diseases (J40-J47)	20	0.4	0.3
12	Diabetes mellitus (E10-E14)	18	0.3	*
13	Septicemia (A40-A41)	17	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
...	All other causes (Residual)	599	10.5	8.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	6,546	100.0	80.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,321	35.5	28.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,028	15.7	12.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	649	9.9	7.9
4	Malignant neoplasms (C00-C97)	609	9.3	7.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	378	5.8	4.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	182	2.8	2.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	103	1.6	1.3
8	Cerebrovascular diseases (I60-I69)	100	1.5	1.2
9	Diabetes mellitus (E10-E14)	59	0.9	0.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	58	0.9	0.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	0.9	0.7
12	Septicemia (A40-A41)	41	0.6	0.5
13	Influenza and pneumonia (J09-J18)	38	0.6	0.5
14	Legal intervention (Y35,Y89.0)	36	0.5	0.4
15	Chronic lower respiratory diseases (J40-J47)	35	0.5	0.4
...	All other causes (Residual)	853	13.0	10.4
Hispanic, both sexes, 35-44 years				
...	All causes	8,695	100.0	127.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,958	22.5	28.6
2	Malignant neoplasms (C00-C97)	1,485	17.1	21.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	962	11.1	14.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	505	5.8	7.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	502	5.8	7.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	499	5.7	7.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	462	5.3	6.8
8	Cerebrovascular diseases (I60-I69)	301	3.5	4.4
9	Diabetes mellitus (E10-E14)	228	2.6	3.3
10	Viral hepatitis (B15-B19)	101	1.2	1.5
11	Influenza and pneumonia (J09-J18)	85	1.0	1.2
12	Septicemia (A40-A41)	81	0.9	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	0.9	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	0.5	0.6
15	Pregnancy, childbirth and the puerperium (O00-O99)	39	0.4	0.6
...	All other causes (Residual)	1,369	15.7	20.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	14,400	100.0	303.2
1	Malignant neoplasms (C00-C97)	3,554	24.7	74.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,317	16.1	48.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,659	11.5	34.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,167	8.1	24.6
5	Cerebrovascular diseases (I60-I69)	656	4.6	13.8
6	Diabetes mellitus (E10-E14)	631	4.4	13.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	515	3.6	10.8
8	Viral hepatitis (B15-B19)	392	2.7	8.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	391	2.7	8.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	253	1.8	5.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	230	1.6	4.8
12	Septicemia (A40-A41)	182	1.3	3.8
13	Chronic lower respiratory diseases (J40-J47)	141	1.0	3.0
14	Influenza and pneumonia (J09-J18)	130	0.9	2.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	112	0.8	2.4
...	All other causes (Residual)	2,070	14.4	43.6
Hispanic, both sexes, 55-64 years				
...	All causes	17,424	100.0	647.4
1	Malignant neoplasms (C00-C97)	5,326	30.6	197.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,751	21.5	139.4
3	Diabetes mellitus (E10-E14)	1,131	6.5	42.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	954	5.5	35.4
5	Cerebrovascular diseases (I60-I69)	911	5.2	33.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	795	4.6	29.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	371	2.1	13.8
8	Viral hepatitis (B15-B19)	358	2.1	13.3
9	Chronic lower respiratory diseases (J40-J47)	297	1.7	11.0
10	Septicemia (A40-A41)	253	1.5	9.4
11	Influenza and pneumonia (J09-J18)	226	1.3	8.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	192	1.1	7.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	191	1.1	7.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	170	1.0	6.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	0.5	3.5
...	All other causes (Residual)	2,403	13.8	89.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	21,127	100.0	1,477.1
1	Malignant neoplasms (C00-C97)	6,472	30.6	452.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,033	23.8	351.9
3	Diabetes mellitus (E10-E14)	1,535	7.3	107.3
4	Cerebrovascular diseases (I60-I69)	1,172	5.5	81.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	700	3.3	48.9
6	Chronic lower respiratory diseases (J40-J47)	645	3.1	45.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	611	2.9	42.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	527	2.5	36.8
9	Influenza and pneumonia (J09-J18)	393	1.9	27.5
10	Septicemia (A40-A41)	375	1.8	26.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	239	1.1	16.7
12	Alzheimer's disease (G30)	181	0.9	12.7
13	Viral hepatitis (B15-B19)	157	0.7	11.0
14	Parkinson's disease (G20-G21)	118	0.6	8.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	106	0.5	7.4
...	All other causes (Residual)	2,863	13.6	200.2
Hispanic, both sexes, 75-84 years				
...	All causes	29,490	100.0	3,678.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,114	27.5	1,012.1
2	Malignant neoplasms (C00-C97)	6,683	22.7	833.6
3	Cerebrovascular diseases (I60-I69)	2,022	6.9	252.2
4	Diabetes mellitus (E10-E14)	1,864	6.3	232.5
5	Chronic lower respiratory diseases (J40-J47)	1,321	4.5	164.8
6	Alzheimer's disease (G30)	867	2.9	108.1
7	Influenza and pneumonia (J09-J18)	806	2.7	100.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	787	2.7	98.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	616	2.1	76.8
10	Septicemia (A40-A41)	498	1.7	62.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	420	1.4	52.4
12	Parkinson's disease (G20-G21)	411	1.4	51.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	385	1.3	48.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	198	0.7	24.7
15	Pneumonitis due to solids and liquids (J69)	155	0.5	19.3
...	All other causes (Residual)	4,343	14.7	541.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	23,890	100.0	8,542.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,149	34.1	2,913.8
2	Malignant neoplasms (C00-C97)	2,871	12.0	1,026.6
3	Cerebrovascular diseases (I60-I69)	1,826	7.6	652.9
4	Alzheimer's disease (G30)	1,372	5.7	490.6
5	Chronic lower respiratory diseases (J40-J47)	999	4.2	357.2
6	Influenza and pneumonia (J09-J18)	952	4.0	340.4
7	Diabetes mellitus (E10-E14)	944	4.0	337.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	532	2.2	190.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	439	1.8	157.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	425	1.8	152.0
11	Septicemia (A40-A41)	347	1.5	124.1
12	Parkinson's disease (G20-G21)	214	0.9	76.5
13	Pneumonitis due to solids and liquids (J69)	210	0.9	75.1
14	Atherosclerosis (I70)	141	0.6	50.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	130	0.5	46.5
...	All other causes (Residual)	4,339	18.2	1,551.5
Hispanic, male, all ages				
...	All causes	75,708	100.0	321.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,657	20.7	66.6
2	Malignant neoplasms (C00-C97)	14,493	19.1	61.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,844	11.7	37.6
4	Cerebrovascular diseases (I60-I69)	3,319	4.4	14.1
5	Diabetes mellitus (E10-E14)	3,199	4.2	13.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,926	3.9	12.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,799	3.7	11.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,078	2.7	8.8
9	Chronic lower respiratory diseases (J40-J47)	1,894	2.5	8.1
10	Certain conditions originating in the perinatal period (P00-P96)	1,643	2.2	7.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,363	1.8	5.8
12	Influenza and pneumonia (J09-J18)	1,361	1.8	5.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,180	1.6	5.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,063	1.4	4.5
15	Septicemia (A40-A41)	944	1.2	4.0
...	All other causes (Residual)	12,945	17.1	55.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	556	100.0	28.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	40.5	11.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	13.7	3.8
3	Malignant neoplasms (C00-C97)	38	6.8	1.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	5.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.1	*
6	Influenza and pneumonia (J09-J18)	13	2.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.3	*
8	Septicemia (A40-A41)	6	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	0.9	*
10	Cerebrovascular diseases (I60-I69)	4	0.7	*
10	Pneumonitis due to solids and liquids (J69)	4	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
13	Meningococcal infection (A39)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Hernia (K40-K46)	1	0.2	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	118	21.2	5.9
Hispanic, male, 5-14 years				
...	All causes	663	100.0	15.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	244	36.8	5.8
2	Malignant neoplasms (C00-C97)	114	17.2	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	7.2	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	5.3	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.5	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	2.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.7	*
7	Influenza and pneumonia (J09-J18)	11	1.7	*
9	Septicemia (A40-A41)	10	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	10	1.5	*
11	Cerebrovascular diseases (I60-I69)	8	1.2	*
12	Anemias (D50-D64)	4	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	125	18.9	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,506	100.0	115.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,035	45.2	52.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,171	26.0	30.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	451	10.0	11.5
4	Malignant neoplasms (C00-C97)	209	4.6	5.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	2.3	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	0.9	1.0
7	Legal intervention (Y35,Y89.0)	24	0.5	0.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.5	0.6
9	Cerebrovascular diseases (I60-I69)	20	0.4	0.5
10	Diabetes mellitus (E10-E14)	14	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.2	*
12	Septicemia (A40-A41)	10	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
15	Aortic aneurysm and dissection (I71)	6	0.1	*
15	Influenza and pneumonia (J09-J18)	6	0.1	*
...	All other causes (Residual)	366	8.1	9.4
Hispanic, male, 25-34 years				
...	All causes	4,956	100.0	110.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,939	39.1	43.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	898	18.1	19.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	548	11.1	12.2
4	Malignant neoplasms (C00-C97)	282	5.7	6.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	277	5.6	6.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	135	2.7	3.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	81	1.6	1.8
8	Cerebrovascular diseases (I60-I69)	60	1.2	1.3
9	Diabetes mellitus (E10-E14)	39	0.8	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	0.7	0.8
11	Legal intervention (Y35,Y89.0)	36	0.7	0.8
12	Chronic lower respiratory diseases (J40-J47)	24	0.5	0.5
13	Influenza and pneumonia (J09-J18)	22	0.4	0.5
14	Septicemia (A40-A41)	19	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.3	*
...	All other causes (Residual)	543	11.0	12.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	6,037	100.0	166.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,558	25.8	42.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	743	12.3	20.5
3	Malignant neoplasms (C00-C97)	633	10.5	17.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	417	6.9	11.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	406	6.7	11.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	396	6.6	10.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	382	6.3	10.5
8	Cerebrovascular diseases (I60-I69)	194	3.2	5.3
9	Diabetes mellitus (E10-E14)	152	2.5	4.2
10	Viral hepatitis (B15-B19)	79	1.3	2.2
11	Influenza and pneumonia (J09-J18)	59	1.0	1.6
12	Septicemia (A40-A41)	49	0.8	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.7	1.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	0.4	0.7
14	Aortic aneurysm and dissection (I71)	26	0.4	0.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.4	0.7
...	All other causes (Residual)	846	14.0	23.3
Hispanic, male, 45-54 years				
...	All causes	9,634	100.0	399.2
1	Malignant neoplasms (C00-C97)	1,791	18.6	74.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,770	18.4	73.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,315	13.6	54.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	929	9.6	38.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	408	4.2	16.9
6	Cerebrovascular diseases (I60-I69)	399	4.1	16.5
7	Diabetes mellitus (E10-E14)	383	4.0	15.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	317	3.3	13.1
9	Viral hepatitis (B15-B19)	289	3.0	12.0
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	209	2.2	8.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	1.5	6.0
12	Septicemia (A40-A41)	112	1.2	4.6
13	Chronic lower respiratory diseases (J40-J47)	84	0.9	3.5
14	Influenza and pneumonia (J09-J18)	79	0.8	3.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	0.7	2.8
...	All other causes (Residual)	1,338	13.9	55.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	10,764	100.0	831.4
1	Malignant neoplasms (C00-C97)	2,873	26.7	221.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,614	24.3	201.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	723	6.7	55.8
4	Diabetes mellitus (E10-E14)	661	6.1	51.1
5	Cerebrovascular diseases (I60-I69)	556	5.2	42.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	556	5.2	42.9
7	Viral hepatitis (B15-B19)	259	2.4	20.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	207	1.9	16.0
9	Chronic lower respiratory diseases (J40-J47)	173	1.6	13.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	162	1.5	12.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	145	1.3	11.2
12	Septicemia (A40-A41)	144	1.3	11.1
13	Influenza and pneumonia (J09-J18)	143	1.3	11.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	0.9	7.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	0.7	5.8
...	All other causes (Residual)	1,372	12.7	106.0
Hispanic, male, 65-74 years				
...	All causes	11,978	100.0	1,862.7
1	Malignant neoplasms (C00-C97)	3,603	30.1	560.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,067	25.6	477.0
3	Diabetes mellitus (E10-E14)	818	6.8	127.2
4	Cerebrovascular diseases (I60-I69)	608	5.1	94.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	441	3.7	68.6
6	Chronic lower respiratory diseases (J40-J47)	387	3.2	60.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	352	2.9	54.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	311	2.6	48.4
9	Influenza and pneumonia (J09-J18)	216	1.8	33.6
10	Septicemia (A40-A41)	202	1.7	31.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	130	1.1	20.2
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	0.7	13.7
13	Alzheimer's disease (G30)	85	0.7	13.2
14	Viral hepatitis (B15-B19)	82	0.7	12.8
15	Parkinson's disease (G20-G21)	70	0.6	10.9
...	All other causes (Residual)	1,518	12.7	236.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	14,438	100.0	4,364.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,080	28.3	1,233.4
2	Malignant neoplasms (C00-C97)	3,548	24.6	1,072.6
3	Cerebrovascular diseases (I60-I69)	872	6.0	263.6
4	Diabetes mellitus (E10-E14)	810	5.6	244.9
5	Chronic lower respiratory diseases (J40-J47)	718	5.0	217.1
6	Influenza and pneumonia (J09-J18)	419	2.9	126.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	391	2.7	118.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	336	2.3	101.6
9	Alzheimer's disease (G30)	315	2.2	95.2
10	Parkinson's disease (G20-G21)	245	1.7	74.1
11	Septicemia (A40-A41)	244	1.7	73.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	195	1.4	59.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	160	1.1	48.4
14	Pneumonitis due to solids and liquids (J69)	87	0.6	26.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	84	0.6	25.4
...	All other causes (Residual)	1,934	13.4	584.7
Hispanic, male, 85 years and over				
...	All causes	8,809	100.0	8,953.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,913	33.1	2,960.8
2	Malignant neoplasms (C00-C97)	1,395	15.8	1,417.9
3	Cerebrovascular diseases (I60-I69)	585	6.6	594.6
4	Chronic lower respiratory diseases (J40-J47)	454	5.2	461.5
5	Alzheimer's disease (G30)	384	4.4	390.3
6	Influenza and pneumonia (J09-J18)	364	4.1	370.0
7	Diabetes mellitus (E10-E14)	319	3.6	324.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	220	2.5	223.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	2.1	186.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	149	1.7	151.4
11	Septicemia (A40-A41)	112	1.3	113.8
12	Parkinson's disease (G20-G21)	105	1.2	106.7
13	Pneumonitis due to solids and liquids (J69)	99	1.1	100.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.5	45.7
14	Aortic aneurysm and dissection (I71)	45	0.5	45.7
...	All other causes (Residual)	1,437	16.3	1,460.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	59,811	100.0	272.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,364	22.3	60.8
2	Malignant neoplasms (C00-C97)	13,167	22.0	59.9
3	Cerebrovascular diseases (I60-I69)	3,759	6.3	17.1
4	Diabetes mellitus (E10-E14)	3,218	5.4	14.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,879	4.8	13.1
6	Alzheimer's disease (G30)	1,670	2.8	7.6
7	Chronic lower respiratory diseases (J40-J47)	1,637	2.7	7.4
8	Influenza and pneumonia (J09-J18)	1,374	2.3	6.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,328	2.2	6.0
10	Certain conditions originating in the perinatal period (P00-P96)	1,303	2.2	5.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,114	1.9	5.1
12	Septicemia (A40-A41)	950	1.6	4.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	928	1.6	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	756	1.3	3.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	540	0.9	2.5
...	All other causes (Residual)	11,824	19.8	53.8
Hispanic, female, 1-4 years				
...	All causes	453	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	27.2	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	12.4	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	10.2	2.4
4	Malignant neoplasms (C00-C97)	30	6.6	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.9	1.2
6	Septicemia (A40-A41)	12	2.6	*
7	Influenza and pneumonia (J09-J18)	11	2.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	11	2.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	2.2	*
10	Cerebrovascular diseases (I60-I69)	4	0.9	*
10	Acute bronchitis and bronchiolitis (J20-J21)	4	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (Residual)	110	24.3	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	494	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	28.9	3.6
2	Malignant neoplasms (C00-C97)	114	23.1	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	7.7	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	5.7	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.4	*
6	Influenza and pneumonia (J09-J18)	9	1.8	*
7	Cerebrovascular diseases (I60-I69)	8	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	7	1.4	*
9	Septicemia (A40-A41)	6	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	1.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.8	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Meningitis (G00,G03)	2	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
...	All other causes (Residual)	102	20.6	2.5
Hispanic, female, 15-24 years				
...	All causes	1,180	100.0	33.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	476	40.3	13.5
2	Malignant neoplasms (C00-C97)	136	11.5	3.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	124	10.5	3.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	76	6.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.5	1.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	28	2.4	0.8
7	Cerebrovascular diseases (I60-I69)	17	1.4	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	1.4	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	9	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.8	*
12	Septicemia (A40-A41)	7	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
12	Influenza and pneumonia (J09-J18)	7	0.6	*
15	Anemias (D50-D64)	5	0.4	*
15	Pneumonitis due to solids and liquids (J69)	5	0.4	*
...	All other causes (Residual)	208	17.6	5.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,590	100.0	43.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	382	24.0	10.4
2	Malignant neoplasms (C00-C97)	327	20.6	8.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	8.2	3.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	6.4	2.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	6.4	2.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	58	3.6	1.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	47	3.0	1.3
8	Cerebrovascular diseases (I60-I69)	40	2.5	1.1
9	Septicemia (A40-A41)	22	1.4	0.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	22	1.4	0.6
11	Diabetes mellitus (E10-E14)	20	1.3	0.5
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.2	*
13	Influenza and pneumonia (J09-J18)	16	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.0	*
15	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
...	All other causes (Residual)	278	17.5	7.6
Hispanic, female, 35-44 years				
...	All causes	2,658	100.0	82.7
1	Malignant neoplasms (C00-C97)	852	32.1	26.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	400	15.0	12.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	219	8.2	6.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	109	4.1	3.4
5	Cerebrovascular diseases (I60-I69)	107	4.0	3.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	96	3.6	3.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	3.1	2.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	80	3.0	2.5
9	Diabetes mellitus (E10-E14)	76	2.9	2.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	39	1.5	1.2
11	Septicemia (A40-A41)	32	1.2	1.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.1	0.9
13	Influenza and pneumonia (J09-J18)	26	1.0	0.8
14	Viral hepatitis (B15-B19)	22	0.8	0.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.7	*
...	All other causes (Residual)	471	17.7	14.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	4,766	100.0	204.0
1	Malignant neoplasms (C00-C97)	1,763	37.0	75.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	547	11.5	23.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	344	7.2	14.7
4	Cerebrovascular diseases (I60-I69)	257	5.4	11.0
5	Diabetes mellitus (E10-E14)	248	5.2	10.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	238	5.0	10.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	107	2.2	4.6
8	Viral hepatitis (B15-B19)	103	2.2	4.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	1.8	3.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	1.6	3.2
11	Septicemia (A40-A41)	70	1.5	3.0
12	Chronic lower respiratory diseases (J40-J47)	57	1.2	2.4
13	Influenza and pneumonia (J09-J18)	51	1.1	2.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	0.9	1.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	0.9	1.9
...	All other causes (Residual)	732	15.4	31.3
Hispanic, female, 55-64 years				
...	All causes	6,660	100.0	476.9
1	Malignant neoplasms (C00-C97)	2,453	36.8	175.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,137	17.1	81.4
3	Diabetes mellitus (E10-E14)	470	7.1	33.7
4	Cerebrovascular diseases (I60-I69)	355	5.3	25.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	3.6	17.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	231	3.5	16.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	2.5	11.7
8	Chronic lower respiratory diseases (J40-J47)	124	1.9	8.9
9	Septicemia (A40-A41)	109	1.6	7.8
10	Viral hepatitis (B15-B19)	99	1.5	7.1
11	Influenza and pneumonia (J09-J18)	83	1.2	5.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.0	4.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	47	0.7	3.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.5	2.6
15	Alzheimer's disease (G30)	30	0.5	2.1
...	All other causes (Residual)	1,014	15.2	72.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	9,149	100.0	1,162.1
1	Malignant neoplasms (C00-C97)	2,869	31.4	364.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,966	21.5	249.7
3	Diabetes mellitus (E10-E14)	717	7.8	91.1
4	Cerebrovascular diseases (I60-I69)	564	6.2	71.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	300	3.3	38.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	259	2.8	32.9
7	Chronic lower respiratory diseases (J40-J47)	258	2.8	32.8
8	Influenza and pneumonia (J09-J18)	177	1.9	22.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	175	1.9	22.2
10	Septicemia (A40-A41)	173	1.9	22.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	109	1.2	13.8
12	Alzheimer's disease (G30)	96	1.0	12.2
13	Viral hepatitis (B15-B19)	75	0.8	9.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.6	6.7
15	Parkinson's disease (G20-G21)	48	0.5	6.1
...	All other causes (Residual)	1,310	14.3	166.4
Hispanic, female, 75-84 years				
...	All causes	15,052	100.0	3,196.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,034	26.8	856.6
2	Malignant neoplasms (C00-C97)	3,135	20.8	665.7
3	Cerebrovascular diseases (I60-I69)	1,150	7.6	244.2
4	Diabetes mellitus (E10-E14)	1,054	7.0	223.8
5	Chronic lower respiratory diseases (J40-J47)	603	4.0	128.0
6	Alzheimer's disease (G30)	552	3.7	117.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	396	2.6	84.1
8	Influenza and pneumonia (J09-J18)	387	2.6	82.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	280	1.9	59.5
10	Septicemia (A40-A41)	254	1.7	53.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	225	1.5	47.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	225	1.5	47.8
13	Parkinson's disease (G20-G21)	166	1.1	35.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	114	0.8	24.2
15	Pneumonitis due to solids and liquids (J69)	68	0.5	14.4
...	All other causes (Residual)	2,409	16.0	511.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	15,081	100.0	8,318.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,236	34.7	2,888.2
2	Malignant neoplasms (C00-C97)	1,476	9.8	814.2
3	Cerebrovascular diseases (I60-I69)	1,241	8.2	684.5
4	Alzheimer's disease (G30)	988	6.6	545.0
5	Diabetes mellitus (E10-E14)	625	4.1	344.8
6	Influenza and pneumonia (J09-J18)	588	3.9	324.3
7	Chronic lower respiratory diseases (J40-J47)	545	3.6	300.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	312	2.1	172.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	290	1.9	160.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	1.6	133.5
11	Septicemia (A40-A41)	235	1.6	129.6
12	Pneumonitis due to solids and liquids (J69)	111	0.7	61.2
13	Parkinson's disease (G20-G21)	109	0.7	60.1
14	Atherosclerosis (I70)	99	0.7	54.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	0.6	46.9
...	All other causes (Residual)	2,899	19.2	1,599.1
Non-Hispanic, white, both sexes, all ages				
...	All causes	1,939,606	100.0	964.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	502,683	25.9	249.9
2	Malignant neoplasms (C00-C97)	456,576	23.5	226.9
3	Chronic lower respiratory diseases (J40-J47)	114,480	5.9	56.9
4	Cerebrovascular diseases (I60-I69)	107,678	5.6	53.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94,584	4.9	47.0
6	Alzheimer's disease (G30)	66,453	3.4	33.0
7	Diabetes mellitus (E10-E14)	50,046	2.6	24.9
8	Influenza and pneumonia (J09-J18)	43,219	2.2	21.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,219	1.8	17.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,897	1.5	14.4
11	Septicemia (A40-A41)	25,856	1.3	12.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	21,598	1.1	10.7
13	Parkinson's disease (G20-G21)	18,098	0.9	9.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,216	0.9	8.6
15	Pneumonitis due to solids and liquids (J69)	14,680	0.8	7.3
...	All other causes (Residual)	343,323	17.7	170.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,325	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	835	35.9	9.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	266	11.4	2.9
3	Malignant neoplasms (C00-C97)	224	9.6	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	6.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	3.2	0.8
6	Influenza and pneumonia (J09-J18)	50	2.2	0.5
7	Septicemia (A40-A41)	42	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.4	0.4
9	Cerebrovascular diseases (I60-I69)	26	1.1	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	25	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	18	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.6	*
14	Meningitis (G00,G03)	10	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.2	*
...	All other causes (Residual)	544	23.4	6.0
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	3,362	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,250	37.2	5.3
2	Malignant neoplasms (C00-C97)	559	16.6	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	202	6.0	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	3.8	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	3.5	0.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	3.2	0.5
7	Influenza and pneumonia (J09-J18)	58	1.7	0.2
8	Cerebrovascular diseases (I60-I69)	47	1.4	0.2
9	Septicemia (A40-A41)	43	1.3	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	39	1.2	0.2
12	Meningitis (G00,G03)	16	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
14	Anemias (D50-D64)	13	0.4	*
15	Diabetes mellitus (E10-E14)	12	0.4	*
...	All other causes (Residual)	712	21.2	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	19,607	100.0	74.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,032	56.3	41.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,932	15.0	11.1
3	Malignant neoplasms (C00-C97)	958	4.9	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	937	4.8	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	565	2.9	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	245	1.2	0.9
7	Influenza and pneumonia (J09-J18)	116	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	102	0.5	0.4
9	Diabetes mellitus (E10-E14)	92	0.5	0.3
10	Septicemia (A40-A41)	89	0.5	0.3
11	Chronic lower respiratory diseases (J40-J47)	59	0.3	0.2
11	Pregnancy, childbirth and the puerperium (O00-O99)	59	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	30	0.2	0.1
...	All other causes (Residual)	2,306	11.8	8.8
Non-Hispanic, white, both sexes, 25-34 years				
...	All causes	24,757	100.0	102.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,347	41.8	42.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,892	15.7	16.1
3	Malignant neoplasms (C00-C97)	2,060	8.3	8.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,708	6.9	7.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,054	4.3	4.4
6	Diabetes mellitus (E10-E14)	321	1.3	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	267	1.1	1.1
8	Cerebrovascular diseases (I60-I69)	231	0.9	1.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	206	0.8	0.9
10	Influenza and pneumonia (J09-J18)	189	0.8	0.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	187	0.8	0.8
12	Septicemia (A40-A41)	153	0.6	0.6
13	Pregnancy, childbirth and the puerperium (O00-O99)	136	0.5	0.6
14	Chronic lower respiratory diseases (J40-J47)	129	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	114	0.5	0.5
...	All other causes (Residual)	3,763	15.2	15.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-44 years				
...	All causes	50,966	100.0	181.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,270	24.1	43.7
2	Malignant neoplasms (C00-C97)	8,830	17.3	31.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,458	14.6	26.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,535	10.9	19.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,726	3.4	6.1
6	Diabetes mellitus (E10-E14)	1,118	2.2	4.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,072	2.1	3.8
8	Cerebrovascular diseases (I60-I69)	1,068	2.1	3.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,019	2.0	3.6
10	Chronic lower respiratory diseases (J40-J47)	539	1.1	1.9
11	Septicemia (A40-A41)	525	1.0	1.9
12	Influenza and pneumonia (J09-J18)	496	1.0	1.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	376	0.7	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	334	0.7	1.2
15	Viral hepatitis (B15-B19)	288	0.6	1.0
...	All other causes (Residual)	8,312	16.3	29.6
Non-Hispanic, white, both sexes, 45-54 years				
...	All causes	127,898	100.0	403.8
1	Malignant neoplasms (C00-C97)	36,052	28.2	113.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,823	20.2	81.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,345	12.0	48.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,866	5.4	21.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,851	4.6	18.5
6	Diabetes mellitus (E10-E14)	3,505	2.7	11.1
7	Cerebrovascular diseases (I60-I69)	3,493	2.7	11.0
8	Chronic lower respiratory diseases (J40-J47)	3,246	2.5	10.2
9	Viral hepatitis (B15-B19)	1,957	1.5	6.2
10	Septicemia (A40-A41)	1,445	1.1	4.6
11	Influenza and pneumonia (J09-J18)	1,270	1.0	4.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,243	1.0	3.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,120	0.9	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,018	0.8	3.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	558	0.4	1.8
...	All other causes (Residual)	19,106	14.9	60.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-64 years				
...	All causes	213,876	100.0	849.6
1	Malignant neoplasms (C00-C97)	80,103	37.5	318.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,803	22.4	189.9
3	Chronic lower respiratory diseases (J40-J47)	11,100	5.2	44.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,402	4.4	37.3
5	Diabetes mellitus (E10-E14)	7,272	3.4	28.9
6	Cerebrovascular diseases (I60-I69)	6,533	3.1	26.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,927	2.8	23.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,593	2.1	18.2
9	Septicemia (A40-A41)	2,868	1.3	11.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,525	1.2	10.0
11	Influenza and pneumonia (J09-J18)	2,253	1.1	9.0
12	Viral hepatitis (B15-B19)	1,539	0.7	6.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,149	0.5	4.6
14	Aortic aneurysm and dissection (I71)	1,141	0.5	4.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,106	0.5	4.4
...	All other causes (Residual)	28,562	13.4	113.5
Non-Hispanic, white, both sexes, 65-74 years				
...	All causes	306,342	100.0	2,006.2
1	Malignant neoplasms (C00-C97)	112,431	36.7	736.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,564	22.4	449.0
3	Chronic lower respiratory diseases (J40-J47)	25,720	8.4	168.4
4	Cerebrovascular diseases (I60-I69)	12,861	4.2	84.2
5	Diabetes mellitus (E10-E14)	10,128	3.3	66.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,997	2.3	45.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,054	1.6	33.1
8	Septicemia (A40-A41)	4,475	1.5	29.3
9	Influenza and pneumonia (J09-J18)	4,161	1.4	27.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	3,930	1.3	25.7
11	Alzheimer's disease (G30)	3,380	1.1	22.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,211	0.7	14.5
13	Aortic aneurysm and dissection (I71)	2,208	0.7	14.5
14	Parkinson's disease (G20-G21)	2,001	0.7	13.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,980	0.6	13.0
...	All other causes (Residual)	40,241	13.1	263.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-84 years				
...	All causes	549,752	100.0	5,113.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	142,967	26.0	1,329.7
2	Malignant neoplasms (C00-C97)	138,954	25.3	1,292.4
3	Chronic lower respiratory diseases (J40-J47)	43,637	7.9	405.9
4	Cerebrovascular diseases (I60-I69)	34,331	6.2	319.3
5	Alzheimer's disease (G30)	20,248	3.7	188.3
6	Diabetes mellitus (E10-E14)	15,567	2.8	144.8
7	Influenza and pneumonia (J09-J18)	12,274	2.2	114.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,954	2.2	111.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,505	2.1	107.0
10	Parkinson's disease (G20-G21)	8,421	1.5	78.3
11	Septicemia (A40-A41)	8,194	1.5	76.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,735	0.9	44.0
13	Pneumonitis due to solids and liquids (J69)	4,489	0.8	41.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,106	0.7	38.2
15	Aortic aneurysm and dissection (I71)	3,831	0.7	35.6
...	All other causes (Residual)	84,539	15.4	786.3
Non-Hispanic, white, both sexes, 85 years and over				
...	All causes	627,642	100.0	13,413.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	207,410	33.0	4,432.5
2	Malignant neoplasms (C00-C97)	76,361	12.2	1,631.9
3	Cerebrovascular diseases (I60-I69)	48,918	7.8	1,045.4
4	Alzheimer's disease (G30)	42,105	6.7	899.8
5	Chronic lower respiratory diseases (J40-J47)	29,973	4.8	640.6
6	Influenza and pneumonia (J09-J18)	22,262	3.5	475.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,501	2.3	309.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,436	2.1	287.1
9	Diabetes mellitus (E10-E14)	12,025	1.9	257.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,616	1.4	184.1
11	Septicemia (A40-A41)	7,924	1.3	169.3
12	Pneumonitis due to solids and liquids (J69)	7,579	1.2	162.0
13	Parkinson's disease (G20-G21)	7,284	1.2	155.7
14	Atherosclerosis (I70)	4,112	0.7	87.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,798	0.6	81.2
...	All other causes (Residual)	121,338	19.3	2,593.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	948,662	100.0	960.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251,229	26.5	254.3
2	Malignant neoplasms (C00-C97)	237,672	25.1	240.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59,274	6.2	60.0
4	Chronic lower respiratory diseases (J40-J47)	53,989	5.7	54.7
5	Cerebrovascular diseases (I60-I69)	41,416	4.4	41.9
6	Diabetes mellitus (E10-E14)	25,584	2.7	25.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22,660	2.4	22.9
8	Alzheimer's disease (G30)	19,378	2.0	19.6
9	Influenza and pneumonia (J09-J18)	19,361	2.0	19.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,896	1.8	17.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	14,007	1.5	14.2
12	Septicemia (A40-A41)	11,653	1.2	11.8
13	Parkinson's disease (G20-G21)	10,457	1.1	10.6
14	Pneumonitis due to solids and liquids (J69)	7,713	0.8	7.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,480	0.7	6.6
...	All other causes (Residual)	150,893	15.9	152.8
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,316	100.0	28.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	39.6	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	10.3	2.9
3	Malignant neoplasms (C00-C97)	111	8.4	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	6.6	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	2.3	0.6
6	Septicemia (A40-A41)	27	2.1	0.6
7	Influenza and pneumonia (J09-J18)	21	1.6	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	14	1.1	*
10	Cerebrovascular diseases (I60-I69)	12	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.8	*
12	Meningitis (G00,G03)	6	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	4	0.3	*
...	All other causes (Residual)	307	23.3	6.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,945	100.0	16.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	763	39.2	6.3
2	Malignant neoplasms (C00-C97)	293	15.1	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	5.7	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	78	4.0	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	3.5	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	2.9	0.5
7	Septicemia (A40-A41)	29	1.5	0.2
8	Chronic lower respiratory diseases (J40-J47)	26	1.3	0.2
9	Influenza and pneumonia (J09-J18)	25	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	24	1.2	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.0	0.2
12	Anemias (D50-D64)	9	0.5	*
13	Diabetes mellitus (E10-E14)	8	0.4	*
13	Pneumonitis due to solids and liquids (J69)	8	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.4	*
...	All other causes (Residual)	418	21.5	3.5
Non-Hispanic, white, male, 15-24 years				
...	All causes	14,129	100.0	104.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,137	57.6	60.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,460	17.4	18.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	658	4.7	4.9
4	Malignant neoplasms (C00-C97)	576	4.1	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	343	2.4	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	1.0	1.0
7	Influenza and pneumonia (J09-J18)	68	0.5	0.5
8	Diabetes mellitus (E10-E14)	60	0.4	0.4
9	Cerebrovascular diseases (I60-I69)	56	0.4	0.4
10	Septicemia (A40-A41)	49	0.3	0.4
11	Chronic lower respiratory diseases (J40-J47)	38	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.2	0.2
12	Pneumonitis due to solids and liquids (J69)	22	0.2	0.2
12	Legal intervention (Y35,Y89.0)	22	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.1	0.1
...	All other causes (Residual)	1,462	10.3	10.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-34 years				
...	All causes	17,138	100.0	140.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,709	45.0	63.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,143	18.3	25.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,163	6.8	9.6
4	Malignant neoplasms (C00-C97)	1,052	6.1	8.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	755	4.4	6.2
6	Diabetes mellitus (E10-E14)	196	1.1	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	148	0.9	1.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	135	0.8	1.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	130	0.8	1.1
10	Cerebrovascular diseases (I60-I69)	107	0.6	0.9
11	Influenza and pneumonia (J09-J18)	106	0.6	0.9
12	Septicemia (A40-A41)	84	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	65	0.4	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	0.4	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.3	0.4
...	All other causes (Residual)	2,230	13.0	18.3
Non-Hispanic, white, male, 35-44 years				
...	All causes	32,139	100.0	228.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,537	26.6	60.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,318	16.5	37.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,209	13.1	29.9
4	Malignant neoplasms (C00-C97)	3,897	12.1	27.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,097	3.4	7.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	852	2.7	6.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	711	2.2	5.1
8	Diabetes mellitus (E10-E14)	709	2.2	5.0
9	Cerebrovascular diseases (I60-I69)	562	1.7	4.0
10	Septicemia (A40-A41)	269	0.8	1.9
11	Influenza and pneumonia (J09-J18)	266	0.8	1.9
12	Chronic lower respiratory diseases (J40-J47)	256	0.8	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	0.6	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	0.6	1.3
15	Viral hepatitis (B15-B19)	181	0.6	1.3
...	All other causes (Residual)	4,888	15.2	34.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-54 years				
...	All causes	79,931	100.0	508.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,107	23.9	121.6
2	Malignant neoplasms (C00-C97)	18,358	23.0	116.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,398	13.0	66.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,119	6.4	32.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,081	5.1	26.0
6	Diabetes mellitus (E10-E14)	2,214	2.8	14.1
7	Cerebrovascular diseases (I60-I69)	1,936	2.4	12.3
8	Chronic lower respiratory diseases (J40-J47)	1,686	2.1	10.7
9	Viral hepatitis (B15-B19)	1,396	1.7	8.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,026	1.3	6.5
11	Septicemia (A40-A41)	762	1.0	4.8
12	Influenza and pneumonia (J09-J18)	727	0.9	4.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	696	0.9	4.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	647	0.8	4.1
15	Aortic aneurysm and dissection (I71)	368	0.5	2.3
...	All other causes (Residual)	11,410	14.3	72.6
Non-Hispanic, white, male, 55-64 years				
...	All causes	129,977	100.0	1,057.5
1	Malignant neoplasms (C00-C97)	43,953	33.8	357.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,001	26.2	276.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,410	4.9	52.2
4	Chronic lower respiratory diseases (J40-J47)	5,771	4.4	47.0
5	Diabetes mellitus (E10-E14)	4,443	3.4	36.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,242	3.3	34.5
7	Cerebrovascular diseases (I60-I69)	3,692	2.8	30.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,450	2.7	28.1
9	Septicemia (A40-A41)	1,471	1.1	12.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,449	1.1	11.8
11	Influenza and pneumonia (J09-J18)	1,341	1.0	10.9
12	Viral hepatitis (B15-B19)	1,144	0.9	9.3
13	Aortic aneurysm and dissection (I71)	872	0.7	7.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	731	0.6	5.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	656	0.5	5.3
...	All other causes (Residual)	16,351	12.6	133.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-74 years				
...	All causes	172,867	100.0	2,432.7
1	Malignant neoplasms (C00-C97)	61,631	35.7	867.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,414	25.1	610.9
3	Chronic lower respiratory diseases (J40-J47)	12,998	7.5	182.9
4	Cerebrovascular diseases (I60-I69)	6,668	3.9	93.8
5	Diabetes mellitus (E10-E14)	5,840	3.4	82.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,447	2.6	62.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,718	1.6	38.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2,435	1.4	34.3
9	Septicemia (A40-A41)	2,292	1.3	32.3
10	Influenza and pneumonia (J09-J18)	2,260	1.3	31.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,808	1.0	25.4
12	Aortic aneurysm and dissection (I71)	1,462	0.8	20.6
13	Alzheimer's disease (G30)	1,436	0.8	20.2
14	Parkinson's disease (G20-G21)	1,306	0.8	18.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,124	0.7	15.8
...	All other causes (Residual)	21,028	12.2	295.9
Non-Hispanic, white, male, 75-84 years				
...	All causes	272,398	100.0	6,152.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,600	27.4	1,685.0
2	Malignant neoplasms (C00-C97)	73,175	26.9	1,652.8
3	Chronic lower respiratory diseases (J40-J47)	20,997	7.7	474.3
4	Cerebrovascular diseases (I60-I69)	14,446	5.3	326.3
5	Diabetes mellitus (E10-E14)	7,625	2.8	172.2
6	Alzheimer's disease (G30)	7,290	2.7	164.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,317	2.3	142.7
8	Influenza and pneumonia (J09-J18)	6,098	2.2	137.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,013	2.2	135.8
10	Parkinson's disease (G20-G21)	5,154	1.9	116.4
11	Septicemia (A40-A41)	3,829	1.4	86.5
12	Pneumonitis due to solids and liquids (J69)	2,534	0.9	57.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,285	0.8	51.6
14	Aortic aneurysm and dissection (I71)	2,084	0.8	47.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,981	0.7	44.7
...	All other causes (Residual)	37,970	13.9	857.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85 years and over				
...	All causes	219,419	100.0	14,588.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,076	33.3	4,858.5
2	Malignant neoplasms (C00-C97)	34,600	15.8	2,300.4
3	Cerebrovascular diseases (I60-I69)	13,873	6.3	922.4
4	Chronic lower respiratory diseases (J40-J47)	12,131	5.5	806.5
5	Alzheimer's disease (G30)	10,324	4.7	686.4
6	Influenza and pneumonia (J09-J18)	8,390	3.8	557.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,746	2.6	382.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,660	2.6	376.3
9	Diabetes mellitus (E10-E14)	4,484	2.0	298.1
10	Parkinson's disease (G20-G21)	3,745	1.7	249.0
11	Pneumonitis due to solids and liquids (J69)	3,563	1.6	236.9
12	Septicemia (A40-A41)	2,789	1.3	185.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,254	1.0	149.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,542	0.7	102.5
15	Aortic aneurysm and dissection (I71)	1,326	0.6	88.2
...	All other causes (Residual)	35,916	16.4	2,387.9
Non-Hispanic, white, female, all ages				
...	All causes	990,944	100.0	967.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251,454	25.4	245.5
2	Malignant neoplasms (C00-C97)	218,904	22.1	213.7
3	Cerebrovascular diseases (I60-I69)	66,262	6.7	64.7
4	Chronic lower respiratory diseases (J40-J47)	60,491	6.1	59.1
5	Alzheimer's disease (G30)	47,075	4.8	46.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,310	3.6	34.5
7	Diabetes mellitus (E10-E14)	24,462	2.5	23.9
8	Influenza and pneumonia (J09-J18)	23,858	2.4	23.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,323	1.7	16.9
10	Septicemia (A40-A41)	14,203	1.4	13.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,736	1.1	10.5
12	Parkinson's disease (G20-G21)	7,641	0.8	7.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,591	0.8	7.4
14	Pneumonitis due to solids and liquids (J69)	6,967	0.7	6.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,237	0.6	6.1
...	All other causes (Residual)	192,430	19.4	187.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	1,009	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	31.1	7.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	12.9	2.9
3	Malignant neoplasms (C00-C97)	113	11.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	5.9	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.5	1.0
6	Influenza and pneumonia (J09-J18)	29	2.9	0.7
7	Septicemia (A40-A41)	15	1.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.4	*
8	Cerebrovascular diseases (I60-I69)	14	1.4	*
10	Certain conditions originating in the perinatal period (P00-P96)	11	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
14	Meningitis (G00,G03)	4	0.4	*
15	Anemias (D50-D64)	2	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.2	*
...	All other causes (Residual)	234	23.2	5.3
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,417	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	487	34.4	4.2
2	Malignant neoplasms (C00-C97)	266	18.8	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	6.4	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	4.4	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	4.2	0.5
6	Influenza and pneumonia (J09-J18)	33	2.3	0.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	2.0	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.6	0.2
8	Cerebrovascular diseases (I60-I69)	23	1.6	0.2
10	Septicemia (A40-A41)	14	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	13	0.9	*
12	Meningitis (G00,G03)	10	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.6	*
14	Anemias (D50-D64)	4	0.3	*
14	Diabetes mellitus (E10-E14)	4	0.3	*
...	All other causes (Residual)	290	20.5	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	5,478	100.0	42.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,895	52.8	22.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	472	8.6	3.7
3	Malignant neoplasms (C00-C97)	382	7.0	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	279	5.1	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	222	4.1	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	2.0	0.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	59	1.1	0.5
8	Influenza and pneumonia (J09-J18)	48	0.9	0.4
9	Cerebrovascular diseases (I60-I69)	46	0.8	0.4
10	Septicemia (A40-A41)	40	0.7	0.3
11	Diabetes mellitus (E10-E14)	32	0.6	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.4	0.2
13	Chronic lower respiratory diseases (J40-J47)	21	0.4	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.4	0.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	11	0.2	*
...	All other causes (Residual)	819	15.0	6.4
Non-Hispanic, white, female, 25-34 years				
...	All causes	7,619	100.0	63.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,638	34.6	22.0
2	Malignant neoplasms (C00-C97)	1,008	13.2	8.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	749	9.8	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	545	7.2	4.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	299	3.9	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	136	1.8	1.1
7	Diabetes mellitus (E10-E14)	125	1.6	1.0
8	Cerebrovascular diseases (I60-I69)	124	1.6	1.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	1.6	1.0
10	Influenza and pneumonia (J09-J18)	83	1.1	0.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	71	0.9	0.6
12	Septicemia (A40-A41)	69	0.9	0.6
13	Chronic lower respiratory diseases (J40-J47)	64	0.8	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	57	0.7	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	0.7	0.4
...	All other causes (Residual)	1,482	19.5	12.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-44 years				
...	All causes	18,827	100.0	134.4
1	Malignant neoplasms (C00-C97)	4,933	26.2	35.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,733	19.8	26.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,140	11.4	15.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,326	7.0	9.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	629	3.3	4.5
6	Cerebrovascular diseases (I60-I69)	506	2.7	3.6
7	Diabetes mellitus (E10-E14)	409	2.2	2.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	361	1.9	2.6
9	Chronic lower respiratory diseases (J40-J47)	283	1.5	2.0
10	Septicemia (A40-A41)	256	1.4	1.8
11	Influenza and pneumonia (J09-J18)	230	1.2	1.6
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	1.0	1.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	167	0.9	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	0.7	1.0
15	Viral hepatitis (B15-B19)	107	0.6	0.8
...	All other causes (Residual)	3,424	18.2	24.4
Non-Hispanic, white, female, 45-54 years				
...	All causes	47,967	100.0	300.5
1	Malignant neoplasms (C00-C97)	17,694	36.9	110.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,716	14.0	42.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,947	10.3	31.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,770	3.7	11.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,747	3.6	10.9
6	Chronic lower respiratory diseases (J40-J47)	1,560	3.3	9.8
7	Cerebrovascular diseases (I60-I69)	1,557	3.2	9.8
8	Diabetes mellitus (E10-E14)	1,291	2.7	8.1
9	Septicemia (A40-A41)	683	1.4	4.3
10	Viral hepatitis (B15-B19)	561	1.2	3.5
11	Influenza and pneumonia (J09-J18)	543	1.1	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	473	1.0	3.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	322	0.7	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	252	0.5	1.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	225	0.5	1.4
...	All other causes (Residual)	7,626	15.9	47.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-64 years				
...	All causes	83,899	100.0	651.3
1	Malignant neoplasms (C00-C97)	36,150	43.1	280.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,802	16.5	107.1
3	Chronic lower respiratory diseases (J40-J47)	5,329	6.4	41.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,992	3.6	23.2
5	Cerebrovascular diseases (I60-I69)	2,841	3.4	22.1
6	Diabetes mellitus (E10-E14)	2,829	3.4	22.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,685	2.0	13.1
8	Septicemia (A40-A41)	1,397	1.7	10.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,143	1.4	8.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,076	1.3	8.4
11	Influenza and pneumonia (J09-J18)	912	1.1	7.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	450	0.5	3.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	418	0.5	3.2
14	Viral hepatitis (B15-B19)	395	0.5	3.1
15	Alzheimer's disease (G30)	344	0.4	2.7
...	All other causes (Residual)	12,136	14.5	94.2
Non-Hispanic, white, female, 65-74 years				
...	All causes	133,475	100.0	1,634.9
1	Malignant neoplasms (C00-C97)	50,800	38.1	622.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,150	18.8	308.1
3	Chronic lower respiratory diseases (J40-J47)	12,722	9.5	155.8
4	Cerebrovascular diseases (I60-I69)	6,193	4.6	75.9
5	Diabetes mellitus (E10-E14)	4,288	3.2	52.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,550	1.9	31.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,336	1.8	28.6
8	Septicemia (A40-A41)	2,183	1.6	26.7
9	Alzheimer's disease (G30)	1,944	1.5	23.8
10	Influenza and pneumonia (J09-J18)	1,901	1.4	23.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,495	1.1	18.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	952	0.7	11.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	856	0.6	10.5
14	Aortic aneurysm and dissection (I71)	746	0.6	9.1
15	Parkinson's disease (G20-G21)	695	0.5	8.5
...	All other causes (Residual)	18,664	14.0	228.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-84 years				
...	All causes	277,354	100.0	4,385.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,367	24.6	1,081.0
2	Malignant neoplasms (C00-C97)	65,779	23.7	1,040.1
3	Chronic lower respiratory diseases (J40-J47)	22,640	8.2	358.0
4	Cerebrovascular diseases (I60-I69)	19,885	7.2	314.4
5	Alzheimer's disease (G30)	12,958	4.7	204.9
6	Diabetes mellitus (E10-E14)	7,942	2.9	125.6
7	Influenza and pneumonia (J09-J18)	6,176	2.2	97.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,637	2.0	89.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,492	2.0	86.8
10	Septicemia (A40-A41)	4,365	1.6	69.0
11	Parkinson's disease (G20-G21)	3,267	1.2	51.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,754	1.0	43.5
13	Pneumonitis due to solids and liquids (J69)	1,955	0.7	30.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,821	0.7	28.8
15	Aortic aneurysm and dissection (I71)	1,747	0.6	27.6
...	All other causes (Residual)	46,569	16.8	736.3
Non-Hispanic, white, female, 85 years and over				
...	All causes	408,223	100.0	12,856.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	134,334	32.9	4,230.8
2	Malignant neoplasms (C00-C97)	41,761	10.2	1,315.2
3	Cerebrovascular diseases (I60-I69)	35,045	8.6	1,103.7
4	Alzheimer's disease (G30)	31,781	7.8	1,000.9
5	Chronic lower respiratory diseases (J40-J47)	17,842	4.4	561.9
6	Influenza and pneumonia (J09-J18)	13,872	3.4	436.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,841	2.2	278.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,690	1.9	242.2
9	Diabetes mellitus (E10-E14)	7,541	1.8	237.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,362	1.6	200.4
11	Septicemia (A40-A41)	5,135	1.3	161.7
12	Pneumonitis due to solids and liquids (J69)	4,016	1.0	126.5
13	Parkinson's disease (G20-G21)	3,539	0.9	111.5
14	Atherosclerosis (I70)	2,889	0.7	91.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,256	0.6	71.1
...	All other causes (Residual)	85,319	20.9	2,687.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	286,366	100.0	750.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,443	24.6	184.7
2	Malignant neoplasms (C00-C97)	63,441	22.2	166.3
3	Cerebrovascular diseases (I60-I69)	16,934	5.9	44.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,332	4.7	34.9
5	Diabetes mellitus (E10-E14)	12,343	4.3	32.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,746	3.1	22.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,318	2.9	21.8
8	Chronic lower respiratory diseases (J40-J47)	7,830	2.7	20.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,398	2.2	16.8
10	Septicemia (A40-A41)	6,241	2.2	16.4
11	Influenza and pneumonia (J09-J18)	5,091	1.8	13.3
12	Certain conditions originating in the perinatal period (P00-P96)	4,868	1.7	12.8
13	Alzheimer's disease (G30)	4,729	1.7	12.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,597	1.6	12.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,525	0.9	6.6
...	All other causes (Residual)	50,530	17.6	132.5
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,102	100.0	43.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	318	28.9	12.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	14.5	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	10.5	4.6
4	Malignant neoplasms (C00-C97)	51	4.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.8	1.7
6	Chronic lower respiratory diseases (J40-J47)	29	2.6	1.2
7	Influenza and pneumonia (J09-J18)	28	2.5	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	26	2.4	1.0
9	Septicemia (A40-A41)	16	1.5	*
9	Cerebrovascular diseases (I60-I69)	16	1.5	*
11	Anemias (D50-D64)	12	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
...	All other causes (Residual)	262	23.8	10.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,335	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	452	33.9	7.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	10.3	2.2
3	Malignant neoplasms (C00-C97)	134	10.0	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	5.8	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	4.7	1.0
6	Chronic lower respiratory diseases (J40-J47)	59	4.4	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	3.1	0.7
8	Influenza and pneumonia (J09-J18)	21	1.6	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.5	0.3
10	Anemias (D50-D64)	13	1.0	*
10	Cerebrovascular diseases (I60-I69)	13	1.0	*
12	Septicemia (A40-A41)	10	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.6	*
14	Diabetes mellitus (E10-E14)	6	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
...	All other causes (Residual)	268	20.1	4.3
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	7,237	100.0	112.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,162	43.7	49.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,699	23.5	26.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	396	5.5	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	344	4.8	5.4
5	Malignant neoplasms (C00-C97)	258	3.6	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	109	1.5	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	1.1	1.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	68	0.9	1.1
9	Anemias (D50-D64)	66	0.9	1.0
10	Chronic lower respiratory diseases (J40-J47)	65	0.9	1.0
11	Diabetes mellitus (E10-E14)	55	0.8	0.9
12	Septicemia (A40-A41)	47	0.6	0.7
12	Legal intervention (Y35,Y89.0)	47	0.6	0.7
14	Cerebrovascular diseases (I60-I69)	45	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.4	0.4
...	All other causes (Residual)	767	10.6	11.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-34 years				
...	All causes	9,521	100.0	176.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,524	26.5	46.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,748	18.4	32.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	985	10.3	18.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	701	7.4	13.0
5	Malignant neoplasms (C00-C97)	612	6.4	11.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	471	4.9	8.7
7	Diabetes mellitus (E10-E14)	212	2.2	3.9
8	Cerebrovascular diseases (I60-I69)	148	1.6	2.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	1.2	2.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	106	1.1	2.0
11	Anemias (D50-D64)	101	1.1	1.9
12	Chronic lower respiratory diseases (J40-J47)	93	1.0	1.7
13	Influenza and pneumonia (J09-J18)	92	1.0	1.7
14	Septicemia (A40-A41)	89	0.9	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	0.9	1.5
...	All other causes (Residual)	1,447	15.2	26.8
Non-Hispanic, black, both sexes, 35-44 years				
...	All causes	16,797	100.0	310.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,962	17.6	54.8
2	Malignant neoplasms (C00-C97)	2,332	13.9	43.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,118	12.6	39.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,990	11.8	36.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,355	8.1	25.1
6	Cerebrovascular diseases (I60-I69)	648	3.9	12.0
7	Diabetes mellitus (E10-E14)	554	3.3	10.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	415	2.5	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	329	2.0	6.1
10	Septicemia (A40-A41)	273	1.6	5.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	200	1.2	3.7
12	Chronic lower respiratory diseases (J40-J47)	193	1.1	3.6
13	Influenza and pneumonia (J09-J18)	175	1.0	3.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	174	1.0	3.2
15	Anemias (D50-D64)	139	0.8	2.6
...	All other causes (Residual)	2,940	17.5	54.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-54 years				
...	All causes	36,465	100.0	713.2
1	Malignant neoplasms (C00-C97)	8,754	24.0	171.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,285	22.7	162.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,724	7.5	53.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,318	6.4	45.3
5	Cerebrovascular diseases (I60-I69)	1,926	5.3	37.7
6	Diabetes mellitus (E10-E14)	1,412	3.9	27.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	843	2.3	16.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	815	2.2	15.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	796	2.2	15.6
10	Septicemia (A40-A41)	715	2.0	14.0
11	Chronic lower respiratory diseases (J40-J47)	669	1.8	13.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	531	1.5	10.4
13	Influenza and pneumonia (J09-J18)	434	1.2	8.5
14	Viral hepatitis (B15-B19)	375	1.0	7.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	317	0.9	6.2
...	All other causes (Residual)	5,551	15.2	108.6
Non-Hispanic, black, both sexes, 55-64 years				
...	All causes	47,199	100.0	1,446.2
1	Malignant neoplasms (C00-C97)	14,622	31.0	448.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,156	25.8	372.5
3	Cerebrovascular diseases (I60-I69)	2,572	5.4	78.8
4	Diabetes mellitus (E10-E14)	2,460	5.2	75.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,584	3.4	48.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,377	2.9	42.2
7	Chronic lower respiratory diseases (J40-J47)	1,166	2.5	35.7
8	Septicemia (A40-A41)	996	2.1	30.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	980	2.1	30.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	857	1.8	26.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	745	1.6	22.8
12	Influenza and pneumonia (J09-J18)	567	1.2	17.4
13	Viral hepatitis (B15-B19)	399	0.8	12.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	316	0.7	9.7
15	Aortic aneurysm and dissection (I71)	227	0.5	7.0
...	All other causes (Residual)	6,175	13.1	189.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-74 years				
...	All causes	50,890	100.0	2,847.8
1	Malignant neoplasms (C00-C97)	15,887	31.2	889.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,507	26.5	755.8
3	Cerebrovascular diseases (I60-I69)	3,224	6.3	180.4
4	Diabetes mellitus (E10-E14)	2,843	5.6	159.1
5	Chronic lower respiratory diseases (J40-J47)	1,886	3.7	105.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,824	3.6	102.1
7	Septicemia (A40-A41)	1,314	2.6	73.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	900	1.8	50.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	822	1.6	46.0
10	Influenza and pneumonia (J09-J18)	791	1.6	44.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	399	0.8	22.3
12	Alzheimer's disease (G30)	357	0.7	20.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	249	0.5	13.9
14	Aortic aneurysm and dissection (I71)	236	0.5	13.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	226	0.4	12.6
...	All other causes (Residual)	6,425	12.6	359.5
Non-Hispanic, black, both sexes, 75-84 years				
...	All causes	58,398	100.0	5,822.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,359	28.0	1,631.2
2	Malignant neoplasms (C00-C97)	14,142	24.2	1,410.1
3	Cerebrovascular diseases (I60-I69)	4,346	7.4	433.3
4	Diabetes mellitus (E10-E14)	3,031	5.2	302.2
5	Chronic lower respiratory diseases (J40-J47)	2,315	4.0	230.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,046	3.5	204.0
7	Alzheimer's disease (G30)	1,620	2.8	161.5
8	Septicemia (A40-A41)	1,500	2.6	149.6
9	Influenza and pneumonia (J09-J18)	1,298	2.2	129.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,085	1.9	108.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	756	1.3	75.4
12	Pneumonitis due to solids and liquids (J69)	436	0.7	43.5
13	Parkinson's disease (G20-G21)	358	0.6	35.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	306	0.5	30.5
15	Aortic aneurysm and dissection (I71)	267	0.5	26.6
...	All other causes (Residual)	8,533	14.6	850.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85 years and over				
...	All causes	48,762	100.0	12,468.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,601	32.0	3,989.1
2	Malignant neoplasms (C00-C97)	6,636	13.6	1,696.8
3	Cerebrovascular diseases (I60-I69)	3,961	8.1	1,012.8
4	Alzheimer's disease (G30)	2,703	5.5	691.1
5	Diabetes mellitus (E10-E14)	1,767	3.6	451.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,736	3.6	443.9
7	Influenza and pneumonia (J09-J18)	1,588	3.3	406.0
8	Chronic lower respiratory diseases (J40-J47)	1,341	2.8	342.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,180	2.4	301.7
10	Septicemia (A40-A41)	1,171	2.4	299.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	611	1.3	156.2
12	Pneumonitis due to solids and liquids (J69)	499	1.0	127.6
13	Atherosclerosis (I70)	277	0.6	70.8
14	Parkinson's disease (G20-G21)	259	0.5	66.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	198	0.4	50.6
...	All other causes (Residual)	9,234	18.9	2,361.1
Non-Hispanic, black, male, all ages				
...	All causes	146,474	100.0	805.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,243	24.1	193.7
2	Malignant neoplasms (C00-C97)	32,766	22.4	180.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,093	6.2	50.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,477	5.1	41.1
5	Cerebrovascular diseases (I60-I69)	7,470	5.1	41.1
6	Diabetes mellitus (E10-E14)	5,434	3.7	29.9
7	Chronic lower respiratory diseases (J40-J47)	4,167	2.8	22.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,127	2.8	22.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,730	2.5	20.5
10	Septicemia (A40-A41)	2,807	1.9	15.4
11	Certain conditions originating in the perinatal period (P00-P96)	2,734	1.9	15.0
12	Influenza and pneumonia (J09-J18)	2,464	1.7	13.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,963	1.3	10.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,662	1.1	9.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,571	1.1	8.6
...	All other causes (Residual)	23,766	16.2	130.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	609	100.0	47.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	30.5	14.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	14.4	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	9.4	4.4
4	Malignant neoplasms (C00-C97)	31	5.1	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.6	2.2
6	Influenza and pneumonia (J09-J18)	15	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	13	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	2.0	*
9	Septicemia (A40-A41)	11	1.8	*
10	Anemias (D50-D64)	4	0.7	*
10	Cerebrovascular diseases (I60-I69)	4	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
13	Meningitis (G00,G03)	3	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	144	23.6	11.2
Non-Hispanic, black, male, 5-14 years				
...	All causes	801	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	34.3	8.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	11.7	3.0
3	Malignant neoplasms (C00-C97)	70	8.7	2.2
4	Chronic lower respiratory diseases (J40-J47)	40	5.0	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	4.6	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.4	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	3.6	0.9
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.7	*
9	Influenza and pneumonia (J09-J18)	13	1.6	*
10	Anemias (D50-D64)	8	1.0	*
11	Septicemia (A40-A41)	4	0.5	*
11	Meningitis (G00,G03)	4	0.5	*
13	Cerebrovascular diseases (I60-I69)	3	0.4	*
13	Diseases of appendix (K35-K38)	3	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (Residual)	169	21.1	5.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,635	100.0	173.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,871	50.9	88.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,267	22.5	38.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	343	6.1	10.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	233	4.1	7.2
5	Malignant neoplasms (C00-C97)	142	2.5	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	56	1.0	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	0.9	1.6
8	Legal intervention (Y35,Y89.0)	45	0.8	1.4
9	Chronic lower respiratory diseases (J40-J47)	44	0.8	1.4
10	Anemias (D50-D64)	36	0.6	1.1
11	Cerebrovascular diseases (I60-I69)	30	0.5	0.9
12	Septicemia (A40-A41)	22	0.4	0.7
13	Diabetes mellitus (E10-E14)	21	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.2	*
15	Influenza and pneumonia (J09-J18)	12	0.2	*
...	All other causes (Residual)	447	7.9	13.7
Non-Hispanic, black, male, 25-34 years				
...	All causes	6,538	100.0	250.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,249	34.4	86.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,342	20.5	51.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	623	9.5	23.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	399	6.1	15.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	372	5.7	14.3
6	Malignant neoplasms (C00-C97)	260	4.0	10.0
7	Diabetes mellitus (E10-E14)	116	1.8	4.4
8	Cerebrovascular diseases (I60-I69)	71	1.1	2.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	0.9	2.3
10	Chronic lower respiratory diseases (J40-J47)	56	0.9	2.1
11	Anemias (D50-D64)	55	0.8	2.1
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	0.8	2.0
13	Influenza and pneumonia (J09-J18)	51	0.8	2.0
14	Legal intervention (Y35,Y89.0)	41	0.6	1.6
15	Septicemia (A40-A41)	40	0.6	1.5
...	All other causes (Residual)	751	11.5	28.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-44 years				
...	All causes	9,981	100.0	393.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,929	19.3	76.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,531	15.3	60.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,217	12.2	48.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,090	10.9	43.0
5	Malignant neoplasms (C00-C97)	895	9.0	35.3
6	Cerebrovascular diseases (I60-I69)	330	3.3	13.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	329	3.3	13.0
8	Diabetes mellitus (E10-E14)	309	3.1	12.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	176	1.8	6.9
10	Septicemia (A40-A41)	126	1.3	5.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	121	1.2	4.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	100	1.0	3.9
12	Influenza and pneumonia (J09-J18)	100	1.0	3.9
14	Chronic lower respiratory diseases (J40-J47)	99	1.0	3.9
15	Aortic aneurysm and dissection (I71)	86	0.9	3.4
...	All other causes (Residual)	1,543	15.5	60.8
Non-Hispanic, black, male, 45-54 years				
...	All causes	21,260	100.0	900.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,251	24.7	222.4
2	Malignant neoplasms (C00-C97)	4,334	20.4	183.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,919	9.0	81.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,533	7.2	64.9
5	Cerebrovascular diseases (I60-I69)	994	4.7	42.1
6	Diabetes mellitus (E10-E14)	784	3.7	33.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	617	2.9	26.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	552	2.6	23.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	430	2.0	18.2
10	Septicemia (A40-A41)	376	1.8	15.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	305	1.4	12.9
12	Chronic lower respiratory diseases (J40-J47)	303	1.4	12.8
13	Influenza and pneumonia (J09-J18)	267	1.3	11.3
14	Viral hepatitis (B15-B19)	249	1.2	10.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	242	1.1	10.2
...	All other causes (Residual)	3,104	14.6	131.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-64 years				
...	All causes	27,823	100.0	1,913.7
1	Malignant neoplasms (C00-C97)	8,113	29.2	558.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,673	27.6	527.7
3	Cerebrovascular diseases (I60-I69)	1,486	5.3	102.2
4	Diabetes mellitus (E10-E14)	1,276	4.6	87.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,138	4.1	78.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	731	2.6	50.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	723	2.6	49.7
8	Chronic lower respiratory diseases (J40-J47)	663	2.4	45.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	598	2.1	41.1
10	Septicemia (A40-A41)	524	1.9	36.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	402	1.4	27.6
12	Influenza and pneumonia (J09-J18)	328	1.2	22.6
13	Viral hepatitis (B15-B19)	287	1.0	19.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	266	1.0	18.3
15	Aortic aneurysm and dissection (I71)	144	0.5	9.9
...	All other causes (Residual)	3,471	12.5	238.7
Non-Hispanic, black, male, 65-74 years				
...	All causes	27,377	100.0	3,678.3
1	Malignant neoplasms (C00-C97)	8,650	31.6	1,162.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,505	27.4	1,008.4
3	Cerebrovascular diseases (I60-I69)	1,692	6.2	227.3
4	Diabetes mellitus (E10-E14)	1,308	4.8	175.7
5	Chronic lower respiratory diseases (J40-J47)	1,055	3.9	141.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	865	3.2	116.2
7	Septicemia (A40-A41)	658	2.4	88.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	594	2.2	79.8
9	Influenza and pneumonia (J09-J18)	451	1.6	60.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	408	1.5	54.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	271	1.0	36.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	170	0.6	22.8
13	Alzheimer's disease (G30)	150	0.5	20.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	135	0.5	18.1
15	Pneumonitis due to solids and liquids (J69)	129	0.5	17.3
...	All other causes (Residual)	3,336	12.2	448.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-84 years				
...	All causes	26,523	100.0	7,295.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,379	27.8	2,029.6
2	Malignant neoplasms (C00-C97)	7,178	27.1	1,974.3
3	Cerebrovascular diseases (I60-I69)	1,823	6.9	501.4
4	Chronic lower respiratory diseases (J40-J47)	1,283	4.8	352.9
5	Diabetes mellitus (E10-E14)	1,153	4.3	317.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	857	3.2	235.7
7	Influenza and pneumonia (J09-J18)	634	2.4	174.4
8	Septicemia (A40-A41)	600	2.3	165.0
9	Alzheimer's disease (G30)	537	2.0	147.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	431	1.6	118.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	392	1.5	107.8
12	Pneumonitis due to solids and liquids (J69)	242	0.9	66.6
13	Parkinson's disease (G20-G21)	199	0.8	54.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	140	0.5	38.5
15	Aortic aneurysm and dissection (I71)	116	0.4	31.9
...	All other causes (Residual)	3,559	13.4	978.9
Non-Hispanic, black, male, 85 years and over				
...	All causes	15,115	100.0	13,189.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,506	29.8	3,931.8
2	Malignant neoplasms (C00-C97)	3,085	20.4	2,691.9
3	Cerebrovascular diseases (I60-I69)	1,012	6.7	883.0
4	Chronic lower respiratory diseases (J40-J47)	599	4.0	522.7
5	Alzheimer's disease (G30)	585	3.9	510.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	574	3.8	500.9
7	Influenza and pneumonia (J09-J18)	551	3.6	480.8
8	Diabetes mellitus (E10-E14)	463	3.1	404.0
9	Septicemia (A40-A41)	386	2.6	336.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	288	1.9	251.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	1.5	200.7
12	Pneumonitis due to solids and liquids (J69)	187	1.2	163.2
13	Parkinson's disease (G20-G21)	115	0.8	100.3
14	Atherosclerosis (I70)	74	0.5	64.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.5	63.7
...	All other causes (Residual)	2,387	15.8	2,082.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	139,892	100.0	701.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,200	25.2	176.4
2	Malignant neoplasms (C00-C97)	30,675	21.9	153.7
3	Cerebrovascular diseases (I60-I69)	9,464	6.8	47.4
4	Diabetes mellitus (E10-E14)	6,909	4.9	34.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,588	3.3	23.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,239	3.0	21.2
7	Chronic lower respiratory diseases (J40-J47)	3,663	2.6	18.4
8	Alzheimer's disease (G30)	3,436	2.5	17.2
9	Septicemia (A40-A41)	3,434	2.5	17.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,634	1.9	13.2
11	Influenza and pneumonia (J09-J18)	2,627	1.9	13.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,271	1.6	11.4
13	Certain conditions originating in the perinatal period (P00-P96)	2,134	1.5	10.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,269	0.9	6.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	863	0.6	4.3
...	All other causes (Residual)	26,486	18.9	132.7
Non-Hispanic, black, female, 1-4 years				
...	All causes	493	100.0	39.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	26.8	10.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	14.6	5.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	12.0	4.8
4	Malignant neoplasms (C00-C97)	20	4.1	1.6
5	Chronic lower respiratory diseases (J40-J47)	16	3.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	14	2.8	*
8	Influenza and pneumonia (J09-J18)	13	2.6	*
9	Cerebrovascular diseases (I60-I69)	12	2.4	*
10	Anemias (D50-D64)	8	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
12	Septicemia (A40-A41)	5	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	115	23.3	9.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	534	100.0	17.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	33.1	5.8
2	Malignant neoplasms (C00-C97)	64	12.0	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	8.2	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	7.5	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.2	0.9
6	Chronic lower respiratory diseases (J40-J47)	19	3.6	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.2	*
8	Cerebrovascular diseases (I60-I69)	10	1.9	*
9	Influenza and pneumonia (J09-J18)	8	1.5	*
10	Septicemia (A40-A41)	6	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
13	Anemias (D50-D64)	5	0.9	*
14	Diabetes mellitus (E10-E14)	4	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
...	All other causes (Residual)	101	18.9	3.3
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,602	100.0	50.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	432	27.0	13.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	291	18.2	9.2
3	Malignant neoplasms (C00-C97)	116	7.2	3.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	6.9	3.5
5	Pregnancy, childbirth and the puerperium (O00-O99)	68	4.2	2.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	53	3.3	1.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	3.3	1.7
8	Diabetes mellitus (E10-E14)	34	2.1	1.1
9	Anemias (D50-D64)	30	1.9	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	1.8	0.9
11	Septicemia (A40-A41)	25	1.6	0.8
12	Chronic lower respiratory diseases (J40-J47)	21	1.3	0.7
13	Cerebrovascular diseases (I60-I69)	15	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.9	*
15	Influenza and pneumonia (J09-J18)	11	0.7	*
...	All other causes (Residual)	299	18.7	9.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-34 years				
...	All causes	2,983	100.0	106.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	406	13.6	14.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	362	12.1	13.0
3	Malignant neoplasms (C00-C97)	352	11.8	12.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	329	11.0	11.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	275	9.2	9.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	106	3.6	3.8
7	Diabetes mellitus (E10-E14)	96	3.2	3.4
8	Cerebrovascular diseases (I60-I69)	77	2.6	2.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	2.4	2.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	1.7	1.8
11	Septicemia (A40-A41)	49	1.6	1.8
12	Anemias (D50-D64)	46	1.5	1.6
13	Influenza and pneumonia (J09-J18)	41	1.4	1.5
14	Chronic lower respiratory diseases (J40-J47)	37	1.2	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	1.0	1.0
...	All other causes (Residual)	655	22.0	23.4
Non-Hispanic, black, female, 35-44 years				
...	All causes	6,816	100.0	237.9
1	Malignant neoplasms (C00-C97)	1,437	21.1	50.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,033	15.2	36.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	773	11.3	27.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	587	8.6	20.5
5	Cerebrovascular diseases (I60-I69)	318	4.7	11.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	265	3.9	9.3
7	Diabetes mellitus (E10-E14)	245	3.6	8.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	2.2	5.3
9	Septicemia (A40-A41)	147	2.2	5.1
10	Chronic lower respiratory diseases (J40-J47)	94	1.4	3.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	1.3	3.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	79	1.2	2.8
13	Influenza and pneumonia (J09-J18)	75	1.1	2.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	74	1.1	2.6
15	Anemias (D50-D64)	64	0.9	2.2
...	All other causes (Residual)	1,386	20.3	48.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-54 years				
...	All causes	15,205	100.0	552.5
1	Malignant neoplasms (C00-C97)	4,420	29.1	160.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,034	20.0	110.2
3	Cerebrovascular diseases (I60-I69)	932	6.1	33.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	805	5.3	29.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	785	5.2	28.5
6	Diabetes mellitus (E10-E14)	628	4.1	22.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	385	2.5	14.0
8	Chronic lower respiratory diseases (J40-J47)	366	2.4	13.3
9	Septicemia (A40-A41)	339	2.2	12.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	291	1.9	10.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	226	1.5	8.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	179	1.2	6.5
13	Influenza and pneumonia (J09-J18)	167	1.1	6.1
14	Viral hepatitis (B15-B19)	126	0.8	4.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	0.5	2.7
...	All other causes (Residual)	2,447	16.1	88.9
Non-Hispanic, black, female, 55-64 years				
...	All causes	19,376	100.0	1,070.7
1	Malignant neoplasms (C00-C97)	6,509	33.6	359.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,483	23.1	247.7
3	Diabetes mellitus (E10-E14)	1,184	6.1	65.4
4	Cerebrovascular diseases (I60-I69)	1,086	5.6	60.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	654	3.4	36.1
6	Chronic lower respiratory diseases (J40-J47)	503	2.6	27.8
7	Septicemia (A40-A41)	472	2.4	26.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	446	2.3	24.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	343	1.8	19.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	259	1.3	14.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	249	1.3	13.8
12	Influenza and pneumonia (J09-J18)	239	1.2	13.2
13	Viral hepatitis (B15-B19)	112	0.6	6.2
14	Aortic aneurysm and dissection (I71)	83	0.4	4.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.4	3.8
...	All other causes (Residual)	2,685	13.9	148.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-74 years				
...	All causes	23,513	100.0	2,255.0
1	Malignant neoplasms (C00-C97)	7,237	30.8	694.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,002	25.5	575.6
3	Diabetes mellitus (E10-E14)	1,535	6.5	147.2
4	Cerebrovascular diseases (I60-I69)	1,532	6.5	146.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	959	4.1	92.0
6	Chronic lower respiratory diseases (J40-J47)	831	3.5	79.7
7	Septicemia (A40-A41)	656	2.8	62.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	414	1.8	39.7
9	Influenza and pneumonia (J09-J18)	340	1.4	32.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	306	1.3	29.3
11	Alzheimer's disease (G30)	207	0.9	19.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	128	0.5	12.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	114	0.5	10.9
14	Aortic aneurysm and dissection (I71)	109	0.5	10.5
15	Pneumonitis due to solids and liquids (J69)	76	0.3	7.3
...	All other causes (Residual)	3,067	13.0	294.1
Non-Hispanic, black, female, 75-84 years				
...	All causes	31,875	100.0	4,985.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,980	28.2	1,404.6
2	Malignant neoplasms (C00-C97)	6,964	21.8	1,089.3
3	Cerebrovascular diseases (I60-I69)	2,523	7.9	394.6
4	Diabetes mellitus (E10-E14)	1,878	5.9	293.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,189	3.7	186.0
6	Alzheimer's disease (G30)	1,083	3.4	169.4
7	Chronic lower respiratory diseases (J40-J47)	1,032	3.2	161.4
8	Septicemia (A40-A41)	900	2.8	140.8
9	Influenza and pneumonia (J09-J18)	664	2.1	103.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	654	2.1	102.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	364	1.1	56.9
12	Pneumonitis due to solids and liquids (J69)	194	0.6	30.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	166	0.5	26.0
14	Parkinson's disease (G20-G21)	159	0.5	24.9
15	Aortic aneurysm and dissection (I71)	151	0.5	23.6
...	All other causes (Residual)	4,974	15.6	778.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85 years and over				
...	All causes	33,647	100.0	12,169.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,095	33.0	4,012.8
2	Malignant neoplasms (C00-C97)	3,551	10.6	1,284.3
3	Cerebrovascular diseases (I60-I69)	2,949	8.8	1,066.6
4	Alzheimer's disease (G30)	2,118	6.3	766.0
5	Diabetes mellitus (E10-E14)	1,304	3.9	471.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,162	3.5	420.3
7	Influenza and pneumonia (J09-J18)	1,037	3.1	375.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	892	2.7	322.6
9	Septicemia (A40-A41)	785	2.3	283.9
10	Chronic lower respiratory diseases (J40-J47)	742	2.2	268.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	381	1.1	137.8
12	Pneumonitis due to solids and liquids (J69)	312	0.9	112.8
13	Atherosclerosis (I70)	203	0.6	73.4
14	Parkinson's disease (G20-G21)	144	0.4	52.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	125	0.4	45.2
...	All other causes (Residual)	6,847	20.3	2,476.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable