

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	144,490	100.0	286.2
1	Malignant neoplasms (C00-C97)	31,119	21.5	61.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,006	20.8	59.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,476	7.3	20.8
4	Cerebrovascular diseases (I60-I69)	7,274	5.0	14.4
5	Diabetes mellitus (E10-E14)	6,556	4.5	13.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,348	3.0	8.6
7	Chronic lower respiratory diseases (J40-J47)	4,172	2.9	8.3
8	Alzheimer's disease (G30)	3,427	2.4	6.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,252	2.3	6.4
10	Influenza and pneumonia (J09-J18)	3,025	2.1	6.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,890	2.0	5.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,661	1.8	5.3
13	Certain conditions originating in the perinatal period (P00-P96)	2,529	1.8	5.0
14	Septicemia (A40-A41)	2,035	1.4	4.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,786	1.2	3.5
...	All other causes (Residual)	28,934	20.0	57.3
Hispanic, both sexes, 1-4 years				
...	All causes	930	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	35.5	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	13.1	3.0
3	Malignant neoplasms (C00-C97)	75	8.1	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	6.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.0	0.9
6	Influenza and pneumonia (J09-J18)	21	2.3	0.5
7	Septicemia (A40-A41)	18	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
9	Cerebrovascular diseases (I60-I69)	8	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.8	*
12	Meningococcal infection (A39)	5	0.5	*
12	Pneumonitis due to solids and liquids (J69)	5	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	216	23.2	5.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	951	100.0	10.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	252	26.5	2.7
2	Malignant neoplasms (C00-C97)	212	22.3	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	5.7	0.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	5.3	0.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	4.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	2.7	0.3
7	Cerebrovascular diseases (I60-I69)	23	2.4	0.2
8	Chronic lower respiratory diseases (J40-J47)	20	2.1	0.2
9	Septicemia (A40-A41)	14	1.5	*
10	Influenza and pneumonia (J09-J18)	13	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
13	Anemias (D50-D64)	3	0.3	*
13	Meningitis (G00,G03)	3	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Pneumonitis due to solids and liquids (J69)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	215	22.6	2.3
Hispanic, both sexes, 15-24 years				
...	All causes	4,795	100.0	54.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,891	39.4	21.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,022	21.3	11.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	628	13.1	7.1
4	Malignant neoplasms (C00-C97)	344	7.2	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	2.6	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	1.2	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	32	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	29	0.6	0.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.5	0.3
10	Influenza and pneumonia (J09-J18)	24	0.5	0.3
11	Septicemia (A40-A41)	21	0.4	0.2
12	Diabetes mellitus (E10-E14)	18	0.4	*
13	Legal intervention (Y35,Y89.0)	16	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	15	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
...	All other causes (Residual)	531	11.1	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	6,022	100.0	71.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,969	32.7	23.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	846	14.0	10.0
3	Malignant neoplasms (C00-C97)	652	10.8	7.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	588	9.8	7.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	372	6.2	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	132	2.2	1.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	112	1.9	1.3
8	Cerebrovascular diseases (I60-I69)	91	1.5	1.1
9	Pregnancy, childbirth and the puerperium (O00-O99)	71	1.2	0.8
10	Diabetes mellitus (E10-E14)	63	1.0	0.7
11	Influenza and pneumonia (J09-J18)	59	1.0	0.7
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	0.7	0.5
13	Septicemia (A40-A41)	44	0.7	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	0.6	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.6	0.4
...	All other causes (Residual)	904	15.0	10.7
Hispanic, both sexes, 35-44 years				
...	All causes	8,142	100.0	111.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,719	21.1	23.6
2	Malignant neoplasms (C00-C97)	1,454	17.9	19.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	919	11.3	12.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	505	6.2	6.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	458	5.6	6.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	415	5.1	5.7
7	Cerebrovascular diseases (I60-I69)	304	3.7	4.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	296	3.6	4.1
9	Diabetes mellitus (E10-E14)	242	3.0	3.3
10	Influenza and pneumonia (J09-J18)	97	1.2	1.3
11	Viral hepatitis (B15-B19)	84	1.0	1.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	1.0	1.1
13	Septicemia (A40-A41)	72	0.9	1.0
14	Chronic lower respiratory diseases (J40-J47)	55	0.7	0.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	47	0.6	0.6
...	All other causes (Residual)	1,396	17.1	19.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	14,915	100.0	273.0
1	Malignant neoplasms (C00-C97)	3,924	26.3	71.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,478	16.6	45.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,512	10.1	27.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,238	8.3	22.7
5	Cerebrovascular diseases (I60-I69)	641	4.3	11.7
6	Diabetes mellitus (E10-E14)	595	4.0	10.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	422	2.8	7.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	411	2.8	7.5
9	Viral hepatitis (B15-B19)	367	2.5	6.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	274	1.8	5.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	223	1.5	4.1
12	Septicemia (A40-A41)	200	1.3	3.7
13	Chronic lower respiratory diseases (J40-J47)	173	1.2	3.2
14	Influenza and pneumonia (J09-J18)	158	1.1	2.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	98	0.7	1.8
...	All other causes (Residual)	2,201	14.8	40.3
Hispanic, both sexes, 55-64 years				
...	All causes	20,066	100.0	624.4
1	Malignant neoplasms (C00-C97)	6,246	31.1	194.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,067	20.3	126.5
3	Diabetes mellitus (E10-E14)	1,239	6.2	38.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,195	6.0	37.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	952	4.7	29.6
6	Cerebrovascular diseases (I60-I69)	861	4.3	26.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	485	2.4	15.1
8	Viral hepatitis (B15-B19)	426	2.1	13.3
9	Chronic lower respiratory diseases (J40-J47)	393	2.0	12.2
10	Septicemia (A40-A41)	318	1.6	9.9
11	Influenza and pneumonia (J09-J18)	287	1.4	8.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	253	1.3	7.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	223	1.1	6.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	188	0.9	5.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	0.5	2.9
...	All other causes (Residual)	2,840	14.2	88.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	22,962	100.0	1,392.7
1	Malignant neoplasms (C00-C97)	7,302	31.8	442.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,059	22.0	306.8
3	Diabetes mellitus (E10-E14)	1,530	6.7	92.8
4	Cerebrovascular diseases (I60-I69)	1,174	5.1	71.2
5	Chronic lower respiratory diseases (J40-J47)	762	3.3	46.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	730	3.2	44.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	700	3.0	42.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	550	2.4	33.4
9	Influenza and pneumonia (J09-J18)	500	2.2	30.3
10	Septicemia (A40-A41)	381	1.7	23.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	279	1.2	16.9
12	Alzheimer's disease (G30)	237	1.0	14.4
13	Viral hepatitis (B15-B19)	146	0.6	8.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	124	0.5	7.5
14	Parkinson's disease (G20-G21)	124	0.5	7.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	124	0.5	7.5
...	All other causes (Residual)	3,240	14.1	196.5
Hispanic, both sexes, 75-84 years				
...	All causes	31,364	100.0	3,637.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,946	25.3	921.5
2	Malignant neoplasms (C00-C97)	7,295	23.3	846.0
3	Cerebrovascular diseases (I60-I69)	2,069	6.6	239.9
4	Diabetes mellitus (E10-E14)	1,747	5.6	202.6
5	Chronic lower respiratory diseases (J40-J47)	1,394	4.4	161.7
6	Alzheimer's disease (G30)	1,088	3.5	126.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	909	2.9	105.4
8	Influenza and pneumonia (J09-J18)	836	2.7	97.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	607	1.9	70.4
10	Septicemia (A40-A41)	538	1.7	62.4
11	Parkinson's disease (G20-G21)	468	1.5	54.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	467	1.5	54.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	466	1.5	54.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	222	0.7	25.7
15	Pneumonitis due to solids and liquids (J69)	201	0.6	23.3
...	All other causes (Residual)	5,111	16.3	592.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	29,166	100.0	10,777.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,913	30.6	3,293.7
2	Malignant neoplasms (C00-C97)	3,599	12.3	1,330.0
3	Cerebrovascular diseases (I60-I69)	2,049	7.0	757.2
4	Alzheimer's disease (G30)	2,045	7.0	755.7
5	Chronic lower respiratory diseases (J40-J47)	1,311	4.5	484.5
6	Diabetes mellitus (E10-E14)	1,119	3.8	413.5
7	Influenza and pneumonia (J09-J18)	996	3.4	368.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	769	2.6	284.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	574	2.0	212.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	540	1.9	199.5
11	Septicemia (A40-A41)	385	1.3	142.3
12	Parkinson's disease (G20-G21)	357	1.2	131.9
13	Pneumonitis due to solids and liquids (J69)	268	0.9	99.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	165	0.6	61.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	141	0.5	52.1
...	All other causes (Residual)	5,935	20.3	2,193.2
Hispanic, male, all ages				
...	All causes	79,622	100.0	310.8
1	Malignant neoplasms (C00-C97)	16,450	20.7	64.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,421	20.6	64.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,594	9.5	29.6
4	Cerebrovascular diseases (I60-I69)	3,382	4.2	13.2
5	Diabetes mellitus (E10-E14)	3,372	4.2	13.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,067	3.9	12.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,435	3.1	9.5
8	Chronic lower respiratory diseases (J40-J47)	2,174	2.7	8.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,168	2.7	8.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,670	2.1	6.5
11	Influenza and pneumonia (J09-J18)	1,565	2.0	6.1
12	Certain conditions originating in the perinatal period (P00-P96)	1,456	1.8	5.7
13	Alzheimer's disease (G30)	1,171	1.5	4.6
14	Septicemia (A40-A41)	1,012	1.3	4.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	910	1.1	3.6
...	All other causes (Residual)	14,775	18.6	57.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	524	100.0	25.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	39.9	10.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	11.3	2.8
3	Malignant neoplasms (C00-C97)	42	8.0	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	8.0	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.1	*
6	Influenza and pneumonia (J09-J18)	13	2.5	*
7	Septicemia (A40-A41)	12	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
10	Pneumonitis due to solids and liquids (J69)	3	0.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
12	Meningitis (G00,G03)	2	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
15	Meningococcal infection (A39)	1	0.2	*
15	Cerebrovascular diseases (I60-I69)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	104	19.8	5.0
Hispanic, male, 5-14 years				
...	All causes	543	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	27.6	3.2
2	Malignant neoplasms (C00-C97)	112	20.6	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	6.4	0.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	5.5	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	4.1	0.5
6	Chronic lower respiratory diseases (J40-J47)	18	3.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.6	*
8	Cerebrovascular diseases (I60-I69)	13	2.4	*
9	Septicemia (A40-A41)	9	1.7	*
9	Influenza and pneumonia (J09-J18)	9	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
13	Anemias (D50-D64)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Meningitis (G00,G03)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (Residual)	114	21.0	2.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,689	100.0	79.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,478	40.1	31.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	914	24.8	19.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	497	13.5	10.7
4	Malignant neoplasms (C00-C97)	225	6.1	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	2.5	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	0.9	0.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.5	0.4
8	Cerebrovascular diseases (I60-I69)	19	0.5	*
9	Influenza and pneumonia (J09-J18)	17	0.5	*
10	Legal intervention (Y35,Y89.0)	16	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.3	*
12	Septicemia (A40-A41)	9	0.2	*
12	Diabetes mellitus (E10-E14)	9	0.2	*
12	Aortic aneurysm and dissection (I71)	9	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	8	0.2	*
...	All other causes (Residual)	330	8.9	7.1
Hispanic, male, 25-34 years				
...	All causes	4,461	100.0	100.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,588	35.6	35.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	741	16.6	16.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	504	11.3	11.4
4	Malignant neoplasms (C00-C97)	316	7.1	7.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	296	6.6	6.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	107	2.4	2.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	93	2.1	2.1
8	Cerebrovascular diseases (I60-I69)	59	1.3	1.3
9	Diabetes mellitus (E10-E14)	41	0.9	0.9
10	Influenza and pneumonia (J09-J18)	36	0.8	0.8
11	Legal intervention (Y35,Y89.0)	27	0.6	0.6
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.6	0.6
13	Septicemia (A40-A41)	23	0.5	0.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	0.5
13	Aortic aneurysm and dissection (I71)	23	0.5	0.5
...	All other causes (Residual)	558	12.5	12.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,460	100.0	146.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,348	24.7	36.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	696	12.7	18.6
3	Malignant neoplasms (C00-C97)	618	11.3	16.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	407	7.5	10.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	366	6.7	9.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	333	6.1	8.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	230	4.2	6.2
8	Cerebrovascular diseases (I60-I69)	191	3.5	5.1
9	Diabetes mellitus (E10-E14)	145	2.7	3.9
10	Influenza and pneumonia (J09-J18)	67	1.2	1.8
11	Viral hepatitis (B15-B19)	63	1.2	1.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	0.9	1.3
13	Septicemia (A40-A41)	44	0.8	1.2
14	Chronic lower respiratory diseases (J40-J47)	32	0.6	0.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	0.5	0.7
...	All other causes (Residual)	845	15.5	22.6
Hispanic, male, 45-54 years				
...	All causes	9,627	100.0	351.9
1	Malignant neoplasms (C00-C97)	1,906	19.8	69.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,809	18.8	66.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,124	11.7	41.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	945	9.8	34.5
5	Cerebrovascular diseases (I60-I69)	391	4.1	14.3
6	Diabetes mellitus (E10-E14)	376	3.9	13.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	347	3.6	12.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	333	3.5	12.2
9	Viral hepatitis (B15-B19)	284	3.0	10.4
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	219	2.3	8.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	134	1.4	4.9
12	Septicemia (A40-A41)	121	1.3	4.4
13	Influenza and pneumonia (J09-J18)	92	1.0	3.4
14	Chronic lower respiratory diseases (J40-J47)	84	0.9	3.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	62	0.6	2.3
...	All other causes (Residual)	1,400	14.5	51.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	12,509	100.0	815.1
1	Malignant neoplasms (C00-C97)	3,460	27.7	225.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,853	22.8	185.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	908	7.3	59.2
4	Diabetes mellitus (E10-E14)	761	6.1	49.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	673	5.4	43.9
6	Cerebrovascular diseases (I60-I69)	490	3.9	31.9
7	Viral hepatitis (B15-B19)	290	2.3	18.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	283	2.3	18.4
9	Chronic lower respiratory diseases (J40-J47)	225	1.8	14.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	205	1.6	13.4
11	Septicemia (A40-A41)	180	1.4	11.7
12	Influenza and pneumonia (J09-J18)	159	1.3	10.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	156	1.2	10.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	142	1.1	9.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	76	0.6	5.0
...	All other causes (Residual)	1,648	13.2	107.4
Hispanic, male, 65-74 years				
...	All causes	13,040	100.0	1,775.0
1	Malignant neoplasms (C00-C97)	4,055	31.1	552.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,119	23.9	424.5
3	Diabetes mellitus (E10-E14)	825	6.3	112.3
4	Cerebrovascular diseases (I60-I69)	620	4.8	84.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	453	3.5	61.7
6	Chronic lower respiratory diseases (J40-J47)	424	3.3	57.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	357	2.7	48.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	356	2.7	48.5
9	Influenza and pneumonia (J09-J18)	288	2.2	39.2
10	Septicemia (A40-A41)	195	1.5	26.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	135	1.0	18.4
12	Alzheimer's disease (G30)	99	0.8	13.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	97	0.7	13.2
14	Viral hepatitis (B15-B19)	86	0.7	11.7
15	Parkinson's disease (G20-G21)	84	0.6	11.4
...	All other causes (Residual)	1,847	14.2	251.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	15,723	100.0	4,461.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,091	26.0	1,160.9
2	Malignant neoplasms (C00-C97)	3,942	25.1	1,118.7
3	Cerebrovascular diseases (I60-I69)	939	6.0	266.5
4	Diabetes mellitus (E10-E14)	826	5.3	234.4
5	Chronic lower respiratory diseases (J40-J47)	750	4.8	212.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	461	2.9	130.8
7	Influenza and pneumonia (J09-J18)	450	2.9	127.7
8	Alzheimer's disease (G30)	431	2.7	122.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	345	2.2	97.9
10	Parkinson's disease (G20-G21)	264	1.7	74.9
11	Septicemia (A40-A41)	251	1.6	71.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	240	1.5	68.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	199	1.3	56.5
14	Pneumonitis due to solids and liquids (J69)	114	0.7	32.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	108	0.7	30.6
...	All other causes (Residual)	2,312	14.7	656.1
Hispanic, male, 85 years and over				
...	All causes	11,171	100.0	11,779.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,393	30.4	3,577.9
2	Malignant neoplasms (C00-C97)	1,765	15.8	1,861.2
3	Cerebrovascular diseases (I60-I69)	644	5.8	679.1
4	Alzheimer's disease (G30)	618	5.5	651.7
5	Chronic lower respiratory diseases (J40-J47)	602	5.4	634.8
6	Influenza and pneumonia (J09-J18)	415	3.7	437.6
7	Diabetes mellitus (E10-E14)	389	3.5	410.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	344	3.1	362.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	2.1	246.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	185	1.7	195.1
11	Parkinson's disease (G20-G21)	184	1.6	194.0
12	Septicemia (A40-A41)	144	1.3	151.8
13	Pneumonitis due to solids and liquids (J69)	137	1.2	144.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.7	86.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	59	0.5	62.2
...	All other causes (Residual)	1,976	17.7	2,083.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	64,868	100.0	260.9
1	Malignant neoplasms (C00-C97)	14,669	22.6	59.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,585	20.9	54.6
3	Cerebrovascular diseases (I60-I69)	3,892	6.0	15.7
4	Diabetes mellitus (E10-E14)	3,184	4.9	12.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,882	4.4	11.6
6	Alzheimer's disease (G30)	2,256	3.5	9.1
7	Chronic lower respiratory diseases (J40-J47)	1,998	3.1	8.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,582	2.4	6.4
9	Influenza and pneumonia (J09-J18)	1,460	2.3	5.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,281	2.0	5.2
11	Certain conditions originating in the perinatal period (P00-P96)	1,073	1.7	4.3
12	Septicemia (A40-A41)	1,023	1.6	4.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	948	1.5	3.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	883	1.4	3.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	493	0.8	2.0
...	All other causes (Residual)	13,659	21.1	54.9
Hispanic, female, 1-4 years				
...	All causes	406	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	29.8	6.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	15.5	3.1
3	Malignant neoplasms (C00-C97)	33	8.1	1.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	5.2	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	3.7	*
6	Influenza and pneumonia (J09-J18)	8	2.0	*
7	Cerebrovascular diseases (I60-I69)	7	1.7	*
8	Septicemia (A40-A41)	6	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.2	*
11	Meningococcal infection (A39)	4	1.0	*
12	Anemias (D50-D64)	3	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.5	*
14	Pneumonitis due to solids and liquids (J69)	2	0.5	*
...	All other causes (Residual)	108	26.6	5.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	408	100.0	8.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	25.0	2.2
2	Malignant neoplasms (C00-C97)	100	24.5	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	5.4	0.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	4.9	0.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	4.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.9	*
7	Cerebrovascular diseases (I60-I69)	10	2.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.5	*
9	Septicemia (A40-A41)	5	1.2	*
10	Influenza and pneumonia (J09-J18)	4	1.0	*
11	Diabetes mellitus (E10-E14)	2	0.5	*
11	Meningitis (G00,G03)	2	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
11	Pneumonitis due to solids and liquids (J69)	2	0.5	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Meningococcal infection (A39)	1	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (Residual)	93	22.8	2.0
Hispanic, female, 15-24 years				
...	All causes	1,106	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	413	37.3	9.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	131	11.8	3.1
3	Malignant neoplasms (C00-C97)	119	10.8	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	108	9.8	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	2.9	0.8
5	Pregnancy, childbirth and the puerperium (O00-O99)	32	2.9	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	2.3	0.6
8	Septicemia (A40-A41)	12	1.1	*
9	Cerebrovascular diseases (I60-I69)	10	0.9	*
10	Diabetes mellitus (E10-E14)	9	0.8	*
11	Influenza and pneumonia (J09-J18)	7	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.4	*
...	All other causes (Residual)	181	16.4	4.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,561	100.0	38.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	381	24.4	9.5
2	Malignant neoplasms (C00-C97)	336	21.5	8.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	6.7	2.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	5.4	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	4.9	1.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	71	4.5	1.8
7	Cerebrovascular diseases (I60-I69)	32	2.0	0.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	25	1.6	0.6
9	Influenza and pneumonia (J09-J18)	23	1.5	0.6
10	Diabetes mellitus (E10-E14)	22	1.4	0.5
11	Septicemia (A40-A41)	21	1.3	0.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.3	0.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.2	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.8	*
...	All other causes (Residual)	315	20.2	7.8
Hispanic, female, 35-44 years				
...	All causes	2,682	100.0	75.2
1	Malignant neoplasms (C00-C97)	836	31.2	23.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	13.8	10.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	223	8.3	6.3
4	Cerebrovascular diseases (I60-I69)	113	4.2	3.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	98	3.7	2.7
6	Diabetes mellitus (E10-E14)	97	3.6	2.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	92	3.4	2.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	3.1	2.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	66	2.5	1.9
10	Pregnancy, childbirth and the puerperium (O00-O99)	37	1.4	1.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	1.2	0.9
12	Influenza and pneumonia (J09-J18)	30	1.1	0.8
13	Septicemia (A40-A41)	28	1.0	0.8
14	Chronic lower respiratory diseases (J40-J47)	23	0.9	0.6
15	Viral hepatitis (B15-B19)	21	0.8	0.6
...	All other causes (Residual)	533	19.9	15.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	5,288	100.0	193.9
1	Malignant neoplasms (C00-C97)	2,018	38.2	74.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	669	12.7	24.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	388	7.3	14.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	293	5.5	10.7
5	Cerebrovascular diseases (I60-I69)	250	4.7	9.2
6	Diabetes mellitus (E10-E14)	219	4.1	8.0
7	Chronic lower respiratory diseases (J40-J47)	89	1.7	3.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	89	1.7	3.3
9	Viral hepatitis (B15-B19)	83	1.6	3.0
10	Septicemia (A40-A41)	79	1.5	2.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	78	1.5	2.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	1.4	2.7
13	Influenza and pneumonia (J09-J18)	66	1.2	2.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	1.0	2.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.8	1.5
...	All other causes (Residual)	796	15.1	29.2
Hispanic, female, 55-64 years				
...	All causes	7,557	100.0	450.1
1	Malignant neoplasms (C00-C97)	2,786	36.9	165.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,214	16.1	72.3
3	Diabetes mellitus (E10-E14)	478	6.3	28.5
4	Cerebrovascular diseases (I60-I69)	371	4.9	22.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	287	3.8	17.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	3.7	16.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	2.7	12.0
8	Chronic lower respiratory diseases (J40-J47)	168	2.2	10.0
9	Septicemia (A40-A41)	138	1.8	8.2
10	Viral hepatitis (B15-B19)	136	1.8	8.1
11	Influenza and pneumonia (J09-J18)	128	1.7	7.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	1.1	4.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	0.6	2.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.5	2.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.4	1.9
...	All other causes (Residual)	1,169	15.5	69.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	9,922	100.0	1,085.5
1	Malignant neoplasms (C00-C97)	3,247	32.7	355.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,940	19.6	212.2
3	Diabetes mellitus (E10-E14)	705	7.1	77.1
4	Cerebrovascular diseases (I60-I69)	554	5.6	60.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	3.5	37.5
6	Chronic lower respiratory diseases (J40-J47)	338	3.4	37.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	277	2.8	30.3
8	Influenza and pneumonia (J09-J18)	212	2.1	23.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	2.0	21.2
10	Septicemia (A40-A41)	186	1.9	20.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	144	1.5	15.8
12	Alzheimer's disease (G30)	138	1.4	15.1
13	Viral hepatitis (B15-B19)	60	0.6	6.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.5	5.9
15	Parkinson's disease (G20-G21)	40	0.4	4.4
...	All other causes (Residual)	1,490	15.0	163.0
Hispanic, female, 75-84 years				
...	All causes	15,641	100.0	3,067.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,855	24.6	756.0
2	Malignant neoplasms (C00-C97)	3,353	21.4	657.6
3	Cerebrovascular diseases (I60-I69)	1,130	7.2	221.6
4	Diabetes mellitus (E10-E14)	921	5.9	180.6
5	Alzheimer's disease (G30)	657	4.2	128.8
6	Chronic lower respiratory diseases (J40-J47)	644	4.1	126.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	448	2.9	87.9
8	Influenza and pneumonia (J09-J18)	386	2.5	75.7
9	Septicemia (A40-A41)	287	1.8	56.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	268	1.7	52.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	262	1.7	51.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	226	1.4	44.3
13	Parkinson's disease (G20-G21)	204	1.3	40.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	114	0.7	22.4
15	Pneumonitis due to solids and liquids (J69)	87	0.6	17.1
...	All other causes (Residual)	2,799	17.9	548.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	17,995	100.0	10,237.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,520	30.7	3,140.3
2	Malignant neoplasms (C00-C97)	1,834	10.2	1,043.4
3	Alzheimer's disease (G30)	1,427	7.9	811.8
4	Cerebrovascular diseases (I60-I69)	1,405	7.8	799.3
5	Diabetes mellitus (E10-E14)	730	4.1	415.3
6	Chronic lower respiratory diseases (J40-J47)	709	3.9	403.3
7	Influenza and pneumonia (J09-J18)	581	3.2	330.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	425	2.4	241.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	389	2.2	221.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	306	1.7	174.1
11	Septicemia (A40-A41)	241	1.3	137.1
12	Parkinson's disease (G20-G21)	173	1.0	98.4
13	Pneumonitis due to solids and liquids (J69)	131	0.7	74.5
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	85	0.5	48.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	83	0.5	47.2
...	All other causes (Residual)	3,956	22.0	2,250.6
Non-Hispanic white, both sexes, all ages				
...	All causes	1,969,916	100.0	984.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	483,973	24.6	241.8
2	Malignant neoplasms (C00-C97)	460,567	23.4	230.1
3	Chronic lower respiratory diseases (J40-J47)	122,887	6.2	61.4
4	Cerebrovascular diseases (I60-I69)	101,849	5.2	50.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94,420	4.8	47.2
6	Alzheimer's disease (G30)	73,502	3.7	36.7
7	Diabetes mellitus (E10-E14)	47,746	2.4	23.9
8	Influenza and pneumonia (J09-J18)	40,244	2.0	20.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,963	1.9	18.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32,010	1.6	16.0
11	Septicemia (A40-A41)	25,951	1.3	13.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	23,655	1.2	11.8
13	Parkinson's disease (G20-G21)	19,818	1.0	9.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,878	1.0	9.4
15	Pneumonitis due to solids and liquids (J69)	14,539	0.7	7.3
...	All other causes (Residual)	372,914	18.9	186.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 1-4 years				
...	All causes	2,139	100.0	24.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	744	34.8	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	229	10.7	2.6
3	Malignant neoplasms (C00-C97)	196	9.2	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	7.4	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	2.8	0.7
6	Influenza and pneumonia (J09-J18)	38	1.8	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.7	0.4
8	Septicemia (A40-A41)	31	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	27	1.3	0.3
10	Cerebrovascular diseases (I60-I69)	23	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	17	0.8	*
12	Meningitis (G00,G03)	13	0.6	*
13	Anemias (D50-D64)	11	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
...	All other causes (Residual)	539	25.2	6.2
Non-Hispanic white, both sexes, 5-14 years				
...	All causes	2,910	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,000	34.4	4.3
2	Malignant neoplasms (C00-C97)	500	17.2	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	166	5.7	0.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	164	5.6	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	3.2	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.1	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	1.6	0.2
8	Influenza and pneumonia (J09-J18)	42	1.4	0.2
9	Cerebrovascular diseases (I60-I69)	39	1.3	0.2
10	Chronic lower respiratory diseases (J40-J47)	34	1.2	0.1
11	Septicemia (A40-A41)	31	1.1	0.1
12	Diabetes mellitus (E10-E14)	17	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	16	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.4	*
...	All other causes (Residual)	646	22.2	2.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 15-24 years				
...	All causes	16,847	100.0	66.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,507	50.5	33.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,190	18.9	12.6
3	Malignant neoplasms (C00-C97)	901	5.3	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	748	4.4	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	520	3.1	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	245	1.5	1.0
7	Influenza and pneumonia (J09-J18)	110	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	100	0.6	0.4
9	Diabetes mellitus (E10-E14)	95	0.6	0.4
10	Septicemia (A40-A41)	76	0.5	0.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	73	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	68	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.2	0.1
15	Pneumonitis due to solids and liquids (J69)	26	0.2	0.1
...	All other causes (Residual)	2,102	12.5	8.3
Non-Hispanic white, both sexes, 25-34 years				
...	All causes	25,486	100.0	105.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,425	40.9	43.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,335	17.0	18.0
3	Malignant neoplasms (C00-C97)	2,154	8.5	8.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,804	7.1	7.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	888	3.5	3.7
6	Diabetes mellitus (E10-E14)	325	1.3	1.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	282	1.1	1.2
8	Cerebrovascular diseases (I60-I69)	260	1.0	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	254	1.0	1.1
10	Influenza and pneumonia (J09-J18)	226	0.9	0.9
11	Septicemia (A40-A41)	167	0.7	0.7
12	Pregnancy, childbirth and the puerperium (O00-O99)	149	0.6	0.6
13	Chronic lower respiratory diseases (J40-J47)	134	0.5	0.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	125	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	0.4	0.4
...	All other causes (Residual)	3,853	15.1	16.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 35-44 years				
...	All causes	44,999	100.0	176.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,980	24.4	43.0
2	Malignant neoplasms (C00-C97)	7,637	17.0	29.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,566	14.6	25.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,359	11.9	21.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,574	3.5	6.2
6	Diabetes mellitus (E10-E14)	966	2.1	3.8
7	Cerebrovascular diseases (I60-I69)	915	2.0	3.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	871	1.9	3.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	504	1.1	2.0
10	Influenza and pneumonia (J09-J18)	492	1.1	1.9
11	Septicemia (A40-A41)	456	1.0	1.8
12	Chronic lower respiratory diseases (J40-J47)	455	1.0	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	334	0.7	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	334	0.7	1.3
15	Viral hepatitis (B15-B19)	200	0.4	0.8
...	All other causes (Residual)	7,356	16.3	28.8
Non-Hispanic white, both sexes, 45-54 years				
...	All causes	128,034	100.0	407.2
1	Malignant neoplasms (C00-C97)	35,789	28.0	113.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,082	19.6	79.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,259	11.9	48.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,755	6.1	24.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,229	4.9	19.8
6	Chronic lower respiratory diseases (J40-J47)	3,555	2.8	11.3
7	Diabetes mellitus (E10-E14)	3,383	2.6	10.8
8	Cerebrovascular diseases (I60-I69)	3,229	2.5	10.3
9	Viral hepatitis (B15-B19)	1,616	1.3	5.1
10	Septicemia (A40-A41)	1,459	1.1	4.6
11	Influenza and pneumonia (J09-J18)	1,307	1.0	4.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,117	0.9	3.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,025	0.8	3.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	949	0.7	3.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	712	0.6	2.3
...	All other causes (Residual)	19,568	15.3	62.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 55-64 years				
...	All causes	229,265	100.0	834.2
1	Malignant neoplasms (C00-C97)	83,251	36.3	302.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,276	21.5	179.3
3	Chronic lower respiratory diseases (J40-J47)	12,142	5.3	44.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,831	4.7	39.4
5	Diabetes mellitus (E10-E14)	7,464	3.3	27.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	7,245	3.2	26.4
7	Cerebrovascular diseases (I60-I69)	6,664	2.9	24.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,734	2.5	20.9
9	Septicemia (A40-A41)	3,065	1.3	11.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,891	1.3	10.5
11	Influenza and pneumonia (J09-J18)	2,605	1.1	9.5
12	Viral hepatitis (B15-B19)	2,149	0.9	7.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,485	0.6	5.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,114	0.5	4.1
15	Aortic aneurysm and dissection (I71)	940	0.4	3.4
...	All other causes (Residual)	32,409	14.1	117.9
Non-Hispanic white, both sexes, 65-74 years				
...	All causes	319,805	100.0	1,876.2
1	Malignant neoplasms (C00-C97)	116,480	36.4	683.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,089	21.3	399.5
3	Chronic lower respiratory diseases (J40-J47)	28,358	8.9	166.4
4	Cerebrovascular diseases (I60-I69)	12,673	4.0	74.3
5	Diabetes mellitus (E10-E14)	9,791	3.1	57.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,581	2.4	44.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,605	1.8	32.9
8	Septicemia (A40-A41)	4,743	1.5	27.8
9	Influenza and pneumonia (J09-J18)	4,461	1.4	26.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,429	1.4	26.0
11	Alzheimer's disease (G30)	3,562	1.1	20.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,683	0.8	15.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,274	0.7	13.3
14	Parkinson's disease (G20-G21)	2,224	0.7	13.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,141	0.7	12.6
...	All other causes (Residual)	44,711	14.0	262.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 75-84 years				
...	All causes	520,983	100.0	4,886.8
1	Malignant neoplasms (C00-C97)	131,485	25.2	1,233.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	125,926	24.2	1,181.2
3	Chronic lower respiratory diseases (J40-J47)	43,827	8.4	411.1
4	Cerebrovascular diseases (I60-I69)	30,451	5.8	285.6
5	Alzheimer's disease (G30)	20,934	4.0	196.4
6	Diabetes mellitus (E10-E14)	13,543	2.6	127.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,028	2.3	112.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,557	2.2	108.4
9	Influenza and pneumonia (J09-J18)	10,775	2.1	101.1
10	Parkinson's disease (G20-G21)	8,794	1.7	82.5
11	Septicemia (A40-A41)	7,755	1.5	72.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,892	0.9	45.9
13	Pneumonitis due to solids and liquids (J69)	4,268	0.8	40.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,076	0.8	38.2
15	Aortic aneurysm and dissection (I71)	2,989	0.6	28.0
...	All other causes (Residual)	87,683	16.8	822.5
Non-Hispanic white, both sexes, 85 years and over				
...	All causes	668,361	100.0	14,286.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	206,389	30.9	4,411.5
2	Malignant neoplasms (C00-C97)	82,136	12.3	1,755.6
3	Alzheimer's disease (G30)	48,261	7.2	1,031.6
4	Cerebrovascular diseases (I60-I69)	47,441	7.1	1,014.0
5	Chronic lower respiratory diseases (J40-J47)	34,281	5.1	732.7
6	Influenza and pneumonia (J09-J18)	20,117	3.0	430.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,492	2.5	352.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,272	2.3	326.4
9	Diabetes mellitus (E10-E14)	12,160	1.8	259.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,315	1.4	199.1
11	Parkinson's disease (G20-G21)	8,314	1.2	177.7
12	Septicemia (A40-A41)	8,094	1.2	173.0
13	Pneumonitis due to solids and liquids (J69)	7,464	1.1	159.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,193	0.6	89.6
15	Atherosclerosis (I70)	3,742	0.6	80.0
...	All other causes (Residual)	144,690	21.6	3,092.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, all ages				
...	All causes	971,604	100.0	987.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,765	25.5	251.8
2	Malignant neoplasms (C00-C97)	241,816	24.9	245.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57,747	5.9	58.7
4	Chronic lower respiratory diseases (J40-J47)	57,390	5.9	58.3
5	Cerebrovascular diseases (I60-I69)	40,040	4.1	40.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,238	2.6	25.7
7	Diabetes mellitus (E10-E14)	25,134	2.6	25.5
8	Alzheimer's disease (G30)	22,278	2.3	22.6
9	Influenza and pneumonia (J09-J18)	18,657	1.9	19.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,503	1.9	18.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,270	1.6	15.5
12	Septicemia (A40-A41)	11,963	1.2	12.2
13	Parkinson's disease (G20-G21)	11,627	1.2	11.8
14	Pneumonitis due to solids and liquids (J69)	7,862	0.8	8.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,476	0.8	7.6
...	All other causes (Residual)	162,838	16.8	165.5
Non-Hispanic white, male, 1-4 years				
...	All causes	1,219	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	456	37.4	10.3
2	Malignant neoplasms (C00-C97)	109	8.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	8.3	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	7.4	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.0	0.8
6	Influenza and pneumonia (J09-J18)	20	1.6	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.4	*
7	Cerebrovascular diseases (I60-I69)	17	1.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	17	1.4	*
10	Septicemia (A40-A41)	13	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	10	0.8	*
12	Meningitis (G00,G03)	9	0.7	*
13	Anemias (D50-D64)	8	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
...	All other causes (Residual)	304	24.9	6.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 5-14 years				
...	All causes	1,691	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	639	37.8	5.4
2	Malignant neoplasms (C00-C97)	277	16.4	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	128	7.6	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	4.4	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	3.3	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	2.4	0.3
7	Cerebrovascular diseases (I60-I69)	23	1.4	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.1	*
9	Septicemia (A40-A41)	18	1.1	*
10	Influenza and pneumonia (J09-J18)	15	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	12	0.7	*
13	Diabetes mellitus (E10-E14)	11	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	356	21.1	3.0
Non-Hispanic white, male, 15-24 years				
...	All causes	12,079	100.0	93.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,172	51.1	47.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,638	21.8	20.4
3	Malignant neoplasms (C00-C97)	565	4.7	4.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	528	4.4	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	382	3.2	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	150	1.2	1.2
7	Cerebrovascular diseases (I60-I69)	61	0.5	0.5
8	Influenza and pneumonia (J09-J18)	59	0.5	0.5
9	Diabetes mellitus (E10-E14)	54	0.4	0.4
10	Chronic lower respiratory diseases (J40-J47)	42	0.3	0.3
11	Septicemia (A40-A41)	38	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.3	0.2
13	Legal intervention (Y35,Y89.0)	24	0.2	0.2
14	Aortic aneurysm and dissection (I71)	18	0.1	*
15	Pneumonitis due to solids and liquids (J69)	17	0.1	*
...	All other causes (Residual)	1,299	10.8	10.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 25-34 years				
...	All causes	17,483	100.0	143.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,638	43.7	62.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,480	19.9	28.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,250	7.1	10.3
4	Malignant neoplasms (C00-C97)	1,106	6.3	9.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	605	3.5	5.0
6	Diabetes mellitus (E10-E14)	186	1.1	1.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	172	1.0	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	0.8	1.2
9	Influenza and pneumonia (J09-J18)	124	0.7	1.0
10	Cerebrovascular diseases (I60-I69)	116	0.7	1.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	88	0.5	0.7
12	Septicemia (A40-A41)	83	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	75	0.4	0.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.3	0.4
...	All other causes (Residual)	2,298	13.1	18.9
Non-Hispanic white, male, 35-44 years				
...	All causes	28,070	100.0	219.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,397	26.4	57.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,687	16.7	36.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,100	14.6	32.0
4	Malignant neoplasms (C00-C97)	3,462	12.3	27.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,012	3.6	7.9
6	Diabetes mellitus (E10-E14)	573	2.0	4.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	570	2.0	4.4
8	Cerebrovascular diseases (I60-I69)	474	1.7	3.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	393	1.4	3.1
10	Influenza and pneumonia (J09-J18)	259	0.9	2.0
11	Septicemia (A40-A41)	235	0.8	1.8
12	Chronic lower respiratory diseases (J40-J47)	202	0.7	1.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	0.7	1.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	178	0.6	1.4
15	Viral hepatitis (B15-B19)	130	0.5	1.0
...	All other causes (Residual)	4,207	15.0	32.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 45-54 years				
...	All causes	79,295	100.0	508.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,293	23.1	117.2
2	Malignant neoplasms (C00-C97)	18,285	23.1	117.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,096	12.7	64.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,924	7.5	38.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,212	5.3	27.0
6	Diabetes mellitus (E10-E14)	2,206	2.8	14.1
7	Cerebrovascular diseases (I60-I69)	1,811	2.3	11.6
8	Chronic lower respiratory diseases (J40-J47)	1,648	2.1	10.6
9	Viral hepatitis (B15-B19)	1,108	1.4	7.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	877	1.1	5.6
11	Septicemia (A40-A41)	790	1.0	5.1
12	Influenza and pneumonia (J09-J18)	713	0.9	4.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	668	0.8	4.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	645	0.8	4.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	471	0.6	3.0
...	All other causes (Residual)	11,548	14.6	74.0
Non-Hispanic white, male, 55-64 years				
...	All causes	140,552	100.0	1,046.2
1	Malignant neoplasms (C00-C97)	46,240	32.9	344.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,191	25.0	261.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,360	5.2	54.8
4	Chronic lower respiratory diseases (J40-J47)	6,252	4.4	46.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,112	3.6	38.1
6	Diabetes mellitus (E10-E14)	4,694	3.3	34.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,355	3.1	32.4
8	Cerebrovascular diseases (I60-I69)	3,814	2.7	28.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,659	1.2	12.3
10	Septicemia (A40-A41)	1,625	1.2	12.1
11	Viral hepatitis (B15-B19)	1,570	1.1	11.7
12	Influenza and pneumonia (J09-J18)	1,528	1.1	11.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	971	0.7	7.2
14	Aortic aneurysm and dissection (I71)	708	0.5	5.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	665	0.5	5.0
...	All other causes (Residual)	18,808	13.4	140.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 65-74 years				
...	All causes	181,567	100.0	2,256.9
1	Malignant neoplasms (C00-C97)	64,238	35.4	798.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,618	24.0	542.2
3	Chronic lower respiratory diseases (J40-J47)	14,286	7.9	177.6
4	Cerebrovascular diseases (I60-I69)	6,648	3.7	82.6
5	Diabetes mellitus (E10-E14)	5,689	3.1	70.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,768	2.6	59.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,108	1.7	38.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2,751	1.5	34.2
9	Influenza and pneumonia (J09-J18)	2,502	1.4	31.1
10	Septicemia (A40-A41)	2,435	1.3	30.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,182	1.2	27.1
12	Alzheimer's disease (G30)	1,573	0.9	19.6
13	Parkinson's disease (G20-G21)	1,480	0.8	18.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,247	0.7	15.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,236	0.7	15.4
...	All other causes (Residual)	23,806	13.1	295.9
Non-Hispanic white, male, 75-84 years				
...	All causes	261,723	100.0	5,770.3
1	Malignant neoplasms (C00-C97)	69,597	26.6	1,534.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	67,646	25.8	1,491.4
3	Chronic lower respiratory diseases (J40-J47)	20,862	8.0	460.0
4	Cerebrovascular diseases (I60-I69)	13,091	5.0	288.6
5	Alzheimer's disease (G30)	7,887	3.0	173.9
6	Diabetes mellitus (E10-E14)	6,963	2.7	153.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,363	2.4	140.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,135	2.3	135.3
9	Parkinson's disease (G20-G21)	5,556	2.1	122.5
10	Influenza and pneumonia (J09-J18)	5,506	2.1	121.4
11	Septicemia (A40-A41)	3,676	1.4	81.0
12	Pneumonitis due to solids and liquids (J69)	2,480	0.9	54.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,241	0.9	49.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,035	0.8	44.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,639	0.6	36.1
...	All other causes (Residual)	40,046	15.3	882.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 85 years and over				
...	All causes	241,741	100.0	15,816.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,521	31.7	5,006.6
2	Malignant neoplasms (C00-C97)	37,916	15.7	2,480.8
3	Chronic lower respiratory diseases (J40-J47)	13,991	5.8	915.4
4	Cerebrovascular diseases (I60-I69)	13,958	5.8	913.2
5	Alzheimer's disease (G30)	12,503	5.2	818.0
6	Influenza and pneumonia (J09-J18)	7,891	3.3	516.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,663	2.8	435.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,529	2.7	427.2
9	Diabetes mellitus (E10-E14)	4,757	2.0	311.2
10	Parkinson's disease (G20-G21)	4,269	1.8	279.3
11	Pneumonitis due to solids and liquids (J69)	3,683	1.5	241.0
12	Septicemia (A40-A41)	3,003	1.2	196.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,629	1.1	172.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,858	0.8	121.6
15	Atherosclerosis (I70)	1,193	0.5	78.1
...	All other causes (Residual)	44,377	18.4	2,903.5
Non-Hispanic white, female, all ages				
...	All causes	998,312	100.0	981.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	236,208	23.7	232.2
2	Malignant neoplasms (C00-C97)	218,751	21.9	215.0
3	Chronic lower respiratory diseases (J40-J47)	65,497	6.6	64.4
4	Cerebrovascular diseases (I60-I69)	61,809	6.2	60.8
5	Alzheimer's disease (G30)	51,224	5.1	50.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36,673	3.7	36.0
7	Diabetes mellitus (E10-E14)	22,612	2.3	22.2
8	Influenza and pneumonia (J09-J18)	21,587	2.2	21.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,460	1.8	18.1
10	Septicemia (A40-A41)	13,988	1.4	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,402	1.1	11.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,385	0.8	8.2
13	Parkinson's disease (G20-G21)	8,191	0.8	8.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,772	0.7	6.7
15	Pneumonitis due to solids and liquids (J69)	6,677	0.7	6.6
...	All other causes (Residual)	210,076	21.0	206.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 1-4 years				
...	All causes	920	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	31.3	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	128	13.9	3.0
3	Malignant neoplasms (C00-C97)	87	9.5	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	7.5	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.5	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	2.2	0.5
7	Septicemia (A40-A41)	18	2.0	*
7	Influenza and pneumonia (J09-J18)	18	2.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	10	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	7	0.8	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
12	Meningitis (G00,G03)	4	0.4	*
13	Meningococcal infection (A39)	3	0.3	*
13	Anemias (D50-D64)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	233	25.3	5.5
Non-Hispanic white, female, 5-14 years				
...	All causes	1,219	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	361	29.6	3.2
2	Malignant neoplasms (C00-C97)	223	18.3	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	7.4	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.0	0.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	3.2	0.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	3.1	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	2.3	0.2
8	Influenza and pneumonia (J09-J18)	27	2.2	0.2
9	Chronic lower respiratory diseases (J40-J47)	20	1.6	0.2
10	Cerebrovascular diseases (I60-I69)	16	1.3	*
11	Septicemia (A40-A41)	13	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
14	Diabetes mellitus (E10-E14)	6	0.5	*
15	Anemias (D50-D64)	5	0.4	*
...	All other causes (Residual)	289	23.7	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 15-24 years				
...	All causes	4,768	100.0	38.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,335	49.0	18.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	552	11.6	4.4
3	Malignant neoplasms (C00-C97)	336	7.0	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	220	4.6	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	138	2.9	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	2.0	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	73	1.5	0.6
8	Influenza and pneumonia (J09-J18)	51	1.1	0.4
9	Diabetes mellitus (E10-E14)	41	0.9	0.3
10	Cerebrovascular diseases (I60-I69)	39	0.8	0.3
11	Septicemia (A40-A41)	38	0.8	0.3
12	Chronic lower respiratory diseases (J40-J47)	26	0.5	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.4	*
15	Anemias (D50-D64)	11	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.2	*
...	All other causes (Residual)	761	16.0	6.1
Non-Hispanic white, female, 25-34 years				
...	All causes	8,003	100.0	66.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,787	34.8	23.3
2	Malignant neoplasms (C00-C97)	1,048	13.1	8.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	855	10.7	7.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	554	6.9	4.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	283	3.5	2.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	149	1.9	1.2
7	Cerebrovascular diseases (I60-I69)	144	1.8	1.2
8	Diabetes mellitus (E10-E14)	139	1.7	1.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	110	1.4	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	1.3	0.9
11	Influenza and pneumonia (J09-J18)	102	1.3	0.9
12	Septicemia (A40-A41)	84	1.0	0.7
13	Chronic lower respiratory diseases (J40-J47)	59	0.7	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	0.6	0.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	37	0.5	0.3
...	All other causes (Residual)	1,492	18.6	12.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 35-44 years				
...	All causes	16,929	100.0	133.1
1	Malignant neoplasms (C00-C97)	4,175	24.7	32.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,583	21.2	28.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,879	11.1	14.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,259	7.4	9.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	562	3.3	4.4
6	Cerebrovascular diseases (I60-I69)	441	2.6	3.5
7	Diabetes mellitus (E10-E14)	393	2.3	3.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	301	1.8	2.4
9	Chronic lower respiratory diseases (J40-J47)	253	1.5	2.0
10	Influenza and pneumonia (J09-J18)	233	1.4	1.8
11	Septicemia (A40-A41)	221	1.3	1.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	0.9	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	143	0.8	1.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	111	0.7	0.9
15	Pregnancy, childbirth and the puerperium (O00-O99)	87	0.5	0.7
...	All other causes (Residual)	3,132	18.5	24.6
Non-Hispanic white, female, 45-54 years				
...	All causes	48,739	100.0	307.7
1	Malignant neoplasms (C00-C97)	17,504	35.9	110.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,789	13.9	42.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,163	10.6	32.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,017	4.1	12.7
5	Chronic lower respiratory diseases (J40-J47)	1,907	3.9	12.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,831	3.8	11.6
7	Cerebrovascular diseases (I60-I69)	1,418	2.9	9.0
8	Diabetes mellitus (E10-E14)	1,177	2.4	7.4
9	Septicemia (A40-A41)	669	1.4	4.2
10	Influenza and pneumonia (J09-J18)	594	1.2	3.8
11	Viral hepatitis (B15-B19)	508	1.0	3.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	449	0.9	2.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	304	0.6	1.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	242	0.5	1.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	241	0.5	1.5
...	All other causes (Residual)	7,926	16.3	50.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 55-64 years				
...	All causes	88,713	100.0	631.5
1	Malignant neoplasms (C00-C97)	37,011	41.7	263.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,085	15.9	100.3
3	Chronic lower respiratory diseases (J40-J47)	5,890	6.6	41.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,471	3.9	24.7
5	Cerebrovascular diseases (I60-I69)	2,850	3.2	20.3
6	Diabetes mellitus (E10-E14)	2,770	3.1	19.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,133	2.4	15.2
8	Septicemia (A40-A41)	1,440	1.6	10.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,379	1.6	9.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,232	1.4	8.8
11	Influenza and pneumonia (J09-J18)	1,077	1.2	7.7
12	Viral hepatitis (B15-B19)	579	0.7	4.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	514	0.6	3.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	449	0.5	3.2
15	Alzheimer's disease (G30)	362	0.4	2.6
...	All other causes (Residual)	13,471	15.2	95.9
Non-Hispanic white, female, 65-74 years				
...	All causes	138,238	100.0	1,535.9
1	Malignant neoplasms (C00-C97)	52,242	37.8	580.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,471	17.7	271.9
3	Chronic lower respiratory diseases (J40-J47)	14,072	10.2	156.4
4	Cerebrovascular diseases (I60-I69)	6,025	4.4	66.9
5	Diabetes mellitus (E10-E14)	4,102	3.0	45.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,813	2.0	31.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,497	1.8	27.7
8	Septicemia (A40-A41)	2,308	1.7	25.6
9	Alzheimer's disease (G30)	1,989	1.4	22.1
10	Influenza and pneumonia (J09-J18)	1,959	1.4	21.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,678	1.2	18.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,038	0.8	11.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	894	0.6	9.9
14	Parkinson's disease (G20-G21)	744	0.5	8.3
15	Pneumonitis due to solids and liquids (J69)	615	0.4	6.8
...	All other causes (Residual)	20,791	15.0	231.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 75-84 years				
...	All causes	259,260	100.0	4,232.6
1	Malignant neoplasms (C00-C97)	61,888	23.9	1,010.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	58,280	22.5	951.5
3	Chronic lower respiratory diseases (J40-J47)	22,965	8.9	374.9
4	Cerebrovascular diseases (I60-I69)	17,360	6.7	283.4
5	Alzheimer's disease (G30)	13,047	5.0	213.0
6	Diabetes mellitus (E10-E14)	6,580	2.5	107.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,665	2.2	92.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,422	2.1	88.5
9	Influenza and pneumonia (J09-J18)	5,269	2.0	86.0
10	Septicemia (A40-A41)	4,079	1.6	66.6
11	Parkinson's disease (G20-G21)	3,238	1.2	52.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,857	1.1	46.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,835	0.7	30.0
14	Pneumonitis due to solids and liquids (J69)	1,788	0.7	29.2
15	Aortic aneurysm and dissection (I71)	1,371	0.5	22.4
...	All other causes (Residual)	47,616	18.4	777.4
Non-Hispanic white, female, 85 years and over				
...	All causes	426,620	100.0	13,543.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	129,868	30.4	4,122.8
2	Malignant neoplasms (C00-C97)	44,220	10.4	1,403.8
3	Alzheimer's disease (G30)	35,758	8.4	1,135.2
4	Cerebrovascular diseases (I60-I69)	33,483	7.8	1,063.0
5	Chronic lower respiratory diseases (J40-J47)	20,290	4.8	644.1
6	Influenza and pneumonia (J09-J18)	12,226	2.9	388.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,963	2.3	316.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,609	2.0	273.3
9	Diabetes mellitus (E10-E14)	7,403	1.7	235.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,686	1.6	212.3
11	Septicemia (A40-A41)	5,091	1.2	161.6
12	Parkinson's disease (G20-G21)	4,045	0.9	128.4
13	Pneumonitis due to solids and liquids (J69)	3,781	0.9	120.0
14	Atherosclerosis (I70)	2,549	0.6	80.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,335	0.5	74.1
...	All other causes (Residual)	100,313	23.5	3,184.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, all ages				
...	All causes	283,438	100.0	718.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,215	24.1	173.0
2	Malignant neoplasms (C00-C97)	65,276	23.0	165.5
3	Cerebrovascular diseases (I60-I69)	15,833	5.6	40.1
4	Diabetes mellitus (E10-E14)	11,996	4.2	30.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,853	4.2	30.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,769	3.1	22.2
7	Chronic lower respiratory diseases (J40-J47)	8,614	3.0	21.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,679	2.7	19.5
9	Septicemia (A40-A41)	5,958	2.1	15.1
10	Alzheimer's disease (G30)	5,169	1.8	13.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,059	1.8	12.8
12	Influenza and pneumonia (J09-J18)	4,854	1.7	12.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,598	1.6	11.7
14	Certain conditions originating in the perinatal period (P00-P96)	3,846	1.4	9.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,601	0.9	6.6
...	All other causes (Residual)	53,118	18.7	134.7
Non-Hispanic black, both sexes, 1-4 years				
...	All causes	993	100.0	40.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	249	25.1	10.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	15.4	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	12.5	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	5.0	2.0
5	Malignant neoplasms (C00-C97)	47	4.7	1.9
6	Influenza and pneumonia (J09-J18)	25	2.5	1.0
7	Chronic lower respiratory diseases (J40-J47)	19	1.9	*
8	Certain conditions originating in the perinatal period (P00-P96)	15	1.5	*
9	Anemias (D50-D64)	14	1.4	*
10	Cerebrovascular diseases (I60-I69)	13	1.3	*
11	Septicemia (A40-A41)	11	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	250	25.2	10.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 5-14 years				
...	All causes	1,118	100.0	18.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	312	27.9	5.1
2	Malignant neoplasms (C00-C97)	142	12.7	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	8.9	1.6
4	Chronic lower respiratory diseases (J40-J47)	71	6.4	1.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	5.5	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	5.4	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.8	0.7
8	Cerebrovascular diseases (I60-I69)	21	1.9	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.5	*
10	Septicemia (A40-A41)	14	1.3	*
10	Anemias (D50-D64)	14	1.3	*
12	Influenza and pneumonia (J09-J18)	11	1.0	*
13	Diabetes mellitus (E10-E14)	6	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (Residual)	235	21.0	3.8
Non-Hispanic black, both sexes, 15-24 years				
...	All causes	6,546	100.0	98.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,762	42.2	41.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,426	21.8	21.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	454	6.9	6.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	327	5.0	4.9
5	Malignant neoplasms (C00-C97)	264	4.0	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	109	1.7	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	1.4	1.3
8	Anemias (D50-D64)	79	1.2	1.2
9	Chronic lower respiratory diseases (J40-J47)	60	0.9	0.9
10	Diabetes mellitus (E10-E14)	51	0.8	0.8
11	Pregnancy, childbirth and the puerperium (O00-O99)	50	0.8	0.8
12	Cerebrovascular diseases (I60-I69)	49	0.7	0.7
13	Septicemia (A40-A41)	41	0.6	0.6
14	Influenza and pneumonia (J09-J18)	39	0.6	0.6
15	Legal intervention (Y35,Y89.0)	28	0.4	0.4
...	All other causes (Residual)	718	11.0	10.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 25-34 years				
...	All causes	8,786	100.0	161.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,366	26.9	43.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,580	18.0	29.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	905	10.3	16.7
4	Malignant neoplasms (C00-C97)	575	6.5	10.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	494	5.6	9.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	471	5.4	8.7
7	Diabetes mellitus (E10-E14)	206	2.3	3.8
8	Cerebrovascular diseases (I60-I69)	137	1.6	2.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	118	1.3	2.2
10	Anemias (D50-D64)	107	1.2	2.0
11	Chronic lower respiratory diseases (J40-J47)	96	1.1	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	1.1	1.7
13	Septicemia (A40-A41)	85	1.0	1.6
14	Influenza and pneumonia (J09-J18)	82	0.9	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	0.8	1.3
...	All other causes (Residual)	1,401	15.9	25.8
Non-Hispanic black, both sexes, 35-44 years				
...	All causes	13,807	100.0	261.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,688	19.5	51.0
2	Malignant neoplasms (C00-C97)	2,085	15.1	39.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,568	11.4	29.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,065	7.7	20.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,065	7.7	20.2
6	Cerebrovascular diseases (I60-I69)	553	4.0	10.5
7	Diabetes mellitus (E10-E14)	507	3.7	9.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	435	3.2	8.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	279	2.0	5.3
10	Septicemia (A40-A41)	210	1.5	4.0
11	Chronic lower respiratory diseases (J40-J47)	179	1.3	3.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	178	1.3	3.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	172	1.2	3.3
14	Influenza and pneumonia (J09-J18)	143	1.0	2.7
15	Anemias (D50-D64)	130	0.9	2.5
...	All other causes (Residual)	2,550	18.5	48.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 45-54 years				
...	All causes	33,698	100.0	610.9
1	Malignant neoplasms (C00-C97)	8,467	25.1	153.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,026	23.8	145.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,274	6.7	41.2
4	Cerebrovascular diseases (I60-I69)	1,710	5.1	31.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,610	4.8	29.2
6	Diabetes mellitus (E10-E14)	1,385	4.1	25.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	804	2.4	14.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	792	2.4	14.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	682	2.0	12.4
10	Chronic lower respiratory diseases (J40-J47)	642	1.9	11.6
11	Septicemia (A40-A41)	605	1.8	11.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	528	1.6	9.6
13	Influenza and pneumonia (J09-J18)	388	1.2	7.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	332	1.0	6.0
15	Viral hepatitis (B15-B19)	310	0.9	5.6
...	All other causes (Residual)	5,143	15.3	93.2
Non-Hispanic black, both sexes, 55-64 years				
...	All causes	51,288	100.0	1,318.8
1	Malignant neoplasms (C00-C97)	16,221	31.6	417.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,686	24.7	326.2
3	Cerebrovascular diseases (I60-I69)	2,637	5.1	67.8
4	Diabetes mellitus (E10-E14)	2,464	4.8	63.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,753	3.4	45.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,527	3.0	39.3
7	Chronic lower respiratory diseases (J40-J47)	1,449	2.8	37.3
8	Septicemia (A40-A41)	1,099	2.1	28.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,014	2.0	26.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,013	2.0	26.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	872	1.7	22.4
12	Influenza and pneumonia (J09-J18)	588	1.1	15.1
13	Viral hepatitis (B15-B19)	495	1.0	12.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	317	0.6	8.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	203	0.4	5.2
...	All other causes (Residual)	6,950	13.6	178.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 65-74 years				
...	All causes	51,930	100.0	2,582.2
1	Malignant neoplasms (C00-C97)	16,467	31.7	818.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,021	25.1	647.5
3	Cerebrovascular diseases (I60-I69)	3,096	6.0	153.9
4	Diabetes mellitus (E10-E14)	2,745	5.3	136.5
5	Chronic lower respiratory diseases (J40-J47)	2,160	4.2	107.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,946	3.7	96.8
7	Septicemia (A40-A41)	1,258	2.4	62.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	940	1.8	46.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	900	1.7	44.8
10	Influenza and pneumonia (J09-J18)	858	1.7	42.7
11	Alzheimer's disease (G30)	402	0.8	20.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	382	0.7	19.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	254	0.5	12.6
14	Pneumonitis due to solids and liquids (J69)	231	0.4	11.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	230	0.4	11.4
...	All other causes (Residual)	7,040	13.6	350.1
Non-Hispanic black, both sexes, 75-84 years				
...	All causes	56,942	100.0	5,477.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,233	26.8	1,465.4
2	Malignant neoplasms (C00-C97)	13,908	24.4	1,338.0
3	Cerebrovascular diseases (I60-I69)	3,892	6.8	374.4
4	Diabetes mellitus (E10-E14)	2,796	4.9	269.0
5	Chronic lower respiratory diseases (J40-J47)	2,312	4.1	222.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,208	3.9	212.4
7	Alzheimer's disease (G30)	1,650	2.9	158.7
8	Septicemia (A40-A41)	1,415	2.5	136.1
9	Influenza and pneumonia (J09-J18)	1,212	2.1	116.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,115	2.0	107.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	813	1.4	78.2
12	Pneumonitis due to solids and liquids (J69)	400	0.7	38.5
13	Parkinson's disease (G20-G21)	320	0.6	30.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	288	0.5	27.7
15	Aortic aneurysm and dissection (I71)	202	0.4	19.4
...	All other causes (Residual)	9,178	16.1	882.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 85 years and over				
...	All causes	51,242	100.0	13,385.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,122	29.5	3,950.3
2	Malignant neoplasms (C00-C97)	7,091	13.8	1,852.4
3	Cerebrovascular diseases (I60-I69)	3,680	7.2	961.3
4	Alzheimer's disease (G30)	3,057	6.0	798.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,860	3.6	485.9
6	Diabetes mellitus (E10-E14)	1,832	3.6	478.6
7	Chronic lower respiratory diseases (J40-J47)	1,617	3.2	422.4
8	Influenza and pneumonia (J09-J18)	1,437	2.8	375.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,364	2.7	356.3
10	Septicemia (A40-A41)	1,133	2.2	296.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	638	1.2	166.7
12	Pneumonitis due to solids and liquids (J69)	453	0.9	118.3
13	Parkinson's disease (G20-G21)	276	0.5	72.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	245	0.5	64.0
15	Atherosclerosis (I70)	229	0.4	59.8
...	All other causes (Residual)	11,208	21.9	2,927.8
Non-Hispanic black, male, all ages				
...	All causes	143,824	100.0	764.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,597	24.1	183.9
2	Malignant neoplasms (C00-C97)	33,607	23.4	178.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,920	5.5	42.1
4	Cerebrovascular diseases (I60-I69)	6,890	4.8	36.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,580	4.6	35.0
6	Diabetes mellitus (E10-E14)	5,572	3.9	29.6
7	Chronic lower respiratory diseases (J40-J47)	4,480	3.1	23.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,984	2.8	21.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,000	2.1	15.9
10	Septicemia (A40-A41)	2,673	1.9	14.2
11	Influenza and pneumonia (J09-J18)	2,340	1.6	12.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,196	1.5	11.7
13	Certain conditions originating in the perinatal period (P00-P96)	2,116	1.5	11.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,712	1.2	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,694	1.2	9.0
...	All other causes (Residual)	24,463	17.0	130.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 1-4 years				
...	All causes	570	100.0	45.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	27.9	12.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	91	16.0	7.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	10.0	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.9	2.2
5	Malignant neoplasms (C00-C97)	26	4.6	2.1
6	Influenza and pneumonia (J09-J18)	15	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	10	1.8	*
8	Anemias (D50-D64)	9	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
12	Septicemia (A40-A41)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
14	Meningococcal infection (A39)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Hernia (K40-K46)	2	0.4	*
...	All other causes (Residual)	141	24.7	11.2
Non-Hispanic black, male, 5-14 years				
...	All causes	649	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	31.9	6.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	10.3	2.1
3	Malignant neoplasms (C00-C97)	62	9.6	2.0
4	Chronic lower respiratory diseases (J40-J47)	51	7.9	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	5.9	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	5.2	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	3.5	0.7
8	Cerebrovascular diseases (I60-I69)	14	2.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.4	*
10	Septicemia (A40-A41)	7	1.1	*
10	Influenza and pneumonia (J09-J18)	7	1.1	*
12	Anemias (D50-D64)	6	0.9	*
13	Diabetes mellitus (E10-E14)	3	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	116	17.9	3.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 15-24 years				
...	All causes	5,033	100.0	150.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,499	49.7	74.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,053	20.9	31.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	385	7.6	11.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	204	4.1	6.1
5	Malignant neoplasms (C00-C97)	148	2.9	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	70	1.4	2.1
7	Anemias (D50-D64)	55	1.1	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	1.1	1.6
9	Chronic lower respiratory diseases (J40-J47)	41	0.8	1.2
10	Legal intervention (Y35,Y89.0)	27	0.5	0.8
11	Septicemia (A40-A41)	26	0.5	0.8
11	Diabetes mellitus (E10-E14)	26	0.5	0.8
13	Influenza and pneumonia (J09-J18)	24	0.5	0.7
14	Cerebrovascular diseases (I60-I69)	21	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.2	*
...	All other causes (Residual)	389	7.7	11.7
Non-Hispanic black, male, 25-34 years				
...	All causes	5,971	100.0	230.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,099	35.2	81.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,157	19.4	44.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	569	9.5	22.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	425	7.1	16.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	274	4.6	10.6
6	Malignant neoplasms (C00-C97)	250	4.2	9.7
7	Diabetes mellitus (E10-E14)	112	1.9	4.3
8	Cerebrovascular diseases (I60-I69)	75	1.3	2.9
9	Chronic lower respiratory diseases (J40-J47)	68	1.1	2.6
10	Anemias (D50-D64)	51	0.9	2.0
11	Influenza and pneumonia (J09-J18)	48	0.8	1.9
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	0.7	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.7	1.6
14	Septicemia (A40-A41)	35	0.6	1.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.5	1.1
...	All other causes (Residual)	696	11.7	26.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 35-44 years				
...	All causes	7,975	100.0	321.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,648	20.7	66.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,079	13.5	43.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	890	11.2	35.8
4	Malignant neoplasms (C00-C97)	784	9.8	31.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	647	8.1	26.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	340	4.3	13.7
7	Diabetes mellitus (E10-E14)	300	3.8	12.1
8	Cerebrovascular diseases (I60-I69)	287	3.6	11.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	1.9	6.2
10	Septicemia (A40-A41)	104	1.3	4.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	104	1.3	4.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	1.2	3.9
13	Chronic lower respiratory diseases (J40-J47)	89	1.1	3.6
14	Influenza and pneumonia (J09-J18)	84	1.1	3.4
15	Aortic aneurysm and dissection (I71)	81	1.0	3.3
...	All other causes (Residual)	1,287	16.1	51.8
Non-Hispanic black, male, 45-54 years				
...	All causes	19,146	100.0	739.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,106	26.7	197.1
2	Malignant neoplasms (C00-C97)	4,034	21.1	155.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,537	8.0	59.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,071	5.6	41.3
5	Cerebrovascular diseases (I60-I69)	904	4.7	34.9
6	Diabetes mellitus (E10-E14)	786	4.1	30.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	544	2.8	21.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	513	2.7	19.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	395	2.1	15.2
10	Septicemia (A40-A41)	314	1.6	12.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	287	1.5	11.1
12	Chronic lower respiratory diseases (J40-J47)	283	1.5	10.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	259	1.4	10.0
14	Influenza and pneumonia (J09-J18)	226	1.2	8.7
15	Viral hepatitis (B15-B19)	208	1.1	8.0
...	All other causes (Residual)	2,679	14.0	103.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 55-64 years				
...	All causes	30,138	100.0	1,705.0
1	Malignant neoplasms (C00-C97)	9,019	29.9	510.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,929	26.3	448.6
3	Cerebrovascular diseases (I60-I69)	1,519	5.0	85.9
4	Diabetes mellitus (E10-E14)	1,339	4.4	75.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,255	4.2	71.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	842	2.8	47.6
7	Chronic lower respiratory diseases (J40-J47)	787	2.6	44.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	720	2.4	40.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	706	2.3	39.9
10	Septicemia (A40-A41)	576	1.9	32.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	494	1.6	27.9
12	Viral hepatitis (B15-B19)	346	1.1	19.6
13	Influenza and pneumonia (J09-J18)	336	1.1	19.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	239	0.8	13.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	0.6	9.4
...	All other causes (Residual)	3,864	12.8	218.6
Non-Hispanic black, male, 65-74 years				
...	All causes	28,060	100.0	3,274.7
1	Malignant neoplasms (C00-C97)	9,004	32.1	1,050.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,406	26.4	864.3
3	Cerebrovascular diseases (I60-I69)	1,607	5.7	187.5
4	Diabetes mellitus (E10-E14)	1,328	4.7	155.0
5	Chronic lower respiratory diseases (J40-J47)	1,204	4.3	140.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	965	3.4	112.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	589	2.1	68.7
8	Septicemia (A40-A41)	587	2.1	68.5
9	Influenza and pneumonia (J09-J18)	484	1.7	56.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	479	1.7	55.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	248	0.9	28.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	182	0.6	21.2
13	Alzheimer's disease (G30)	168	0.6	19.6
14	Pneumonitis due to solids and liquids (J69)	142	0.5	16.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	110	0.4	12.8
...	All other causes (Residual)	3,557	12.7	415.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 75-84 years				
...	All causes	26,281	100.0	6,849.1
1	Malignant neoplasms (C00-C97)	7,159	27.2	1,865.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,033	26.8	1,832.9
3	Cerebrovascular diseases (I60-I69)	1,561	5.9	406.8
4	Chronic lower respiratory diseases (J40-J47)	1,215	4.6	316.6
5	Diabetes mellitus (E10-E14)	1,161	4.4	302.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	953	3.6	248.4
7	Septicemia (A40-A41)	618	2.4	161.1
8	Alzheimer's disease (G30)	606	2.3	157.9
9	Influenza and pneumonia (J09-J18)	582	2.2	151.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	446	1.7	116.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	438	1.7	114.1
12	Pneumonitis due to solids and liquids (J69)	230	0.9	59.9
13	Parkinson's disease (G20-G21)	189	0.7	49.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	122	0.5	31.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	90	0.3	23.5
...	All other causes (Residual)	3,878	14.8	1,010.6
Non-Hispanic black, male, 85 years and over				
...	All causes	16,055	100.0	14,974.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,572	28.5	4,264.2
2	Malignant neoplasms (C00-C97)	3,117	19.4	2,907.2
3	Cerebrovascular diseases (I60-I69)	879	5.5	819.8
4	Chronic lower respiratory diseases (J40-J47)	725	4.5	676.2
5	Alzheimer's disease (G30)	670	4.2	624.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	606	3.8	565.2
7	Diabetes mellitus (E10-E14)	514	3.2	479.4
8	Influenza and pneumonia (J09-J18)	492	3.1	458.9
9	Septicemia (A40-A41)	357	2.2	333.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	355	2.2	331.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	251	1.6	234.1
12	Pneumonitis due to solids and liquids (J69)	166	1.0	154.8
13	Parkinson's disease (G20-G21)	129	0.8	120.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.5	76.5
15	Atherosclerosis (I70)	60	0.4	56.0
...	All other causes (Residual)	3,080	19.2	2,872.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, all ages				
...	All causes	139,614	100.0	676.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,618	24.1	163.0
2	Malignant neoplasms (C00-C97)	31,669	22.7	153.5
3	Cerebrovascular diseases (I60-I69)	8,943	6.4	43.4
4	Diabetes mellitus (E10-E14)	6,424	4.6	31.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,785	3.4	23.2
6	Chronic lower respiratory diseases (J40-J47)	4,134	3.0	20.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,933	2.8	19.1
8	Alzheimer's disease (G30)	3,698	2.6	17.9
9	Septicemia (A40-A41)	3,285	2.4	15.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,863	2.1	13.9
11	Influenza and pneumonia (J09-J18)	2,514	1.8	12.2
12	Certain conditions originating in the perinatal period (P00-P96)	1,730	1.2	8.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,598	1.1	7.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,099	0.8	5.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	907	0.6	4.4
...	All other causes (Residual)	28,414	20.4	137.8
Non-Hispanic black, female, 1-4 years				
...	All causes	423	100.0	34.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	21.3	7.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	15.8	5.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	14.7	5.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	5.2	1.8
5	Malignant neoplasms (C00-C97)	21	5.0	1.7
6	Influenza and pneumonia (J09-J18)	10	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.1	*
8	Cerebrovascular diseases (I60-I69)	8	1.9	*
9	Septicemia (A40-A41)	7	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	7	1.7	*
11	Anemias (D50-D64)	5	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Meningitis (G00,G03)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	104	24.6	8.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 5-14 years				
...	All causes	469	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	22.4	3.5
2	Malignant neoplasms (C00-C97)	80	17.1	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.8	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	6.0	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.7	0.7
6	Chronic lower respiratory diseases (J40-J47)	20	4.3	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	4.3	0.7
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.7	*
8	Anemias (D50-D64)	8	1.7	*
10	Septicemia (A40-A41)	7	1.5	*
10	Cerebrovascular diseases (I60-I69)	7	1.5	*
12	Influenza and pneumonia (J09-J18)	4	0.9	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
...	All other causes (Residual)	116	24.7	3.8
Non-Hispanic black, female, 15-24 years				
...	All causes	1,513	100.0	45.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	373	24.7	11.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	263	17.4	7.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	8.1	3.7
4	Malignant neoplasms (C00-C97)	116	7.7	3.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	4.6	2.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	50	3.3	1.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	39	2.6	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	2.4	1.1
9	Cerebrovascular diseases (I60-I69)	28	1.9	0.8
10	Diabetes mellitus (E10-E14)	25	1.7	0.8
11	Anemias (D50-D64)	24	1.6	0.7
12	Chronic lower respiratory diseases (J40-J47)	19	1.3	*
13	Septicemia (A40-A41)	15	1.0	*
13	Influenza and pneumonia (J09-J18)	15	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.0	*
...	All other causes (Residual)	303	20.0	9.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 25-34 years				
...	All causes	2,815	100.0	99.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	423	15.0	14.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	336	11.9	11.8
3	Malignant neoplasms (C00-C97)	325	11.5	11.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	267	9.5	9.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	197	7.0	6.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	118	4.2	4.2
7	Diabetes mellitus (E10-E14)	94	3.3	3.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	2.5	2.4
9	Cerebrovascular diseases (I60-I69)	62	2.2	2.2
10	Anemias (D50-D64)	56	2.0	2.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	1.8	1.8
12	Septicemia (A40-A41)	50	1.8	1.8
13	Influenza and pneumonia (J09-J18)	34	1.2	1.2
14	Chronic lower respiratory diseases (J40-J47)	28	1.0	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	1.0	1.0
...	All other causes (Residual)	676	24.0	23.8
Non-Hispanic black, female, 35-44 years				
...	All causes	5,832	100.0	209.1
1	Malignant neoplasms (C00-C97)	1,301	22.3	46.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,040	17.8	37.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	489	8.4	17.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	418	7.2	15.0
5	Cerebrovascular diseases (I60-I69)	266	4.6	9.5
6	Diabetes mellitus (E10-E14)	207	3.5	7.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	175	3.0	6.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	125	2.1	4.5
9	Septicemia (A40-A41)	106	1.8	3.8
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	95	1.6	3.4
11	Chronic lower respiratory diseases (J40-J47)	90	1.5	3.2
12	Pregnancy, childbirth and the puerperium (O00-O99)	78	1.3	2.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	75	1.3	2.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	74	1.3	2.7
15	Anemias (D50-D64)	64	1.1	2.3
...	All other causes (Residual)	1,229	21.1	44.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 45-54 years				
...	All causes	14,552	100.0	497.4
1	Malignant neoplasms (C00-C97)	4,433	30.5	151.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,920	20.1	99.8
3	Cerebrovascular diseases (I60-I69)	806	5.5	27.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	737	5.1	25.2
5	Diabetes mellitus (E10-E14)	599	4.1	20.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	539	3.7	18.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	397	2.7	13.6
8	Chronic lower respiratory diseases (J40-J47)	359	2.5	12.3
9	Septicemia (A40-A41)	291	2.0	9.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	291	2.0	9.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	241	1.7	8.2
12	Influenza and pneumonia (J09-J18)	162	1.1	5.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	0.9	4.7
14	Viral hepatitis (B15-B19)	102	0.7	3.5
15	Anemias (D50-D64)	74	0.5	2.5
...	All other causes (Residual)	2,463	16.9	84.2
Non-Hispanic black, female, 55-64 years				
...	All causes	21,150	100.0	996.9
1	Malignant neoplasms (C00-C97)	7,202	34.1	339.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,757	22.5	224.2
3	Diabetes mellitus (E10-E14)	1,125	5.3	53.0
4	Cerebrovascular diseases (I60-I69)	1,118	5.3	52.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	685	3.2	32.3
6	Chronic lower respiratory diseases (J40-J47)	662	3.1	31.2
7	Septicemia (A40-A41)	523	2.5	24.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	498	2.4	23.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	378	1.8	17.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	308	1.5	14.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	293	1.4	13.8
12	Influenza and pneumonia (J09-J18)	252	1.2	11.9
13	Viral hepatitis (B15-B19)	149	0.7	7.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.4	4.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	0.4	3.7
...	All other causes (Residual)	3,031	14.3	142.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 65-74 years				
...	All causes	23,870	100.0	2,068.1
1	Malignant neoplasms (C00-C97)	7,463	31.3	646.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,615	23.5	486.5
3	Cerebrovascular diseases (I60-I69)	1,489	6.2	129.0
4	Diabetes mellitus (E10-E14)	1,417	5.9	122.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	981	4.1	85.0
6	Chronic lower respiratory diseases (J40-J47)	956	4.0	82.8
7	Septicemia (A40-A41)	671	2.8	58.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	461	1.9	39.9
9	Influenza and pneumonia (J09-J18)	374	1.6	32.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	311	1.3	26.9
11	Alzheimer's disease (G30)	234	1.0	20.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	134	0.6	11.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	120	0.5	10.4
14	Pneumonitis due to solids and liquids (J69)	89	0.4	7.7
15	Viral hepatitis (B15-B19)	73	0.3	6.3
...	All other causes (Residual)	3,482	14.6	301.7
Non-Hispanic black, female, 75-84 years				
...	All causes	30,661	100.0	4,675.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,200	26.7	1,250.4
2	Malignant neoplasms (C00-C97)	6,749	22.0	1,029.2
3	Cerebrovascular diseases (I60-I69)	2,331	7.6	355.5
4	Diabetes mellitus (E10-E14)	1,635	5.3	249.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,255	4.1	191.4
6	Chronic lower respiratory diseases (J40-J47)	1,097	3.6	167.3
7	Alzheimer's disease (G30)	1,044	3.4	159.2
8	Septicemia (A40-A41)	797	2.6	121.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	669	2.2	102.0
10	Influenza and pneumonia (J09-J18)	630	2.1	96.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	375	1.2	57.2
12	Pneumonitis due to solids and liquids (J69)	170	0.6	25.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	166	0.5	25.3
14	Parkinson's disease (G20-G21)	131	0.4	20.0
15	Aortic aneurysm and dissection (I71)	120	0.4	18.3
...	All other causes (Residual)	5,292	17.3	807.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 85 years and over				
...	All causes	35,187	100.0	12,767.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,550	30.0	3,828.1
2	Malignant neoplasms (C00-C97)	3,974	11.3	1,442.0
3	Cerebrovascular diseases (I60-I69)	2,801	8.0	1,016.4
4	Alzheimer's disease (G30)	2,387	6.8	866.1
5	Diabetes mellitus (E10-E14)	1,318	3.7	478.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,254	3.6	455.0
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,009	2.9	366.1
8	Influenza and pneumonia (J09-J18)	945	2.7	342.9
9	Chronic lower respiratory diseases (J40-J47)	892	2.5	323.7
10	Septicemia (A40-A41)	776	2.2	281.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	387	1.1	140.4
12	Pneumonitis due to solids and liquids (J69)	287	0.8	104.1
13	Atherosclerosis (I70)	169	0.5	61.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	163	0.5	59.1
15	Parkinson's disease (G20-G21)	147	0.4	53.3
...	All other causes (Residual)	8,128	23.1	2,949.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable