

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	163,241	100.0	301.9
1	Malignant neoplasms (C00-C97)	35,147	21.5	65.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,243	20.4	61.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,015	7.4	22.2
4	Cerebrovascular diseases (I60-I69)	8,127	5.0	15.0
5	Diabetes mellitus (E10-E14)	7,632	4.7	14.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	5,141	3.1	9.5
7	Chronic lower respiratory diseases (J40-J47)	4,827	3.0	8.9
8	Alzheimer's disease (G30)	4,127	2.5	7.6
9	Influenza and pneumonia (J09-J18)	3,592	2.2	6.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,083	1.9	5.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,865	1.8	5.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,571	1.6	4.8
13	Septicemia (A40-A41)	2,435	1.5	4.5
14	Certain conditions originating in the perinatal period (P00-P96)	2,428	1.5	4.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,092	1.3	3.9
...	All other causes (Residual)	33,916	20.8	62.7
Hispanic, both sexes, 1-4 years				
...	All causes	854	100.0	20.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	32.6	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	14.3	3.0
3	Malignant neoplasms (C00-C97)	90	10.5	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	6.2	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.7	0.6
6	Influenza and pneumonia (J09-J18)	19	2.2	*
7	Septicemia (A40-A41)	9	1.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.1	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.1	*
11	Anemias (D50-D64)	5	0.6	*
11	Meningitis (G00,G03)	5	0.6	*
13	Cerebrovascular diseases (I60-I69)	4	0.5	*
14	Diseases of appendix (K35-K38)	3	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	213	24.9	5.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,073	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	26.0	2.8
2	Malignant neoplasms (C00-C97)	209	19.5	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	7.4	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	6.6	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	5.5	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.9	0.3
7	Cerebrovascular diseases (I60-I69)	26	2.4	0.3
8	Influenza and pneumonia (J09-J18)	22	2.1	0.2
8	Chronic lower respiratory diseases (J40-J47)	22	2.1	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.1	*
11	Septicemia (A40-A41)	10	0.9	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
15	Diabetes mellitus (E10-E14)	4	0.4	*
...	All other causes (Residual)	232	21.6	2.3
Hispanic, both sexes, 15-24 years				
...	All causes	4,657	100.0	50.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,944	41.7	21.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	849	18.2	9.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	635	13.6	6.8
4	Malignant neoplasms (C00-C97)	339	7.3	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	2.6	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.2	0.6
7	Cerebrovascular diseases (I60-I69)	32	0.7	0.3
8	Influenza and pneumonia (J09-J18)	27	0.6	0.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	27	0.6	0.3
10	Septicemia (A40-A41)	23	0.5	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.5	0.2
12	Diabetes mellitus (E10-E14)	18	0.4	*
13	Legal intervention (Y35,Y89.0)	17	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.3	*
15	Chronic lower respiratory diseases (J40-J47)	14	0.3	*
...	All other causes (Residual)	520	11.2	5.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	6,341	100.0	72.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,203	34.7	25.1
2	Malignant neoplasms (C00-C97)	724	11.4	8.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	724	11.4	8.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	687	10.8	7.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	375	5.9	4.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	128	2.0	1.5
7	Cerebrovascular diseases (I60-I69)	107	1.7	1.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	103	1.6	1.2
9	Diabetes mellitus (E10-E14)	88	1.4	1.0
10	Influenza and pneumonia (J09-J18)	70	1.1	0.8
11	Pregnancy, childbirth and the puerperium (O00-O99)	53	0.8	0.6
12	Septicemia (A40-A41)	50	0.8	0.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	0.8	0.5
14	Chronic lower respiratory diseases (J40-J47)	41	0.6	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.5	0.4
...	All other causes (Residual)	906	14.3	10.3
Hispanic, both sexes, 35-44 years				
...	All causes	8,698	100.0	111.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,914	22.0	24.6
2	Malignant neoplasms (C00-C97)	1,619	18.6	20.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	992	11.4	12.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	525	6.0	6.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	483	5.6	6.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	414	4.8	5.3
7	Cerebrovascular diseases (I60-I69)	307	3.5	3.9
8	Diabetes mellitus (E10-E14)	266	3.1	3.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	192	2.2	2.5
10	Influenza and pneumonia (J09-J18)	123	1.4	1.6
11	Septicemia (A40-A41)	92	1.1	1.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	1.0	1.1
13	Viral hepatitis (B15-B19)	60	0.7	0.8
14	Pregnancy, childbirth and the puerperium (O00-O99)	56	0.6	0.7
15	Chronic lower respiratory diseases (J40-J47)	54	0.6	0.7
...	All other causes (Residual)	1,514	17.4	19.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	16,394	100.0	269.6
1	Malignant neoplasms (C00-C97)	4,119	25.1	67.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,620	16.0	43.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,860	11.3	30.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,402	8.6	23.1
5	Diabetes mellitus (E10-E14)	734	4.5	12.1
6	Cerebrovascular diseases (I60-I69)	696	4.2	11.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	441	2.7	7.3
8	Viral hepatitis (B15-B19)	351	2.1	5.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	325	2.0	5.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	256	1.6	4.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	245	1.5	4.0
12	Septicemia (A40-A41)	231	1.4	3.8
13	Influenza and pneumonia (J09-J18)	224	1.4	3.7
14	Chronic lower respiratory diseases (J40-J47)	158	1.0	2.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	126	0.8	2.1
...	All other causes (Residual)	2,606	15.9	42.9
Hispanic, both sexes, 55-64 years				
...	All causes	23,326	100.0	610.7
1	Malignant neoplasms (C00-C97)	7,261	31.1	190.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,525	19.4	118.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,467	6.3	38.4
4	Diabetes mellitus (E10-E14)	1,425	6.1	37.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,262	5.4	33.0
6	Cerebrovascular diseases (I60-I69)	1,012	4.3	26.5
7	Viral hepatitis (B15-B19)	514	2.2	13.5
8	Chronic lower respiratory diseases (J40-J47)	471	2.0	12.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	433	1.9	11.3
10	Influenza and pneumonia (J09-J18)	378	1.6	9.9
11	Septicemia (A40-A41)	369	1.6	9.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	279	1.2	7.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	229	1.0	6.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	196	0.8	5.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	103	0.4	2.7
...	All other causes (Residual)	3,402	14.6	89.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	26,684	100.0	1,342.8
1	Malignant neoplasms (C00-C97)	8,355	31.3	420.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,873	22.0	295.5
3	Diabetes mellitus (E10-E14)	1,751	6.6	88.1
4	Cerebrovascular diseases (I60-I69)	1,321	5.0	66.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	923	3.5	46.4
6	Chronic lower respiratory diseases (J40-J47)	857	3.2	43.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	693	2.6	34.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	676	2.5	34.0
9	Influenza and pneumonia (J09-J18)	579	2.2	29.1
10	Septicemia (A40-A41)	444	1.7	22.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	317	1.2	16.0
12	Alzheimer's disease (G30)	297	1.1	14.9
13	Viral hepatitis (B15-B19)	211	0.8	10.6
14	Parkinson's disease (G20-G21)	202	0.8	10.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	141	0.5	7.1
...	All other causes (Residual)	4,044	15.2	203.5
Hispanic, both sexes, 75-84 years				
...	All causes	34,766	100.0	3,524.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,372	24.1	848.8
2	Malignant neoplasms (C00-C97)	8,083	23.2	819.5
3	Cerebrovascular diseases (I60-I69)	2,210	6.4	224.1
4	Diabetes mellitus (E10-E14)	1,948	5.6	197.5
5	Chronic lower respiratory diseases (J40-J47)	1,567	4.5	158.9
6	Alzheimer's disease (G30)	1,229	3.5	124.6
7	Influenza and pneumonia (J09-J18)	900	2.6	91.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	842	2.4	85.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	697	2.0	70.7
10	Septicemia (A40-A41)	646	1.9	65.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	567	1.6	57.5
12	Parkinson's disease (G20-G21)	561	1.6	56.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	554	1.6	56.2
14	Pneumonitis due to solids and liquids (J69)	242	0.7	24.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	230	0.7	23.3
...	All other causes (Residual)	6,118	17.6	620.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	35,690	100.0	9,986.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,254	28.7	2,869.3
2	Malignant neoplasms (C00-C97)	4,336	12.1	1,213.3
3	Alzheimer's disease (G30)	2,535	7.1	709.4
4	Cerebrovascular diseases (I60-I69)	2,397	6.7	670.7
5	Chronic lower respiratory diseases (J40-J47)	1,629	4.6	455.8
6	Diabetes mellitus (E10-E14)	1,397	3.9	390.9
7	Influenza and pneumonia (J09-J18)	1,215	3.4	340.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	788	2.2	220.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	728	2.0	203.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	716	2.0	200.4
11	Septicemia (A40-A41)	526	1.5	147.2
12	Parkinson's disease (G20-G21)	488	1.4	136.6
13	Pneumonitis due to solids and liquids (J69)	325	0.9	90.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	193	0.5	54.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	175	0.5	49.0
...	All other causes (Residual)	7,988	22.4	2,235.2
Hispanic, male, all ages				
...	All causes	88,880	100.0	323.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,377	20.7	66.9
2	Malignant neoplasms (C00-C97)	18,371	20.7	66.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,760	9.9	31.9
4	Diabetes mellitus (E10-E14)	3,934	4.4	14.3
5	Cerebrovascular diseases (I60-I69)	3,841	4.3	14.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,552	4.0	12.9
7	Chronic lower respiratory diseases (J40-J47)	2,536	2.9	9.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,279	2.6	8.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,132	2.4	7.8
10	Influenza and pneumonia (J09-J18)	1,816	2.0	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,583	1.8	5.8
12	Certain conditions originating in the perinatal period (P00-P96)	1,320	1.5	4.8
13	Alzheimer's disease (G30)	1,298	1.5	4.7
14	Septicemia (A40-A41)	1,226	1.4	4.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	972	1.1	3.5
...	All other causes (Residual)	16,883	19.0	61.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	483	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	37.7	8.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	13.0	3.0
3	Malignant neoplasms (C00-C97)	48	9.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	6.4	1.5
5	Influenza and pneumonia (J09-J18)	9	1.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	1.7	*
7	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
8	Septicemia (A40-A41)	4	0.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	4	0.8	*
10	Meningitis (G00,G03)	3	0.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
11	Anemias (D50-D64)	2	0.4	*
11	Diseases of appendix (K35-K38)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Cerebrovascular diseases (I60-I69)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
...	All other causes (Residual)	113	23.4	5.4
Hispanic, male, 5-14 years				
...	All causes	591	100.0	11.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	29.3	3.4
2	Malignant neoplasms (C00-C97)	106	17.9	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.4	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	6.1	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	5.6	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.0	*
7	Chronic lower respiratory diseases (J40-J47)	15	2.5	*
8	Cerebrovascular diseases (I60-I69)	10	1.7	*
9	Influenza and pneumonia (J09-J18)	7	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
11	Septicemia (A40-A41)	5	0.8	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.8	*
13	Anemias (D50-D64)	2	0.3	*
13	Meningitis (G00,G03)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	131	22.2	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,516	100.0	72.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,513	43.0	31.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	740	21.0	15.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	487	13.9	10.1
4	Malignant neoplasms (C00-C97)	209	5.9	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	86	2.4	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	0.9	0.7
7	Legal intervention (Y35,Y89.0)	17	0.5	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.4	*
9	Cerebrovascular diseases (I60-I69)	15	0.4	*
11	Influenza and pneumonia (J09-J18)	14	0.4	*
12	Septicemia (A40-A41)	13	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	10	0.3	*
14	Diabetes mellitus (E10-E14)	9	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.2	*
...	All other causes (Residual)	332	9.4	6.9
Hispanic, male, 25-34 years				
...	All causes	4,632	100.0	99.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,771	38.2	38.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	616	13.3	13.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	572	12.3	12.3
4	Malignant neoplasms (C00-C97)	363	7.8	7.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	261	5.6	5.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	94	2.0	2.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	84	1.8	1.8
8	Cerebrovascular diseases (I60-I69)	69	1.5	1.5
9	Diabetes mellitus (E10-E14)	57	1.2	1.2
10	Influenza and pneumonia (J09-J18)	41	0.9	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	0.7	0.7
11	Legal intervention (Y35,Y89.0)	31	0.7	0.7
13	Septicemia (A40-A41)	30	0.6	0.6
14	Chronic lower respiratory diseases (J40-J47)	23	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.5	0.5
...	All other causes (Residual)	567	12.2	12.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,728	100.0	143.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,479	25.8	37.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	758	13.2	19.0
3	Malignant neoplasms (C00-C97)	655	11.4	16.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	421	7.3	10.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	374	6.5	9.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	335	5.8	8.4
7	Cerebrovascular diseases (I60-I69)	183	3.2	4.6
8	Diabetes mellitus (E10-E14)	166	2.9	4.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	157	2.7	3.9
10	Influenza and pneumonia (J09-J18)	79	1.4	2.0
11	Septicemia (A40-A41)	54	0.9	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	0.9	1.3
13	Viral hepatitis (B15-B19)	45	0.8	1.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	0.5	0.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.5	0.7
...	All other causes (Residual)	912	15.9	22.8
Hispanic, male, 45-54 years				
...	All causes	10,683	100.0	348.0
1	Malignant neoplasms (C00-C97)	2,039	19.1	66.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,950	18.3	63.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,430	13.4	46.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,074	10.1	35.0
5	Diabetes mellitus (E10-E14)	480	4.5	15.6
6	Cerebrovascular diseases (I60-I69)	423	4.0	13.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	334	3.1	10.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	264	2.5	8.6
9	Viral hepatitis (B15-B19)	262	2.5	8.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	216	2.0	7.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	1.4	4.8
12	Influenza and pneumonia (J09-J18)	145	1.4	4.7
13	Septicemia (A40-A41)	127	1.2	4.1
14	Chronic lower respiratory diseases (J40-J47)	97	0.9	3.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	0.7	2.6
...	All other causes (Residual)	1,614	15.1	52.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	14,674	100.0	797.4
1	Malignant neoplasms (C00-C97)	4,001	27.3	217.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,276	22.3	178.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,106	7.5	60.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	944	6.4	51.3
5	Diabetes mellitus (E10-E14)	823	5.6	44.7
6	Cerebrovascular diseases (I60-I69)	623	4.2	33.9
7	Viral hepatitis (B15-B19)	368	2.5	20.0
8	Chronic lower respiratory diseases (J40-J47)	266	1.8	14.5
9	Influenza and pneumonia (J09-J18)	239	1.6	13.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	233	1.6	12.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	225	1.5	12.2
12	Septicemia (A40-A41)	204	1.4	11.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	153	1.0	8.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	139	0.9	7.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	0.6	4.5
...	All other causes (Residual)	1,991	13.6	108.2
Hispanic, male, 65-74 years				
...	All causes	15,355	100.0	1,710.4
1	Malignant neoplasms (C00-C97)	4,629	30.1	515.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,740	24.4	416.6
3	Diabetes mellitus (E10-E14)	958	6.2	106.7
4	Cerebrovascular diseases (I60-I69)	737	4.8	82.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	563	3.7	62.7
6	Chronic lower respiratory diseases (J40-J47)	513	3.3	57.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	462	3.0	51.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	344	2.2	38.3
9	Influenza and pneumonia (J09-J18)	337	2.2	37.5
10	Septicemia (A40-A41)	241	1.6	26.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	184	1.2	20.5
12	Alzheimer's disease (G30)	145	0.9	16.2
13	Parkinson's disease (G20-G21)	137	0.9	15.3
14	Viral hepatitis (B15-B19)	122	0.8	13.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	0.7	11.9
...	All other causes (Residual)	2,136	13.9	237.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	17,160	100.0	4,218.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,302	25.1	1,057.5
2	Malignant neoplasms (C00-C97)	4,277	24.9	1,051.3
3	Cerebrovascular diseases (I60-I69)	985	5.7	242.1
4	Diabetes mellitus (E10-E14)	903	5.3	222.0
5	Chronic lower respiratory diseases (J40-J47)	833	4.9	204.8
6	Alzheimer's disease (G30)	450	2.6	110.6
6	Influenza and pneumonia (J09-J18)	450	2.6	110.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	431	2.5	105.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	378	2.2	92.9
10	Parkinson's disease (G20-G21)	333	1.9	81.9
11	Septicemia (A40-A41)	315	1.8	77.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	268	1.6	65.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	260	1.5	63.9
14	Pneumonitis due to solids and liquids (J69)	142	0.8	34.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	122	0.7	30.0
...	All other causes (Residual)	2,711	15.8	666.4
Hispanic, male, 85 years and over				
...	All causes	13,454	100.0	10,596.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,944	29.3	3,106.2
2	Malignant neoplasms (C00-C97)	2,040	15.2	1,606.6
3	Cerebrovascular diseases (I60-I69)	788	5.9	620.6
4	Chronic lower respiratory diseases (J40-J47)	744	5.5	586.0
5	Alzheimer's disease (G30)	673	5.0	530.0
6	Diabetes mellitus (E10-E14)	536	4.0	422.1
7	Influenza and pneumonia (J09-J18)	472	3.5	371.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	335	2.5	263.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	330	2.5	259.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	253	1.9	199.3
11	Parkinson's disease (G20-G21)	251	1.9	197.7
12	Septicemia (A40-A41)	211	1.6	166.2
13	Pneumonitis due to solids and liquids (J69)	145	1.1	114.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.6	59.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	73	0.5	57.5
...	All other causes (Residual)	2,583	19.2	2,034.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	74,361	100.0	279.4
1	Malignant neoplasms (C00-C97)	16,776	22.6	63.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,866	20.0	55.9
3	Cerebrovascular diseases (I60-I69)	4,286	5.8	16.1
4	Diabetes mellitus (E10-E14)	3,698	5.0	13.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,255	4.4	12.2
6	Alzheimer's disease (G30)	2,829	3.8	10.6
7	Chronic lower respiratory diseases (J40-J47)	2,291	3.1	8.6
8	Influenza and pneumonia (J09-J18)	1,776	2.4	6.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,589	2.1	6.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,500	2.0	5.6
11	Septicemia (A40-A41)	1,209	1.6	4.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,120	1.5	4.2
13	Certain conditions originating in the perinatal period (P00-P96)	1,108	1.5	4.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	813	1.1	3.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	586	0.8	2.2
...	All other causes (Residual)	16,659	22.4	62.6
Hispanic, female, 1-4 years				
...	All causes	371	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	25.9	4.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	15.9	2.9
3	Malignant neoplasms (C00-C97)	42	11.3	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	5.9	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	4.0	*
6	Influenza and pneumonia (J09-J18)	10	2.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.9	*
8	Septicemia (A40-A41)	5	1.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	5	1.3	*
10	Anemias (D50-D64)	3	0.8	*
10	Cerebrovascular diseases (I60-I69)	3	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
13	Meningitis (G00,G03)	2	0.5	*
14	Shigellosis and amebiasis (A03,A06)	1	0.3	*
14	Tuberculosis (A16-A19)	1	0.3	*
14	Meningococcal infection (A39)	1	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
14	Diseases of appendix (K35-K38)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	93	25.1	4.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	482	100.0	9.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	22.0	2.2
2	Malignant neoplasms (C00-C97)	103	21.4	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	8.5	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	7.3	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	5.4	0.5
6	Cerebrovascular diseases (I60-I69)	16	3.3	*
7	Influenza and pneumonia (J09-J18)	15	3.1	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.7	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
11	Septicemia (A40-A41)	5	1.0	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.8	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	97	20.1	2.0
Hispanic, female, 15-24 years				
...	All causes	1,141	100.0	25.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	431	37.8	9.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	13.0	3.3
3	Malignant neoplasms (C00-C97)	130	11.4	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	9.6	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.0	0.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	27	2.4	0.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	2.0	0.5
8	Cerebrovascular diseases (I60-I69)	17	1.5	*
9	Influenza and pneumonia (J09-J18)	13	1.1	*
10	Septicemia (A40-A41)	10	0.9	*
11	Diabetes mellitus (E10-E14)	9	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
13	Anemias (D50-D64)	5	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	169	14.8	3.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,709	100.0	41.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	432	25.3	10.5
2	Malignant neoplasms (C00-C97)	361	21.1	8.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	6.7	2.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	114	6.7	2.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	108	6.3	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	53	3.1	1.3
7	Cerebrovascular diseases (I60-I69)	38	2.2	0.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	34	2.0	0.8
9	Diabetes mellitus (E10-E14)	31	1.8	0.8
10	Influenza and pneumonia (J09-J18)	29	1.7	0.7
11	Septicemia (A40-A41)	20	1.2	0.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	19	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	18	1.1	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.7	*
...	All other causes (Residual)	308	18.0	7.5
Hispanic, female, 35-44 years				
...	All causes	2,970	100.0	78.3
1	Malignant neoplasms (C00-C97)	964	32.5	25.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	435	14.6	11.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	234	7.9	6.2
4	Cerebrovascular diseases (I60-I69)	124	4.2	3.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	109	3.7	2.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	104	3.5	2.7
7	Diabetes mellitus (E10-E14)	100	3.4	2.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	2.7	2.1
9	Pregnancy, childbirth and the puerperium (O00-O99)	56	1.9	1.5
10	Influenza and pneumonia (J09-J18)	44	1.5	1.2
11	Septicemia (A40-A41)	38	1.3	1.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	35	1.2	0.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.2	0.9
14	Chronic lower respiratory diseases (J40-J47)	27	0.9	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.6	*
...	All other causes (Residual)	569	19.2	15.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	5,711	100.0	189.6
1	Malignant neoplasms (C00-C97)	2,080	36.4	69.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	670	11.7	22.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	430	7.5	14.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	328	5.7	10.9
5	Cerebrovascular diseases (I60-I69)	273	4.8	9.1
6	Diabetes mellitus (E10-E14)	254	4.4	8.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	1.9	3.6
8	Septicemia (A40-A41)	104	1.8	3.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	97	1.7	3.2
10	Viral hepatitis (B15-B19)	89	1.6	3.0
11	Influenza and pneumonia (J09-J18)	79	1.4	2.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.1	2.0
12	Chronic lower respiratory diseases (J40-J47)	61	1.1	2.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	0.8	1.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	0.7	1.3
...	All other causes (Residual)	992	17.4	32.9
Hispanic, female, 55-64 years				
...	All causes	8,652	100.0	437.2
1	Malignant neoplasms (C00-C97)	3,260	37.7	164.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,249	14.4	63.1
3	Diabetes mellitus (E10-E14)	602	7.0	30.4
4	Cerebrovascular diseases (I60-I69)	389	4.5	19.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	361	4.2	18.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	318	3.7	16.1
7	Chronic lower respiratory diseases (J40-J47)	205	2.4	10.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	200	2.3	10.1
9	Septicemia (A40-A41)	165	1.9	8.3
10	Viral hepatitis (B15-B19)	146	1.7	7.4
11	Influenza and pneumonia (J09-J18)	139	1.6	7.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	90	1.0	4.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	0.6	2.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	43	0.5	2.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.5	2.0
...	All other causes (Residual)	1,392	16.1	70.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	11,329	100.0	1,039.9
1	Malignant neoplasms (C00-C97)	3,726	32.9	342.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,133	18.8	195.8
3	Diabetes mellitus (E10-E14)	793	7.0	72.8
4	Cerebrovascular diseases (I60-I69)	584	5.2	53.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	360	3.2	33.0
6	Chronic lower respiratory diseases (J40-J47)	344	3.0	31.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	332	2.9	30.5
8	Influenza and pneumonia (J09-J18)	242	2.1	22.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	2.0	21.2
10	Septicemia (A40-A41)	203	1.8	18.6
11	Alzheimer's disease (G30)	152	1.3	14.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	133	1.2	12.2
13	Viral hepatitis (B15-B19)	89	0.8	8.2
14	Parkinson's disease (G20-G21)	65	0.6	6.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	61	0.5	5.6
...	All other causes (Residual)	1,881	16.6	172.7
Hispanic, female, 75-84 years				
...	All causes	17,606	100.0	3,037.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,070	23.1	702.3
2	Malignant neoplasms (C00-C97)	3,806	21.6	656.7
3	Cerebrovascular diseases (I60-I69)	1,225	7.0	211.4
4	Diabetes mellitus (E10-E14)	1,045	5.9	180.3
5	Alzheimer's disease (G30)	779	4.4	134.4
6	Chronic lower respiratory diseases (J40-J47)	734	4.2	126.6
7	Influenza and pneumonia (J09-J18)	450	2.6	77.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	411	2.3	70.9
9	Septicemia (A40-A41)	331	1.9	57.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	319	1.8	55.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	299	1.7	51.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	294	1.7	50.7
13	Parkinson's disease (G20-G21)	228	1.3	39.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	108	0.6	18.6
15	Pneumonitis due to solids and liquids (J69)	100	0.6	17.3
...	All other causes (Residual)	3,407	19.4	587.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	22,236	100.0	9,651.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,310	28.4	2,738.8
2	Malignant neoplasms (C00-C97)	2,296	10.3	996.6
3	Alzheimer's disease (G30)	1,862	8.4	808.2
4	Cerebrovascular diseases (I60-I69)	1,609	7.2	698.4
5	Chronic lower respiratory diseases (J40-J47)	885	4.0	384.1
6	Diabetes mellitus (E10-E14)	861	3.9	373.7
7	Influenza and pneumonia (J09-J18)	743	3.3	322.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	535	2.4	232.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	398	1.8	172.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	381	1.7	165.4
11	Septicemia (A40-A41)	315	1.4	136.7
12	Parkinson's disease (G20-G21)	237	1.1	102.9
13	Pneumonitis due to solids and liquids (J69)	180	0.8	78.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	117	0.5	50.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	102	0.5	44.3
...	All other causes (Residual)	5,405	24.3	2,346.0
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,052,660	100.0	1,021.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	488,817	23.8	243.3
2	Malignant neoplasms (C00-C97)	462,850	22.5	230.4
3	Chronic lower respiratory diseases (J40-J47)	131,667	6.4	65.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100,773	4.9	50.2
5	Cerebrovascular diseases (I60-I69)	99,759	4.9	49.7
6	Alzheimer's disease (G30)	73,234	3.6	36.4
7	Diabetes mellitus (E10-E14)	51,297	2.5	25.5
8	Influenza and pneumonia (J09-J18)	45,350	2.2	22.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34,253	1.7	17.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,173	1.7	17.0
11	Septicemia (A40-A41)	28,431	1.4	14.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26,692	1.3	13.3
13	Parkinson's disease (G20-G21)	22,241	1.1	11.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22,043	1.1	11.0
15	Pneumonitis due to solids and liquids (J69)	15,630	0.8	7.8
...	All other causes (Residual)	415,450	20.2	206.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,977	100.0	23.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	675	34.1	8.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	11.5	2.7
3	Malignant neoplasms (C00-C97)	160	8.1	1.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	128	6.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	4.1	1.0
6	Influenza and pneumonia (J09-J18)	53	2.7	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	1.5	0.4
8	Chronic lower respiratory diseases (J40-J47)	24	1.2	0.3
9	Septicemia (A40-A41)	22	1.1	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	22	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	15	0.8	*
12	Meningitis (G00,G03)	9	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	9	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
15	Anemias (D50-D64)	6	0.3	*
...	All other causes (Residual)	508	25.7	6.1
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,824	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	838	29.7	3.7
2	Malignant neoplasms (C00-C97)	512	18.1	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	244	8.6	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	5.9	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	4.3	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	2.7	0.3
7	Influenza and pneumonia (J09-J18)	71	2.5	0.3
8	Cerebrovascular diseases (I60-I69)	42	1.5	0.2
9	Chronic lower respiratory diseases (J40-J47)	38	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.2	0.2
11	Septicemia (A40-A41)	28	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	19	0.7	*
13	Diabetes mellitus (E10-E14)	18	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
15	Meningitis (G00,G03)	7	0.2	*
15	Pneumonitis due to solids and liquids (J69)	7	0.2	*
...	All other causes (Residual)	591	20.9	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	16,092	100.0	64.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,820	48.6	31.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,388	21.1	13.6
3	Malignant neoplasms (C00-C97)	804	5.0	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	687	4.3	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	485	3.0	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	1.4	0.9
7	Influenza and pneumonia (J09-J18)	132	0.8	0.5
8	Diabetes mellitus (E10-E14)	95	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	76	0.5	0.3
10	Cerebrovascular diseases (I60-I69)	75	0.5	0.3
11	Septicemia (A40-A41)	70	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	54	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.3	0.2
14	Legal intervention (Y35,Y89.0)	34	0.2	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.2	0.1
...	All other causes (Residual)	2,070	12.9	8.3
Non-Hispanic, white, both sexes, 25-34 years				
...	All causes	27,583	100.0	110.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,594	42.0	46.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,725	17.1	18.9
3	Malignant neoplasms (C00-C97)	2,071	7.5	8.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,773	6.4	7.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	924	3.3	3.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	416	1.5	1.7
7	Diabetes mellitus (E10-E14)	342	1.2	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	290	1.1	1.2
9	Influenza and pneumonia (J09-J18)	266	1.0	1.1
10	Cerebrovascular diseases (I60-I69)	237	0.9	0.9
11	Pregnancy, childbirth and the puerperium (O00-O99)	209	0.8	0.8
12	Septicemia (A40-A41)	176	0.6	0.7
13	Chronic lower respiratory diseases (J40-J47)	143	0.5	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	0.4	0.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	116	0.4	0.5
...	All other causes (Residual)	4,181	15.2	16.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-44 years				
...	All causes	43,931	100.0	181.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,120	25.3	45.8
2	Malignant neoplasms (C00-C97)	7,101	16.2	29.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,141	14.0	25.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,298	12.1	21.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,600	3.6	6.6
6	Diabetes mellitus (E10-E14)	1,062	2.4	4.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	832	1.9	3.4
8	Cerebrovascular diseases (I60-I69)	758	1.7	3.1
9	Influenza and pneumonia (J09-J18)	558	1.3	2.3
10	Chronic lower respiratory diseases (J40-J47)	477	1.1	2.0
11	Septicemia (A40-A41)	469	1.1	1.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	345	0.8	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	311	0.7	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	266	0.6	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	171	0.4	0.7
...	All other causes (Residual)	7,422	16.9	30.6
Non-Hispanic, white, both sexes, 45-54 years				
...	All causes	122,531	100.0	415.4
1	Malignant neoplasms (C00-C97)	32,434	26.5	110.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,715	19.4	80.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,377	12.5	52.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,527	6.1	25.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,218	5.1	21.1
6	Chronic lower respiratory diseases (J40-J47)	3,708	3.0	12.6
7	Diabetes mellitus (E10-E14)	3,486	2.8	11.8
8	Cerebrovascular diseases (I60-I69)	2,913	2.4	9.9
9	Septicemia (A40-A41)	1,477	1.2	5.0
9	Influenza and pneumonia (J09-J18)	1,477	1.2	5.0
11	Viral hepatitis (B15-B19)	1,255	1.0	4.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,032	0.8	3.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	895	0.7	3.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	820	0.7	2.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	786	0.6	2.7
...	All other causes (Residual)	19,411	15.8	65.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-64 years				
...	All causes	245,518	100.0	849.7
1	Malignant neoplasms (C00-C97)	84,411	34.4	292.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,348	20.9	177.7
3	Chronic lower respiratory diseases (J40-J47)	13,388	5.5	46.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,901	5.3	44.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	8,814	3.6	30.5
6	Diabetes mellitus (E10-E14)	8,176	3.3	28.3
7	Cerebrovascular diseases (I60-I69)	6,995	2.8	24.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,411	2.6	22.2
9	Septicemia (A40-A41)	3,665	1.5	12.7
10	Influenza and pneumonia (J09-J18)	3,337	1.4	11.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,793	1.1	9.7
12	Viral hepatitis (B15-B19)	2,629	1.1	9.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,890	0.8	6.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,134	0.5	3.9
15	Aortic aneurysm and dissection (I71)	956	0.4	3.3
...	All other causes (Residual)	36,670	14.9	126.9
Non-Hispanic, white, both sexes, 65-74 years				
...	All causes	355,949	100.0	1,811.8
1	Malignant neoplasms (C00-C97)	124,636	35.0	634.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,980	21.1	381.7
3	Chronic lower respiratory diseases (J40-J47)	31,661	8.9	161.2
4	Cerebrovascular diseases (I60-I69)	13,353	3.8	68.0
5	Diabetes mellitus (E10-E14)	11,629	3.3	59.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,806	2.5	44.8
7	Septicemia (A40-A41)	5,666	1.6	28.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,655	1.6	28.8
9	Influenza and pneumonia (J09-J18)	5,515	1.5	28.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5,393	1.5	27.5
11	Alzheimer's disease (G30)	3,759	1.1	19.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,468	1.0	17.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,764	0.8	14.1
14	Parkinson's disease (G20-G21)	2,742	0.8	14.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,297	0.6	11.7
...	All other causes (Residual)	53,625	15.1	273.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-84 years				
...	All causes	512,252	100.0	4,769.4
1	Malignant neoplasms (C00-C97)	126,182	24.6	1,174.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	119,291	23.3	1,110.7
3	Chronic lower respiratory diseases (J40-J47)	44,318	8.7	412.6
4	Cerebrovascular diseases (I60-I69)	28,507	5.6	265.4
5	Alzheimer's disease (G30)	19,636	3.8	182.8
6	Diabetes mellitus (E10-E14)	13,725	2.7	127.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,453	2.4	115.9
8	Influenza and pneumonia (J09-J18)	11,162	2.2	103.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,948	1.9	92.6
10	Parkinson's disease (G20-G21)	9,307	1.8	86.7
11	Septicemia (A40-A41)	7,929	1.5	73.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,105	1.0	47.5
13	Pneumonitis due to solids and liquids (J69)	4,119	0.8	38.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,911	0.8	36.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,115	0.6	29.0
...	All other causes (Residual)	93,544	18.3	871.0
Non-Hispanic, white, both sexes, 85 years and over				
...	All causes	713,390	100.0	14,176.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	210,771	29.5	4,188.5
2	Malignant neoplasms (C00-C97)	84,491	11.8	1,679.0
3	Alzheimer's disease (G30)	49,031	6.9	974.4
4	Cerebrovascular diseases (I60-I69)	46,814	6.6	930.3
5	Chronic lower respiratory diseases (J40-J47)	37,849	5.3	752.2
6	Influenza and pneumonia (J09-J18)	22,713	3.2	451.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,639	2.6	370.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,281	2.0	283.8
9	Diabetes mellitus (E10-E14)	12,756	1.8	253.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,273	1.6	224.0
11	Parkinson's disease (G20-G21)	9,657	1.4	191.9
12	Septicemia (A40-A41)	8,871	1.2	176.3
13	Pneumonitis due to solids and liquids (J69)	8,118	1.1	161.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,546	0.6	90.3
15	Atherosclerosis (I70)	3,397	0.5	67.5
...	All other causes (Residual)	170,183	23.9	3,381.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	1,021,135	100.0	1,032.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	256,294	25.1	259.0
2	Malignant neoplasms (C00-C97)	244,646	24.0	247.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61,391	6.0	62.1
4	Chronic lower respiratory diseases (J40-J47)	61,106	6.0	61.8
5	Cerebrovascular diseases (I60-I69)	40,350	4.0	40.8
6	Diabetes mellitus (E10-E14)	27,807	2.7	28.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,621	2.6	26.9
8	Alzheimer's disease (G30)	22,345	2.2	22.6
9	Influenza and pneumonia (J09-J18)	21,059	2.1	21.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	17,290	1.7	17.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,213	1.7	17.4
12	Parkinson's disease (G20-G21)	13,370	1.3	13.5
13	Septicemia (A40-A41)	13,364	1.3	13.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,955	0.9	9.1
15	Pneumonitis due to solids and liquids (J69)	8,534	0.8	8.6
...	All other causes (Residual)	180,790	17.7	182.7
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,149	100.0	26.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	437	38.0	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	8.8	2.4
3	Malignant neoplasms (C00-C97)	96	8.4	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	5.8	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	4.2	1.1
6	Influenza and pneumonia (J09-J18)	31	2.7	0.7
7	Certain conditions originating in the perinatal period (P00-P96)	18	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
9	Septicemia (A40-A41)	13	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	13	1.1	*
11	Cerebrovascular diseases (I60-I69)	8	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
13	Anemias (D50-D64)	4	0.3	*
13	Meningitis (G00,G03)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	285	24.8	6.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,657	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	519	31.3	4.5
2	Malignant neoplasms (C00-C97)	266	16.1	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	169	10.2	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	5.7	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	4.3	0.6
6	Influenza and pneumonia (J09-J18)	41	2.5	0.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	2.2	0.3
8	Chronic lower respiratory diseases (J40-J47)	24	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	21	1.3	0.2
11	Septicemia (A40-A41)	16	1.0	*
12	Certain conditions originating in the perinatal period (P00-P96)	11	0.7	*
13	Diabetes mellitus (E10-E14)	10	0.6	*
14	Meningitis (G00,G03)	6	0.4	*
14	Aortic aneurysm and dissection (I71)	6	0.4	*
...	All other causes (Residual)	342	20.6	3.0
Non-Hispanic, white, male, 15-24 years				
...	All causes	11,461	100.0	89.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,695	49.7	44.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,732	23.8	21.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	496	4.3	3.9
4	Malignant neoplasms (C00-C97)	456	4.0	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	315	2.7	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	1.2	1.1
7	Influenza and pneumonia (J09-J18)	79	0.7	0.6
8	Diabetes mellitus (E10-E14)	58	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	46	0.4	0.4
10	Septicemia (A40-A41)	37	0.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.3	0.3
12	Chronic lower respiratory diseases (J40-J47)	32	0.3	0.3
13	Legal intervention (Y35,Y89.0)	30	0.3	0.2
14	Aortic aneurysm and dissection (I71)	16	0.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.1	*
...	All other causes (Residual)	1,278	11.2	10.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-34 years				
...	All causes	18,904	100.0	149.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,491	44.9	67.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,729	19.7	29.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,219	6.4	9.7
4	Malignant neoplasms (C00-C97)	1,071	5.7	8.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	657	3.5	5.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	264	1.4	2.1
7	Diabetes mellitus (E10-E14)	215	1.1	1.7
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	0.9	1.3
9	Influenza and pneumonia (J09-J18)	144	0.8	1.1
10	Cerebrovascular diseases (I60-I69)	131	0.7	1.0
11	Septicemia (A40-A41)	101	0.5	0.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	88	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	79	0.4	0.6
14	Legal intervention (Y35,Y89.0)	66	0.3	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	0.3	0.5
...	All other causes (Residual)	2,424	12.8	19.2
Non-Hispanic, white, male, 35-44 years				
...	All causes	27,197	100.0	223.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,461	27.4	61.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,276	15.7	35.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,013	14.8	32.9
4	Malignant neoplasms (C00-C97)	3,127	11.5	25.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,018	3.7	8.3
6	Diabetes mellitus (E10-E14)	664	2.4	5.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	556	2.0	4.6
8	Cerebrovascular diseases (I60-I69)	423	1.6	3.5
9	Influenza and pneumonia (J09-J18)	287	1.1	2.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	277	1.0	2.3
11	Septicemia (A40-A41)	219	0.8	1.8
12	Chronic lower respiratory diseases (J40-J47)	203	0.7	1.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	172	0.6	1.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	0.5	1.1
15	Aortic aneurysm and dissection (I71)	121	0.4	1.0
...	All other causes (Residual)	4,240	15.6	34.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-54 years				
...	All causes	74,958	100.0	511.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,177	22.9	117.2
2	Malignant neoplasms (C00-C97)	16,402	21.9	111.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,052	13.4	68.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,564	7.4	38.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,074	5.4	27.8
6	Diabetes mellitus (E10-E14)	2,209	2.9	15.1
7	Chronic lower respiratory diseases (J40-J47)	1,689	2.3	11.5
8	Cerebrovascular diseases (I60-I69)	1,610	2.1	11.0
9	Viral hepatitis (B15-B19)	874	1.2	6.0
10	Influenza and pneumonia (J09-J18)	818	1.1	5.6
11	Septicemia (A40-A41)	783	1.0	5.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	674	0.9	4.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	606	0.8	4.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	600	0.8	4.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	536	0.7	3.7
...	All other causes (Residual)	11,290	15.1	77.0
Non-Hispanic, white, male, 55-64 years				
...	All causes	150,822	100.0	1,069.0
1	Malignant neoplasms (C00-C97)	46,963	31.1	332.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,711	24.3	260.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,659	5.7	61.4
4	Chronic lower respiratory diseases (J40-J47)	6,946	4.6	49.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,248	4.1	44.3
6	Diabetes mellitus (E10-E14)	5,152	3.4	36.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,820	3.2	34.2
8	Cerebrovascular diseases (I60-I69)	4,094	2.7	29.0
9	Septicemia (A40-A41)	1,998	1.3	14.2
10	Influenza and pneumonia (J09-J18)	1,903	1.3	13.5
11	Viral hepatitis (B15-B19)	1,895	1.3	13.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,596	1.1	11.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,257	0.8	8.9
14	Aortic aneurysm and dissection (I71)	724	0.5	5.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	657	0.4	4.7
...	All other causes (Residual)	21,199	14.1	150.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-74 years				
...	All causes	202,897	100.0	2,174.3
1	Malignant neoplasms (C00-C97)	68,797	33.9	737.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,545	23.9	520.2
3	Chronic lower respiratory diseases (J40-J47)	15,677	7.7	168.0
4	Cerebrovascular diseases (I60-I69)	7,144	3.5	76.6
5	Diabetes mellitus (E10-E14)	6,891	3.4	73.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,602	2.8	60.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,455	1.7	37.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,052	1.5	32.7
9	Influenza and pneumonia (J09-J18)	3,048	1.5	32.7
10	Septicemia (A40-A41)	2,955	1.5	31.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,806	1.4	30.1
12	Parkinson's disease (G20-G21)	1,841	0.9	19.7
13	Alzheimer's disease (G30)	1,614	0.8	17.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,497	0.7	16.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,384	0.7	14.8
...	All other causes (Residual)	28,589	14.1	306.4
Non-Hispanic, white, male, 75-84 years				
...	All causes	261,077	100.0	5,582.7
1	Malignant neoplasms (C00-C97)	67,568	25.9	1,444.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	65,848	25.2	1,408.1
3	Chronic lower respiratory diseases (J40-J47)	21,123	8.1	451.7
4	Cerebrovascular diseases (I60-I69)	12,721	4.9	272.0
5	Alzheimer's disease (G30)	7,526	2.9	160.9
6	Diabetes mellitus (E10-E14)	7,316	2.8	156.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,725	2.6	143.8
8	Parkinson's disease (G20-G21)	5,960	2.3	127.4
9	Influenza and pneumonia (J09-J18)	5,731	2.2	122.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,291	2.0	113.1
11	Septicemia (A40-A41)	3,771	1.4	80.6
12	Pneumonitis due to solids and liquids (J69)	2,386	0.9	51.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,209	0.8	47.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,204	0.8	47.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,846	0.7	39.5
...	All other causes (Residual)	42,852	16.4	916.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85 years and over				
...	All causes	264,933	100.0	15,485.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,020	31.0	4,794.2
2	Malignant neoplasms (C00-C97)	39,871	15.0	2,330.5
3	Chronic lower respiratory diseases (J40-J47)	15,315	5.8	895.2
4	Cerebrovascular diseases (I60-I69)	14,124	5.3	825.6
5	Alzheimer's disease (G30)	12,880	4.9	752.9
6	Influenza and pneumonia (J09-J18)	8,935	3.4	522.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,408	2.8	433.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,431	2.4	375.9
9	Diabetes mellitus (E10-E14)	5,286	2.0	309.0
10	Parkinson's disease (G20-G21)	5,209	2.0	304.5
11	Pneumonitis due to solids and liquids (J69)	4,108	1.6	240.1
12	Septicemia (A40-A41)	3,439	1.3	201.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,310	1.2	193.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,084	0.8	121.8
15	Aortic aneurysm and dissection (I71)	1,095	0.4	64.0
...	All other causes (Residual)	53,418	20.2	3,122.4
Non-Hispanic, white, female, all ages				
...	All causes	1,031,525	100.0	1,011.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	232,523	22.5	228.0
2	Malignant neoplasms (C00-C97)	218,204	21.2	214.0
3	Chronic lower respiratory diseases (J40-J47)	70,561	6.8	69.2
4	Cerebrovascular diseases (I60-I69)	59,409	5.8	58.3
5	Alzheimer's disease (G30)	50,889	4.9	49.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39,382	3.8	38.6
7	Influenza and pneumonia (J09-J18)	24,291	2.4	23.8
8	Diabetes mellitus (E10-E14)	23,490	2.3	23.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,960	1.6	16.6
10	Septicemia (A40-A41)	15,067	1.5	14.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,088	1.3	12.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,402	0.9	9.2
13	Parkinson's disease (G20-G21)	8,871	0.9	8.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,632	0.7	7.5
15	Pneumonitis due to solids and liquids (J69)	7,096	0.7	7.0
...	All other causes (Residual)	234,660	22.7	230.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	828	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	28.7	5.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	15.2	3.1
3	Malignant neoplasms (C00-C97)	64	7.7	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	7.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	4.1	0.8
6	Influenza and pneumonia (J09-J18)	22	2.7	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.3	*
9	Septicemia (A40-A41)	9	1.1	*
10	Cerebrovascular diseases (I60-I69)	7	0.8	*
11	Meningitis (G00,G03)	5	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (Residual)	222	26.8	5.5
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,167	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	319	27.3	2.9
2	Malignant neoplasms (C00-C97)	246	21.1	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	6.4	0.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	6.2	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	4.2	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	3.3	0.4
7	Influenza and pneumonia (J09-J18)	30	2.6	0.3
8	Cerebrovascular diseases (I60-I69)	21	1.8	0.2
9	Chronic lower respiratory diseases (J40-J47)	14	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.1	*
11	Septicemia (A40-A41)	12	1.0	*
12	Diabetes mellitus (E10-E14)	8	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (Residual)	253	21.7	2.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,631	100.0	38.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,125	45.9	17.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	656	14.2	5.4
3	Malignant neoplasms (C00-C97)	348	7.5	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	191	4.1	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	3.7	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	1.8	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	76	1.6	0.6
8	Influenza and pneumonia (J09-J18)	53	1.1	0.4
9	Diabetes mellitus (E10-E14)	37	0.8	0.3
10	Septicemia (A40-A41)	33	0.7	0.3
11	Cerebrovascular diseases (I60-I69)	29	0.6	0.2
12	Chronic lower respiratory diseases (J40-J47)	22	0.5	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.2	*
...	All other causes (Residual)	765	16.5	6.3
Non-Hispanic, white, female, 25-34 years				
...	All causes	8,679	100.0	70.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,103	35.8	25.1
2	Malignant neoplasms (C00-C97)	1,000	11.5	8.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	996	11.5	8.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	554	6.4	4.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	267	3.1	2.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	209	2.4	1.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	152	1.8	1.2
8	Diabetes mellitus (E10-E14)	127	1.5	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	1.5	1.0
10	Influenza and pneumonia (J09-J18)	122	1.4	1.0
11	Cerebrovascular diseases (I60-I69)	106	1.2	0.9
12	Septicemia (A40-A41)	75	0.9	0.6
13	Chronic lower respiratory diseases (J40-J47)	64	0.7	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	0.7	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.5	0.4
...	All other causes (Residual)	1,672	19.3	13.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-44 years				
...	All causes	16,734	100.0	138.6
1	Malignant neoplasms (C00-C97)	3,974	23.7	32.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,659	21.9	30.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,865	11.1	15.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,285	7.7	10.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	582	3.5	4.8
6	Diabetes mellitus (E10-E14)	398	2.4	3.3
7	Cerebrovascular diseases (I60-I69)	335	2.0	2.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	276	1.6	2.3
9	Chronic lower respiratory diseases (J40-J47)	274	1.6	2.3
10	Influenza and pneumonia (J09-J18)	271	1.6	2.2
11	Septicemia (A40-A41)	250	1.5	2.1
12	Pregnancy, childbirth and the puerperium (O00-O99)	140	0.8	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	139	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	0.8	1.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.4	0.6
...	All other causes (Residual)	3,091	18.5	25.6
Non-Hispanic, white, female, 45-54 years				
...	All causes	47,573	100.0	320.5
1	Malignant neoplasms (C00-C97)	16,032	33.7	108.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,538	13.7	44.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,325	11.2	35.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,144	4.5	14.4
5	Chronic lower respiratory diseases (J40-J47)	2,019	4.2	13.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,963	4.1	13.2
7	Cerebrovascular diseases (I60-I69)	1,303	2.7	8.8
8	Diabetes mellitus (E10-E14)	1,277	2.7	8.6
9	Septicemia (A40-A41)	694	1.5	4.7
10	Influenza and pneumonia (J09-J18)	659	1.4	4.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	432	0.9	2.9
12	Viral hepatitis (B15-B19)	381	0.8	2.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	289	0.6	1.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	260	0.5	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	250	0.5	1.7
...	All other causes (Residual)	8,007	16.8	53.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-64 years				
...	All causes	94,696	100.0	640.5
1	Malignant neoplasms (C00-C97)	37,448	39.5	253.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,637	15.5	99.0
3	Chronic lower respiratory diseases (J40-J47)	6,442	6.8	43.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,242	4.5	28.7
5	Diabetes mellitus (E10-E14)	3,024	3.2	20.5
6	Cerebrovascular diseases (I60-I69)	2,901	3.1	19.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,566	2.7	17.4
8	Septicemia (A40-A41)	1,667	1.8	11.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,591	1.7	10.8
10	Influenza and pneumonia (J09-J18)	1,434	1.5	9.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,197	1.3	8.1
12	Viral hepatitis (B15-B19)	734	0.8	5.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	633	0.7	4.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	477	0.5	3.2
15	Alzheimer's disease (G30)	435	0.5	2.9
...	All other causes (Residual)	15,268	16.1	103.3
Non-Hispanic, white, female, 65-74 years				
...	All causes	153,052	100.0	1,483.9
1	Malignant neoplasms (C00-C97)	55,839	36.5	541.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,435	17.3	256.3
3	Chronic lower respiratory diseases (J40-J47)	15,984	10.4	155.0
4	Cerebrovascular diseases (I60-I69)	6,209	4.1	60.2
5	Diabetes mellitus (E10-E14)	4,738	3.1	45.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,204	2.1	31.1
7	Septicemia (A40-A41)	2,711	1.8	26.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,603	1.7	25.2
9	Influenza and pneumonia (J09-J18)	2,467	1.6	23.9
10	Alzheimer's disease (G30)	2,145	1.4	20.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,938	1.3	18.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,267	0.8	12.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	913	0.6	8.9
14	Parkinson's disease (G20-G21)	901	0.6	8.7
15	Pneumonitis due to solids and liquids (J69)	703	0.5	6.8
...	All other causes (Residual)	24,995	16.3	242.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-84 years				
...	All causes	251,175	100.0	4,142.2
1	Malignant neoplasms (C00-C97)	58,614	23.3	966.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,443	21.3	881.3
3	Chronic lower respiratory diseases (J40-J47)	23,195	9.2	382.5
4	Cerebrovascular diseases (I60-I69)	15,786	6.3	260.3
5	Alzheimer's disease (G30)	12,110	4.8	199.7
6	Diabetes mellitus (E10-E14)	6,409	2.6	105.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,728	2.3	94.5
8	Influenza and pneumonia (J09-J18)	5,431	2.2	89.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,657	1.9	76.8
10	Septicemia (A40-A41)	4,158	1.7	68.6
11	Parkinson's disease (G20-G21)	3,347	1.3	55.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,901	1.2	47.8
13	Pneumonitis due to solids and liquids (J69)	1,733	0.7	28.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,702	0.7	28.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,406	0.6	23.2
...	All other causes (Residual)	50,555	20.1	833.7
Non-Hispanic, white, female, 85 years and over				
...	All causes	448,457	100.0	13,502.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	128,751	28.7	3,876.5
2	Malignant neoplasms (C00-C97)	44,620	9.9	1,343.5
3	Alzheimer's disease (G30)	36,151	8.1	1,088.5
4	Cerebrovascular diseases (I60-I69)	32,690	7.3	984.3
5	Chronic lower respiratory diseases (J40-J47)	22,534	5.0	678.5
6	Influenza and pneumonia (J09-J18)	13,778	3.1	414.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,231	2.5	338.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,963	1.8	239.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,850	1.8	236.4
10	Diabetes mellitus (E10-E14)	7,470	1.7	224.9
11	Septicemia (A40-A41)	5,432	1.2	163.6
12	Parkinson's disease (G20-G21)	4,448	1.0	133.9
13	Pneumonitis due to solids and liquids (J69)	4,010	0.9	120.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,462	0.5	74.1
15	Atherosclerosis (I70)	2,315	0.5	69.7
...	All other causes (Residual)	116,752	26.0	3,515.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	299,227	100.0	733.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,102	23.8	174.3
2	Malignant neoplasms (C00-C97)	67,272	22.5	164.9
3	Cerebrovascular diseases (I60-I69)	16,111	5.4	39.5
4	Diabetes mellitus (E10-E14)	13,239	4.4	32.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,169	4.4	32.3
6	Chronic lower respiratory diseases (J40-J47)	9,816	3.3	24.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,340	2.8	20.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,950	2.7	19.5
9	Septicemia (A40-A41)	6,193	2.1	15.2
10	Alzheimer's disease (G30)	5,664	1.9	13.9
11	Influenza and pneumonia (J09-J18)	5,494	1.8	13.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,400	1.8	13.2
13	Certain conditions originating in the perinatal period (P00-P96)	3,854	1.3	9.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,673	1.2	9.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,998	1.0	7.3
...	All other causes (Residual)	58,952	19.7	144.5
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	963	100.0	39.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	285	29.6	11.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	13.8	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	9.1	3.6
4	Malignant neoplasms (C00-C97)	62	6.4	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	6.0	2.4
6	Chronic lower respiratory diseases (J40-J47)	26	2.7	1.1
7	Influenza and pneumonia (J09-J18)	21	2.2	0.9
8	Septicemia (A40-A41)	18	1.9	*
9	Cerebrovascular diseases (I60-I69)	15	1.6	*
10	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
11	Anemias (D50-D64)	8	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
13	Meningitis (G00,G03)	5	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
...	All other causes (Residual)	219	22.7	9.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,132	100.0	18.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	324	28.6	5.2
2	Malignant neoplasms (C00-C97)	119	10.5	1.9
3	Chronic lower respiratory diseases (J40-J47)	86	7.6	1.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	7.2	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	7.1	1.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	4.9	0.9
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.3	0.8
8	Influenza and pneumonia (J09-J18)	27	2.4	0.4
9	Cerebrovascular diseases (I60-I69)	18	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.2	*
11	Septicemia (A40-A41)	13	1.1	*
12	Anemias (D50-D64)	11	1.0	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	239	21.1	3.9
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	6,423	100.0	94.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,693	41.9	39.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,376	21.4	20.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	513	8.0	7.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	287	4.5	4.2
5	Malignant neoplasms (C00-C97)	239	3.7	3.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	85	1.3	1.2
7	Chronic lower respiratory diseases (J40-J47)	80	1.2	1.2
8	Diabetes mellitus (E10-E14)	73	1.1	1.1
9	Anemias (D50-D64)	70	1.1	1.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	69	1.1	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	0.9	0.8
12	Legal intervention (Y35,Y89.0)	43	0.7	0.6
13	Cerebrovascular diseases (I60-I69)	38	0.6	0.6
14	Septicemia (A40-A41)	36	0.6	0.5
15	Influenza and pneumonia (J09-J18)	31	0.5	0.5
...	All other causes (Residual)	733	11.4	10.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-34 years				
...	All causes	9,382	100.0	162.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,445	26.1	42.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,764	18.8	30.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	971	10.3	16.8
4	Malignant neoplasms (C00-C97)	639	6.8	11.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	544	5.8	9.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	391	4.2	6.8
7	Diabetes mellitus (E10-E14)	221	2.4	3.8
8	Pregnancy, childbirth and the puerperium (O00-O99)	149	1.6	2.6
9	Cerebrovascular diseases (I60-I69)	142	1.5	2.5
10	Anemias (D50-D64)	113	1.2	2.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	1.1	1.7
12	Chronic lower respiratory diseases (J40-J47)	99	1.1	1.7
13	Influenza and pneumonia (J09-J18)	96	1.0	1.7
14	Septicemia (A40-A41)	89	0.9	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	0.8	1.3
...	All other causes (Residual)	1,546	16.5	26.8
Non-Hispanic, black, both sexes, 35-44 years				
...	All causes	13,579	100.0	260.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,770	20.4	53.2
2	Malignant neoplasms (C00-C97)	1,892	13.9	36.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,747	12.9	33.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,225	9.0	23.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	675	5.0	13.0
6	Diabetes mellitus (E10-E14)	531	3.9	10.2
7	Cerebrovascular diseases (I60-I69)	485	3.6	9.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	426	3.1	8.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	242	1.8	4.6
10	Septicemia (A40-A41)	225	1.7	4.3
11	Chronic lower respiratory diseases (J40-J47)	201	1.5	3.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	191	1.4	3.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	176	1.3	3.4
14	Influenza and pneumonia (J09-J18)	157	1.2	3.0
15	Aortic aneurysm and dissection (I71)	101	0.7	1.9
...	All other causes (Residual)	2,535	18.7	48.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-54 years				
...	All causes	31,760	100.0	581.9
1	Malignant neoplasms (C00-C97)	7,589	23.9	139.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,585	23.9	139.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,399	7.6	44.0
4	Cerebrovascular diseases (I60-I69)	1,455	4.6	26.7
5	Diabetes mellitus (E10-E14)	1,401	4.4	25.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,162	3.7	21.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	761	2.4	13.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	735	2.3	13.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	675	2.1	12.4
10	Chronic lower respiratory diseases (J40-J47)	651	2.0	11.9
11	Septicemia (A40-A41)	636	2.0	11.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	548	1.7	10.0
13	Influenza and pneumonia (J09-J18)	448	1.4	8.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	360	1.1	6.6
15	Viral hepatitis (B15-B19)	212	0.7	3.9
...	All other causes (Residual)	5,143	16.2	94.2
Non-Hispanic, black, both sexes, 55-64 years				
...	All causes	57,480	100.0	1,301.1
1	Malignant neoplasms (C00-C97)	17,531	30.5	396.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,154	24.6	320.4
3	Diabetes mellitus (E10-E14)	2,863	5.0	64.8
4	Cerebrovascular diseases (I60-I69)	2,805	4.9	63.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,306	4.0	52.2
6	Chronic lower respiratory diseases (J40-J47)	1,785	3.1	40.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,532	2.7	34.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,257	2.2	28.5
9	Septicemia (A40-A41)	1,167	2.0	26.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	965	1.7	21.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	905	1.6	20.5
12	Influenza and pneumonia (J09-J18)	854	1.5	19.3
13	Viral hepatitis (B15-B19)	636	1.1	14.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	350	0.6	7.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	245	0.4	5.5
...	All other causes (Residual)	8,125	14.1	183.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-74 years				
...	All causes	57,508	100.0	2,464.8
1	Malignant neoplasms (C00-C97)	17,778	30.9	762.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,392	25.0	616.8
3	Cerebrovascular diseases (I60-I69)	3,188	5.5	136.6
4	Diabetes mellitus (E10-E14)	3,143	5.5	134.7
5	Chronic lower respiratory diseases (J40-J47)	2,496	4.3	107.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,883	3.3	80.7
7	Septicemia (A40-A41)	1,337	2.3	57.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,085	1.9	46.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,070	1.9	45.9
10	Influenza and pneumonia (J09-J18)	1,038	1.8	44.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	511	0.9	21.9
12	Alzheimer's disease (G30)	416	0.7	17.8
13	Pneumonitis due to solids and liquids (J69)	306	0.5	13.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	303	0.5	13.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	247	0.4	10.6
...	All other causes (Residual)	8,315	14.5	356.4
Non-Hispanic, black, both sexes, 75-84 years				
...	All causes	59,224	100.0	5,270.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,199	25.7	1,352.7
2	Malignant neoplasms (C00-C97)	14,147	23.9	1,259.1
3	Cerebrovascular diseases (I60-I69)	4,073	6.9	362.5
4	Diabetes mellitus (E10-E14)	2,987	5.0	265.8
5	Chronic lower respiratory diseases (J40-J47)	2,543	4.3	226.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,118	3.6	188.5
7	Alzheimer's disease (G30)	1,750	3.0	155.8
8	Septicemia (A40-A41)	1,430	2.4	127.3
9	Influenza and pneumonia (J09-J18)	1,221	2.1	108.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,212	2.0	107.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	793	1.3	70.6
12	Pneumonitis due to solids and liquids (J69)	436	0.7	38.8
13	Parkinson's disease (G20-G21)	435	0.7	38.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	304	0.5	27.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	208	0.4	18.5
...	All other causes (Residual)	10,368	17.5	922.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85 years and over				
...	All causes	54,975	100.0	12,659.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,540	28.3	3,578.4
2	Malignant neoplasms (C00-C97)	7,267	13.2	1,673.4
3	Cerebrovascular diseases (I60-I69)	3,852	7.0	887.0
4	Alzheimer's disease (G30)	3,411	6.2	785.4
5	Diabetes mellitus (E10-E14)	2,014	3.7	463.8
6	Chronic lower respiratory diseases (J40-J47)	1,838	3.3	423.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,746	3.2	402.0
8	Influenza and pneumonia (J09-J18)	1,531	2.8	352.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,411	2.6	324.9
10	Septicemia (A40-A41)	1,188	2.2	273.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	692	1.3	159.3
12	Pneumonitis due to solids and liquids (J69)	497	0.9	114.4
13	Parkinson's disease (G20-G21)	355	0.6	81.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	239	0.4	55.0
15	Atherosclerosis (I70)	196	0.4	45.1
...	All other causes (Residual)	13,198	24.0	3,039.1
Non-Hispanic, black, male, all ages				
...	All causes	152,661	100.0	782.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,562	23.9	187.4
2	Malignant neoplasms (C00-C97)	34,319	22.5	175.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,845	5.8	45.3
4	Cerebrovascular diseases (I60-I69)	7,266	4.8	37.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,851	4.5	35.1
6	Diabetes mellitus (E10-E14)	6,298	4.1	32.3
7	Chronic lower respiratory diseases (J40-J47)	5,019	3.3	25.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,944	2.6	20.2
9	Septicemia (A40-A41)	2,854	1.9	14.6
10	Influenza and pneumonia (J09-J18)	2,657	1.7	13.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,467	1.6	12.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,429	1.6	12.4
13	Certain conditions originating in the perinatal period (P00-P96)	2,102	1.4	10.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,915	1.3	9.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,850	1.2	9.5
...	All other causes (Residual)	27,283	17.9	139.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	538	100.0	43.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	30.7	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	12.6	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	9.3	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	6.7	2.9
5	Malignant neoplasms (C00-C97)	33	6.1	2.7
6	Chronic lower respiratory diseases (J40-J47)	15	2.8	*
7	Septicemia (A40-A41)	12	2.2	*
7	Cerebrovascular diseases (I60-I69)	12	2.2	*
7	Influenza and pneumonia (J09-J18)	12	2.2	*
10	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
11	Anemias (D50-D64)	4	0.7	*
11	Meningitis (G00,G03)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	115	21.4	9.3
Non-Hispanic, black, male, 5-14 years				
...	All causes	653	100.0	20.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	32.2	6.7
2	Malignant neoplasms (C00-C97)	65	10.0	2.1
3	Chronic lower respiratory diseases (J40-J47)	54	8.3	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	8.1	1.7
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	5.7	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	5.1	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	4.4	0.9
8	Influenza and pneumonia (J09-J18)	18	2.8	*
9	Cerebrovascular diseases (I60-I69)	10	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.4	*
11	Septicemia (A40-A41)	8	1.2	*
12	Anemias (D50-D64)	5	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	113	17.3	3.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	4,957	100.0	143.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,450	49.4	71.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	995	20.1	28.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	421	8.5	12.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	191	3.9	5.5
5	Malignant neoplasms (C00-C97)	128	2.6	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.2	1.8
7	Chronic lower respiratory diseases (J40-J47)	55	1.1	1.6
8	Diabetes mellitus (E10-E14)	48	1.0	1.4
9	Anemias (D50-D64)	44	0.9	1.3
10	Legal intervention (Y35,Y89.0)	43	0.9	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	0.7	1.0
12	Cerebrovascular diseases (I60-I69)	21	0.4	0.6
13	Septicemia (A40-A41)	19	0.4	*
14	Influenza and pneumonia (J09-J18)	16	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
...	All other causes (Residual)	421	8.5	12.2
Non-Hispanic, black, male, 25-34 years				
...	All causes	6,489	100.0	233.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,174	33.5	78.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,307	20.1	46.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	629	9.7	22.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	454	7.0	16.3
5	Malignant neoplasms (C00-C97)	307	4.7	11.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	270	4.2	9.7
7	Diabetes mellitus (E10-E14)	117	1.8	4.2
8	Cerebrovascular diseases (I60-I69)	73	1.1	2.6
9	Anemias (D50-D64)	58	0.9	2.1
10	Influenza and pneumonia (J09-J18)	54	0.8	1.9
11	Chronic lower respiratory diseases (J40-J47)	53	0.8	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	0.8	1.8
13	Septicemia (A40-A41)	41	0.6	1.5
14	Legal intervention (Y35,Y89.0)	40	0.6	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	0.6	1.4
...	All other causes (Residual)	824	12.7	29.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-44 years				
...	All causes	8,105	100.0	330.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,793	22.1	73.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,224	15.1	49.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,056	13.0	43.0
4	Malignant neoplasms (C00-C97)	708	8.7	28.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	384	4.7	15.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	346	4.3	14.1
7	Diabetes mellitus (E10-E14)	319	3.9	13.0
8	Cerebrovascular diseases (I60-I69)	264	3.3	10.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	1.7	5.7
10	Chronic lower respiratory diseases (J40-J47)	117	1.4	4.8
11	Septicemia (A40-A41)	110	1.4	4.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	109	1.3	4.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	97	1.2	4.0
14	Influenza and pneumonia (J09-J18)	79	1.0	3.2
15	Aortic aneurysm and dissection (I71)	76	0.9	3.1
...	All other causes (Residual)	1,284	15.8	52.3
Non-Hispanic, black, male, 45-54 years				
...	All causes	18,072	100.0	705.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,754	26.3	185.6
2	Malignant neoplasms (C00-C97)	3,526	19.5	137.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,625	9.0	63.4
4	Cerebrovascular diseases (I60-I69)	818	4.5	31.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	767	4.2	29.9
6	Diabetes mellitus (E10-E14)	766	4.2	29.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	599	3.3	23.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	448	2.5	17.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	404	2.2	15.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	340	1.9	13.3
11	Septicemia (A40-A41)	335	1.9	13.1
12	Chronic lower respiratory diseases (J40-J47)	306	1.7	11.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	277	1.5	10.8
14	Influenza and pneumonia (J09-J18)	266	1.5	10.4
15	Viral hepatitis (B15-B19)	139	0.8	5.4
...	All other causes (Residual)	2,702	15.0	105.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-64 years				
...	All causes	33,644	100.0	1,672.2
1	Malignant neoplasms (C00-C97)	9,611	28.6	477.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,870	26.4	440.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,673	5.0	83.2
4	Cerebrovascular diseases (I60-I69)	1,618	4.8	80.4
5	Diabetes mellitus (E10-E14)	1,599	4.8	79.5
6	Chronic lower respiratory diseases (J40-J47)	906	2.7	45.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	869	2.6	43.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	866	2.6	43.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	674	2.0	33.5
10	Septicemia (A40-A41)	598	1.8	29.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	519	1.5	25.8
12	Influenza and pneumonia (J09-J18)	477	1.4	23.7
13	Viral hepatitis (B15-B19)	431	1.3	21.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	276	0.8	13.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	186	0.6	9.2
...	All other causes (Residual)	4,471	13.3	222.2
Non-Hispanic, black, male, 65-74 years				
...	All causes	31,433	100.0	3,137.2
1	Malignant neoplasms (C00-C97)	9,676	30.8	965.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,254	26.3	823.8
3	Cerebrovascular diseases (I60-I69)	1,697	5.4	169.4
4	Diabetes mellitus (E10-E14)	1,572	5.0	156.9
5	Chronic lower respiratory diseases (J40-J47)	1,394	4.4	139.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	905	2.9	90.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	717	2.3	71.6
8	Septicemia (A40-A41)	670	2.1	66.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	577	1.8	57.6
10	Influenza and pneumonia (J09-J18)	570	1.8	56.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	356	1.1	35.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	205	0.7	20.5
13	Pneumonitis due to solids and liquids (J69)	188	0.6	18.8
14	Alzheimer's disease (G30)	184	0.6	18.4
15	Viral hepatitis (B15-B19)	147	0.5	14.7
...	All other causes (Residual)	4,321	13.7	431.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-84 years				
...	All causes	27,582	100.0	6,501.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,165	26.0	1,688.9
2	Malignant neoplasms (C00-C97)	7,098	25.7	1,673.1
3	Cerebrovascular diseases (I60-I69)	1,686	6.1	397.4
4	Chronic lower respiratory diseases (J40-J47)	1,314	4.8	309.7
5	Diabetes mellitus (E10-E14)	1,270	4.6	299.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	939	3.4	221.3
7	Alzheimer's disease (G30)	648	2.3	152.7
8	Septicemia (A40-A41)	619	2.2	145.9
9	Influenza and pneumonia (J09-J18)	574	2.1	135.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	526	1.9	124.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	450	1.6	106.1
12	Parkinson's disease (G20-G21)	281	1.0	66.2
13	Pneumonitis due to solids and liquids (J69)	239	0.9	56.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	147	0.5	34.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	98	0.4	23.1
...	All other causes (Residual)	4,528	16.4	1,067.3
Non-Hispanic, black, male, 85 years and over				
...	All causes	17,477	100.0	13,959.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,788	27.4	3,824.4
2	Malignant neoplasms (C00-C97)	3,165	18.1	2,528.0
3	Cerebrovascular diseases (I60-I69)	1,044	6.0	833.9
4	Chronic lower respiratory diseases (J40-J47)	799	4.6	638.2
5	Alzheimer's disease (G30)	763	4.4	609.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	613	3.5	489.6
7	Diabetes mellitus (E10-E14)	604	3.5	482.4
8	Influenza and pneumonia (J09-J18)	551	3.2	440.1
9	Septicemia (A40-A41)	412	2.4	329.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	361	2.1	288.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	265	1.5	211.7
12	Pneumonitis due to solids and liquids (J69)	215	1.2	171.7
13	Parkinson's disease (G20-G21)	165	0.9	131.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	92	0.5	73.5
15	Atherosclerosis (I70)	53	0.3	42.3
...	All other causes (Residual)	3,587	20.5	2,865.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	146,566	100.0	688.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,540	23.6	162.2
2	Malignant neoplasms (C00-C97)	32,953	22.5	154.8
3	Cerebrovascular diseases (I60-I69)	8,845	6.0	41.5
4	Diabetes mellitus (E10-E14)	6,941	4.7	32.6
5	Chronic lower respiratory diseases (J40-J47)	4,797	3.3	22.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,396	3.0	20.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,324	3.0	20.3
8	Alzheimer's disease (G30)	4,018	2.7	18.9
9	Septicemia (A40-A41)	3,339	2.3	15.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,933	2.0	13.8
11	Influenza and pneumonia (J09-J18)	2,837	1.9	13.3
12	Certain conditions originating in the perinatal period (P00-P96)	1,752	1.2	8.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,244	0.8	5.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,099	0.7	5.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,083	0.7	5.1
...	All other causes (Residual)	31,465	21.5	147.8
Non-Hispanic, black, female, 1-4 years				
...	All causes	425	100.0	35.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	28.2	10.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	15.3	5.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	8.9	3.2
4	Malignant neoplasms (C00-C97)	29	6.8	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	5.2	1.8
6	Chronic lower respiratory diseases (J40-J47)	11	2.6	*
7	Influenza and pneumonia (J09-J18)	9	2.1	*
8	Septicemia (A40-A41)	6	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
11	Anemias (D50-D64)	4	0.9	*
12	Cerebrovascular diseases (I60-I69)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	98	23.1	8.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	479	100.0	15.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	23.8	3.7
2	Malignant neoplasms (C00-C97)	54	11.3	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	9.0	1.4
4	Chronic lower respiratory diseases (J40-J47)	32	6.7	1.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	5.8	0.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	4.6	0.7
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	4.2	0.7
8	Influenza and pneumonia (J09-J18)	9	1.9	*
9	Cerebrovascular diseases (I60-I69)	8	1.7	*
10	Anemias (D50-D64)	6	1.3	*
11	Septicemia (A40-A41)	5	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
14	Meningitis (G00,G03)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	122	25.5	4.0
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,466	100.0	43.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	381	26.0	11.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	243	16.6	7.2
3	Malignant neoplasms (C00-C97)	111	7.6	3.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	6.5	2.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	6.3	2.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	69	4.7	2.0
7	Anemias (D50-D64)	26	1.8	0.8
8	Diabetes mellitus (E10-E14)	25	1.7	0.7
8	Chronic lower respiratory diseases (J40-J47)	25	1.7	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	24	1.6	0.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	1.5	0.7
12	Septicemia (A40-A41)	17	1.2	*
12	Cerebrovascular diseases (I60-I69)	17	1.2	*
14	Influenza and pneumonia (J09-J18)	15	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
...	All other causes (Residual)	296	20.2	8.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-34 years				
...	All causes	2,893	100.0	97.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	457	15.8	15.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	342	11.8	11.5
3	Malignant neoplasms (C00-C97)	332	11.5	11.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	271	9.4	9.1
5	Pregnancy, childbirth and the puerperium (O00-O99)	149	5.2	5.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	121	4.2	4.1
7	Diabetes mellitus (E10-E14)	104	3.6	3.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	3.1	3.0
9	Cerebrovascular diseases (I60-I69)	69	2.4	2.3
10	Anemias (D50-D64)	55	1.9	1.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	1.8	1.7
12	Septicemia (A40-A41)	48	1.7	1.6
13	Chronic lower respiratory diseases (J40-J47)	46	1.6	1.5
14	Influenza and pneumonia (J09-J18)	42	1.5	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	1.2	1.1
...	All other causes (Residual)	682	23.6	22.9
Non-Hispanic, black, female, 35-44 years				
...	All causes	5,474	100.0	198.6
1	Malignant neoplasms (C00-C97)	1,184	21.6	43.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	977	17.8	35.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	523	9.6	19.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	291	5.3	10.6
5	Cerebrovascular diseases (I60-I69)	221	4.0	8.0
6	Diabetes mellitus (E10-E14)	212	3.9	7.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	169	3.1	6.1
8	Septicemia (A40-A41)	115	2.1	4.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	1.9	3.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	94	1.7	3.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	86	1.6	3.1
12	Chronic lower respiratory diseases (J40-J47)	84	1.5	3.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	1.5	2.9
14	Influenza and pneumonia (J09-J18)	78	1.4	2.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	1.2	2.4
...	All other causes (Residual)	1,190	21.7	43.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-54 years				
...	All causes	13,688	100.0	472.6
1	Malignant neoplasms (C00-C97)	4,063	29.7	140.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,831	20.7	97.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	774	5.7	26.7
4	Cerebrovascular diseases (I60-I69)	637	4.7	22.0
5	Diabetes mellitus (E10-E14)	635	4.6	21.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	395	2.9	13.6
7	Chronic lower respiratory diseases (J40-J47)	345	2.5	11.9
8	Septicemia (A40-A41)	301	2.2	10.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	287	2.1	9.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	271	2.0	9.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	208	1.5	7.2
12	Influenza and pneumonia (J09-J18)	182	1.3	6.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	162	1.2	5.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	0.6	2.9
15	Viral hepatitis (B15-B19)	73	0.5	2.5
...	All other causes (Residual)	2,441	17.8	84.3
Non-Hispanic, black, female, 55-64 years				
...	All causes	23,836	100.0	990.8
1	Malignant neoplasms (C00-C97)	7,920	33.2	329.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,284	22.2	219.6
3	Diabetes mellitus (E10-E14)	1,264	5.3	52.5
4	Cerebrovascular diseases (I60-I69)	1,187	5.0	49.3
5	Chronic lower respiratory diseases (J40-J47)	879	3.7	36.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	663	2.8	27.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	633	2.7	26.3
8	Septicemia (A40-A41)	569	2.4	23.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	391	1.6	16.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	386	1.6	16.0
11	Influenza and pneumonia (J09-J18)	377	1.6	15.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	291	1.2	12.1
13	Viral hepatitis (B15-B19)	205	0.9	8.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.4	3.9
15	Pneumonitis due to solids and liquids (J69)	83	0.3	3.5
...	All other causes (Residual)	3,610	15.1	150.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-74 years				
...	All causes	26,075	100.0	1,958.7
1	Malignant neoplasms (C00-C97)	8,102	31.1	608.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,138	23.5	461.1
3	Diabetes mellitus (E10-E14)	1,571	6.0	118.0
4	Cerebrovascular diseases (I60-I69)	1,491	5.7	112.0
5	Chronic lower respiratory diseases (J40-J47)	1,102	4.2	82.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	978	3.8	73.5
7	Septicemia (A40-A41)	667	2.6	50.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	508	1.9	38.2
9	Influenza and pneumonia (J09-J18)	468	1.8	35.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	353	1.4	26.5
11	Alzheimer's disease (G30)	232	0.9	17.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	155	0.6	11.6
13	Pneumonitis due to solids and liquids (J69)	118	0.5	8.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	102	0.4	7.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	98	0.4	7.4
...	All other causes (Residual)	3,992	15.3	299.9
Non-Hispanic, black, female, 75-84 years				
...	All causes	31,642	100.0	4,524.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,034	25.4	1,148.8
2	Malignant neoplasms (C00-C97)	7,049	22.3	1,007.9
3	Cerebrovascular diseases (I60-I69)	2,387	7.5	341.3
4	Diabetes mellitus (E10-E14)	1,717	5.4	245.5
5	Chronic lower respiratory diseases (J40-J47)	1,229	3.9	175.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,179	3.7	168.6
7	Alzheimer's disease (G30)	1,102	3.5	157.6
8	Septicemia (A40-A41)	811	2.6	116.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	686	2.2	98.1
10	Influenza and pneumonia (J09-J18)	647	2.0	92.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	343	1.1	49.0
12	Pneumonitis due to solids and liquids (J69)	197	0.6	28.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	157	0.5	22.4
14	Parkinson's disease (G20-G21)	154	0.5	22.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	110	0.3	15.7
...	All other causes (Residual)	5,840	18.5	835.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85 years and over				
...	All causes	37,498	100.0	12,132.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,752	28.7	3,478.7
2	Malignant neoplasms (C00-C97)	4,102	10.9	1,327.2
3	Cerebrovascular diseases (I60-I69)	2,808	7.5	908.5
4	Alzheimer's disease (G30)	2,648	7.1	856.7
5	Diabetes mellitus (E10-E14)	1,410	3.8	456.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,133	3.0	366.6
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,050	2.8	339.7
8	Chronic lower respiratory diseases (J40-J47)	1,039	2.8	336.2
9	Influenza and pneumonia (J09-J18)	980	2.6	317.1
10	Septicemia (A40-A41)	776	2.1	251.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	427	1.1	138.2
12	Pneumonitis due to solids and liquids (J69)	282	0.8	91.2
13	Parkinson's disease (G20-G21)	190	0.5	61.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	147	0.4	47.6
15	Atherosclerosis (I70)	143	0.4	46.3
...	All other causes (Residual)	9,611	25.6	3,109.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013