

A revised mortality data file for 2014 was released on April 3, 2017 to include corrections from Tennessee and Massachusetts affecting 125 records previously coded to Accidental discharge of firearms (W32-W34). After the revision, 50% of the changed records are now classified as Assault (homicide) by discharge of firearms (*U01.4 and X93-X95); 42% as Intentional self-harm (suicide) by discharge of firearms (X72-X74); 4% as Discharge of firearms, undetermined intent (Y22-Y24); and the remaining 4% to various other causes. For more information, please see "[Deaths: Final Data for 2014](#)." This table was updated on May 3, 2017 to correct errors. The shaded areas in this table indicate the changes that were made.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	169,387	100.0	305.8
1	Malignant neoplasms (C00-C97)	36,447	21.5	65.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,021	20.1	61.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,413	7.3	22.4
4	Cerebrovascular diseases (I60-I69)	8,713	5.1	15.7
5	Diabetes mellitus (E10-E14)	7,795	4.6	14.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	5,658	3.3	10.2
7	Alzheimer's disease (G30)	4,934	2.9	8.9
8	Chronic lower respiratory diseases (J40-J47)	4,795	2.8	8.7
9	Influenza and pneumonia (J09-J18)	3,875	2.3	7.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,273	1.9	5.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,246	1.9	5.9
12	Septicemia (A40-A41)	2,613	1.5	4.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,610	1.5	4.7
14	Certain conditions originating in the perinatal period (P00-P96)	2,422	1.4	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,139	1.3	3.9
...	All other causes (residual)	34,433	20.3	62.2
Hispanic, both sexes, 1-4 years				
...	All causes	769	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	28.7	5.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	12.6	2.4
3	Malignant neoplasms (C00-C97)	81	10.5	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	9.1	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.8	0.7
6	Influenza and pneumonia (J09-J18)	20	2.6	0.5
7	Chronic lower respiratory diseases (J40-J47)	10	1.3	*
7	Certain conditions originating in the perinatal period (P00-P96)	10	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.0	*
10	Cerebrovascular diseases (I60-I69)	7	0.9	*
11	Septicemia (A40-A41)	5	0.7	*
12	Nutritional deficiencies (E40-E64)	4	0.5	*
12	Meningitis (G00,G03)	4	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (residual)	196	25.5	4.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,116	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	28.0	3.1
2	Malignant neoplasms (C00-C97)	219	19.6	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	7.2	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	5.9	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	4.9	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	2.4	0.3
6	Cerebrovascular diseases (I60-I69)	27	2.4	0.3
8	Chronic lower respiratory diseases (J40-J47)	24	2.2	0.2
9	Influenza and pneumonia (J09-J18)	23	2.1	0.2
10	Septicemia (A40-A41)	16	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
13	Anemias (D50-D64)	5	0.4	*
14	Diabetes mellitus (E10-E14)	4	0.4	*
14	Meningitis (G00,G03)	4	0.4	*
...	All other causes (residual)	233	20.9	2.3
Hispanic, both sexes, 15-24 years				
...	All causes	4,905	100.0	52.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,044	41.7	21.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	812	16.6	8.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	720	14.7	7.6
4	Malignant neoplasms (C00-C97)	364	7.4	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	128	2.6	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	1.3	0.7
7	Cerebrovascular diseases (I60-I69)	34	0.7	0.4
8	Influenza and pneumonia (J09-J18)	31	0.6	0.3
9	Pregnancy, childbirth and the puerperium (O00-O99)	29	0.6	0.3
10	Septicemia (A40-A41)	24	0.5	0.3
11	Chronic lower respiratory diseases (J40-J47)	21	0.4	0.2
12	Legal intervention (Y35,Y89.0)	20	0.4	0.2
13	Diabetes mellitus (E10-E14)	18	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
...	All other causes (residual)	567	11.6	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	6,575	100.0	74.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,343	35.6	26.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	743	11.3	8.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	711	10.8	8.0
4	Malignant neoplasms (C00-C97)	631	9.6	7.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	395	6.0	4.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	151	2.3	1.7
7	Cerebrovascular diseases (I60-I69)	112	1.7	1.3
8	Influenza and pneumonia (J09-J18)	102	1.6	1.1
9	Diabetes mellitus (E10-E14)	101	1.5	1.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	99	1.5	1.1
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	0.9	0.7
12	Pregnancy, childbirth and the puerperium (O00-O99)	56	0.9	0.6
13	Septicemia (A40-A41)	53	0.8	0.6
14	Chronic lower respiratory diseases (J40-J47)	42	0.6	0.5
15	Legal intervention (Y35,Y89.0)	33	0.5	0.4
...	All other causes (residual)	943	14.3	10.6
Hispanic, both sexes, 35-44 years				
...	All causes	9,168	100.0	114.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,980	21.6	24.8
2	Malignant neoplasms (C00-C97)	1,698	18.5	21.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,034	11.3	12.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	605	6.6	7.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	533	5.8	6.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	490	5.3	6.1
7	Cerebrovascular diseases (I60-I69)	318	3.5	4.0
8	Diabetes mellitus (E10-E14)	259	2.8	3.2
9	Influenza and pneumonia (J09-J18)	193	2.1	2.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	173	1.9	2.2
11	Septicemia (A40-A41)	123	1.3	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	1.0	1.1
13	Pregnancy, childbirth and the puerperium (O00-O99)	61	0.7	0.8
14	Viral hepatitis (B15-B19)	58	0.6	0.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	51	0.6	0.6
...	All other causes (residual)	1,504	16.4	18.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	16,705	100.0	263.9
1	Malignant neoplasms (C00-C97)	4,233	25.3	66.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,714	16.2	42.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,872	11.2	29.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,422	8.5	22.5
5	Diabetes mellitus (E10-E14)	780	4.7	12.3
6	Cerebrovascular diseases (I60-I69)	666	4.0	10.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	511	3.1	8.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	331	2.0	5.2
9	Viral hepatitis (B15-B19)	299	1.8	4.7
10	Influenza and pneumonia (J09-J18)	280	1.7	4.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	249	1.5	3.9
12	Septicemia (A40-A41)	247	1.5	3.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	242	1.4	3.8
14	Chronic lower respiratory diseases (J40-J47)	159	1.0	2.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	146	0.9	2.3
...	All other causes (residual)	2,554	15.3	40.3
Hispanic, both sexes, 55-64 years				
...	All causes	24,701	100.0	609.4
1	Malignant neoplasms (C00-C97)	7,488	30.3	184.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,788	19.4	118.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,744	7.1	43.0
4	Diabetes mellitus (E10-E14)	1,496	6.1	36.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,250	5.1	30.8
6	Cerebrovascular diseases (I60-I69)	1,010	4.1	24.9
7	Influenza and pneumonia (J09-J18)	536	2.2	13.2
8	Viral hepatitis (B15-B19)	480	1.9	11.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	472	1.9	11.6
10	Septicemia (A40-A41)	462	1.9	11.4
11	Chronic lower respiratory diseases (J40-J47)	437	1.8	10.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	315	1.3	7.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	254	1.0	6.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	199	0.8	4.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	0.5	2.9
...	All other causes (residual)	3,651	14.8	90.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	28,036	100.0	1,316.5
1	Malignant neoplasms (C00-C97)	8,715	31.1	409.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,988	21.4	281.2
3	Diabetes mellitus (E10-E14)	1,824	6.5	85.7
4	Cerebrovascular diseases (I60-I69)	1,380	4.9	64.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,071	3.8	50.3
6	Chronic lower respiratory diseases (J40-J47)	895	3.2	42.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	759	2.7	35.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	732	2.6	34.4
9	Influenza and pneumonia (J09-J18)	585	2.1	27.5
10	Septicemia (A40-A41)	503	1.8	23.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	365	1.3	17.1
12	Alzheimer's disease (G30)	336	1.2	15.8
13	Parkinson's disease (G20-G21)	201	0.7	9.4
14	Viral hepatitis (B15-B19)	199	0.7	9.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	0.6	7.3
...	All other causes (residual)	4,328	15.4	203.2
Hispanic, both sexes, 75-84 years				
...	All causes	35,342	100.0	3,415.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,352	23.6	807.2
2	Malignant neoplasms (C00-C97)	8,254	23.4	797.7
3	Cerebrovascular diseases (I60-I69)	2,385	6.7	230.5
4	Diabetes mellitus (E10-E14)	1,977	5.6	191.1
5	Chronic lower respiratory diseases (J40-J47)	1,552	4.4	150.0
6	Alzheimer's disease (G30)	1,438	4.1	139.0
7	Influenza and pneumonia (J09-J18)	915	2.6	88.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	907	2.6	87.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	742	2.1	71.7
10	Septicemia (A40-A41)	633	1.8	61.2
11	Parkinson's disease (G20-G21)	564	1.6	54.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	558	1.6	53.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	554	1.6	53.5
14	Pneumonitis due to solids and liquids (J69)	241	0.7	23.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	229	0.6	22.1
...	All other causes (residual)	6,041	17.1	583.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	37,292	100.0	9,635.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,502	28.2	2,713.4
2	Malignant neoplasms (C00-C97)	4,753	12.7	1,228.0
3	Alzheimer's disease (G30)	3,090	8.3	798.4
4	Cerebrovascular diseases (I60-I69)	2,759	7.4	712.8
5	Chronic lower respiratory diseases (J40-J47)	1,612	4.3	416.5
6	Diabetes mellitus (E10-E14)	1,334	3.6	344.7
7	Influenza and pneumonia (J09-J18)	1,155	3.1	298.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	767	2.1	198.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	747	2.0	193.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	736	2.0	190.2
11	Parkinson's disease (G20-G21)	547	1.5	141.3
12	Septicemia (A40-A41)	508	1.4	131.3
13	Pneumonitis due to solids and liquids (J69)	322	0.9	83.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	216	0.6	55.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	175	0.5	45.2
...	All other causes (residual)	8,069	21.6	2,084.8
Hispanic, male, all ages				
...	All causes	92,474	100.0	330.1
1	Malignant neoplasms (C00-C97)	19,040	20.6	68.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,884	20.4	67.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,969	9.7	32.0
4	Diabetes mellitus (E10-E14)	4,149	4.5	14.8
5	Cerebrovascular diseases (I60-I69)	4,092	4.4	14.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,911	4.2	14.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,584	2.8	9.2
8	Chronic lower respiratory diseases (J40-J47)	2,433	2.6	8.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,122	2.3	7.6
10	Influenza and pneumonia (J09-J18)	1,975	2.1	7.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,700	1.8	6.1
12	Alzheimer's disease (G30)	1,600	1.7	5.7
13	Certain conditions originating in the perinatal period (P00-P96)	1,369	1.5	4.9
14	Septicemia (A40-A41)	1,319	1.4	4.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,016	1.1	3.6
...	All other causes (residual)	17,311	18.7	61.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	421	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	32.8	6.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	11.2	2.2
3	Malignant neoplasms (C00-C97)	37	8.8	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	7.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.3	*
6	Influenza and pneumonia (J09-J18)	9	2.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	7	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
9	Cerebrovascular diseases (I60-I69)	5	1.2	*
10	Nutritional deficiencies (E40-E64)	3	0.7	*
11	Septicemia (A40-A41)	2	0.5	*
11	Pneumonitis due to solids and liquids (J69)	2	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
13	Aortic aneurysm and dissection (I71)	1	0.2	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	111	26.4	5.3
Hispanic, male, 5-14 years				
...	All causes	638	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	30.3	3.8
2	Malignant neoplasms (C00-C97)	132	20.7	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	6.7	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	5.5	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	3.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	14	2.2	*
8	Cerebrovascular diseases (I60-I69)	12	1.9	*
9	Septicemia (A40-A41)	11	1.7	*
9	Influenza and pneumonia (J09-J18)	11	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
12	Anemias (D50-D64)	4	0.6	*
13	Nutritional deficiencies (E40-E64)	3	0.5	*
13	Meningitis (G00,G03)	3	0.5	*
15	Diseases of appendix (K35-K38)	2	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (residual)	129	20.2	2.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,669	100.0	75.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,572	42.8	32.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	691	18.8	14.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	563	15.3	11.6
4	Malignant neoplasms (C00-C97)	237	6.5	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	92	2.5	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.0	0.8
7	Cerebrovascular diseases (I60-I69)	21	0.6	0.4
8	Influenza and pneumonia (J09-J18)	19	0.5	*
8	Legal intervention (Y35,Y89.0)	19	0.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.4	*
11	Septicemia (A40-A41)	13	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	12	0.3	*
13	Diabetes mellitus (E10-E14)	10	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.2	*
15	Anemias (D50-D64)	5	0.1	*
15	Aortic aneurysm and dissection (I71)	5	0.1	*
15	Pneumonitis due to solids and liquids (J69)	5	0.1	*
...	All other causes (residual)	346	9.4	7.1
Hispanic, male, 25-34 years				
...	All causes	4,836	100.0	103.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,867	38.6	40.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	611	12.6	13.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	604	12.5	12.9
4	Malignant neoplasms (C00-C97)	329	6.8	7.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	313	6.5	6.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	114	2.4	2.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	71	1.5	1.5
8	Cerebrovascular diseases (I60-I69)	70	1.4	1.5
9	Influenza and pneumonia (J09-J18)	64	1.3	1.4
10	Diabetes mellitus (E10-E14)	61	1.3	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	0.7	0.8
12	Legal intervention (Y35,Y89.0)	32	0.7	0.7
13	Septicemia (A40-A41)	28	0.6	0.6
14	Chronic lower respiratory diseases (J40-J47)	24	0.5	0.5
15	Aortic aneurysm and dissection (I71)	20	0.4	0.4
...	All other causes (residual)	593	12.3	12.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	6,099	100.0	149.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,555	25.5	38.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	756	12.4	18.5
3	Malignant neoplasms (C00-C97)	704	11.5	17.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	497	8.1	12.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	409	6.7	10.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	405	6.6	9.9
7	Cerebrovascular diseases (I60-I69)	193	3.2	4.7
8	Diabetes mellitus (E10-E14)	161	2.6	3.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	137	2.2	3.4
10	Influenza and pneumonia (J09-J18)	122	2.0	3.0
11	Septicemia (A40-A41)	79	1.3	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	0.9	1.3
13	Viral hepatitis (B15-B19)	38	0.6	0.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	0.6	0.9
15	Legal intervention (Y35,Y89.0)	33	0.5	0.8
...	All other causes (residual)	917	15.0	22.5
Hispanic, male, 45-54 years				
...	All causes	10,894	100.0	341.0
1	Malignant neoplasms (C00-C97)	2,080	19.1	65.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,045	18.8	64.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,403	12.9	43.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,095	10.1	34.3
5	Diabetes mellitus (E10-E14)	515	4.7	16.1
6	Cerebrovascular diseases (I60-I69)	425	3.9	13.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	400	3.7	12.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	271	2.5	8.5
9	Viral hepatitis (B15-B19)	215	2.0	6.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	199	1.8	6.2
11	Influenza and pneumonia (J09-J18)	176	1.6	5.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	1.5	5.0
13	Septicemia (A40-A41)	152	1.4	4.8
14	Chronic lower respiratory diseases (J40-J47)	96	0.9	3.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	95	0.9	3.0
...	All other causes (residual)	1,568	14.4	49.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	15,398	100.0	787.7
1	Malignant neoplasms (C00-C97)	4,062	26.4	207.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,314	21.5	169.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,321	8.6	67.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	924	6.0	47.3
5	Diabetes mellitus (E10-E14)	910	5.9	46.6
6	Cerebrovascular diseases (I60-I69)	649	4.2	33.2
7	Viral hepatitis (B15-B19)	345	2.2	17.6
8	Influenza and pneumonia (J09-J18)	301	2.0	15.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	271	1.8	13.9
10	Septicemia (A40-A41)	258	1.7	13.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	241	1.6	12.3
12	Chronic lower respiratory diseases (J40-J47)	240	1.6	12.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	169	1.1	8.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	166	1.1	8.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	0.6	4.5
...	All other causes (residual)	2,139	13.9	109.4
Hispanic, male, 65-74 years				
...	All causes	15,948	100.0	1,655.1
1	Malignant neoplasms (C00-C97)	4,798	30.1	497.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,773	23.7	391.6
3	Diabetes mellitus (E10-E14)	986	6.2	102.3
4	Cerebrovascular diseases (I60-I69)	742	4.7	77.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	646	4.1	67.0
6	Chronic lower respiratory diseases (J40-J47)	523	3.3	54.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	502	3.1	52.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	379	2.4	39.3
9	Influenza and pneumonia (J09-J18)	323	2.0	33.5
10	Septicemia (A40-A41)	274	1.7	28.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	203	1.3	21.1
12	Alzheimer's disease (G30)	144	0.9	14.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	127	0.8	13.2
14	Parkinson's disease (G20-G21)	123	0.8	12.8
15	Viral hepatitis (B15-B19)	112	0.7	11.6
...	All other causes (residual)	2,293	14.4	238.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	17,544	100.0	4,103.3
1	Malignant neoplasms (C00-C97)	4,389	25.0	1,026.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,360	24.9	1,019.7
3	Cerebrovascular diseases (I60-I69)	1,051	6.0	245.8
4	Diabetes mellitus (E10-E14)	967	5.5	226.2
5	Chronic lower respiratory diseases (J40-J47)	786	4.5	183.8
6	Alzheimer's disease (G30)	544	3.1	127.2
7	Influenza and pneumonia (J09-J18)	452	2.6	105.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	449	2.6	105.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	394	2.2	92.2
10	Parkinson's disease (G20-G21)	334	1.9	78.1
11	Septicemia (A40-A41)	284	1.6	66.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	260	1.5	60.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	259	1.5	60.6
14	Pneumonitis due to solids and liquids (J69)	144	0.8	33.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	127	0.7	29.7
...	All other causes (residual)	2,744	15.6	641.8
Hispanic, male, 85 years and over				
...	All causes	14,395	100.0	10,318.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,168	29.0	2,987.5
2	Malignant neoplasms (C00-C97)	2,267	15.7	1,624.9
3	Cerebrovascular diseases (I60-I69)	916	6.4	656.6
4	Alzheimer's disease (G30)	877	6.1	628.6
5	Chronic lower respiratory diseases (J40-J47)	707	4.9	506.8
6	Diabetes mellitus (E10-E14)	538	3.7	385.6
7	Influenza and pneumonia (J09-J18)	478	3.3	342.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	354	2.5	253.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	331	2.3	237.3
10	Parkinson's disease (G20-G21)	263	1.8	188.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	239	1.7	171.3
12	Septicemia (A40-A41)	195	1.4	139.8
13	Pneumonitis due to solids and liquids (J69)	167	1.2	119.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	92	0.6	65.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	63	0.4	45.2
...	All other causes (residual)	2,740	19.0	1,964.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	76,913	100.0	281.0
1	Malignant neoplasms (C00-C97)	17,407	22.6	63.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,137	19.7	55.3
3	Cerebrovascular diseases (I60-I69)	4,621	6.0	16.9
4	Diabetes mellitus (E10-E14)	3,646	4.7	13.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,444	4.5	12.6
6	Alzheimer's disease (G30)	3,334	4.3	12.2
7	Chronic lower respiratory diseases (J40-J47)	2,362	3.1	8.6
8	Influenza and pneumonia (J09-J18)	1,900	2.5	6.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,747	2.3	6.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,573	2.0	5.7
11	Septicemia (A40-A41)	1,294	1.7	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,123	1.5	4.1
13	Certain conditions originating in the perinatal period (P00-P96)	1,053	1.4	3.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	873	1.1	3.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	662	0.9	2.4
...	All other causes (residual)	16,737	21.8	61.2
Hispanic, female, 1-4 years				
...	All causes	348	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	23.9	4.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	14.4	2.5
3	Malignant neoplasms (C00-C97)	44	12.6	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	11.2	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	4.3	*
6	Influenza and pneumonia (J09-J18)	11	3.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	2.0	*
8	Meningitis (G00,G03)	4	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.1	*
10	Septicemia (A40-A41)	3	0.9	*
10	Acute bronchitis and bronchiolitis (J20-J21)	3	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	3	0.9	*
13	Cerebrovascular diseases (I60-I69)	2	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
15	Anemias (D50-D64)	1	0.3	*
15	Diabetes mellitus (E10-E14)	1	0.3	*
15	Nutritional deficiencies (E40-E64)	1	0.3	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.3	*
15	Hernia (K40-K46)	1	0.3	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	71	20.4	3.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	478	100.0	9.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	120	25.1	2.4
2	Malignant neoplasms (C00-C97)	87	18.2	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	7.7	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.7	0.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	6.5	0.6
6	Cerebrovascular diseases (I60-I69)	15	3.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.7	*
8	Influenza and pneumonia (J09-J18)	12	2.5	*
9	Chronic lower respiratory diseases (J40-J47)	10	2.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
12	Septicemia (A40-A41)	5	1.0	*
13	Diabetes mellitus (E10-E14)	4	0.8	*
14	Hernia (K40-K46)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (residual)	95	19.9	1.9
Hispanic, female, 15-24 years				
...	All causes	1,236	100.0	27.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	472	38.2	10.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	157	12.7	3.4
3	Malignant neoplasms (C00-C97)	127	10.3	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	9.8	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	2.9	0.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	29	2.3	0.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	2.3	0.6
8	Cerebrovascular diseases (I60-I69)	13	1.1	*
9	Influenza and pneumonia (J09-J18)	12	1.0	*
10	Septicemia (A40-A41)	11	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
12	Diabetes mellitus (E10-E14)	8	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
14	Anemias (D50-D64)	5	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (residual)	197	15.9	4.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,739	100.0	41.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	476	27.4	11.3
2	Malignant neoplasms (C00-C97)	302	17.4	7.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	139	8.0	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	5.8	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	4.7	1.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	56	3.2	1.3
7	Cerebrovascular diseases (I60-I69)	42	2.4	1.0
8	Diabetes mellitus (E10-E14)	40	2.3	0.9
9	Influenza and pneumonia (J09-J18)	38	2.2	0.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	37	2.1	0.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	28	1.6	0.7
12	Septicemia (A40-A41)	25	1.4	0.6
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	1.4	0.6
14	Chronic lower respiratory diseases (J40-J47)	18	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.8	*
...	All other causes (residual)	317	18.2	7.5
Hispanic, female, 35-44 years				
...	All causes	3,069	100.0	78.5
1	Malignant neoplasms (C00-C97)	994	32.4	25.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	425	13.8	10.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	278	9.1	7.1
4	Cerebrovascular diseases (I60-I69)	125	4.1	3.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	124	4.0	3.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	108	3.5	2.8
7	Diabetes mellitus (E10-E14)	98	3.2	2.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	85	2.8	2.2
9	Influenza and pneumonia (J09-J18)	71	2.3	1.8
10	Pregnancy, childbirth and the puerperium (O00-O99)	61	2.0	1.6
11	Septicemia (A40-A41)	44	1.4	1.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	36	1.2	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.1	0.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.8	0.7
15	Viral hepatitis (B15-B19)	20	0.7	0.5
...	All other causes (residual)	541	17.6	13.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	5,811	100.0	185.3
1	Malignant neoplasms (C00-C97)	2,153	37.1	68.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	669	11.5	21.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	469	8.1	15.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	327	5.6	10.4
5	Diabetes mellitus (E10-E14)	265	4.6	8.4
6	Cerebrovascular diseases (I60-I69)	241	4.1	7.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	111	1.9	3.5
8	Influenza and pneumonia (J09-J18)	104	1.8	3.3
9	Septicemia (A40-A41)	95	1.6	3.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	1.5	2.9
11	Viral hepatitis (B15-B19)	84	1.4	2.7
12	Chronic lower respiratory diseases (J40-J47)	63	1.1	2.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	60	1.0	1.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	51	0.9	1.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	0.7	1.4
...	All other causes (residual)	986	17.0	31.4
Hispanic, female, 55-64 years				
...	All causes	9,303	100.0	443.3
1	Malignant neoplasms (C00-C97)	3,426	36.8	163.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,474	15.8	70.2
3	Diabetes mellitus (E10-E14)	586	6.3	27.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	423	4.5	20.2
5	Cerebrovascular diseases (I60-I69)	361	3.9	17.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	326	3.5	15.5
7	Influenza and pneumonia (J09-J18)	235	2.5	11.2
8	Septicemia (A40-A41)	204	2.2	9.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	2.2	9.6
10	Chronic lower respiratory diseases (J40-J47)	197	2.1	9.4
11	Viral hepatitis (B15-B19)	135	1.5	6.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	85	0.9	4.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	0.8	3.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.5	2.4
15	Pneumonitis due to solids and liquids (J69)	35	0.4	1.7
...	All other causes (residual)	1,491	16.0	71.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	12,088	100.0	1,036.7
1	Malignant neoplasms (C00-C97)	3,917	32.4	335.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,215	18.3	190.0
3	Diabetes mellitus (E10-E14)	838	6.9	71.9
4	Cerebrovascular diseases (I60-I69)	638	5.3	54.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	425	3.5	36.4
6	Chronic lower respiratory diseases (J40-J47)	372	3.1	31.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	353	2.9	30.3
8	Influenza and pneumonia (J09-J18)	262	2.2	22.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	257	2.1	22.0
10	Septicemia (A40-A41)	229	1.9	19.6
11	Alzheimer's disease (G30)	192	1.6	16.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	162	1.3	13.9
13	Viral hepatitis (B15-B19)	87	0.7	7.5
14	Parkinson's disease (G20-G21)	78	0.6	6.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.5	5.1
...	All other causes (residual)	2,003	16.6	171.8
Hispanic, female, 75-84 years				
...	All causes	17,798	100.0	2,931.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,992	22.4	657.5
2	Malignant neoplasms (C00-C97)	3,865	21.7	636.6
3	Cerebrovascular diseases (I60-I69)	1,334	7.5	219.7
4	Diabetes mellitus (E10-E14)	1,010	5.7	166.3
5	Alzheimer's disease (G30)	894	5.0	147.2
6	Chronic lower respiratory diseases (J40-J47)	766	4.3	126.2
7	Influenza and pneumonia (J09-J18)	463	2.6	76.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	458	2.6	75.4
9	Septicemia (A40-A41)	349	2.0	57.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	348	2.0	57.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	298	1.7	49.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	295	1.7	48.6
13	Parkinson's disease (G20-G21)	230	1.3	37.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	102	0.6	16.8
15	Pneumonitis due to solids and liquids (J69)	97	0.5	16.0
...	All other causes (residual)	3,297	18.5	543.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	22,897	100.0	9,250.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,334	27.7	2,558.9
2	Malignant neoplasms (C00-C97)	2,486	10.9	1,004.3
3	Alzheimer's disease (G30)	2,213	9.7	894.0
4	Cerebrovascular diseases (I60-I69)	1,843	8.0	744.6
5	Chronic lower respiratory diseases (J40-J47)	905	4.0	365.6
6	Diabetes mellitus (E10-E14)	796	3.5	321.6
7	Influenza and pneumonia (J09-J18)	677	3.0	273.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	508	2.2	205.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	413	1.8	166.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	405	1.8	163.6
11	Septicemia (A40-A41)	313	1.4	126.4
12	Parkinson's disease (G20-G21)	284	1.2	114.7
13	Pneumonitis due to solids and liquids (J69)	155	0.7	62.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	124	0.5	50.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	112	0.5	45.2
...	All other causes (residual)	5,329	23.3	2,152.9
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,066,949	100.0	1,028.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	489,926	23.7	243.7
2	Malignant neoplasms (C00-C97)	466,271	22.6	231.9
3	Chronic lower respiratory diseases (J40-J47)	129,519	6.3	64.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104,682	5.1	52.1
5	Cerebrovascular diseases (I60-I69)	102,326	5.0	50.9
6	Alzheimer's disease (G30)	80,014	3.9	39.8
7	Diabetes mellitus (E10-E14)	51,944	2.5	25.8
8	Influenza and pneumonia (J09-J18)	43,377	2.1	21.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35,444	1.7	17.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,691	1.7	17.3
11	Septicemia (A40-A41)	28,868	1.4	14.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	27,830	1.3	13.8
13	Parkinson's disease (G20-G21)	22,917	1.1	11.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21,511	1.0	10.7
15	Pneumonitis due to solids and liquids (J69)	15,644	0.8	7.8
...	All other causes (residual)	411,985	19.9	204.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,876	100.0	22.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	655	34.9	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	9.6	2.2
3	Malignant neoplasms (C00-C97)	176	9.4	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	7.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	3.0	0.7
6	Influenza and pneumonia (J09-J18)	53	2.8	0.6
7	Septicemia (A40-A41)	32	1.7	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.1	0.2
9	Chronic lower respiratory diseases (J40-J47)	18	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	14	0.7	*
11	Cerebrovascular diseases (I60-I69)	13	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
13	Meningitis (G00,G03)	8	0.4	*
14	Anemias (D50-D64)	6	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.3	*
...	All other causes (residual)	493	26.3	5.9
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,684	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	769	28.7	3.5
2	Malignant neoplasms (C00-C97)	451	16.8	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	285	10.6	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	176	6.6	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	106	3.9	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.4	0.4
7	Influenza and pneumonia (J09-J18)	52	1.9	0.2
8	Cerebrovascular diseases (I60-I69)	42	1.6	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	1.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	37	1.4	0.2
11	Septicemia (A40-A41)	26	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	15	0.6	*
13	Diabetes mellitus (E10-E14)	12	0.4	*
14	Anemias (D50-D64)	8	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.2	*
...	All other causes (residual)	562	20.9	2.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	16,040	100.0	64.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,743	48.3	31.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,503	21.8	14.1
3	Malignant neoplasms (C00-C97)	859	5.4	3.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	607	3.8	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	493	3.1	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	222	1.4	0.9
7	Influenza and pneumonia (J09-J18)	123	0.8	0.5
8	Diabetes mellitus (E10-E14)	101	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	88	0.5	0.4
10	Chronic lower respiratory diseases (J40-J47)	75	0.5	0.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	71	0.4	0.3
12	Septicemia (A40-A41)	65	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.2	0.1
15	Legal intervention (Y35,Y89.0)	27	0.2	0.1
...	All other causes (residual)	1,986	12.4	8.0
Non-Hispanic, white, both sexes, 25-34 years				
...	All causes	28,969	100.0	115.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,348	42.6	49.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,908	16.9	19.5
3	Malignant neoplasms (C00-C97)	2,144	7.4	8.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,794	6.2	7.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	980	3.4	3.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	441	1.5	1.8
7	Diabetes mellitus (E10-E14)	337	1.2	1.3
8	Influenza and pneumonia (J09-J18)	322	1.1	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	288	1.0	1.1
10	Cerebrovascular diseases (I60-I69)	281	1.0	1.1
11	Septicemia (A40-A41)	205	0.7	0.8
12	Chronic lower respiratory diseases (J40-J47)	192	0.7	0.8
13	Pregnancy, childbirth and the puerperium (O00-O99)	188	0.6	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	0.4	0.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	101	0.3	0.4
...	All other causes (residual)	4,325	14.9	17.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-44 years				
...	All causes	44,565	100.0	185.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,591	26.0	48.3
2	Malignant neoplasms (C00-C97)	6,934	15.6	28.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,109	13.7	25.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,372	12.1	22.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,644	3.7	6.9
6	Diabetes mellitus (E10-E14)	1,093	2.5	4.6
7	Cerebrovascular diseases (I60-I69)	818	1.8	3.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	799	1.8	3.3
9	Influenza and pneumonia (J09-J18)	674	1.5	2.8
10	Chronic lower respiratory diseases (J40-J47)	490	1.1	2.0
11	Septicemia (A40-A41)	482	1.1	2.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	334	0.7	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	286	0.6	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	276	0.6	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	170	0.4	0.7
...	All other causes (residual)	7,493	16.8	31.2
Non-Hispanic, white, both sexes, 45-54 years				
...	All causes	120,679	100.0	417.8
1	Malignant neoplasms (C00-C97)	31,132	25.8	107.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,162	19.2	80.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,522	12.9	53.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,594	6.3	26.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,110	5.1	21.2
6	Diabetes mellitus (E10-E14)	3,570	3.0	12.4
7	Chronic lower respiratory diseases (J40-J47)	3,513	2.9	12.2
8	Cerebrovascular diseases (I60-I69)	2,909	2.4	10.1
9	Influenza and pneumonia (J09-J18)	1,837	1.5	6.4
10	Septicemia (A40-A41)	1,597	1.3	5.5
11	Viral hepatitis (B15-B19)	1,213	1.0	4.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,044	0.9	3.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	882	0.7	3.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	760	0.6	2.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	749	0.6	2.6
...	All other causes (residual)	19,085	15.8	66.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-64 years				
...	All causes	252,298	100.0	864.3
1	Malignant neoplasms (C00-C97)	85,576	33.9	293.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,711	20.9	180.6
3	Chronic lower respiratory diseases (J40-J47)	13,905	5.5	47.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,670	5.4	46.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	9,305	3.7	31.9
6	Diabetes mellitus (E10-E14)	8,356	3.3	28.6
7	Cerebrovascular diseases (I60-I69)	7,170	2.8	24.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,755	2.7	23.1
9	Septicemia (A40-A41)	3,777	1.5	12.9
10	Influenza and pneumonia (J09-J18)	3,710	1.5	12.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,833	1.1	9.7
12	Viral hepatitis (B15-B19)	2,638	1.0	9.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,968	0.8	6.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,232	0.5	4.2
15	Aortic aneurysm and dissection (I71)	1,009	0.4	3.5
...	All other causes (residual)	37,683	14.9	129.1
Non-Hispanic, white, both sexes, 65-74 years				
...	All causes	367,779	100.0	1,798.8
1	Malignant neoplasms (C00-C97)	126,710	34.5	619.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,376	21.0	378.5
3	Chronic lower respiratory diseases (J40-J47)	31,561	8.6	154.4
4	Cerebrovascular diseases (I60-I69)	13,917	3.8	68.1
5	Diabetes mellitus (E10-E14)	12,396	3.4	60.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,520	2.6	46.6
7	Septicemia (A40-A41)	5,997	1.6	29.3
8	Influenza and pneumonia (J09-J18)	5,939	1.6	29.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,916	1.6	28.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	5,890	1.6	28.8
11	Alzheimer's disease (G30)	4,263	1.2	20.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,733	1.0	18.3
13	Parkinson's disease (G20-G21)	2,925	0.8	14.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,898	0.8	14.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,440	0.7	11.9
...	All other causes (residual)	56,298	15.3	275.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-84 years				
...	All causes	510,281	100.0	4,700.7
1	Malignant neoplasms (C00-C97)	126,707	24.8	1,167.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	118,301	23.2	1,089.8
3	Chronic lower respiratory diseases (J40-J47)	42,747	8.4	393.8
4	Cerebrovascular diseases (I60-I69)	28,444	5.6	262.0
5	Alzheimer's disease (G30)	21,373	4.2	196.9
6	Diabetes mellitus (E10-E14)	13,556	2.7	124.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,785	2.5	117.8
8	Influenza and pneumonia (J09-J18)	10,444	2.0	96.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,049	2.0	92.6
10	Parkinson's disease (G20-G21)	9,497	1.9	87.5
11	Septicemia (A40-A41)	7,709	1.5	71.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,963	1.0	45.7
13	Pneumonitis due to solids and liquids (J69)	4,132	0.8	38.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,104	0.8	37.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,251	0.6	29.9
...	All other causes (residual)	92,219	18.1	849.5
Non-Hispanic, white, both sexes, 85 years and over				
...	All causes	711,340	100.0	13,977.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	209,684	29.5	4,120.2
2	Malignant neoplasms (C00-C97)	85,539	12.0	1,680.8
3	Alzheimer's disease (G30)	53,634	7.5	1,053.9
4	Cerebrovascular diseases (I60-I69)	48,591	6.8	954.8
5	Chronic lower respiratory diseases (J40-J47)	36,970	5.2	726.4
6	Influenza and pneumonia (J09-J18)	20,152	2.8	396.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,505	2.7	383.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,358	2.0	282.1
9	Diabetes mellitus (E10-E14)	12,520	1.8	246.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,706	1.5	210.4
11	Parkinson's disease (G20-G21)	9,961	1.4	195.7
12	Septicemia (A40-A41)	8,920	1.3	175.3
13	Pneumonitis due to solids and liquids (J69)	7,992	1.1	157.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,752	0.7	93.4
15	Atherosclerosis (I70)	3,170	0.4	62.3
...	All other causes (residual)	164,886	23.2	3,239.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	1,035,345	100.0	1,045.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	258,473	25.0	261.0
2	Malignant neoplasms (C00-C97)	246,872	23.8	249.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63,807	6.2	64.4
4	Chronic lower respiratory diseases (J40-J47)	60,428	5.8	61.0
5	Cerebrovascular diseases (I60-I69)	41,410	4.0	41.8
6	Diabetes mellitus (E10-E14)	28,743	2.8	29.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,410	2.6	27.7
8	Alzheimer's disease (G30)	24,321	2.3	24.6
9	Influenza and pneumonia (J09-J18)	20,633	2.0	20.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,878	1.7	18.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	17,844	1.7	18.0
12	Parkinson's disease (G20-G21)	13,830	1.3	14.0
13	Septicemia (A40-A41)	13,514	1.3	13.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,984	0.9	9.1
15	Pneumonitis due to solids and liquids (J69)	8,623	0.8	8.7
...	All other causes (residual)	182,575	17.6	184.3
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,060	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	36.9	9.2
2	Malignant neoplasms (C00-C97)	110	10.4	2.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	9.5	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	7.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	2.7	0.7
5	Influenza and pneumonia (J09-J18)	29	2.7	0.7
7	Septicemia (A40-A41)	13	1.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	10	0.9	*
10	Cerebrovascular diseases (I60-I69)	7	0.7	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.7	*
12	Meningitis (G00,G03)	4	0.4	*
13	Anemias (D50-D64)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
15	Diseases of appendix (K35-K38)	2	0.2	*
...	All other causes (residual)	259	24.4	6.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,620	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	511	31.5	4.5
2	Malignant neoplasms (C00-C97)	256	15.8	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	191	11.8	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	5.7	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	4.1	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	2.8	0.4
7	Influenza and pneumonia (J09-J18)	26	1.6	0.2
8	Chronic lower respiratory diseases (J40-J47)	25	1.5	0.2
9	Cerebrovascular diseases (I60-I69)	24	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.2	*
11	Septicemia (A40-A41)	10	0.6	*
11	Diabetes mellitus (E10-E14)	10	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	9	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
15	Anemias (D50-D64)	4	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.2	*
...	All other causes (residual)	322	19.9	2.8
Non-Hispanic, white, male, 15-24 years				
...	All causes	11,508	100.0	90.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,707	49.6	44.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,850	24.8	22.4
3	Malignant neoplasms (C00-C97)	509	4.4	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	447	3.9	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	318	2.8	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	1.1	1.0
7	Diabetes mellitus (E10-E14)	69	0.6	0.5
8	Influenza and pneumonia (J09-J18)	66	0.6	0.5
9	Chronic lower respiratory diseases (J40-J47)	49	0.4	0.4
10	Cerebrovascular diseases (I60-I69)	48	0.4	0.4
11	Septicemia (A40-A41)	38	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.2	0.2
12	Legal intervention (Y35,Y89.0)	24	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.1	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.1	*
...	All other causes (residual)	1,203	10.5	9.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-34 years				
...	All causes	19,845	100.0	155.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,991	45.3	70.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,883	19.6	30.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,213	6.1	9.5
4	Malignant neoplasms (C00-C97)	1,097	5.5	8.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	672	3.4	5.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	253	1.3	2.0
7	Diabetes mellitus (E10-E14)	217	1.1	1.7
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	176	0.9	1.4
9	Influenza and pneumonia (J09-J18)	164	0.8	1.3
10	Cerebrovascular diseases (I60-I69)	152	0.8	1.2
11	Chronic lower respiratory diseases (J40-J47)	114	0.6	0.9
12	Septicemia (A40-A41)	104	0.5	0.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	82	0.4	0.6
14	Legal intervention (Y35,Y89.0)	76	0.4	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	0.3	0.5
...	All other causes (residual)	2,583	13.0	20.3
Non-Hispanic, white, male, 35-44 years				
...	All causes	27,413	100.0	227.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,792	28.4	64.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,207	15.3	34.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,995	14.6	33.1
4	Malignant neoplasms (C00-C97)	3,087	11.3	25.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	995	3.6	8.3
6	Diabetes mellitus (E10-E14)	686	2.5	5.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	549	2.0	4.6
8	Cerebrovascular diseases (I60-I69)	442	1.6	3.7
9	Influenza and pneumonia (J09-J18)	356	1.3	3.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	250	0.9	2.1
11	Septicemia (A40-A41)	230	0.8	1.9
12	Chronic lower respiratory diseases (J40-J47)	217	0.8	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	176	0.6	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	143	0.5	1.2
15	Aortic aneurysm and dissection (I71)	130	0.5	1.1
...	All other causes (residual)	4,158	15.2	34.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-54 years				
...	All causes	73,411	100.0	511.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,511	22.5	115.0
2	Malignant neoplasms (C00-C97)	15,578	21.2	108.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,230	13.9	71.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,535	7.5	38.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,992	5.4	27.8
6	Diabetes mellitus (E10-E14)	2,285	3.1	15.9
7	Cerebrovascular diseases (I60-I69)	1,621	2.2	11.3
8	Chronic lower respiratory diseases (J40-J47)	1,584	2.2	11.0
9	Influenza and pneumonia (J09-J18)	984	1.3	6.9
10	Septicemia (A40-A41)	848	1.2	5.9
11	Viral hepatitis (B15-B19)	823	1.1	5.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	653	0.9	4.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	636	0.9	4.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	615	0.8	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	497	0.7	3.5
...	All other causes (residual)	11,019	15.0	76.7
Non-Hispanic, white, male, 55-64 years				
...	All causes	154,680	100.0	1,085.3
1	Malignant neoplasms (C00-C97)	47,744	30.9	335.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,564	24.3	263.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,084	5.9	63.7
4	Chronic lower respiratory diseases (J40-J47)	7,189	4.6	50.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,412	4.1	45.0
6	Diabetes mellitus (E10-E14)	5,313	3.4	37.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,097	3.3	35.8
8	Cerebrovascular diseases (I60-I69)	4,092	2.6	28.7
9	Influenza and pneumonia (J09-J18)	2,111	1.4	14.8
10	Septicemia (A40-A41)	1,990	1.3	14.0
11	Viral hepatitis (B15-B19)	1,868	1.2	13.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,657	1.1	11.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,249	0.8	8.8
14	Aortic aneurysm and dissection (I71)	765	0.5	5.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	743	0.5	5.2
...	All other causes (residual)	21,802	14.1	153.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-74 years				
...	All causes	210,808	100.0	2,170.0
1	Malignant neoplasms (C00-C97)	70,155	33.3	722.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,411	23.9	518.9
3	Chronic lower respiratory diseases (J40-J47)	15,784	7.5	162.5
4	Diabetes mellitus (E10-E14)	7,525	3.6	77.5
5	Cerebrovascular diseases (I60-I69)	7,447	3.5	76.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,969	2.8	61.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,799	1.8	39.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,267	1.5	33.6
9	Influenza and pneumonia (J09-J18)	3,260	1.5	33.6
10	Septicemia (A40-A41)	3,100	1.5	31.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,987	1.4	30.7
12	Parkinson's disease (G20-G21)	1,953	0.9	20.1
13	Alzheimer's disease (G30)	1,837	0.9	18.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,629	0.8	16.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,483	0.7	15.3
...	All other causes (residual)	30,202	14.3	310.9
Non-Hispanic, white, male, 75-84 years				
...	All causes	261,374	100.0	5,499.0
1	Malignant neoplasms (C00-C97)	67,689	25.9	1,424.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	65,695	25.1	1,382.1
3	Chronic lower respiratory diseases (J40-J47)	20,397	7.8	429.1
4	Cerebrovascular diseases (I60-I69)	12,636	4.8	265.8
5	Alzheimer's disease (G30)	7,981	3.1	167.9
6	Diabetes mellitus (E10-E14)	7,322	2.8	154.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,993	2.7	147.1
8	Parkinson's disease (G20-G21)	6,119	2.3	128.7
9	Influenza and pneumonia (J09-J18)	5,397	2.1	113.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,392	2.1	113.4
11	Septicemia (A40-A41)	3,642	1.4	76.6
12	Pneumonitis due to solids and liquids (J69)	2,423	0.9	51.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,371	0.9	49.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,179	0.8	45.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,879	0.7	39.5
...	All other causes (residual)	43,259	16.6	910.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85 years and over				
...	All causes	267,765	100.0	15,286.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,395	30.8	4,703.8
2	Malignant neoplasms (C00-C97)	40,622	15.2	2,319.1
3	Chronic lower respiratory diseases (J40-J47)	15,049	5.6	859.1
4	Cerebrovascular diseases (I60-I69)	14,910	5.6	851.2
5	Alzheimer's disease (G30)	14,189	5.3	810.0
6	Influenza and pneumonia (J09-J18)	8,202	3.1	468.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,804	2.9	445.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,621	2.5	378.0
9	Parkinson's disease (G20-G21)	5,409	2.0	308.8
10	Diabetes mellitus (E10-E14)	5,314	2.0	303.4
11	Pneumonitis due to solids and liquids (J69)	4,117	1.5	235.0
12	Septicemia (A40-A41)	3,506	1.3	200.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,271	1.2	186.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,181	0.8	124.5
15	Aortic aneurysm and dissection (I71)	1,040	0.4	59.4
...	All other causes (residual)	53,135	19.8	3,033.4
Non-Hispanic, white, female, all ages				
...	All causes	1,031,604	100.0	1,011.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	231,453	22.4	226.9
2	Malignant neoplasms (C00-C97)	219,399	21.3	215.1
3	Chronic lower respiratory diseases (J40-J47)	69,091	6.7	67.7
4	Cerebrovascular diseases (I60-I69)	60,916	5.9	59.7
5	Alzheimer's disease (G30)	55,693	5.4	54.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40,875	4.0	40.1
7	Diabetes mellitus (E10-E14)	23,201	2.2	22.7
8	Influenza and pneumonia (J09-J18)	22,744	2.2	22.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,813	1.6	16.5
10	Septicemia (A40-A41)	15,354	1.5	15.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,527	1.2	12.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,986	1.0	9.8
13	Parkinson's disease (G20-G21)	9,087	0.9	8.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,034	0.8	7.9
15	Pneumonitis due to solids and liquids (J69)	7,021	0.7	6.9
...	All other causes (residual)	229,410	22.2	224.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	816	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	264	32.4	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	9.7	2.0
3	Malignant neoplasms (C00-C97)	66	8.1	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	7.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	3.3	0.7
6	Influenza and pneumonia (J09-J18)	24	2.9	0.6
7	Septicemia (A40-A41)	19	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	8	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.9	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.7	*
13	Meningitis (G00,G03)	4	0.5	*
14	Anemias (D50-D64)	3	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.4	*
14	Pneumonitis due to solids and liquids (J69)	3	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	3	0.4	*
...	All other causes (residual)	221	27.1	5.5
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,064	100.0	9.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	24.2	2.4
2	Malignant neoplasms (C00-C97)	195	18.3	1.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	8.8	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	7.9	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	4.1	0.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	3.8	0.4
7	Influenza and pneumonia (J09-J18)	26	2.4	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.8	*
9	Cerebrovascular diseases (I60-I69)	18	1.7	*
10	Septicemia (A40-A41)	16	1.5	*
11	Chronic lower respiratory diseases (J40-J47)	12	1.1	*
12	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
13	Anemias (D50-D64)	4	0.4	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
15	Tuberculosis (A16-A19)	2	0.2	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
...	All other causes (residual)	241	22.7	2.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,532	100.0	37.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,036	44.9	16.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	653	14.4	5.4
3	Malignant neoplasms (C00-C97)	350	7.7	2.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	175	3.9	1.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	3.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	2.1	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	71	1.6	0.6
8	Influenza and pneumonia (J09-J18)	57	1.3	0.5
9	Cerebrovascular diseases (I60-I69)	40	0.9	0.3
10	Diabetes mellitus (E10-E14)	32	0.7	0.3
11	Septicemia (A40-A41)	27	0.6	0.2
12	Chronic lower respiratory diseases (J40-J47)	26	0.6	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.3	*
15	Pneumonitis due to solids and liquids (J69)	11	0.2	*
...	All other causes (residual)	760	16.8	6.3
Non-Hispanic, white, female, 25-34 years				
...	All causes	9,124	100.0	73.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,357	36.8	26.9
2	Malignant neoplasms (C00-C97)	1,047	11.5	8.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,025	11.2	8.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	581	6.4	4.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	308	3.4	2.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	188	2.1	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	188	2.1	1.5
8	Influenza and pneumonia (J09-J18)	158	1.7	1.3
9	Cerebrovascular diseases (I60-I69)	129	1.4	1.0
10	Diabetes mellitus (E10-E14)	120	1.3	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	1.2	0.9
12	Septicemia (A40-A41)	101	1.1	0.8
13	Chronic lower respiratory diseases (J40-J47)	78	0.9	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	0.5	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.4	0.3
...	All other causes (residual)	1,650	18.1	13.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-44 years				
...	All causes	17,152	100.0	143.8
1	Malignant neoplasms (C00-C97)	3,847	22.4	32.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,799	22.1	31.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,902	11.1	15.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,377	8.0	11.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	649	3.8	5.4
6	Diabetes mellitus (E10-E14)	407	2.4	3.4
7	Cerebrovascular diseases (I60-I69)	376	2.2	3.2
8	Influenza and pneumonia (J09-J18)	318	1.9	2.7
9	Chronic lower respiratory diseases (J40-J47)	273	1.6	2.3
10	Septicemia (A40-A41)	252	1.5	2.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	250	1.5	2.1
12	Pregnancy, childbirth and the puerperium (O00-O99)	159	0.9	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	133	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	0.6	0.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	84	0.5	0.7
...	All other causes (residual)	3,216	18.8	27.0
Non-Hispanic, white, female, 45-54 years				
...	All causes	47,268	100.0	325.5
1	Malignant neoplasms (C00-C97)	15,554	32.9	107.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,651	14.1	45.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,292	11.2	36.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,118	4.5	14.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,059	4.4	14.2
6	Chronic lower respiratory diseases (J40-J47)	1,929	4.1	13.3
7	Cerebrovascular diseases (I60-I69)	1,288	2.7	8.9
8	Diabetes mellitus (E10-E14)	1,285	2.7	8.8
9	Influenza and pneumonia (J09-J18)	853	1.8	5.9
10	Septicemia (A40-A41)	749	1.6	5.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	391	0.8	2.7
12	Viral hepatitis (B15-B19)	390	0.8	2.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	285	0.6	2.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	267	0.6	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	252	0.5	1.7
...	All other causes (residual)	7,905	16.7	54.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-64 years				
...	All causes	97,618	100.0	653.5
1	Malignant neoplasms (C00-C97)	37,832	38.8	253.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,147	15.5	101.4
3	Chronic lower respiratory diseases (J40-J47)	6,716	6.9	45.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,586	4.7	30.7
5	Cerebrovascular diseases (I60-I69)	3,078	3.2	20.6
6	Diabetes mellitus (E10-E14)	3,043	3.1	20.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,893	3.0	19.4
8	Septicemia (A40-A41)	1,787	1.8	12.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,658	1.7	11.1
10	Influenza and pneumonia (J09-J18)	1,599	1.6	10.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,176	1.2	7.9
12	Viral hepatitis (B15-B19)	770	0.8	5.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	719	0.7	4.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	489	0.5	3.3
15	Alzheimer's disease (G30)	395	0.4	2.6
...	All other causes (residual)	15,730	16.1	105.3
Non-Hispanic, white, female, 65-74 years				
...	All causes	156,971	100.0	1,462.9
1	Malignant neoplasms (C00-C97)	56,555	36.0	527.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,965	17.2	251.3
3	Chronic lower respiratory diseases (J40-J47)	15,777	10.1	147.0
4	Cerebrovascular diseases (I60-I69)	6,470	4.1	60.3
5	Diabetes mellitus (E10-E14)	4,871	3.1	45.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,551	2.3	33.1
7	Septicemia (A40-A41)	2,897	1.8	27.0
8	Influenza and pneumonia (J09-J18)	2,679	1.7	25.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,649	1.7	24.7
10	Alzheimer's disease (G30)	2,426	1.5	22.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,091	1.3	19.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,269	0.8	11.8
13	Parkinson's disease (G20-G21)	972	0.6	9.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	957	0.6	8.9
15	Pneumonitis due to solids and liquids (J69)	795	0.5	7.4
...	All other causes (residual)	26,047	16.6	242.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-84 years				
...	All causes	248,907	100.0	4,078.9
1	Malignant neoplasms (C00-C97)	59,018	23.7	967.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,606	21.1	862.1
3	Chronic lower respiratory diseases (J40-J47)	22,350	9.0	366.3
4	Cerebrovascular diseases (I60-I69)	15,808	6.4	259.0
5	Alzheimer's disease (G30)	13,392	5.4	219.5
6	Diabetes mellitus (E10-E14)	6,234	2.5	102.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,792	2.3	94.9
8	Influenza and pneumonia (J09-J18)	5,047	2.0	82.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,657	1.9	76.3
10	Septicemia (A40-A41)	4,067	1.6	66.6
11	Parkinson's disease (G20-G21)	3,378	1.4	55.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,784	1.1	45.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,733	0.7	28.4
14	Pneumonitis due to solids and liquids (J69)	1,709	0.7	28.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,440	0.6	23.6
...	All other causes (residual)	48,892	19.6	801.2
Non-Hispanic, white, female, 85 years and over				
...	All causes	443,575	100.0	13,290.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	127,289	28.7	3,813.8
2	Malignant neoplasms (C00-C97)	44,917	10.1	1,345.8
3	Alzheimer's disease (G30)	39,445	8.9	1,181.9
4	Cerebrovascular diseases (I60-I69)	33,681	7.6	1,009.2
5	Chronic lower respiratory diseases (J40-J47)	21,921	4.9	656.8
6	Influenza and pneumonia (J09-J18)	11,950	2.7	358.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,701	2.6	350.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,737	1.7	231.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,435	1.7	222.8
10	Diabetes mellitus (E10-E14)	7,206	1.6	215.9
11	Septicemia (A40-A41)	5,414	1.2	162.2
12	Parkinson's disease (G20-G21)	4,552	1.0	136.4
13	Pneumonitis due to solids and liquids (J69)	3,875	0.9	116.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,571	0.6	77.0
15	Atherosclerosis (I70)	2,132	0.5	63.9
...	All other causes (residual)	111,749	25.2	3,348.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	303,844	100.0	735.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,894	23.7	174.0
2	Malignant neoplasms (C00-C97)	68,104	22.4	164.8
3	Cerebrovascular diseases (I60-I69)	16,883	5.6	40.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,820	4.5	33.4
5	Diabetes mellitus (E10-E14)	13,264	4.4	32.1
6	Chronic lower respiratory diseases (J40-J47)	9,762	3.2	23.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,496	2.8	20.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,767	2.6	18.8
9	Alzheimer's disease (G30)	6,493	2.1	15.7
10	Septicemia (A40-A41)	6,321	2.1	15.3
11	Influenza and pneumonia (J09-J18)	5,462	1.8	13.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,322	1.8	12.9
13	Certain conditions originating in the perinatal period (P00-P96)	3,810	1.3	9.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,527	1.2	8.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,011	1.0	7.3
...	All other causes (residual)	59,908	19.7	145.0
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	967	100.0	39.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	28.9	11.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	14.8	5.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	10.0	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	5.6	2.2
5	Malignant neoplasms (C00-C97)	50	5.2	2.0
6	Influenza and pneumonia (J09-J18)	28	2.9	1.1
7	Chronic lower respiratory diseases (J40-J47)	20	2.1	0.8
8	Septicemia (A40-A41)	12	1.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
10	Cerebrovascular diseases (I60-I69)	9	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
11	Acute bronchitis and bronchiolitis (J20-J21)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
14	Anemias (D50-D64)	6	0.6	*
15	Meningitis (G00,G03)	4	0.4	*
...	All other causes (residual)	232	24.0	9.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,162	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	327	28.1	5.3
2	Malignant neoplasms (C00-C97)	142	12.2	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	103	8.9	1.7
4	Chronic lower respiratory diseases (J40-J47)	71	6.1	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	6.0	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	5.2	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	4.7	0.9
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.7	0.3
9	Anemias (D50-D64)	17	1.5	*
10	Influenza and pneumonia (J09-J18)	16	1.4	*
11	Septicemia (A40-A41)	15	1.3	*
11	Cerebrovascular diseases (I60-I69)	15	1.3	*
13	Diabetes mellitus (E10-E14)	10	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	9	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (residual)	221	19.0	3.6
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	6,425	100.0	94.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,632	41.0	38.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,466	22.8	21.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	519	8.1	7.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	280	4.4	4.1
5	Malignant neoplasms (C00-C97)	261	4.1	3.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	83	1.3	1.2
7	Chronic lower respiratory diseases (J40-J47)	71	1.1	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	1.0	1.0
9	Anemias (D50-D64)	64	1.0	0.9
10	Diabetes mellitus (E10-E14)	59	0.9	0.9
11	Pregnancy, childbirth and the puerperium (O00-O99)	56	0.9	0.8
12	Cerebrovascular diseases (I60-I69)	48	0.7	0.7
13	Influenza and pneumonia (J09-J18)	36	0.6	0.5
14	Septicemia (A40-A41)	32	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.5	0.4
15	Legal intervention (Y35,Y89.0)	29	0.5	0.4
...	All other causes (residual)	695	10.8	10.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-34 years				
...	All causes	9,373	100.0	157.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,348	25.1	39.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,904	20.3	32.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	998	10.6	16.8
4	Malignant neoplasms (C00-C97)	623	6.6	10.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	553	5.9	9.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	369	3.9	6.2
7	Diabetes mellitus (E10-E14)	233	2.5	3.9
8	Cerebrovascular diseases (I60-I69)	151	1.6	2.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	136	1.5	2.3
10	Anemias (D50-D64)	125	1.3	2.1
11	Chronic lower respiratory diseases (J40-J47)	121	1.3	2.0
12	Influenza and pneumonia (J09-J18)	98	1.0	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	0.9	1.4
14	Septicemia (A40-A41)	81	0.9	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	0.7	1.0
...	All other causes (residual)	1,488	15.9	25.1
Non-Hispanic, black, both sexes, 35-44 years				
...	All causes	13,773	100.0	261.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,746	19.9	52.2
2	Malignant neoplasms (C00-C97)	1,943	14.1	36.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,813	13.2	34.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,201	8.7	22.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	633	4.6	12.0
6	Diabetes mellitus (E10-E14)	544	3.9	10.3
7	Cerebrovascular diseases (I60-I69)	507	3.7	9.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	429	3.1	8.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	1.9	5.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	195	1.4	3.7
10	Influenza and pneumonia (J09-J18)	195	1.4	3.7
12	Septicemia (A40-A41)	187	1.4	3.6
13	Chronic lower respiratory diseases (J40-J47)	185	1.3	3.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	178	1.3	3.4
15	Anemias (D50-D64)	141	1.0	2.7
...	All other causes (residual)	2,611	19.0	49.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-54 years				
...	All causes	31,297	100.0	577.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,572	24.2	139.7
2	Malignant neoplasms (C00-C97)	7,366	23.5	135.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,473	7.9	45.6
4	Cerebrovascular diseases (I60-I69)	1,455	4.6	26.8
5	Diabetes mellitus (E10-E14)	1,426	4.6	26.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,084	3.5	20.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	717	2.3	13.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	702	2.2	13.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	651	2.1	12.0
10	Chronic lower respiratory diseases (J40-J47)	643	2.1	11.9
11	Septicemia (A40-A41)	584	1.9	10.8
12	Influenza and pneumonia (J09-J18)	488	1.6	9.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	476	1.5	8.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	359	1.1	6.6
15	Viral hepatitis (B15-B19)	183	0.6	3.4
...	All other causes (residual)	5,118	16.4	94.4
Non-Hispanic, black, both sexes, 55-64 years				
...	All causes	59,079	100.0	1,296.9
1	Malignant neoplasms (C00-C97)	17,817	30.2	391.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,392	24.4	315.9
3	Cerebrovascular diseases (I60-I69)	2,934	5.0	64.4
4	Diabetes mellitus (E10-E14)	2,865	4.8	62.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,421	4.1	53.1
6	Chronic lower respiratory diseases (J40-J47)	1,805	3.1	39.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,560	2.6	34.2
8	Septicemia (A40-A41)	1,311	2.2	28.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,277	2.2	28.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,006	1.7	22.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	976	1.7	21.4
12	Influenza and pneumonia (J09-J18)	912	1.5	20.0
13	Viral hepatitis (B15-B19)	636	1.1	14.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	349	0.6	7.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	239	0.4	5.2
...	All other causes (residual)	8,579	14.5	188.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-74 years				
...	All causes	60,068	100.0	2,432.6
1	Malignant neoplasms (C00-C97)	18,437	30.7	746.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,881	24.8	602.6
3	Cerebrovascular diseases (I60-I69)	3,476	5.8	140.8
4	Diabetes mellitus (E10-E14)	3,178	5.3	128.7
5	Chronic lower respiratory diseases (J40-J47)	2,551	4.2	103.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,065	3.4	83.6
7	Septicemia (A40-A41)	1,471	2.4	59.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,160	1.9	47.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,004	1.7	40.7
10	Influenza and pneumonia (J09-J18)	986	1.6	39.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	577	1.0	23.4
12	Alzheimer's disease (G30)	486	0.8	19.7
13	Pneumonitis due to solids and liquids (J69)	359	0.6	14.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	306	0.5	12.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	295	0.5	11.9
...	All other causes (residual)	8,836	14.7	357.8
Non-Hispanic, black, both sexes, 75-84 years				
...	All causes	59,483	100.0	5,144.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,295	25.7	1,322.7
2	Malignant neoplasms (C00-C97)	14,072	23.7	1,217.0
3	Cerebrovascular diseases (I60-I69)	4,124	6.9	356.6
4	Diabetes mellitus (E10-E14)	2,969	5.0	256.8
5	Chronic lower respiratory diseases (J40-J47)	2,508	4.2	216.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,064	3.5	178.5
7	Alzheimer's disease (G30)	2,058	3.5	178.0
8	Septicemia (A40-A41)	1,371	2.3	118.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,204	2.0	104.1
10	Influenza and pneumonia (J09-J18)	1,174	2.0	101.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	849	1.4	73.4
12	Parkinson's disease (G20-G21)	483	0.8	41.8
13	Pneumonitis due to solids and liquids (J69)	481	0.8	41.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	308	0.5	26.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	184	0.3	15.9
...	All other causes (residual)	10,339	17.4	894.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85 years and over				
...	All causes	55,499	100.0	12,331.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,498	27.9	3,443.5
2	Malignant neoplasms (C00-C97)	7,383	13.3	1,640.4
3	Cerebrovascular diseases (I60-I69)	4,141	7.5	920.1
4	Alzheimer's disease (G30)	3,844	6.9	854.1
5	Diabetes mellitus (E10-E14)	1,978	3.6	439.5
6	Chronic lower respiratory diseases (J40-J47)	1,779	3.2	395.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,752	3.2	389.3
8	Influenza and pneumonia (J09-J18)	1,458	2.6	324.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,384	2.5	307.5
10	Septicemia (A40-A41)	1,205	2.2	267.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	723	1.3	160.6
12	Pneumonitis due to solids and liquids (J69)	514	0.9	114.2
13	Parkinson's disease (G20-G21)	409	0.7	90.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	280	0.5	62.2
15	Nutritional deficiencies (E40-E64)	194	0.3	43.1
...	All other causes (residual)	12,957	23.3	2,878.9
Non-Hispanic, black, male, all ages				
...	All causes	154,836	100.0	783.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,265	24.1	188.5
2	Malignant neoplasms (C00-C97)	34,539	22.3	174.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,316	6.0	47.1
4	Cerebrovascular diseases (I60-I69)	7,650	4.9	38.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,718	4.3	34.0
6	Diabetes mellitus (E10-E14)	6,369	4.1	32.2
7	Chronic lower respiratory diseases (J40-J47)	4,940	3.2	25.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,983	2.6	20.2
9	Septicemia (A40-A41)	2,938	1.9	14.9
10	Influenza and pneumonia (J09-J18)	2,656	1.7	13.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,440	1.6	12.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,321	1.5	11.7
13	Certain conditions originating in the perinatal period (P00-P96)	2,134	1.4	10.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,876	1.2	9.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,873	1.2	9.5
...	All other causes (residual)	27,818	18.0	140.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	560	100.0	45.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	30.9	13.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	13.9	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	10.0	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.6	2.1
5	Malignant neoplasms (C00-C97)	25	4.5	2.0
6	Influenza and pneumonia (J09-J18)	18	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.6	*
9	Septicemia (A40-A41)	7	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
11	Anemias (D50-D64)	5	0.9	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
14	Meningitis (G00,G03)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (residual)	133	23.8	10.7
Non-Hispanic, black, male, 5-14 years				
...	All causes	701	100.0	22.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	31.2	7.0
2	Malignant neoplasms (C00-C97)	75	10.7	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	9.8	2.2
4	Chronic lower respiratory diseases (J40-J47)	47	6.7	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	5.7	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	5.7	1.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	5.7	1.3
8	Influenza and pneumonia (J09-J18)	9	1.3	*
9	Septicemia (A40-A41)	8	1.1	*
9	Anemias (D50-D64)	8	1.1	*
11	Cerebrovascular diseases (I60-I69)	7	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
13	Diabetes mellitus (E10-E14)	5	0.7	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
15	Diseases of appendix (K35-K38)	2	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	114	16.3	3.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	4,935	100.0	142.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,388	48.4	69.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,092	22.1	31.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	428	8.7	12.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	3.4	4.9
5	Malignant neoplasms (C00-C97)	153	3.1	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	49	1.0	1.4
7	Chronic lower respiratory diseases (J40-J47)	47	1.0	1.4
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	0.7	1.1
9	Diabetes mellitus (E10-E14)	33	0.7	1.0
10	Anemias (D50-D64)	32	0.6	0.9
11	Cerebrovascular diseases (I60-I69)	29	0.6	0.8
11	Legal intervention (Y35,Y89.0)	29	0.6	0.8
13	Influenza and pneumonia (J09-J18)	26	0.5	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
15	Septicemia (A40-A41)	12	0.2	*
...	All other causes (residual)	393	8.0	11.4
Non-Hispanic, black, male, 25-34 years				
...	All causes	6,498	100.0	225.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,099	32.3	72.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,420	21.9	49.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	622	9.6	21.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	466	7.2	16.2
5	Malignant neoplasms (C00-C97)	256	3.9	8.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	247	3.8	8.6
7	Diabetes mellitus (E10-E14)	128	2.0	4.4
8	Cerebrovascular diseases (I60-I69)	89	1.4	3.1
9	Anemias (D50-D64)	78	1.2	2.7
10	Chronic lower respiratory diseases (J40-J47)	71	1.1	2.5
11	Influenza and pneumonia (J09-J18)	47	0.7	1.6
12	Septicemia (A40-A41)	43	0.7	1.5
12	Legal intervention (Y35,Y89.0)	43	0.7	1.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.6	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	0.5	1.1
...	All other causes (residual)	815	12.5	28.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-44 years				
...	All causes	8,066	100.0	325.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,699	21.1	68.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,258	15.6	50.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,037	12.9	41.8
4	Malignant neoplasms (C00-C97)	755	9.4	30.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	360	4.5	14.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	342	4.2	13.8
7	Cerebrovascular diseases (I60-I69)	300	3.7	12.1
8	Diabetes mellitus (E10-E14)	298	3.7	12.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	150	1.9	6.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	128	1.6	5.2
11	Influenza and pneumonia (J09-J18)	100	1.2	4.0
12	Chronic lower respiratory diseases (J40-J47)	99	1.2	4.0
13	Septicemia (A40-A41)	91	1.1	3.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	87	1.1	3.5
15	Aortic aneurysm and dissection (I71)	74	0.9	3.0
...	All other causes (residual)	1,288	16.0	52.0
Non-Hispanic, black, male, 45-54 years				
...	All causes	17,668	100.0	694.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,728	26.8	185.9
2	Malignant neoplasms (C00-C97)	3,392	19.2	133.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,689	9.6	66.4
4	Diabetes mellitus (E10-E14)	815	4.6	32.0
5	Cerebrovascular diseases (I60-I69)	783	4.4	30.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	708	4.0	27.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	583	3.3	22.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	416	2.4	16.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	359	2.0	14.1
10	Septicemia (A40-A41)	297	1.7	11.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	287	1.6	11.3
12	Chronic lower respiratory diseases (J40-J47)	274	1.6	10.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	1.5	10.7
14	Influenza and pneumonia (J09-J18)	254	1.4	10.0
15	Viral hepatitis (B15-B19)	116	0.7	4.6
...	All other causes (residual)	2,695	15.3	105.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-64 years				
...	All causes	34,268	100.0	1,650.4
1	Malignant neoplasms (C00-C97)	9,617	28.1	463.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,020	26.3	434.4
3	Cerebrovascular diseases (I60-I69)	1,699	5.0	81.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,694	4.9	81.6
5	Diabetes mellitus (E10-E14)	1,581	4.6	76.1
6	Chronic lower respiratory diseases (J40-J47)	923	2.7	44.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	875	2.6	42.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	835	2.4	40.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	688	2.0	33.1
10	Septicemia (A40-A41)	678	2.0	32.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	581	1.7	28.0
12	Influenza and pneumonia (J09-J18)	515	1.5	24.8
13	Viral hepatitis (B15-B19)	433	1.3	20.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	280	0.8	13.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	183	0.5	8.8
...	All other causes (residual)	4,666	13.6	224.7
Non-Hispanic, black, male, 65-74 years				
...	All causes	32,962	100.0	3,110.5
1	Malignant neoplasms (C00-C97)	10,028	30.4	946.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,568	26.0	808.5
3	Cerebrovascular diseases (I60-I69)	1,842	5.6	173.8
4	Diabetes mellitus (E10-E14)	1,637	5.0	154.5
5	Chronic lower respiratory diseases (J40-J47)	1,440	4.4	135.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	993	3.0	93.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	770	2.3	72.7
8	Septicemia (A40-A41)	750	2.3	70.8
9	Influenza and pneumonia (J09-J18)	583	1.8	55.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	540	1.6	51.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	403	1.2	38.0
12	Pneumonitis due to solids and liquids (J69)	229	0.7	21.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	211	0.6	19.9
14	Alzheimer's disease (G30)	191	0.6	18.0
15	Viral hepatitis (B15-B19)	164	0.5	15.5
...	All other causes (residual)	4,613	14.0	435.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-84 years				
...	All causes	27,672	100.0	6,287.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,384	26.7	1,677.9
2	Malignant neoplasms (C00-C97)	7,051	25.5	1,602.2
3	Cerebrovascular diseases (I60-I69)	1,768	6.4	401.7
4	Diabetes mellitus (E10-E14)	1,271	4.6	288.8
5	Chronic lower respiratory diseases (J40-J47)	1,255	4.5	285.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	903	3.3	205.2
7	Alzheimer's disease (G30)	717	2.6	162.9
8	Septicemia (A40-A41)	634	2.3	144.1
9	Influenza and pneumonia (J09-J18)	563	2.0	127.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	503	1.8	114.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	478	1.7	108.6
12	Parkinson's disease (G20-G21)	289	1.0	65.7
13	Pneumonitis due to solids and liquids (J69)	252	0.9	57.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	133	0.5	30.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	95	0.3	21.6
...	All other causes (residual)	4,376	15.8	994.4
Non-Hispanic, black, male, 85 years and over				
...	All causes	17,810	100.0	13,532.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,947	27.8	3,759.0
2	Malignant neoplasms (C00-C97)	3,185	17.9	2,420.1
3	Cerebrovascular diseases (I60-I69)	1,115	6.3	847.2
4	Alzheimer's disease (G30)	866	4.9	658.0
5	Chronic lower respiratory diseases (J40-J47)	769	4.3	584.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	628	3.5	477.2
7	Diabetes mellitus (E10-E14)	599	3.4	455.1
8	Influenza and pneumonia (J09-J18)	514	2.9	390.6
9	Septicemia (A40-A41)	391	2.2	297.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	366	2.1	278.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	297	1.7	225.7
12	Parkinson's disease (G20-G21)	209	1.2	158.8
13	Pneumonitis due to solids and liquids (J69)	205	1.2	155.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	88	0.5	66.9
15	Nutritional deficiencies (E40-E64)	70	0.4	53.2
...	All other causes (residual)	3,561	20.0	2,705.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	149,008	100.0	691.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,629	23.2	160.7
2	Malignant neoplasms (C00-C97)	33,565	22.5	155.8
3	Cerebrovascular diseases (I60-I69)	9,233	6.2	42.8
4	Diabetes mellitus (E10-E14)	6,895	4.6	32.0
5	Chronic lower respiratory diseases (J40-J47)	4,822	3.2	22.4
6	Alzheimer's disease (G30)	4,675	3.1	21.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,513	3.0	20.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,504	3.0	20.9
9	Septicemia (A40-A41)	3,383	2.3	15.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,882	1.9	13.4
11	Influenza and pneumonia (J09-J18)	2,806	1.9	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,676	1.1	7.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,206	0.8	5.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,138	0.8	5.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,049	0.7	4.9
...	All other causes (residual)	32,032	21.5	148.6
Non-Hispanic, black, female, 1-4 years				
...	All causes	407	100.0	34.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	26.0	8.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	16.0	5.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	10.1	3.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	6.9	2.3
5	Malignant neoplasms (C00-C97)	25	6.1	2.1
6	Chronic lower respiratory diseases (J40-J47)	11	2.7	*
7	Influenza and pneumonia (J09-J18)	10	2.5	*
8	Septicemia (A40-A41)	5	1.2	*
9	Cerebrovascular diseases (I60-I69)	4	1.0	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	4	1.0	*
11	Acute bronchitis and bronchiolitis (J20-J21)	3	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
13	Salmonella infections (A01-A02)	1	0.2	*
13	Meningococcal infection (A39)	1	0.2	*
13	Malaria (B50-B54)	1	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Nutritional deficiencies (E40-E64)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (residual)	94	23.1	7.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	461	100.0	15.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	108	23.4	3.5
2	Malignant neoplasms (C00-C97)	67	14.5	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	7.4	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	6.5	1.0
5	Chronic lower respiratory diseases (J40-J47)	24	5.2	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.6	0.7
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	3.3	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	3.3	*
9	Anemias (D50-D64)	9	2.0	*
10	Cerebrovascular diseases (I60-I69)	8	1.7	*
11	Septicemia (A40-A41)	7	1.5	*
11	Influenza and pneumonia (J09-J18)	7	1.5	*
13	Diabetes mellitus (E10-E14)	5	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
...	All other causes (residual)	105	22.8	3.4
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,490	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	374	25.1	11.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	244	16.4	7.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	110	7.4	3.3
4	Malignant neoplasms (C00-C97)	108	7.2	3.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	6.1	2.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	56	3.8	1.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	34	2.3	1.0
8	Anemias (D50-D64)	32	2.1	1.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	1.9	0.8
10	Diabetes mellitus (E10-E14)	26	1.7	0.8
11	Chronic lower respiratory diseases (J40-J47)	24	1.6	0.7
12	Septicemia (A40-A41)	20	1.3	0.6
13	Cerebrovascular diseases (I60-I69)	19	1.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.8	*
15	Influenza and pneumonia (J09-J18)	10	0.7	*
...	All other causes (residual)	302	20.3	9.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-34 years				
...	All causes	2,875	100.0	94.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	484	16.8	15.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	376	13.1	12.3
3	Malignant neoplasms (C00-C97)	367	12.8	12.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	249	8.7	8.1
5	Pregnancy, childbirth and the puerperium (O00-O99)	136	4.7	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	122	4.2	4.0
7	Diabetes mellitus (E10-E14)	105	3.7	3.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	3.0	2.8
9	Cerebrovascular diseases (I60-I69)	62	2.2	2.0
10	Influenza and pneumonia (J09-J18)	51	1.8	1.7
11	Chronic lower respiratory diseases (J40-J47)	50	1.7	1.6
12	Anemias (D50-D64)	47	1.6	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.4	1.3
14	Septicemia (A40-A41)	38	1.3	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.0	1.0
...	All other causes (residual)	630	21.9	20.6
Non-Hispanic, black, female, 35-44 years				
...	All causes	5,707	100.0	205.2
1	Malignant neoplasms (C00-C97)	1,188	20.8	42.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,047	18.3	37.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	555	9.7	20.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	273	4.8	9.8
5	Diabetes mellitus (E10-E14)	246	4.3	8.8
6	Cerebrovascular diseases (I60-I69)	207	3.6	7.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	2.9	5.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	2.0	4.1
9	Pregnancy, childbirth and the puerperium (O00-O99)	103	1.8	3.7
10	Septicemia (A40-A41)	96	1.7	3.5
11	Influenza and pneumonia (J09-J18)	95	1.7	3.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	91	1.6	3.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	1.5	3.1
14	Chronic lower respiratory diseases (J40-J47)	86	1.5	3.1
15	Anemias (D50-D64)	68	1.2	2.4
...	All other causes (residual)	1,286	22.5	46.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-54 years				
...	All causes	13,629	100.0	473.9
1	Malignant neoplasms (C00-C97)	3,974	29.2	138.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,844	20.9	98.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	784	5.8	27.3
4	Cerebrovascular diseases (I60-I69)	672	4.9	23.4
5	Diabetes mellitus (E10-E14)	611	4.5	21.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	376	2.8	13.1
7	Chronic lower respiratory diseases (J40-J47)	369	2.7	12.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	292	2.1	10.2
9	Septicemia (A40-A41)	287	2.1	10.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	286	2.1	9.9
11	Influenza and pneumonia (J09-J18)	234	1.7	8.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	189	1.4	6.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	134	1.0	4.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	0.6	3.0
15	Anemias (D50-D64)	76	0.6	2.6
...	All other causes (residual)	2,414	17.7	83.9
Non-Hispanic, black, female, 55-64 years				
...	All causes	24,811	100.0	1,000.8
1	Malignant neoplasms (C00-C97)	8,200	33.0	330.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,372	21.7	216.7
3	Diabetes mellitus (E10-E14)	1,284	5.2	51.8
4	Cerebrovascular diseases (I60-I69)	1,235	5.0	49.8
5	Chronic lower respiratory diseases (J40-J47)	882	3.6	35.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	727	2.9	29.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	685	2.8	27.6
8	Septicemia (A40-A41)	633	2.6	25.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	442	1.8	17.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	425	1.7	17.1
11	Influenza and pneumonia (J09-J18)	397	1.6	16.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	288	1.2	11.6
13	Viral hepatitis (B15-B19)	203	0.8	8.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	96	0.4	3.9
15	Pneumonitis due to solids and liquids (J69)	81	0.3	3.3
...	All other causes (residual)	3,861	15.6	155.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-74 years				
...	All causes	27,106	100.0	1,922.9
1	Malignant neoplasms (C00-C97)	8,409	31.0	596.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,313	23.3	447.9
3	Cerebrovascular diseases (I60-I69)	1,634	6.0	115.9
4	Diabetes mellitus (E10-E14)	1,541	5.7	109.3
5	Chronic lower respiratory diseases (J40-J47)	1,111	4.1	78.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,072	4.0	76.0
7	Septicemia (A40-A41)	721	2.7	51.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	464	1.7	32.9
9	Influenza and pneumonia (J09-J18)	403	1.5	28.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	390	1.4	27.7
11	Alzheimer's disease (G30)	295	1.1	20.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	174	0.6	12.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	158	0.6	11.2
14	Pneumonitis due to solids and liquids (J69)	130	0.5	9.2
15	Aortic aneurysm and dissection (I71)	89	0.3	6.3
...	All other causes (residual)	4,202	15.5	298.1
Non-Hispanic, black, female, 75-84 years				
...	All causes	31,811	100.0	4,441.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,911	24.9	1,104.5
2	Malignant neoplasms (C00-C97)	7,021	22.1	980.3
3	Cerebrovascular diseases (I60-I69)	2,356	7.4	328.9
4	Diabetes mellitus (E10-E14)	1,698	5.3	237.1
5	Alzheimer's disease (G30)	1,341	4.2	187.2
6	Chronic lower respiratory diseases (J40-J47)	1,253	3.9	174.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,161	3.6	162.1
8	Septicemia (A40-A41)	737	2.3	102.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	701	2.2	97.9
10	Influenza and pneumonia (J09-J18)	611	1.9	85.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	1.2	51.8
12	Pneumonitis due to solids and liquids (J69)	229	0.7	32.0
13	Parkinson's disease (G20-G21)	194	0.6	27.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	175	0.6	24.4
15	Aortic aneurysm and dissection (I71)	109	0.3	15.2
...	All other causes (residual)	5,943	18.7	829.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2014

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85 years and over				
...	All causes	37,689	100.0	11,834.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,551	28.0	3,313.2
2	Malignant neoplasms (C00-C97)	4,198	11.1	1,318.2
3	Cerebrovascular diseases (I60-I69)	3,026	8.0	950.2
4	Alzheimer's disease (G30)	2,978	7.9	935.1
5	Diabetes mellitus (E10-E14)	1,379	3.7	433.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,124	3.0	353.0
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,018	2.7	319.7
8	Chronic lower respiratory diseases (J40-J47)	1,010	2.7	317.2
9	Influenza and pneumonia (J09-J18)	944	2.5	296.4
10	Septicemia (A40-A41)	814	2.2	255.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	426	1.1	133.8
12	Pneumonitis due to solids and liquids (J69)	309	0.8	97.0
13	Parkinson's disease (G20-G21)	200	0.5	62.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	192	0.5	60.3
15	Atherosclerosis (I70)	135	0.4	42.4
...	All other causes (residual)	9,385	24.9	2,947.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014