

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	144,490	100.0	286.2
1	Malignant neoplasms (C00-C97)	31,119	21.5	61.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,006	20.8	59.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,476	7.3	20.8
4	Cerebrovascular diseases (I60-I69)	7,274	5.0	14.4
5	Diabetes mellitus (E10-E14)	6,556	4.5	13.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,348	3.0	8.6
7	Chronic lower respiratory diseases (J40-J47)	4,172	2.9	8.3
8	Alzheimer's disease (G30)	3,427	2.4	6.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,252	2.3	6.4
10	Influenza and pneumonia (J09-J18)	3,025	2.1	6.0
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,890	2.0	5.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,661	1.8	5.3
13	Certain conditions originating in the perinatal period (P00-P96)	2,529	1.8	5.0
14	Septicemia (A40-A41)	2,035	1.4	4.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,786	1.2	3.5
...	All other causes (Residual)	28,934	20.0	57.3
Hispanic, both sexes, 1-4 years				
...	All causes	930	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	35.5	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	13.1	3.0
3	Malignant neoplasms (C00-C97)	75	8.1	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	6.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	4.0	0.9
6	Influenza and pneumonia (J09-J18)	21	2.3	0.5
7	Septicemia (A40-A41)	18	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
9	Cerebrovascular diseases (I60-I69)	8	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.8	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.8	*
12	Meningococcal infection (A39)	5	0.5	*
12	Pneumonitis due to solids and liquids (J69)	5	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	216	23.2	5.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	951	100.0	10.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	252	26.5	2.7
2	Malignant neoplasms (C00-C97)	212	22.3	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	5.7	0.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	5.3	0.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	4.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	2.7	0.3
7	Cerebrovascular diseases (I60-I69)	23	2.4	0.2
8	Chronic lower respiratory diseases (J40-J47)	20	2.1	0.2
9	Septicemia (A40-A41)	14	1.5	*
10	Influenza and pneumonia (J09-J18)	13	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
13	Anemias (D50-D64)	3	0.3	*
13	Meningitis (G00,G03)	3	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Pneumonitis due to solids and liquids (J69)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	215	22.6	2.3
Hispanic, both sexes, 15-24 years				
...	All causes	4,795	100.0	54.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,891	39.4	21.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,022	21.3	11.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	628	13.1	7.1
4	Malignant neoplasms (C00-C97)	344	7.2	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	2.6	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	1.2	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	32	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	29	0.6	0.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.5	0.3
10	Influenza and pneumonia (J09-J18)	24	0.5	0.3
11	Septicemia (A40-A41)	21	0.4	0.2
12	Diabetes mellitus (E10-E14)	18	0.4	*
13	Legal intervention (Y35,Y89.0)	16	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	15	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
...	All other causes (Residual)	531	11.1	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	14,164	100.0	90.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,688	26.0	23.4
2	Malignant neoplasms (C00-C97)	2,106	14.9	13.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,291	9.1	8.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,261	8.9	8.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,093	7.7	6.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	570	4.0	3.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	428	3.0	2.7
8	Cerebrovascular diseases (I60-I69)	395	2.8	2.5
9	Diabetes mellitus (E10-E14)	305	2.2	1.9
10	Influenza and pneumonia (J09-J18)	156	1.1	1.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	118	0.8	0.7
12	Septicemia (A40-A41)	116	0.8	0.7
13	Pregnancy, childbirth and the puerperium (O00-O99)	108	0.8	0.7
14	Viral hepatitis (B15-B19)	92	0.6	0.6
15	Chronic lower respiratory diseases (J40-J47)	84	0.6	0.5
...	All other causes (Residual)	2,353	16.6	15.0
Hispanic, both sexes, 45-64 years				
...	All causes	34,981	100.0	403.1
1	Malignant neoplasms (C00-C97)	10,170	29.1	117.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,545	18.7	75.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,464	7.0	28.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,433	7.0	28.0
5	Diabetes mellitus (E10-E14)	1,834	5.2	21.1
6	Cerebrovascular diseases (I60-I69)	1,502	4.3	17.3
7	Viral hepatitis (B15-B19)	793	2.3	9.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	708	2.0	8.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	675	1.9	7.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	599	1.7	6.9
11	Chronic lower respiratory diseases (J40-J47)	566	1.6	6.5
12	Septicemia (A40-A41)	518	1.5	6.0
13	Influenza and pneumonia (J09-J18)	445	1.3	5.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	367	1.0	4.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	321	0.9	3.7
...	All other causes (Residual)	5,041	14.4	58.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	83,492	100.0	3,001.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,918	26.3	788.0
2	Malignant neoplasms (C00-C97)	18,196	21.8	654.2
3	Cerebrovascular diseases (I60-I69)	5,292	6.3	190.2
4	Diabetes mellitus (E10-E14)	4,396	5.3	158.0
5	Chronic lower respiratory diseases (J40-J47)	3,467	4.2	124.6
6	Alzheimer's disease (G30)	3,370	4.0	121.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,378	2.8	85.5
8	Influenza and pneumonia (J09-J18)	2,332	2.8	83.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,697	2.0	61.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,337	1.6	48.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,320	1.6	47.5
12	Septicemia (A40-A41)	1,304	1.6	46.9
13	Parkinson's disease (G20-G21)	949	1.1	34.1
14	Pneumonitis due to solids and liquids (J69)	567	0.7	20.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	511	0.6	18.4
...	All other causes (Residual)	14,458	17.3	519.8
Hispanic, male, all ages				
...	All causes	79,622	100.0	310.8
1	Malignant neoplasms (C00-C97)	16,450	20.7	64.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,421	20.6	64.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,594	9.5	29.6
4	Cerebrovascular diseases (I60-I69)	3,382	4.2	13.2
5	Diabetes mellitus (E10-E14)	3,372	4.2	13.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,067	3.9	12.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,435	3.1	9.5
8	Chronic lower respiratory diseases (J40-J47)	2,174	2.7	8.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,168	2.7	8.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,670	2.1	6.5
11	Influenza and pneumonia (J09-J18)	1,565	2.0	6.1
12	Certain conditions originating in the perinatal period (P00-P96)	1,456	1.8	5.7
13	Alzheimer's disease (G30)	1,171	1.5	4.6
14	Septicemia (A40-A41)	1,012	1.3	4.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	910	1.1	3.6
...	All other causes (Residual)	14,775	18.6	57.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	524	100.0	25.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	39.9	10.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	11.3	2.8
3	Malignant neoplasms (C00-C97)	42	8.0	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	8.0	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.1	*
6	Influenza and pneumonia (J09-J18)	13	2.5	*
7	Septicemia (A40-A41)	12	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
10	Pneumonitis due to solids and liquids (J69)	3	0.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
12	Meningitis (G00,G03)	2	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
15	Meningococcal infection (A39)	1	0.2	*
15	Cerebrovascular diseases (I60-I69)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	104	19.8	5.0
Hispanic, male, 5-14 years				
...	All causes	543	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	27.6	3.2
2	Malignant neoplasms (C00-C97)	112	20.6	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	6.4	0.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	5.5	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	4.1	0.5
6	Chronic lower respiratory diseases (J40-J47)	18	3.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.6	*
8	Cerebrovascular diseases (I60-I69)	13	2.4	*
9	Septicemia (A40-A41)	9	1.7	*
9	Influenza and pneumonia (J09-J18)	9	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
13	Anemias (D50-D64)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Meningitis (G00,G03)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (Residual)	114	21.0	2.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,689	100.0	79.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,478	40.1	31.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	914	24.8	19.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	497	13.5	10.7
4	Malignant neoplasms (C00-C97)	225	6.1	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	2.5	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	0.9	0.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	20	0.5	0.4
8	Cerebrovascular diseases (I60-I69)	19	0.5	*
9	Influenza and pneumonia (J09-J18)	17	0.5	*
10	Legal intervention (Y35,Y89.0)	16	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.3	*
12	Septicemia (A40-A41)	9	0.2	*
12	Diabetes mellitus (E10-E14)	9	0.2	*
12	Aortic aneurysm and dissection (I71)	9	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	8	0.2	*
...	All other causes (Residual)	330	8.9	7.1
Hispanic, male, 25-44 years				
...	All causes	9,921	100.0	121.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,936	29.6	36.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,074	10.8	13.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	992	10.0	12.2
4	Malignant neoplasms (C00-C97)	934	9.4	11.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	911	9.2	11.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	459	4.6	5.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	337	3.4	4.1
8	Cerebrovascular diseases (I60-I69)	250	2.5	3.1
9	Diabetes mellitus (E10-E14)	186	1.9	2.3
10	Influenza and pneumonia (J09-J18)	103	1.0	1.3
11	Viral hepatitis (B15-B19)	70	0.7	0.9
12	Septicemia (A40-A41)	67	0.7	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	0.7	0.8
14	Chronic lower respiratory diseases (J40-J47)	51	0.5	0.6
15	Aortic aneurysm and dissection (I71)	49	0.5	0.6
...	All other causes (Residual)	1,436	14.5	17.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	22,136	100.0	518.3
1	Malignant neoplasms (C00-C97)	5,366	24.2	125.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,662	21.1	109.2
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,853	8.4	43.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,797	8.1	42.1
5	Diabetes mellitus (E10-E14)	1,137	5.1	26.6
6	Cerebrovascular diseases (I60-I69)	881	4.0	20.6
7	Viral hepatitis (B15-B19)	574	2.6	13.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	552	2.5	12.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	489	2.2	11.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	417	1.9	9.8
11	Chronic lower respiratory diseases (J40-J47)	309	1.4	7.2
12	Septicemia (A40-A41)	301	1.4	7.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	295	1.3	6.9
14	Influenza and pneumonia (J09-J18)	251	1.1	5.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	204	0.9	4.8
...	All other causes (Residual)	3,048	13.8	71.4
Hispanic, male, 65 years and over				
...	All causes	39,934	100.0	3,378.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,603	26.6	897.1
2	Malignant neoplasms (C00-C97)	9,762	24.4	826.0
3	Cerebrovascular diseases (I60-I69)	2,203	5.5	186.4
4	Diabetes mellitus (E10-E14)	2,040	5.1	172.6
5	Chronic lower respiratory diseases (J40-J47)	1,776	4.4	150.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,162	2.9	98.3
7	Influenza and pneumonia (J09-J18)	1,153	2.9	97.6
8	Alzheimer's disease (G30)	1,148	2.9	97.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	935	2.3	79.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	752	1.9	63.6
11	Septicemia (A40-A41)	590	1.5	49.9
12	Parkinson's disease (G20-G21)	532	1.3	45.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	519	1.3	43.9
14	Pneumonitis due to solids and liquids (J69)	318	0.8	26.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	260	0.7	22.0
...	All other causes (Residual)	6,181	15.5	523.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	64,868	100.0	260.9
1	Malignant neoplasms (C00-C97)	14,669	22.6	59.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,585	20.9	54.6
3	Cerebrovascular diseases (I60-I69)	3,892	6.0	15.7
4	Diabetes mellitus (E10-E14)	3,184	4.9	12.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,882	4.4	11.6
6	Alzheimer's disease (G30)	2,256	3.5	9.1
7	Chronic lower respiratory diseases (J40-J47)	1,998	3.1	8.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,582	2.4	6.4
9	Influenza and pneumonia (J09-J18)	1,460	2.3	5.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,281	2.0	5.2
11	Certain conditions originating in the perinatal period (P00-P96)	1,073	1.7	4.3
12	Septicemia (A40-A41)	1,023	1.6	4.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	948	1.5	3.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	883	1.4	3.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	493	0.8	2.0
...	All other causes (Residual)	13,659	21.1	54.9
Hispanic, female, 1-4 years				
...	All causes	406	100.0	20.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	29.8	6.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	15.5	3.1
3	Malignant neoplasms (C00-C97)	33	8.1	1.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	5.2	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	3.7	*
6	Influenza and pneumonia (J09-J18)	8	2.0	*
7	Cerebrovascular diseases (I60-I69)	7	1.7	*
8	Septicemia (A40-A41)	6	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.2	*
11	Meningococcal infection (A39)	4	1.0	*
12	Anemias (D50-D64)	3	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.5	*
14	Pneumonitis due to solids and liquids (J69)	2	0.5	*
...	All other causes (Residual)	108	26.6	5.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	408	100.0	8.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	25.0	2.2
2	Malignant neoplasms (C00-C97)	100	24.5	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	5.4	0.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	4.9	0.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	4.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.9	*
7	Cerebrovascular diseases (I60-I69)	10	2.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.5	*
9	Septicemia (A40-A41)	5	1.2	*
10	Influenza and pneumonia (J09-J18)	4	1.0	*
11	Diabetes mellitus (E10-E14)	2	0.5	*
11	Meningitis (G00,G03)	2	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
11	Pneumonitis due to solids and liquids (J69)	2	0.5	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Meningococcal infection (A39)	1	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (Residual)	93	22.8	2.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-24 years				
...	All causes	1,106	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	413	37.3	9.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	131	11.8	3.1
3	Malignant neoplasms (C00-C97)	119	10.8	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	108	9.8	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	2.9	0.8
5	Pregnancy, childbirth and the puerperium (O00-O99)	32	2.9	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	2.3	0.6
8	Septicemia (A40-A41)	12	1.1	*
9	Cerebrovascular diseases (I60-I69)	10	0.9	*
10	Diabetes mellitus (E10-E14)	9	0.8	*
11	Influenza and pneumonia (J09-J18)	7	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.4	*
...	All other causes (Residual)	181	16.4	4.3
Hispanic, female, 25-44 years				
...	All causes	4,243	100.0	56.0
1	Malignant neoplasms (C00-C97)	1,172	27.6	15.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	752	17.7	9.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	299	7.0	3.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	4.4	2.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	182	4.3	2.4
6	Cerebrovascular diseases (I60-I69)	145	3.4	1.9
7	Diabetes mellitus (E10-E14)	119	2.8	1.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	111	2.6	1.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	108	2.5	1.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	91	2.1	1.2
11	Influenza and pneumonia (J09-J18)	53	1.2	0.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	1.2	0.7
13	Septicemia (A40-A41)	49	1.2	0.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	0.9	0.5
15	Chronic lower respiratory diseases (J40-J47)	33	0.8	0.4
...	All other causes (Residual)	853	20.1	11.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-64 years				
...	All causes	12,845	100.0	291.5
1	Malignant neoplasms (C00-C97)	4,804	37.4	109.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,883	14.7	42.7
3	Diabetes mellitus (E10-E14)	697	5.4	15.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	667	5.2	15.1
5	Cerebrovascular diseases (I60-I69)	621	4.8	14.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	580	4.5	13.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	291	2.3	6.6
8	Chronic lower respiratory diseases (J40-J47)	257	2.0	5.8
9	Viral hepatitis (B15-B19)	219	1.7	5.0
10	Septicemia (A40-A41)	217	1.7	4.9
11	Influenza and pneumonia (J09-J18)	194	1.5	4.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	1.0	2.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	117	0.9	2.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	110	0.9	2.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.6	1.8
...	All other causes (Residual)	1,984	15.4	45.0
Hispanic, female, 65 years and over				
...	All causes	43,558	100.0	2,722.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,315	26.0	707.3
2	Malignant neoplasms (C00-C97)	8,434	19.4	527.2
3	Cerebrovascular diseases (I60-I69)	3,089	7.1	193.1
4	Diabetes mellitus (E10-E14)	2,356	5.4	147.3
5	Alzheimer's disease (G30)	2,222	5.1	138.9
6	Chronic lower respiratory diseases (J40-J47)	1,691	3.9	105.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,216	2.8	76.0
8	Influenza and pneumonia (J09-J18)	1,179	2.7	73.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	801	1.8	50.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	762	1.7	47.6
11	Septicemia (A40-A41)	714	1.6	44.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	585	1.3	36.6
13	Parkinson's disease (G20-G21)	417	1.0	26.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	251	0.6	15.7
15	Pneumonitis due to solids and liquids (J69)	249	0.6	15.6
...	All other causes (Residual)	8,277	19.0	517.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, all ages				
...	All causes	1,969,916	100.0	984.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	483,973	24.6	241.8
2	Malignant neoplasms (C00-C97)	460,567	23.4	230.1
3	Chronic lower respiratory diseases (J40-J47)	122,887	6.2	61.4
4	Cerebrovascular diseases (I60-I69)	101,849	5.2	50.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94,420	4.8	47.2
6	Alzheimer's disease (G30)	73,502	3.7	36.7
7	Diabetes mellitus (E10-E14)	47,746	2.4	23.9
8	Influenza and pneumonia (J09-J18)	40,244	2.0	20.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,963	1.9	18.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32,010	1.6	16.0
11	Septicemia (A40-A41)	25,951	1.3	13.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	23,655	1.2	11.8
13	Parkinson's disease (G20-G21)	19,818	1.0	9.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,878	1.0	9.4
15	Pneumonitis due to solids and liquids (J69)	14,539	0.7	7.3
...	All other causes (Residual)	372,914	18.9	186.3
Non-Hispanic white, both sexes, 1-4 years				
...	All causes	2,139	100.0	24.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	744	34.8	8.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	229	10.7	2.6
3	Malignant neoplasms (C00-C97)	196	9.2	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	7.4	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	2.8	0.7
6	Influenza and pneumonia (J09-J18)	38	1.8	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.7	0.4
8	Septicemia (A40-A41)	31	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	27	1.3	0.3
10	Cerebrovascular diseases (I60-I69)	23	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	17	0.8	*
12	Meningitis (G00,G03)	13	0.6	*
13	Anemias (D50-D64)	11	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
...	All other causes (Residual)	539	25.2	6.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 5-14 years				
...	All causes	2,910	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,000	34.4	4.3
2	Malignant neoplasms (C00-C97)	500	17.2	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	166	5.7	0.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	164	5.6	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	3.2	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	3.1	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	1.6	0.2
8	Influenza and pneumonia (J09-J18)	42	1.4	0.2
9	Cerebrovascular diseases (I60-I69)	39	1.3	0.2
10	Chronic lower respiratory diseases (J40-J47)	34	1.2	0.1
11	Septicemia (A40-A41)	31	1.1	0.1
12	Diabetes mellitus (E10-E14)	17	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	16	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.4	*
...	All other causes (Residual)	646	22.2	2.8
Non-Hispanic white, both sexes, 15-24 years				
...	All causes	16,847	100.0	66.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,507	50.5	33.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,190	18.9	12.6
3	Malignant neoplasms (C00-C97)	901	5.3	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	748	4.4	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	520	3.1	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	245	1.5	1.0
7	Influenza and pneumonia (J09-J18)	110	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	100	0.6	0.4
9	Diabetes mellitus (E10-E14)	95	0.6	0.4
10	Septicemia (A40-A41)	76	0.5	0.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	73	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	68	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.2	0.1
15	Pneumonitis due to solids and liquids (J69)	26	0.2	0.1
...	All other causes (Residual)	2,102	12.5	8.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 25-44 years				
...	All causes	70,485	100.0	141.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,405	30.4	43.1
2	Malignant neoplasms (C00-C97)	9,791	13.9	19.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,694	13.8	19.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,370	11.9	16.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,856	2.6	3.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,759	2.5	3.5
7	Diabetes mellitus (E10-E14)	1,291	1.8	2.6
8	Cerebrovascular diseases (I60-I69)	1,175	1.7	2.4
9	Influenza and pneumonia (J09-J18)	718	1.0	1.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	629	0.9	1.3
11	Septicemia (A40-A41)	623	0.9	1.3
12	Chronic lower respiratory diseases (J40-J47)	589	0.8	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	588	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	439	0.6	0.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	270	0.4	0.5
...	All other causes (Residual)	11,288	16.0	22.7
Non-Hispanic white, both sexes, 45-64 years				
...	All causes	357,299	100.0	606.3
1	Malignant neoplasms (C00-C97)	119,040	33.3	202.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,358	20.8	126.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26,090	7.3	44.3
4	Chronic lower respiratory diseases (J40-J47)	15,697	4.4	26.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13,489	3.8	22.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	13,474	3.8	22.9
7	Diabetes mellitus (E10-E14)	10,847	3.0	18.4
8	Cerebrovascular diseases (I60-I69)	9,893	2.8	16.8
9	Septicemia (A40-A41)	4,524	1.3	7.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,008	1.1	6.8
11	Influenza and pneumonia (J09-J18)	3,912	1.1	6.6
12	Viral hepatitis (B15-B19)	3,765	1.1	6.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,197	0.6	3.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,648	0.5	2.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,605	0.4	2.7
...	All other causes (Residual)	52,752	14.8	89.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, both sexes, 65 years and over				
...	All causes	1,509,149	100.0	4,660.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	400,404	26.5	1,236.4
2	Malignant neoplasms (C00-C97)	330,101	21.9	1,019.3
3	Chronic lower respiratory diseases (J40-J47)	106,466	7.1	328.8
4	Cerebrovascular diseases (I60-I69)	90,565	6.0	279.7
5	Alzheimer's disease (G30)	72,757	4.8	224.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36,101	2.4	111.5
7	Diabetes mellitus (E10-E14)	35,494	2.4	109.6
8	Influenza and pneumonia (J09-J18)	35,353	2.3	109.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32,434	2.1	100.2
10	Septicemia (A40-A41)	20,592	1.4	63.6
11	Parkinson's disease (G20-G21)	19,332	1.3	59.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16,481	1.1	50.9
13	Pneumonitis due to solids and liquids (J69)	13,213	0.9	40.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10,410	0.7	32.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	8,301	0.6	25.6
...	All other causes (Residual)	281,145	18.6	868.1
Non-Hispanic white, male, all ages				
...	All causes	971,604	100.0	987.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,765	25.5	251.8
2	Malignant neoplasms (C00-C97)	241,816	24.9	245.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57,747	5.9	58.7
4	Chronic lower respiratory diseases (J40-J47)	57,390	5.9	58.3
5	Cerebrovascular diseases (I60-I69)	40,040	4.1	40.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,238	2.6	25.7
7	Diabetes mellitus (E10-E14)	25,134	2.6	25.5
8	Alzheimer's disease (G30)	22,278	2.3	22.6
9	Influenza and pneumonia (J09-J18)	18,657	1.9	19.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,503	1.9	18.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,270	1.6	15.5
12	Septicemia (A40-A41)	11,963	1.2	12.2
13	Parkinson's disease (G20-G21)	11,627	1.2	11.8
14	Pneumonitis due to solids and liquids (J69)	7,862	0.8	8.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,476	0.8	7.6
...	All other causes (Residual)	162,838	16.8	165.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 1-4 years				
...	All causes	1,219	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	456	37.4	10.3
2	Malignant neoplasms (C00-C97)	109	8.9	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	8.3	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	7.4	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.0	0.8
6	Influenza and pneumonia (J09-J18)	20	1.6	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.4	*
7	Cerebrovascular diseases (I60-I69)	17	1.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	17	1.4	*
10	Septicemia (A40-A41)	13	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	10	0.8	*
12	Meningitis (G00,G03)	9	0.7	*
13	Anemias (D50-D64)	8	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
...	All other causes (Residual)	304	24.9	6.8
Non-Hispanic white, male, 5-14 years				
...	All causes	1,691	100.0	14.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	639	37.8	5.4
2	Malignant neoplasms (C00-C97)	277	16.4	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	128	7.6	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	4.4	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	3.3	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	2.4	0.3
7	Cerebrovascular diseases (I60-I69)	23	1.4	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.1	*
9	Septicemia (A40-A41)	18	1.1	*
10	Influenza and pneumonia (J09-J18)	15	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	12	0.7	*
13	Diabetes mellitus (E10-E14)	11	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	356	21.1	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 15-24 years				
...	All causes	12,079	100.0	93.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,172	51.1	47.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,638	21.8	20.4
3	Malignant neoplasms (C00-C97)	565	4.7	4.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	528	4.4	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	382	3.2	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	150	1.2	1.2
7	Cerebrovascular diseases (I60-I69)	61	0.5	0.5
8	Influenza and pneumonia (J09-J18)	59	0.5	0.5
9	Diabetes mellitus (E10-E14)	54	0.4	0.4
10	Chronic lower respiratory diseases (J40-J47)	42	0.3	0.3
11	Septicemia (A40-A41)	38	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.3	0.2
13	Legal intervention (Y35,Y89.0)	24	0.2	0.2
14	Aortic aneurysm and dissection (I71)	18	0.1	*
15	Pneumonitis due to solids and liquids (J69)	17	0.1	*
...	All other causes (Residual)	1,299	10.8	10.0
Non-Hispanic white, male, 25-44 years				
...	All causes	45,553	100.0	182.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,035	33.0	60.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,580	16.6	30.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,937	13.0	23.8
4	Malignant neoplasms (C00-C97)	4,568	10.0	18.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,184	2.6	4.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,175	2.6	4.7
7	Diabetes mellitus (E10-E14)	759	1.7	3.0
8	Cerebrovascular diseases (I60-I69)	590	1.3	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	481	1.1	1.9
10	Influenza and pneumonia (J09-J18)	383	0.8	1.5
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	337	0.7	1.3
12	Septicemia (A40-A41)	318	0.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	277	0.6	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	231	0.5	0.9
15	Aortic aneurysm and dissection (I71)	175	0.4	0.7
...	All other causes (Residual)	6,523	14.3	26.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, male, 45-64 years				
...	All causes	219,847	100.0	757.0
1	Malignant neoplasms (C00-C97)	64,525	29.3	222.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,484	24.3	184.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,456	7.9	60.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,279	4.7	35.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	9,324	4.2	32.1
6	Chronic lower respiratory diseases (J40-J47)	7,900	3.6	27.2
7	Diabetes mellitus (E10-E14)	6,900	3.1	23.8
8	Cerebrovascular diseases (I60-I69)	5,625	2.6	19.4
9	Viral hepatitis (B15-B19)	2,678	1.2	9.2
10	Septicemia (A40-A41)	2,415	1.1	8.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,327	1.1	8.0
12	Influenza and pneumonia (J09-J18)	2,241	1.0	7.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,442	0.7	5.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,373	0.6	4.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,040	0.5	3.6
...	All other causes (Residual)	30,838	14.0	106.2
Non-Hispanic white, male, 65 years and over				
...	All causes	685,031	100.0	4,855.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	187,785	27.4	1,330.9
2	Malignant neoplasms (C00-C97)	171,751	25.1	1,217.3
3	Chronic lower respiratory diseases (J40-J47)	49,139	7.2	348.3
4	Cerebrovascular diseases (I60-I69)	33,697	4.9	238.8
5	Alzheimer's disease (G30)	21,963	3.2	155.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,660	2.6	125.2
7	Diabetes mellitus (E10-E14)	17,409	2.5	123.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,906	2.3	112.7
9	Influenza and pneumonia (J09-J18)	15,899	2.3	112.7
10	Parkinson's disease (G20-G21)	11,305	1.7	80.1
11	Septicemia (A40-A41)	9,114	1.3	64.6
12	Pneumonitis due to solids and liquids (J69)	7,029	1.0	49.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,900	0.9	41.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,346	0.8	37.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,749	0.7	33.7
...	All other causes (Residual)	110,379	16.1	782.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, all ages				
...	All causes	998,312	100.0	981.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	236,208	23.7	232.2
2	Malignant neoplasms (C00-C97)	218,751	21.9	215.0
3	Chronic lower respiratory diseases (J40-J47)	65,497	6.6	64.4
4	Cerebrovascular diseases (I60-I69)	61,809	6.2	60.8
5	Alzheimer's disease (G30)	51,224	5.1	50.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36,673	3.7	36.0
7	Diabetes mellitus (E10-E14)	22,612	2.3	22.2
8	Influenza and pneumonia (J09-J18)	21,587	2.2	21.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,460	1.8	18.1
10	Septicemia (A40-A41)	13,988	1.4	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,402	1.1	11.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,385	0.8	8.2
13	Parkinson's disease (G20-G21)	8,191	0.8	8.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,772	0.7	6.7
15	Pneumonitis due to solids and liquids (J69)	6,677	0.7	6.6
...	All other causes (Residual)	210,076	21.0	206.5
Non-Hispanic white, female, 1-4 years				
...	All causes	920	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	31.3	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	128	13.9	3.0
3	Malignant neoplasms (C00-C97)	87	9.5	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	7.5	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.5	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	2.2	0.5
7	Septicemia (A40-A41)	18	2.0	*
7	Influenza and pneumonia (J09-J18)	18	2.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	10	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	7	0.8	*
11	Cerebrovascular diseases (I60-I69)	6	0.7	*
12	Meningitis (G00,G03)	4	0.4	*
13	Meningococcal infection (A39)	3	0.3	*
13	Anemias (D50-D64)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	233	25.3	5.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 5-14 years				
...	All causes	1,219	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	361	29.6	3.2
2	Malignant neoplasms (C00-C97)	223	18.3	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	7.4	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.0	0.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	3.2	0.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	3.1	0.3
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	2.3	0.2
8	Influenza and pneumonia (J09-J18)	27	2.2	0.2
9	Chronic lower respiratory diseases (J40-J47)	20	1.6	0.2
10	Cerebrovascular diseases (I60-I69)	16	1.3	*
11	Septicemia (A40-A41)	13	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
14	Diabetes mellitus (E10-E14)	6	0.5	*
15	Anemias (D50-D64)	5	0.4	*
...	All other causes (Residual)	289	23.7	2.6
Non-Hispanic white, female, 15-24 years				
...	All causes	4,768	100.0	38.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,335	49.0	18.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	552	11.6	4.4
3	Malignant neoplasms (C00-C97)	336	7.0	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	220	4.6	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	138	2.9	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	2.0	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	73	1.5	0.6
8	Influenza and pneumonia (J09-J18)	51	1.1	0.4
9	Diabetes mellitus (E10-E14)	41	0.9	0.3
10	Cerebrovascular diseases (I60-I69)	39	0.8	0.3
11	Septicemia (A40-A41)	38	0.8	0.3
12	Chronic lower respiratory diseases (J40-J47)	26	0.5	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.4	*
15	Anemias (D50-D64)	11	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.2	*
...	All other causes (Residual)	761	16.0	6.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 25-44 years				
...	All causes	24,932	100.0	101.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,370	25.5	25.8
2	Malignant neoplasms (C00-C97)	5,223	20.9	21.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,433	9.8	9.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,114	8.5	8.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	672	2.7	2.7
6	Cerebrovascular diseases (I60-I69)	585	2.3	2.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	584	2.3	2.4
8	Diabetes mellitus (E10-E14)	532	2.1	2.2
9	Influenza and pneumonia (J09-J18)	335	1.3	1.4
10	Chronic lower respiratory diseases (J40-J47)	312	1.3	1.3
11	Septicemia (A40-A41)	305	1.2	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	251	1.0	1.0
13	Pregnancy, childbirth and the puerperium (O00-O99)	236	0.9	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	0.8	0.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	148	0.6	0.6
...	All other causes (Residual)	4,624	18.5	18.7
Non-Hispanic white, female, 45-64 years				
...	All causes	137,452	100.0	459.9
1	Malignant neoplasms (C00-C97)	54,515	39.7	182.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,874	15.2	69.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,634	6.3	28.9
4	Chronic lower respiratory diseases (J40-J47)	7,797	5.7	26.1
5	Cerebrovascular diseases (I60-I69)	4,268	3.1	14.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,150	3.0	13.9
7	Diabetes mellitus (E10-E14)	3,947	2.9	13.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,210	2.3	10.7
9	Septicemia (A40-A41)	2,109	1.5	7.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,681	1.2	5.6
11	Influenza and pneumonia (J09-J18)	1,671	1.2	5.6
12	Viral hepatitis (B15-B19)	1,087	0.8	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	755	0.5	2.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	680	0.5	2.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	553	0.4	1.9
...	All other causes (Residual)	21,521	15.7	72.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic white, female, 65 years and over				
...	All causes	824,118	100.0	4,509.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	212,619	25.8	1,163.4
2	Malignant neoplasms (C00-C97)	158,350	19.2	866.5
3	Chronic lower respiratory diseases (J40-J47)	57,327	7.0	313.7
4	Cerebrovascular diseases (I60-I69)	56,868	6.9	311.2
5	Alzheimer's disease (G30)	50,794	6.2	277.9
6	Influenza and pneumonia (J09-J18)	19,454	2.4	106.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,441	2.2	100.9
8	Diabetes mellitus (E10-E14)	18,085	2.2	99.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,528	2.0	90.4
10	Septicemia (A40-A41)	11,478	1.4	62.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,581	1.3	57.9
12	Parkinson's disease (G20-G21)	8,027	1.0	43.9
13	Pneumonitis due to solids and liquids (J69)	6,184	0.8	33.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,064	0.6	27.7
15	Atherosclerosis (I70)	3,611	0.4	19.8
...	All other causes (Residual)	170,707	20.7	934.1
Non-Hispanic black, both sexes, all ages				
...	All causes	283,438	100.0	718.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,215	24.1	173.0
2	Malignant neoplasms (C00-C97)	65,276	23.0	165.5
3	Cerebrovascular diseases (I60-I69)	15,833	5.6	40.1
4	Diabetes mellitus (E10-E14)	11,996	4.2	30.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,853	4.2	30.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,769	3.1	22.2
7	Chronic lower respiratory diseases (J40-J47)	8,614	3.0	21.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,679	2.7	19.5
9	Septicemia (A40-A41)	5,958	2.1	15.1
10	Alzheimer's disease (G30)	5,169	1.8	13.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,059	1.8	12.8
12	Influenza and pneumonia (J09-J18)	4,854	1.7	12.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,598	1.6	11.7
14	Certain conditions originating in the perinatal period (P00-P96)	3,846	1.4	9.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,601	0.9	6.6
...	All other causes (Residual)	53,118	18.7	134.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 1-4 years				
...	All causes	993	100.0	40.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	249	25.1	10.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	15.4	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	12.5	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	5.0	2.0
5	Malignant neoplasms (C00-C97)	47	4.7	1.9
6	Influenza and pneumonia (J09-J18)	25	2.5	1.0
7	Chronic lower respiratory diseases (J40-J47)	19	1.9	*
8	Certain conditions originating in the perinatal period (P00-P96)	15	1.5	*
9	Anemias (D50-D64)	14	1.4	*
10	Cerebrovascular diseases (I60-I69)	13	1.3	*
11	Septicemia (A40-A41)	11	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Meningitis (G00,G03)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	250	25.2	10.1
Non-Hispanic black, both sexes, 5-14 years				
...	All causes	1,118	100.0	18.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	312	27.9	5.1
2	Malignant neoplasms (C00-C97)	142	12.7	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	8.9	1.6
4	Chronic lower respiratory diseases (J40-J47)	71	6.4	1.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	5.5	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	5.4	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.8	0.7
8	Cerebrovascular diseases (I60-I69)	21	1.9	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.5	*
10	Septicemia (A40-A41)	14	1.3	*
10	Anemias (D50-D64)	14	1.3	*
12	Influenza and pneumonia (J09-J18)	11	1.0	*
13	Diabetes mellitus (E10-E14)	6	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (Residual)	235	21.0	3.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 15-24 years				
...	All causes	6,546	100.0	98.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,762	42.2	41.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,426	21.8	21.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	454	6.9	6.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	327	5.0	4.9
5	Malignant neoplasms (C00-C97)	264	4.0	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	109	1.7	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	1.4	1.3
8	Anemias (D50-D64)	79	1.2	1.2
9	Chronic lower respiratory diseases (J40-J47)	60	0.9	0.9
10	Diabetes mellitus (E10-E14)	51	0.8	0.8
11	Pregnancy, childbirth and the puerperium (O00-O99)	50	0.8	0.8
12	Cerebrovascular diseases (I60-I69)	49	0.7	0.7
13	Septicemia (A40-A41)	41	0.6	0.6
14	Influenza and pneumonia (J09-J18)	39	0.6	0.6
15	Legal intervention (Y35,Y89.0)	28	0.4	0.4
...	All other causes (Residual)	718	11.0	10.8
Non-Hispanic black, both sexes, 25-44 years				
...	All causes	22,593	100.0	211.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,593	15.9	33.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,431	15.2	32.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,148	13.9	29.4
4	Malignant neoplasms (C00-C97)	2,660	11.8	24.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,536	6.8	14.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	929	4.1	8.7
7	Diabetes mellitus (E10-E14)	713	3.2	6.7
8	Cerebrovascular diseases (I60-I69)	690	3.1	6.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	372	1.6	3.5
10	Septicemia (A40-A41)	295	1.3	2.8
11	Chronic lower respiratory diseases (J40-J47)	275	1.2	2.6
12	Anemias (D50-D64)	237	1.0	2.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	227	1.0	2.1
14	Influenza and pneumonia (J09-J18)	225	1.0	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	214	0.9	2.0
...	All other causes (Residual)	4,048	17.9	37.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, both sexes, 45-64 years				
...	All causes	84,986	100.0	903.6
1	Malignant neoplasms (C00-C97)	24,688	29.0	262.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,712	24.4	220.2
3	Cerebrovascular diseases (I60-I69)	4,347	5.1	46.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,027	4.7	42.8
5	Diabetes mellitus (E10-E14)	3,849	4.5	40.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,624	3.1	27.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,319	2.7	24.7
8	Chronic lower respiratory diseases (J40-J47)	2,091	2.5	22.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,817	2.1	19.3
10	Septicemia (A40-A41)	1,704	2.0	18.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,400	1.6	14.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	999	1.2	10.6
13	Influenza and pneumonia (J09-J18)	976	1.1	10.4
14	Viral hepatitis (B15-B19)	805	0.9	8.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	535	0.6	5.7
...	All other causes (Residual)	12,093	14.2	128.6
Non-Hispanic black, both sexes, 65 years and over				
...	All causes	160,114	100.0	4,663.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,376	27.1	1,263.3
2	Malignant neoplasms (C00-C97)	37,466	23.4	1,091.2
3	Cerebrovascular diseases (I60-I69)	10,668	6.7	310.7
4	Diabetes mellitus (E10-E14)	7,373	4.6	214.7
5	Chronic lower respiratory diseases (J40-J47)	6,089	3.8	177.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,014	3.8	175.2
7	Alzheimer's disease (G30)	5,109	3.2	148.8
8	Septicemia (A40-A41)	3,806	2.4	110.9
9	Influenza and pneumonia (J09-J18)	3,507	2.2	102.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,419	2.1	99.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,351	1.5	68.5
12	Pneumonitis due to solids and liquids (J69)	1,084	0.7	31.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	763	0.5	22.2
14	Parkinson's disease (G20-G21)	747	0.5	21.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	565	0.4	16.5
...	All other causes (Residual)	27,777	17.3	809.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, all ages				
...	All causes	143,824	100.0	764.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,597	24.1	183.9
2	Malignant neoplasms (C00-C97)	33,607	23.4	178.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,920	5.5	42.1
4	Cerebrovascular diseases (I60-I69)	6,890	4.8	36.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,580	4.6	35.0
6	Diabetes mellitus (E10-E14)	5,572	3.9	29.6
7	Chronic lower respiratory diseases (J40-J47)	4,480	3.1	23.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,984	2.8	21.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,000	2.1	15.9
10	Septicemia (A40-A41)	2,673	1.9	14.2
11	Influenza and pneumonia (J09-J18)	2,340	1.6	12.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,196	1.5	11.7
13	Certain conditions originating in the perinatal period (P00-P96)	2,116	1.5	11.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,712	1.2	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,694	1.2	9.0
...	All other causes (Residual)	24,463	17.0	130.0
Non-Hispanic black, male, 1-4 years				
...	All causes	570	100.0	45.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	27.9	12.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	91	16.0	7.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	10.0	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.9	2.2
5	Malignant neoplasms (C00-C97)	26	4.6	2.1
6	Influenza and pneumonia (J09-J18)	15	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	10	1.8	*
8	Anemias (D50-D64)	9	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.1	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
12	Septicemia (A40-A41)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
14	Meningococcal infection (A39)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Hernia (K40-K46)	2	0.4	*
...	All other causes (Residual)	141	24.7	11.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 5-14 years				
...	All causes	649	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	31.9	6.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	10.3	2.1
3	Malignant neoplasms (C00-C97)	62	9.6	2.0
4	Chronic lower respiratory diseases (J40-J47)	51	7.9	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	5.9	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	5.2	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	3.5	0.7
8	Cerebrovascular diseases (I60-I69)	14	2.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.4	*
10	Septicemia (A40-A41)	7	1.1	*
10	Influenza and pneumonia (J09-J18)	7	1.1	*
12	Anemias (D50-D64)	6	0.9	*
13	Diabetes mellitus (E10-E14)	3	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	116	17.9	3.7
Non-Hispanic black, male, 15-24 years				
...	All causes	5,033	100.0	150.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,499	49.7	74.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,053	20.9	31.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	385	7.6	11.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	204	4.1	6.1
5	Malignant neoplasms (C00-C97)	148	2.9	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	70	1.4	2.1
7	Anemias (D50-D64)	55	1.1	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	1.1	1.6
9	Chronic lower respiratory diseases (J40-J47)	41	0.8	1.2
10	Legal intervention (Y35,Y89.0)	27	0.5	0.8
11	Septicemia (A40-A41)	26	0.5	0.8
11	Diabetes mellitus (E10-E14)	26	0.5	0.8
13	Influenza and pneumonia (J09-J18)	24	0.5	0.7
14	Cerebrovascular diseases (I60-I69)	21	0.4	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.2	*
...	All other causes (Residual)	389	7.7	11.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 25-44 years				
...	All causes	13,946	100.0	275.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,989	21.4	58.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,236	16.0	44.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,217	15.9	43.7
4	Malignant neoplasms (C00-C97)	1,034	7.4	20.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	921	6.6	18.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	765	5.5	15.1
7	Diabetes mellitus (E10-E14)	412	3.0	8.1
8	Cerebrovascular diseases (I60-I69)	362	2.6	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	1.4	3.8
10	Chronic lower respiratory diseases (J40-J47)	157	1.1	3.1
11	Septicemia (A40-A41)	139	1.0	2.7
12	Influenza and pneumonia (J09-J18)	132	0.9	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	126	0.9	2.5
14	Anemias (D50-D64)	117	0.8	2.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	115	0.8	2.3
...	All other causes (Residual)	2,029	14.5	40.0
Non-Hispanic black, male, 45-64 years				
...	All causes	49,284	100.0	1,130.9
1	Malignant neoplasms (C00-C97)	13,053	26.5	299.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,035	26.4	299.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,792	5.7	64.1
4	Cerebrovascular diseases (I60-I69)	2,423	4.9	55.6
5	Diabetes mellitus (E10-E14)	2,125	4.3	48.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,777	3.6	40.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,237	2.5	28.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,233	2.5	28.3
9	Chronic lower respiratory diseases (J40-J47)	1,070	2.2	24.6
10	Septicemia (A40-A41)	890	1.8	20.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	783	1.6	18.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	781	1.6	17.9
13	Influenza and pneumonia (J09-J18)	562	1.1	12.9
14	Viral hepatitis (B15-B19)	554	1.1	12.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	426	0.9	9.8
...	All other causes (Residual)	6,543	13.3	150.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, male, 65 years and over				
...	All causes	70,396	100.0	5,223.0
1	Malignant neoplasms (C00-C97)	19,280	27.4	1,430.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,011	27.0	1,410.5
3	Cerebrovascular diseases (I60-I69)	4,047	5.7	300.3
4	Chronic lower respiratory diseases (J40-J47)	3,144	4.5	233.3
5	Diabetes mellitus (E10-E14)	3,003	4.3	222.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,524	3.6	187.3
7	Septicemia (A40-A41)	1,562	2.2	115.9
8	Influenza and pneumonia (J09-J18)	1,558	2.2	115.6
9	Alzheimer's disease (G30)	1,444	2.1	107.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,280	1.8	95.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,278	1.8	94.8
12	Pneumonitis due to solids and liquids (J69)	538	0.8	39.9
13	Parkinson's disease (G20-G21)	413	0.6	30.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	344	0.5	25.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	314	0.4	23.3
...	All other causes (Residual)	10,656	15.1	790.6
Non-Hispanic black, female, all ages				
...	All causes	139,614	100.0	676.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,618	24.1	163.0
2	Malignant neoplasms (C00-C97)	31,669	22.7	153.5
3	Cerebrovascular diseases (I60-I69)	8,943	6.4	43.4
4	Diabetes mellitus (E10-E14)	6,424	4.6	31.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,785	3.4	23.2
6	Chronic lower respiratory diseases (J40-J47)	4,134	3.0	20.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,933	2.8	19.1
8	Alzheimer's disease (G30)	3,698	2.6	17.9
9	Septicemia (A40-A41)	3,285	2.4	15.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,863	2.1	13.9
11	Influenza and pneumonia (J09-J18)	2,514	1.8	12.2
12	Certain conditions originating in the perinatal period (P00-P96)	1,730	1.2	8.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,598	1.1	7.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,099	0.8	5.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	907	0.6	4.4
...	All other causes (Residual)	28,414	20.4	137.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 1-4 years				
...	All causes	423	100.0	34.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	21.3	7.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	15.8	5.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	14.7	5.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	5.2	1.8
5	Malignant neoplasms (C00-C97)	21	5.0	1.7
6	Influenza and pneumonia (J09-J18)	10	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.1	*
8	Cerebrovascular diseases (I60-I69)	8	1.9	*
9	Septicemia (A40-A41)	7	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	7	1.7	*
11	Anemias (D50-D64)	5	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Meningitis (G00,G03)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	104	24.6	8.5
Non-Hispanic black, female, 5-14 years				
...	All causes	469	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	22.4	3.5
2	Malignant neoplasms (C00-C97)	80	17.1	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.8	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	6.0	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.7	0.7
6	Chronic lower respiratory diseases (J40-J47)	20	4.3	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	4.3	0.7
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.7	*
8	Anemias (D50-D64)	8	1.7	*
10	Septicemia (A40-A41)	7	1.5	*
10	Cerebrovascular diseases (I60-I69)	7	1.5	*
12	Influenza and pneumonia (J09-J18)	4	0.9	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
...	All other causes (Residual)	116	24.7	3.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 15-24 years				
...	All causes	1,513	100.0	45.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	373	24.7	11.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	263	17.4	7.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	8.1	3.7
4	Malignant neoplasms (C00-C97)	116	7.7	3.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	4.6	2.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	50	3.3	1.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	39	2.6	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	2.4	1.1
9	Cerebrovascular diseases (I60-I69)	28	1.9	0.8
10	Diabetes mellitus (E10-E14)	25	1.7	0.8
11	Anemias (D50-D64)	24	1.6	0.7
12	Chronic lower respiratory diseases (J40-J47)	19	1.3	*
13	Septicemia (A40-A41)	15	1.0	*
13	Influenza and pneumonia (J09-J18)	15	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.0	*
...	All other causes (Residual)	303	20.0	9.1
Non-Hispanic black, female, 25-44 years				
...	All causes	8,647	100.0	153.6
1	Malignant neoplasms (C00-C97)	1,626	18.8	28.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,376	15.9	24.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	912	10.5	16.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	615	7.1	10.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	442	5.1	7.9
6	Cerebrovascular diseases (I60-I69)	328	3.8	5.8
7	Diabetes mellitus (E10-E14)	301	3.5	5.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	196	2.3	3.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	2.0	3.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	164	1.9	2.9
11	Septicemia (A40-A41)	156	1.8	2.8
12	Anemias (D50-D64)	120	1.4	2.1
13	Chronic lower respiratory diseases (J40-J47)	118	1.4	2.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	1.2	1.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	99	1.1	1.8
...	All other causes (Residual)	1,916	22.2	34.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2010

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic black, female, 45-64 years				
...	All causes	35,702	100.0	707.4
1	Malignant neoplasms (C00-C97)	11,635	32.6	230.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,677	21.5	152.1
3	Cerebrovascular diseases (I60-I69)	1,924	5.4	38.1
4	Diabetes mellitus (E10-E14)	1,724	4.8	34.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,235	3.5	24.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,082	3.0	21.4
7	Chronic lower respiratory diseases (J40-J47)	1,021	2.9	20.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	847	2.4	16.8
9	Septicemia (A40-A41)	814	2.3	16.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	619	1.7	12.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	584	1.6	11.6
12	Influenza and pneumonia (J09-J18)	414	1.2	8.2
13	Viral hepatitis (B15-B19)	251	0.7	5.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	216	0.6	4.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	152	0.4	3.0
...	All other causes (Residual)	5,507	15.4	109.1
Non-Hispanic black, female, 65 years and over				
...	All causes	89,718	100.0	4,301.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,365	27.2	1,168.2
2	Malignant neoplasms (C00-C97)	18,186	20.3	872.0
3	Cerebrovascular diseases (I60-I69)	6,621	7.4	317.5
4	Diabetes mellitus (E10-E14)	4,370	4.9	209.5
5	Alzheimer's disease (G30)	3,665	4.1	175.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,490	3.9	167.3
7	Chronic lower respiratory diseases (J40-J47)	2,945	3.3	141.2
8	Septicemia (A40-A41)	2,244	2.5	107.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,139	2.4	102.6
10	Influenza and pneumonia (J09-J18)	1,949	2.2	93.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,073	1.2	51.4
12	Pneumonitis due to solids and liquids (J69)	546	0.6	26.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	449	0.5	21.5
14	Parkinson's disease (G20-G21)	334	0.4	16.0
15	Atherosclerosis (I70)	301	0.3	14.4
...	All other causes (Residual)	17,041	19.0	817.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable