

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, all ages				
...	All causes	2,744,248	100.0	849.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	635,260	23.1	196.6
2	Malignant neoplasms (C00-C97)	598,038	21.8	185.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161,374	5.9	49.9
4	Chronic lower respiratory diseases (J40-J47)	154,596	5.6	47.8
5	Cerebrovascular diseases (I60-I69)	142,142	5.2	44.0
6	Alzheimer's disease (G30)	116,103	4.2	35.9
7	Diabetes mellitus (E10-E14)	80,058	2.9	24.8
8	Influenza and pneumonia (J09-J18)	51,537	1.9	15.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,046	1.8	15.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44,965	1.6	13.9
11	Septicemia (A40-A41)	40,613	1.5	12.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	40,545	1.5	12.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33,246	1.2	10.3
14	Parkinson's disease (G20-G21)	29,697	1.1	9.2
15	Pneumonitis due to solids and liquids (J69)	19,715	0.7	6.1
...	All other causes (residual)	546,313	19.9	169.1
All races and origins, both sexes, 1-4 years				
...	All causes	4,045	100.0	25.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,261	31.2	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	433	10.7	2.7
3	Malignant neoplasms (C00-C97)	377	9.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	8.4	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	118	2.9	0.7
6	Influenza and pneumonia (J09-J18)	103	2.5	0.6
7	Septicemia (A40-A41)	70	1.7	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	60	1.5	0.4
9	Cerebrovascular diseases (I60-I69)	55	1.4	0.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	1.2	0.3
12	Anemias (D50-D64)	20	0.5	0.1
13	Acute bronchitis and bronchiolitis (J20-J21)	18	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
15	Meningitis (G00,G03)	14	0.3	*
...	All other causes (residual)	1,062	26.3	6.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 5-14 years				
...	All causes	5,503	100.0	13.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,634	29.7	4.0
2	Malignant neoplasms (C00-C97)	880	16.0	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	443	8.1	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	349	6.3	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	286	5.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	188	3.4	0.5
7	Chronic lower respiratory diseases (J40-J47)	143	2.6	0.3
8	Cerebrovascular diseases (I60-I69)	88	1.6	0.2
9	Influenza and pneumonia (J09-J18)	87	1.6	0.2
10	Septicemia (A40-A41)	71	1.3	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	1.1	0.1
12	Diabetes mellitus (E10-E14)	26	0.5	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	26	0.5	0.1
14	Anemias (D50-D64)	24	0.4	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	23	0.4	0.1
...	All other causes (residual)	1,176	21.4	2.9
All races and origins, both sexes, 15-24 years				
...	All causes	32,575	100.0	74.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,895	42.7	31.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,723	17.6	13.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,172	15.9	11.9
4	Malignant neoplasms (C00-C97)	1,431	4.4	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	949	2.9	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	388	1.2	0.9
7	Diabetes mellitus (E10-E14)	211	0.6	0.5
8	Chronic lower respiratory diseases (J40-J47)	206	0.6	0.5
9	Influenza and pneumonia (J09-J18)	189	0.6	0.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	184	0.6	0.4
11	Cerebrovascular diseases (I60-I69)	148	0.5	0.3
12	Septicemia (A40-A41)	143	0.4	0.3
13	Anemias (D50-D64)	78	0.2	0.2
14	Legal intervention (Y35,Y89.0)	77	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.2	0.2
...	All other causes (residual)	3,711	11.4	8.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 25-34 years				
...	All causes	57,616	100.0	129.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23,984	41.6	53.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,366	12.8	16.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,376	9.3	12.0
4	Malignant neoplasms (C00-C97)	3,791	6.6	8.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,445	6.0	7.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	925	1.6	2.1
7	Diabetes mellitus (E10-E14)	792	1.4	1.8
8	Cerebrovascular diseases (I60-I69)	575	1.0	1.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	546	0.9	1.2
10	Pregnancy, childbirth and the puerperium (O00-O99)	472	0.8	1.1
11	Influenza and pneumonia (J09-J18)	452	0.8	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	436	0.8	1.0
13	Septicemia (A40-A41)	419	0.7	0.9
14	Chronic lower respiratory diseases (J40-J47)	353	0.6	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	0.5	0.6
...	All other causes (residual)	8,419	14.6	18.8
All races and origins, both sexes, 35-44 years				
...	All causes	77,792	100.0	192.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,975	27.0	51.8
2	Malignant neoplasms (C00-C97)	10,903	14.0	26.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,477	13.5	25.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,030	9.0	17.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,369	4.3	8.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,851	3.7	7.0
7	Diabetes mellitus (E10-E14)	2,049	2.6	5.1
8	Cerebrovascular diseases (I60-I69)	1,851	2.4	4.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	971	1.2	2.4
10	Septicemia (A40-A41)	897	1.2	2.2
11	Influenza and pneumonia (J09-J18)	894	1.1	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	743	1.0	1.8
13	Chronic lower respiratory diseases (J40-J47)	692	0.9	1.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	459	0.6	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	421	0.5	1.0
...	All other causes (residual)	13,210	17.0	32.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 45-54 years				
...	All causes	173,516	100.0	405.5
1	Malignant neoplasms (C00-C97)	41,291	23.8	96.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,027	19.6	79.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23,377	13.5	54.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,437	4.9	19.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	8,364	4.8	19.5
6	Diabetes mellitus (E10-E14)	6,267	3.6	14.6
7	Cerebrovascular diseases (I60-I69)	5,353	3.1	12.5
8	Chronic lower respiratory diseases (J40-J47)	4,307	2.5	10.1
9	Septicemia (A40-A41)	2,472	1.4	5.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,152	1.2	5.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,140	1.2	5.0
12	Influenza and pneumonia (J09-J18)	2,133	1.2	5.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,831	1.1	4.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,606	0.9	3.8
15	Viral hepatitis (B15-B19)	1,162	0.7	2.7
...	All other causes (residual)	28,597	16.5	66.8
All races and origins, both sexes, 55-64 years				
...	All causes	366,445	100.0	883.8
1	Malignant neoplasms (C00-C97)	116,364	31.8	280.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,610	21.5	189.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21,860	6.0	52.7
4	Chronic lower respiratory diseases (J40-J47)	17,810	4.9	43.0
5	Diabetes mellitus (E10-E14)	14,251	3.9	34.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	13,448	3.7	32.4
7	Cerebrovascular diseases (I60-I69)	12,310	3.4	29.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,759	2.1	18.7
9	Septicemia (A40-A41)	5,941	1.6	14.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,650	1.5	13.6
11	Influenza and pneumonia (J09-J18)	5,034	1.4	12.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,741	1.0	9.0
13	Viral hepatitis (B15-B19)	2,996	0.8	7.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,803	0.5	4.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,568	0.4	3.8
...	All other causes (residual)	57,300	15.6	138.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 65-74 years				
...	All causes	512,080	100.0	1,788.6
1	Malignant neoplasms (C00-C97)	165,583	32.3	578.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	112,361	21.9	392.5
3	Chronic lower respiratory diseases (J40-J47)	38,403	7.5	134.1
4	Cerebrovascular diseases (I60-I69)	21,768	4.3	76.0
5	Diabetes mellitus (E10-E14)	20,002	3.9	69.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,050	2.7	49.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,904	1.9	34.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	8,797	1.7	30.7
9	Septicemia (A40-A41)	8,767	1.7	30.6
10	Influenza and pneumonia (J09-J18)	8,169	1.6	28.5
11	Alzheimer's disease (G30)	6,761	1.3	23.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,454	1.1	19.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,403	0.9	15.4
14	Parkinson's disease (G20-G21)	4,094	0.8	14.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,122	0.6	10.9
...	All other causes (residual)	80,442	15.7	281.0
All races and origins, both sexes, 75-84 years				
...	All causes	636,916	100.0	4,474.8
1	Malignant neoplasms (C00-C97)	153,961	24.2	1,081.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	147,619	23.2	1,037.1
3	Chronic lower respiratory diseases (J40-J47)	49,413	7.8	347.2
4	Cerebrovascular diseases (I60-I69)	37,789	5.9	265.5
5	Alzheimer's disease (G30)	30,479	4.8	214.1
6	Diabetes mellitus (E10-E14)	19,633	3.1	137.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,759	2.5	110.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,959	2.2	98.1
9	Influenza and pneumonia (J09-J18)	12,596	2.0	88.5
10	Parkinson's disease (G20-G21)	12,165	1.9	85.5
11	Septicemia (A40-A41)	10,522	1.7	73.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,644	1.2	53.7
13	Pneumonitis due to solids and liquids (J69)	5,140	0.8	36.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,699	0.7	33.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,544	0.7	31.9
...	All other causes (residual)	110,994	17.4	779.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 85 years and over				
...	All causes	854,462	100.0	13,392.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,138	28.9	3,873.4
2	Malignant neoplasms (C00-C97)	103,383	12.1	1,620.3
3	Alzheimer's disease (G30)	77,643	9.1	1,216.9
4	Cerebrovascular diseases (I60-I69)	62,073	7.3	972.9
5	Chronic lower respiratory diseases (J40-J47)	43,186	5.1	676.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23,332	2.7	365.7
7	Influenza and pneumonia (J09-J18)	21,714	2.5	340.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,232	2.0	270.1
9	Diabetes mellitus (E10-E14)	16,817	2.0	263.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,185	1.7	222.3
11	Parkinson's disease (G20-G21)	12,670	1.5	198.6
12	Septicemia (A40-A41)	11,116	1.3	174.2
13	Pneumonitis due to solids and liquids (J69)	9,352	1.1	146.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,243	0.6	82.2
15	Nutritional deficiencies (E40-E64)	3,406	0.4	53.4
...	All other causes (residual)	185,972	21.8	2,914.8
All races and origins, male, all ages				
...	All causes	1,400,232	100.0	880.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	339,265	24.2	213.3
2	Malignant neoplasms (C00-C97)	314,571	22.5	197.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103,864	7.4	65.3
4	Chronic lower respiratory diseases (J40-J47)	73,045	5.2	45.9
5	Cerebrovascular diseases (I60-I69)	59,355	4.2	37.3
6	Diabetes mellitus (E10-E14)	43,763	3.1	27.5
7	Alzheimer's disease (G30)	35,372	2.5	22.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34,727	2.5	21.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	25,818	1.8	16.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,399	1.8	16.0
11	Influenza and pneumonia (J09-J18)	25,011	1.8	15.7
12	Septicemia (A40-A41)	19,678	1.4	12.4
13	Parkinson's disease (G20-G21)	17,877	1.3	11.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15,467	1.1	9.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,598	1.0	9.2
...	All other causes (residual)	252,422	18.0	158.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 1-4 years				
...	All causes	2,256	100.0	27.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	776	34.4	9.5
2	Malignant neoplasms (C00-C97)	206	9.1	2.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	206	9.1	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	176	7.8	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	2.7	0.8
6	Influenza and pneumonia (J09-J18)	53	2.3	0.6
7	Septicemia (A40-A41)	37	1.6	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	35	1.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	34	1.5	0.4
10	Cerebrovascular diseases (I60-I69)	28	1.2	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.0	0.3
12	Acute bronchitis and bronchiolitis (J20-J21)	14	0.6	*
13	Anemias (D50-D64)	7	0.3	*
13	Meningitis (G00,G03)	7	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (residual)	586	26.0	7.2
All races and origins, male, 5-14 years				
...	All causes	3,141	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	990	31.5	4.7
2	Malignant neoplasms (C00-C97)	490	15.6	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	8.7	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	187	6.0	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	167	5.3	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	3.2	0.5
7	Chronic lower respiratory diseases (J40-J47)	86	2.7	0.4
8	Cerebrovascular diseases (I60-I69)	55	1.8	0.3
9	Influenza and pneumonia (J09-J18)	49	1.6	0.2
10	Septicemia (A40-A41)	37	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.2	0.2
12	Anemias (D50-D64)	16	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	14	0.4	*
14	Diabetes mellitus (E10-E14)	12	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.4	*
...	All other causes (residual)	618	19.7	3.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 15-24 years				
...	All causes	24,013	100.0	107.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,241	42.6	45.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,575	19.1	20.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,479	18.7	20.1
4	Malignant neoplasms (C00-C97)	902	3.8	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	629	2.6	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	231	1.0	1.0
7	Chronic lower respiratory diseases (J40-J47)	127	0.5	0.6
8	Diabetes mellitus (E10-E14)	121	0.5	0.5
9	Influenza and pneumonia (J09-J18)	105	0.4	0.5
10	Cerebrovascular diseases (I60-I69)	92	0.4	0.4
11	Septicemia (A40-A41)	78	0.3	0.3
12	Legal intervention (Y35,Y89.0)	75	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.2	0.2
14	Anemias (D50-D64)	45	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.2	0.2
...	All other causes (residual)	2,229	9.3	10.0
All races and origins, male, 25-34 years				
...	All causes	40,257	100.0	178.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,737	44.1	78.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,887	14.6	26.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,540	11.3	20.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,304	5.7	10.2
5	Malignant neoplasms (C00-C97)	1,825	4.5	8.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	565	1.4	2.5
7	Diabetes mellitus (E10-E14)	456	1.1	2.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	403	1.0	1.8
9	Cerebrovascular diseases (I60-I69)	331	0.8	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	245	0.6	1.1
11	Influenza and pneumonia (J09-J18)	243	0.6	1.1
12	Septicemia (A40-A41)	209	0.5	0.9
13	Chronic lower respiratory diseases (J40-J47)	194	0.5	0.9
14	Legal intervention (Y35,Y89.0)	173	0.4	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	0.4	0.7
...	All other causes (residual)	4,997	12.4	22.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 35-44 years				
...	All causes	49,215	100.0	244.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,981	30.4	74.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,147	14.5	35.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,294	10.8	26.3
4	Malignant neoplasms (C00-C97)	4,625	9.4	22.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,708	5.5	13.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,761	3.6	8.7
7	Diabetes mellitus (E10-E14)	1,238	2.5	6.1
8	Cerebrovascular diseases (I60-I69)	1,060	2.2	5.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	644	1.3	3.2
10	Influenza and pneumonia (J09-J18)	492	1.0	2.4
11	Septicemia (A40-A41)	461	0.9	2.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	413	0.8	2.0
13	Chronic lower respiratory diseases (J40-J47)	364	0.7	1.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	279	0.6	1.4
15	Aortic aneurysm and dissection (I71)	251	0.5	1.2
...	All other causes (residual)	7,497	15.2	37.2
All races and origins, male, 45-54 years				
...	All causes	105,086	100.0	497.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,754	22.6	112.5
2	Malignant neoplasms (C00-C97)	19,933	19.0	94.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,936	15.2	75.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,198	5.9	29.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,424	5.2	25.7
6	Diabetes mellitus (E10-E14)	3,984	3.8	18.9
7	Cerebrovascular diseases (I60-I69)	2,978	2.8	14.1
8	Chronic lower respiratory diseases (J40-J47)	1,931	1.8	9.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,628	1.5	7.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,344	1.3	6.4
11	Septicemia (A40-A41)	1,327	1.3	6.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,267	1.2	6.0
13	Influenza and pneumonia (J09-J18)	1,157	1.1	5.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,018	1.0	4.8
15	Viral hepatitis (B15-B19)	766	0.7	3.6
...	All other causes (residual)	16,441	15.6	77.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 55-64 years				
...	All causes	222,384	100.0	1,112.0
1	Malignant neoplasms (C00-C97)	63,763	28.7	318.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,258	24.4	271.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,174	6.8	75.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	9,175	4.1	45.9
5	Chronic lower respiratory diseases (J40-J47)	9,064	4.1	45.3
6	Diabetes mellitus (E10-E14)	8,776	3.9	43.9
7	Cerebrovascular diseases (I60-I69)	7,160	3.2	35.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,745	2.6	28.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,238	1.5	16.2
10	Septicemia (A40-A41)	3,222	1.4	16.1
11	Influenza and pneumonia (J09-J18)	2,840	1.3	14.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,391	1.1	12.0
13	Viral hepatitis (B15-B19)	2,089	0.9	10.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,375	0.6	6.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,030	0.5	5.2
...	All other causes (residual)	33,084	14.9	165.4
All races and origins, male, 65-74 years				
...	All causes	292,742	100.0	2,186.1
1	Malignant neoplasms (C00-C97)	91,612	31.3	684.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,829	24.5	536.4
3	Chronic lower respiratory diseases (J40-J47)	19,699	6.7	147.1
4	Diabetes mellitus (E10-E14)	11,676	4.0	87.2
5	Cerebrovascular diseases (I60-I69)	11,585	4.0	86.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,133	3.1	68.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,658	1.9	42.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,307	1.8	39.6
9	Influenza and pneumonia (J09-J18)	4,557	1.6	34.0
10	Septicemia (A40-A41)	4,556	1.6	34.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,463	1.2	25.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,021	1.0	22.6
13	Alzheimer's disease (G30)	2,833	1.0	21.2
14	Parkinson's disease (G20-G21)	2,717	0.9	20.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,822	0.6	13.6
...	All other causes (residual)	43,274	14.8	323.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 75-84 years				
...	All causes	324,819	100.0	5,258.6
1	Malignant neoplasms (C00-C97)	81,820	25.2	1,324.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	81,041	24.9	1,312.0
3	Chronic lower respiratory diseases (J40-J47)	23,659	7.3	383.0
4	Cerebrovascular diseases (I60-I69)	17,108	5.3	277.0
5	Alzheimer's disease (G30)	11,545	3.6	186.9
6	Diabetes mellitus (E10-E14)	10,391	3.2	168.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,662	2.7	140.2
8	Parkinson's disease (G20-G21)	7,795	2.4	126.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,272	2.2	117.7
10	Influenza and pneumonia (J09-J18)	6,498	2.0	105.2
11	Septicemia (A40-A41)	5,084	1.6	82.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,394	1.0	54.9
13	Pneumonitis due to solids and liquids (J69)	2,962	0.9	48.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,659	0.8	43.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,457	0.8	39.8
...	All other causes (residual)	52,472	16.2	849.5
All races and origins, male, 85 years and over				
...	All causes	323,355	100.0	14,533.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	97,970	30.3	4,403.5
2	Malignant neoplasms (C00-C97)	49,356	15.3	2,218.4
3	Alzheimer's disease (G30)	20,477	6.3	920.4
4	Cerebrovascular diseases (I60-I69)	18,876	5.8	848.4
5	Chronic lower respiratory diseases (J40-J47)	17,863	5.5	802.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,506	2.9	427.3
7	Influenza and pneumonia (J09-J18)	8,911	2.8	400.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,687	2.4	345.5
9	Diabetes mellitus (E10-E14)	7,104	2.2	319.3
10	Parkinson's disease (G20-G21)	6,861	2.1	308.4
11	Pneumonitis due to solids and liquids (J69)	4,746	1.5	213.3
12	Septicemia (A40-A41)	4,555	1.4	204.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,405	1.4	198.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,382	0.7	107.1
15	Aortic aneurysm and dissection (I71)	1,121	0.3	50.4
...	All other causes (residual)	61,535	19.0	2,765.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, all ages				
...	All causes	1,344,016	100.0	819.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	295,995	22.0	180.4
2	Malignant neoplasms (C00-C97)	283,467	21.1	172.8
3	Cerebrovascular diseases (I60-I69)	82,787	6.2	50.5
4	Chronic lower respiratory diseases (J40-J47)	81,551	6.1	49.7
5	Alzheimer's disease (G30)	80,731	6.0	49.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57,510	4.3	35.1
7	Diabetes mellitus (E10-E14)	36,295	2.7	22.1
8	Influenza and pneumonia (J09-J18)	26,526	2.0	16.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,647	1.8	15.0
10	Septicemia (A40-A41)	20,935	1.6	12.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,648	1.4	11.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	14,727	1.1	9.0
13	Parkinson's disease (G20-G21)	11,820	0.9	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,238	0.8	6.2
15	Pneumonitis due to solids and liquids (J69)	8,853	0.7	5.4
...	All other causes (residual)	289,286	21.5	176.3
All races and origins, female, 1-4 years				
...	All causes	1,789	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	485	27.1	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	12.7	2.9
3	Malignant neoplasms (C00-C97)	171	9.6	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	9.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	3.1	0.7
6	Influenza and pneumonia (J09-J18)	50	2.8	0.6
7	Septicemia (A40-A41)	33	1.8	0.4
8	Cerebrovascular diseases (I60-I69)	27	1.5	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.5	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	25	1.4	0.3
11	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
12	Anemias (D50-D64)	13	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.6	*
14	Meningitis (G00,G03)	7	0.4	*
15	Diabetes mellitus (E10-E14)	5	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (residual)	469	26.2	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 5-14 years				
...	All causes	2,362	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	644	27.3	3.2
2	Malignant neoplasms (C00-C97)	390	16.5	1.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	171	7.2	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	6.9	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	5.0	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	3.8	0.4
7	Chronic lower respiratory diseases (J40-J47)	57	2.4	0.3
8	Influenza and pneumonia (J09-J18)	38	1.6	0.2
9	Septicemia (A40-A41)	34	1.4	0.2
10	Cerebrovascular diseases (I60-I69)	33	1.4	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.9	0.1
12	Diabetes mellitus (E10-E14)	14	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	12	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.4	*
...	All other causes (residual)	556	23.5	2.8
All races and origins, female, 15-24 years				
...	All causes	8,562	100.0	40.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,654	42.7	17.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,148	13.4	5.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	693	8.1	3.3
4	Malignant neoplasms (C00-C97)	529	6.2	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	320	3.7	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	184	2.1	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	157	1.8	0.7
8	Diabetes mellitus (E10-E14)	90	1.1	0.4
9	Influenza and pneumonia (J09-J18)	84	1.0	0.4
10	Chronic lower respiratory diseases (J40-J47)	79	0.9	0.4
11	Septicemia (A40-A41)	65	0.8	0.3
12	Cerebrovascular diseases (I60-I69)	56	0.7	0.3
13	Anemias (D50-D64)	33	0.4	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	30	0.4	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.3	0.1
...	All other causes (residual)	1,411	16.5	6.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 25-34 years				
...	All causes	17,359	100.0	78.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,247	36.0	28.3
2	Malignant neoplasms (C00-C97)	1,966	11.3	8.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,479	8.5	6.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,141	6.6	5.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	836	4.8	3.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	472	2.7	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	360	2.1	1.6
8	Diabetes mellitus (E10-E14)	336	1.9	1.5
9	Cerebrovascular diseases (I60-I69)	244	1.4	1.1
10	Septicemia (A40-A41)	210	1.2	1.0
11	Influenza and pneumonia (J09-J18)	209	1.2	0.9
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	1.1	0.9
13	Chronic lower respiratory diseases (J40-J47)	159	0.9	0.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	143	0.8	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	0.7	0.5
...	All other causes (residual)	3,249	18.7	14.7
All races and origins, female, 35-44 years				
...	All causes	28,577	100.0	140.7
1	Malignant neoplasms (C00-C97)	6,278	22.0	30.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,994	21.0	29.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,330	11.7	16.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,736	6.1	8.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,090	3.8	5.4
6	Diabetes mellitus (E10-E14)	811	2.8	4.0
7	Cerebrovascular diseases (I60-I69)	791	2.8	3.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	661	2.3	3.3
9	Septicemia (A40-A41)	436	1.5	2.1
10	Influenza and pneumonia (J09-J18)	402	1.4	2.0
11	Pregnancy, childbirth and the puerperium (O00-O99)	351	1.2	1.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	330	1.2	1.6
13	Chronic lower respiratory diseases (J40-J47)	328	1.1	1.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	327	1.1	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	217	0.8	1.1
...	All other causes (residual)	5,495	19.2	27.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 45-54 years				
...	All causes	68,430	100.0	315.6
1	Malignant neoplasms (C00-C97)	21,358	31.2	98.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,273	15.0	47.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,441	10.9	34.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,940	4.3	13.6
5	Chronic lower respiratory diseases (J40-J47)	2,376	3.5	11.0
6	Cerebrovascular diseases (I60-I69)	2,375	3.5	11.0
7	Diabetes mellitus (E10-E14)	2,283	3.3	10.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,239	3.3	10.3
9	Septicemia (A40-A41)	1,145	1.7	5.3
10	Influenza and pneumonia (J09-J18)	976	1.4	4.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	873	1.3	4.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	588	0.9	2.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	524	0.8	2.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	487	0.7	2.2
15	Viral hepatitis (B15-B19)	396	0.6	1.8
...	All other causes (residual)	12,156	17.8	56.1
All races and origins, female, 55-64 years				
...	All causes	144,061	100.0	671.2
1	Malignant neoplasms (C00-C97)	52,601	36.5	245.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,352	16.9	113.5
3	Chronic lower respiratory diseases (J40-J47)	8,746	6.1	40.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,686	4.6	31.1
5	Diabetes mellitus (E10-E14)	5,475	3.8	25.5
6	Cerebrovascular diseases (I60-I69)	5,150	3.6	24.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,273	3.0	19.9
8	Septicemia (A40-A41)	2,719	1.9	12.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,412	1.7	11.2
10	Influenza and pneumonia (J09-J18)	2,194	1.5	10.2
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,014	1.4	9.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,350	0.9	6.3
13	Viral hepatitis (B15-B19)	907	0.6	4.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	694	0.5	3.2
15	Alzheimer's disease (G30)	644	0.4	3.0
...	All other causes (residual)	23,844	16.6	111.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 65-74 years				
...	All causes	219,338	100.0	1,439.3
1	Malignant neoplasms (C00-C97)	73,971	33.7	485.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,532	18.5	266.0
3	Chronic lower respiratory diseases (J40-J47)	18,704	8.5	122.7
4	Cerebrovascular diseases (I60-I69)	10,183	4.6	66.8
5	Diabetes mellitus (E10-E14)	8,326	3.8	54.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,917	2.2	32.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,597	2.1	30.2
8	Septicemia (A40-A41)	4,211	1.9	27.6
9	Alzheimer's disease (G30)	3,928	1.8	25.8
10	Influenza and pneumonia (J09-J18)	3,612	1.6	23.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	3,139	1.4	20.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,433	1.1	16.0
13	Parkinson's disease (G20-G21)	1,377	0.6	9.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,300	0.6	8.5
15	Pneumonitis due to solids and liquids (J69)	1,107	0.5	7.3
...	All other causes (residual)	37,001	16.9	242.8
All races and origins, female, 75-84 years				
...	All causes	312,097	100.0	3,873.8
1	Malignant neoplasms (C00-C97)	72,141	23.1	895.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	66,578	21.3	826.4
3	Chronic lower respiratory diseases (J40-J47)	25,754	8.3	319.7
4	Cerebrovascular diseases (I60-I69)	20,681	6.6	256.7
5	Alzheimer's disease (G30)	18,934	6.1	235.0
6	Diabetes mellitus (E10-E14)	9,242	3.0	114.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,097	2.3	88.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,687	2.1	83.0
9	Influenza and pneumonia (J09-J18)	6,098	2.0	75.7
10	Septicemia (A40-A41)	5,438	1.7	67.5
11	Parkinson's disease (G20-G21)	4,370	1.4	54.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,250	1.4	52.8
13	Pneumonitis due to solids and liquids (J69)	2,178	0.7	27.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2,087	0.7	25.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,040	0.7	25.3
...	All other causes (residual)	58,522	18.8	726.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 85 years and over				
...	All causes	531,107	100.0	12,780.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	149,168	28.1	3,589.7
2	Alzheimer's disease (G30)	57,166	10.8	1,375.7
3	Malignant neoplasms (C00-C97)	54,027	10.2	1,300.1
4	Cerebrovascular diseases (I60-I69)	43,197	8.1	1,039.5
5	Chronic lower respiratory diseases (J40-J47)	25,323	4.8	609.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,826	2.6	332.7
7	Influenza and pneumonia (J09-J18)	12,803	2.4	308.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,780	1.8	235.4
9	Diabetes mellitus (E10-E14)	9,713	1.8	233.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,545	1.8	229.7
11	Septicemia (A40-A41)	6,561	1.2	157.9
12	Parkinson's disease (G20-G21)	5,809	1.1	139.8
13	Pneumonitis due to solids and liquids (J69)	4,606	0.9	110.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,861	0.5	68.8
15	Nutritional deficiencies (E40-E64)	2,312	0.4	55.6
...	All other causes (residual)	124,410	23.4	2,993.9
Hispanic, both sexes, all ages				
...	All causes	188,254	100.0	327.6
1	Malignant neoplasms (C00-C97)	39,263	20.9	68.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,799	20.1	65.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,711	8.3	27.3
4	Cerebrovascular diseases (I60-I69)	10,283	5.5	17.9
5	Diabetes mellitus (E10-E14)	8,546	4.5	14.9
6	Alzheimer's disease (G30)	6,833	3.6	11.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,141	3.3	10.7
8	Chronic lower respiratory diseases (J40-J47)	5,287	2.8	9.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,775	2.0	6.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,668	1.9	6.4
11	Influenza and pneumonia (J09-J18)	3,641	1.9	6.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,187	1.7	5.5
13	Septicemia (A40-A41)	2,718	1.4	4.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,488	1.3	4.3
15	Certain conditions originating in the perinatal period (P00-P96)	2,415	1.3	4.2
...	All other causes (residual)	36,499	19.4	63.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 1-4 years				
...	All causes	799	100.0	19.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	29.2	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	12.4	2.4
3	Malignant neoplasms (C00-C97)	89	11.1	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	7.8	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	3.4	0.7
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	2.0	*
6	Influenza and pneumonia (J09-J18)	16	2.0	*
8	Septicemia (A40-A41)	14	1.8	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.5	*
10	Cerebrovascular diseases (I60-I69)	8	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	8	1.0	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
13	Anemias (D50-D64)	3	0.4	*
14	Meningitis (G00,G03)	2	0.3	*
14	Hernia (K40-K46)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	*
...	All other causes (residual)	200	25.0	4.9
Hispanic, both sexes, 5-14 years				
...	All causes	1,164	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	28.4	3.2
2	Malignant neoplasms (C00-C97)	226	19.4	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	7.5	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	5.2	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	4.7	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.3	0.4
7	Cerebrovascular diseases (I60-I69)	29	2.5	0.3
8	Chronic lower respiratory diseases (J40-J47)	22	1.9	0.2
9	Septicemia (A40-A41)	18	1.5	*
10	Influenza and pneumonia (J09-J18)	14	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.0	*
12	Anemias (D50-D64)	7	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Diseases of appendix (K35-K38)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (residual)	254	21.8	2.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-24 years				
...	All causes	5,722	100.0	59.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,527	44.2	26.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	987	17.2	10.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	841	14.7	8.8
4	Malignant neoplasms (C00-C97)	370	6.5	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	2.1	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	1.5	0.9
7	Cerebrovascular diseases (I60-I69)	34	0.6	0.4
7	Influenza and pneumonia (J09-J18)	34	0.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	30	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	29	0.5	0.3
11	Diabetes mellitus (E10-E14)	25	0.4	0.3
12	Septicemia (A40-A41)	22	0.4	0.2
13	Legal intervention (Y35,Y89.0)	21	0.4	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.2	*
...	All other causes (residual)	572	10.0	6.0
Hispanic, both sexes, 25-34 years				
...	All causes	8,195	100.0	90.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,212	39.2	35.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	964	11.8	10.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	901	11.0	9.9
4	Malignant neoplasms (C00-C97)	706	8.6	7.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	437	5.3	4.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	194	2.4	2.1
7	Cerebrovascular diseases (I60-I69)	108	1.3	1.2
8	Diabetes mellitus (E10-E14)	99	1.2	1.1
9	Pregnancy, childbirth and the puerperium (O00-O99)	90	1.1	1.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	82	1.0	0.9
11	Influenza and pneumonia (J09-J18)	74	0.9	0.8
12	Septicemia (A40-A41)	69	0.8	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	0.8	0.7
14	Chronic lower respiratory diseases (J40-J47)	36	0.4	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.4	0.4
14	Legal intervention (Y35,Y89.0)	36	0.4	0.4
...	All other causes (residual)	1,089	13.3	12.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-44 years				
...	All causes	10,272	100.0	124.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,592	25.2	31.4
2	Malignant neoplasms (C00-C97)	1,793	17.5	21.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,067	10.4	12.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	703	6.8	8.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	622	6.1	7.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	576	5.6	7.0
7	Cerebrovascular diseases (I60-I69)	326	3.2	4.0
8	Diabetes mellitus (E10-E14)	268	2.6	3.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	141	1.4	1.7
10	Influenza and pneumonia (J09-J18)	135	1.3	1.6
11	Septicemia (A40-A41)	102	1.0	1.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	100	1.0	1.2
13	Chronic lower respiratory diseases (J40-J47)	68	0.7	0.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	61	0.6	0.7
15	Pregnancy, childbirth and the puerperium (O00-O99)	53	0.5	0.6
...	All other causes (residual)	1,665	16.2	20.2
Hispanic, both sexes, 45-54 years				
...	All causes	17,753	100.0	265.6
1	Malignant neoplasms (C00-C97)	4,281	24.1	64.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,880	16.2	43.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,371	13.4	35.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,428	8.0	21.4
5	Diabetes mellitus (E10-E14)	815	4.6	12.2
6	Cerebrovascular diseases (I60-I69)	726	4.1	10.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	550	3.1	8.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	287	1.6	4.3
9	Septicemia (A40-A41)	284	1.6	4.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	273	1.5	4.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	258	1.5	3.9
12	Influenza and pneumonia (J09-J18)	242	1.4	3.6
13	Viral hepatitis (B15-B19)	218	1.2	3.3
14	Chronic lower respiratory diseases (J40-J47)	187	1.1	2.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	167	0.9	2.5
...	All other causes (residual)	2,786	15.7	41.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-64 years				
...	All causes	27,587	100.0	617.2
1	Malignant neoplasms (C00-C97)	8,218	29.8	183.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,342	19.4	119.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,848	6.7	41.3
4	Diabetes mellitus (E10-E14)	1,783	6.5	39.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,706	6.2	38.2
6	Cerebrovascular diseases (I60-I69)	1,200	4.3	26.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	604	2.2	13.5
8	Chronic lower respiratory diseases (J40-J47)	532	1.9	11.9
9	Influenza and pneumonia (J09-J18)	439	1.6	9.8
10	Septicemia (A40-A41)	420	1.5	9.4
11	Viral hepatitis (B15-B19)	388	1.4	8.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	337	1.2	7.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	317	1.1	7.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	215	0.8	4.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	134	0.5	3.0
...	All other causes (residual)	4,104	14.9	91.8
Hispanic, both sexes, 65-74 years				
...	All causes	31,550	100.0	1,313.9
1	Malignant neoplasms (C00-C97)	9,410	29.8	391.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,831	21.7	284.5
3	Diabetes mellitus (E10-E14)	2,069	6.6	86.2
4	Cerebrovascular diseases (I60-I69)	1,705	5.4	71.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,115	3.5	46.4
6	Chronic lower respiratory diseases (J40-J47)	989	3.1	41.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	890	2.8	37.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	872	2.8	36.3
9	Influenza and pneumonia (J09-J18)	643	2.0	26.8
10	Septicemia (A40-A41)	569	1.8	23.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	422	1.3	17.6
12	Alzheimer's disease (G30)	416	1.3	17.3
13	Viral hepatitis (B15-B19)	224	0.7	9.3
14	Parkinson's disease (G20-G21)	221	0.7	9.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	164	0.5	6.8
...	All other causes (residual)	5,010	15.9	208.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-84 years				
...	All causes	38,106	100.0	3,397.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,046	23.7	806.5
2	Malignant neoplasms (C00-C97)	8,932	23.4	796.3
3	Cerebrovascular diseases (I60-I69)	2,651	7.0	236.3
4	Diabetes mellitus (E10-E14)	2,031	5.3	181.1
5	Alzheimer's disease (G30)	1,831	4.8	163.2
6	Chronic lower respiratory diseases (J40-J47)	1,610	4.2	143.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	984	2.6	87.7
8	Influenza and pneumonia (J09-J18)	899	2.4	80.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	810	2.1	72.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	709	1.9	63.2
11	Parkinson's disease (G20-G21)	678	1.8	60.4
12	Septicemia (A40-A41)	638	1.7	56.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	570	1.5	50.8
14	Pneumonitis due to solids and liquids (J69)	277	0.7	24.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	254	0.7	22.6
...	All other causes (residual)	6,186	16.2	551.5
Hispanic, both sexes, 85 years and over				
...	All causes	42,276	100.0	9,481.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,945	28.3	2,679.1
2	Malignant neoplasms (C00-C97)	5,218	12.3	1,170.3
3	Alzheimer's disease (G30)	4,493	10.6	1,007.7
4	Cerebrovascular diseases (I60-I69)	3,471	8.2	778.5
5	Chronic lower respiratory diseases (J40-J47)	1,803	4.3	404.4
6	Diabetes mellitus (E10-E14)	1,454	3.4	326.1
7	Influenza and pneumonia (J09-J18)	1,111	2.6	249.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	928	2.2	208.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	875	2.1	196.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	866	2.0	194.2
11	Parkinson's disease (G20-G21)	676	1.6	151.6
12	Septicemia (A40-A41)	537	1.3	120.4
13	Pneumonitis due to solids and liquids (J69)	366	0.9	82.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	242	0.6	54.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	214	0.5	48.0
...	All other causes (residual)	8,077	19.1	1,811.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	103,532	100.0	356.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,945	20.2	72.2
2	Malignant neoplasms (C00-C97)	20,504	19.8	70.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,510	11.1	39.7
4	Cerebrovascular diseases (I60-I69)	4,798	4.6	16.5
5	Diabetes mellitus (E10-E14)	4,603	4.4	15.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,225	4.1	14.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,944	2.8	10.1
8	Chronic lower respiratory diseases (J40-J47)	2,759	2.7	9.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,657	2.6	9.2
10	Alzheimer's disease (G30)	2,151	2.1	7.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,931	1.9	6.7
12	Influenza and pneumonia (J09-J18)	1,877	1.8	6.5
13	Septicemia (A40-A41)	1,392	1.3	4.8
14	Certain conditions originating in the perinatal period (P00-P96)	1,348	1.3	4.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,197	1.2	4.1
...	All other causes (residual)	18,691	18.1	64.4
Hispanic, male, 1-4 years				
...	All causes	443	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	33.0	7.0
2	Malignant neoplasms (C00-C97)	48	10.8	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	10.2	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	7.4	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.2	*
6	Influenza and pneumonia (J09-J18)	10	2.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.8	*
8	Cerebrovascular diseases (I60-I69)	6	1.4	*
9	Septicemia (A40-A41)	5	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
12	Acute bronchitis and bronchiolitis (J20-J21)	3	0.7	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Hernia (K40-K46)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	108	24.4	5.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-14 years				
...	All causes	641	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	29.5	3.6
2	Malignant neoplasms (C00-C97)	125	19.5	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	6.4	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	4.8	0.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	4.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.4	0.4
7	Cerebrovascular diseases (I60-I69)	18	2.8	*
8	Septicemia (A40-A41)	12	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.7	*
11	Influenza and pneumonia (J09-J18)	8	1.2	*
12	Anemias (D50-D64)	3	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
15	Meningococcal infection (A39)	1	0.2	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (residual)	125	19.5	2.4
Hispanic, male, 15-24 years				
...	All causes	4,304	100.0	87.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,897	44.1	38.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	869	20.2	17.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	665	15.5	13.5
4	Malignant neoplasms (C00-C97)	249	5.8	5.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	2.0	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	1.2	1.1
7	Cerebrovascular diseases (I60-I69)	22	0.5	0.4
7	Influenza and pneumonia (J09-J18)	22	0.5	0.4
9	Legal intervention (Y35,Y89.0)	21	0.5	0.4
10	Diabetes mellitus (E10-E14)	17	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	17	0.4	*
12	Septicemia (A40-A41)	12	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.1	*
15	Pneumonitis due to solids and liquids (J69)	5	0.1	*
...	All other causes (residual)	354	8.2	7.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-34 years				
...	All causes	6,059	100.0	127.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,558	42.2	53.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	829	13.7	17.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	740	12.2	15.6
4	Malignant neoplasms (C00-C97)	344	5.7	7.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	313	5.2	6.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	141	2.3	3.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	71	1.2	1.5
8	Cerebrovascular diseases (I60-I69)	65	1.1	1.4
9	Diabetes mellitus (E10-E14)	58	1.0	1.2
10	Influenza and pneumonia (J09-J18)	50	0.8	1.1
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.7	0.8
12	Septicemia (A40-A41)	38	0.6	0.8
13	Legal intervention (Y35,Y89.0)	36	0.6	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.4	0.5
15	Chronic lower respiratory diseases (J40-J47)	20	0.3	0.4
...	All other causes (residual)	733	12.1	15.4
Hispanic, male, 35-44 years				
...	All causes	6,947	100.0	164.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,071	29.8	49.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	795	11.4	18.8
3	Malignant neoplasms (C00-C97)	716	10.3	17.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	570	8.2	13.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	483	7.0	11.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	455	6.5	10.8
7	Cerebrovascular diseases (I60-I69)	209	3.0	5.0
8	Diabetes mellitus (E10-E14)	175	2.5	4.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	110	1.6	2.6
10	Influenza and pneumonia (J09-J18)	92	1.3	2.2
11	Septicemia (A40-A41)	59	0.8	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	0.7	1.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	0.5	0.9
13	Chronic lower respiratory diseases (J40-J47)	36	0.5	0.9
15	Aortic aneurysm and dissection (I71)	34	0.5	0.8
...	All other causes (residual)	1,055	15.2	25.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-54 years				
...	All causes	11,551	100.0	342.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,162	18.7	64.1
2	Malignant neoplasms (C00-C97)	2,049	17.7	60.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,833	15.9	54.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,085	9.4	32.2
5	Diabetes mellitus (E10-E14)	503	4.4	14.9
6	Cerebrovascular diseases (I60-I69)	447	3.9	13.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	446	3.9	13.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	226	2.0	6.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	213	1.8	6.3
10	Septicemia (A40-A41)	182	1.6	5.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	171	1.5	5.1
12	Viral hepatitis (B15-B19)	154	1.3	4.6
13	Influenza and pneumonia (J09-J18)	150	1.3	4.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	0.9	3.0
15	Chronic lower respiratory diseases (J40-J47)	91	0.8	2.7
...	All other causes (residual)	1,738	15.0	51.5
Hispanic, male, 55-64 years				
...	All causes	17,416	100.0	804.1
1	Malignant neoplasms (C00-C97)	4,475	25.7	206.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,820	21.9	176.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,363	7.8	62.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,279	7.3	59.1
5	Diabetes mellitus (E10-E14)	1,085	6.2	50.1
6	Cerebrovascular diseases (I60-I69)	741	4.3	34.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	357	2.0	16.5
8	Chronic lower respiratory diseases (J40-J47)	296	1.7	13.7
9	Viral hepatitis (B15-B19)	282	1.6	13.0
10	Influenza and pneumonia (J09-J18)	281	1.6	13.0
11	Septicemia (A40-A41)	261	1.5	12.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	256	1.5	11.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	197	1.1	9.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	171	1.0	7.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	112	0.6	5.2
...	All other causes (residual)	2,440	14.0	112.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-74 years				
...	All causes	18,211	100.0	1,668.9
1	Malignant neoplasms (C00-C97)	5,224	28.7	478.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,336	23.8	397.4
3	Diabetes mellitus (E10-E14)	1,175	6.5	107.7
4	Cerebrovascular diseases (I60-I69)	946	5.2	86.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	723	4.0	66.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	624	3.4	57.2
7	Chronic lower respiratory diseases (J40-J47)	575	3.2	52.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	450	2.5	41.2
9	Influenza and pneumonia (J09-J18)	360	2.0	33.0
10	Septicemia (A40-A41)	294	1.6	26.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	254	1.4	23.3
12	Alzheimer's disease (G30)	192	1.1	17.6
13	Viral hepatitis (B15-B19)	141	0.8	12.9
14	Parkinson's disease (G20-G21)	129	0.7	11.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	121	0.7	11.1
...	All other causes (residual)	2,667	14.6	244.4
Hispanic, male, 75-84 years				
...	All causes	18,848	100.0	4,054.0
1	Malignant neoplasms (C00-C97)	4,725	25.1	1,016.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,634	24.6	996.7
3	Cerebrovascular diseases (I60-I69)	1,195	6.3	257.0
4	Diabetes mellitus (E10-E14)	988	5.2	212.5
5	Chronic lower respiratory diseases (J40-J47)	881	4.7	189.5
6	Alzheimer's disease (G30)	686	3.6	147.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	484	2.6	104.1
8	Influenza and pneumonia (J09-J18)	440	2.3	94.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	416	2.2	89.5
10	Parkinson's disease (G20-G21)	401	2.1	86.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	353	1.9	75.9
12	Septicemia (A40-A41)	281	1.5	60.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	257	1.4	55.3
14	Pneumonitis due to solids and liquids (J69)	157	0.8	33.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	128	0.7	27.5
...	All other causes (residual)	2,822	15.0	607.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85 years and over				
...	All causes	16,425	100.0	10,139.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,727	28.8	2,918.0
2	Malignant neoplasms (C00-C97)	2,540	15.5	1,568.0
3	Alzheimer's disease (G30)	1,232	7.5	760.5
4	Cerebrovascular diseases (I60-I69)	1,135	6.9	700.6
5	Chronic lower respiratory diseases (J40-J47)	826	5.0	509.9
6	Diabetes mellitus (E10-E14)	600	3.7	370.4
7	Influenza and pneumonia (J09-J18)	442	2.7	272.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	397	2.4	245.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	382	2.3	235.8
10	Parkinson's disease (G20-G21)	348	2.1	214.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	336	2.0	207.4
12	Septicemia (A40-A41)	221	1.3	136.4
13	Pneumonitis due to solids and liquids (J69)	171	1.0	105.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	105	0.6	64.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	98	0.6	60.5
...	All other causes (residual)	2,865	17.4	1,768.6
Hispanic, female, all ages				
...	All causes	84,722	100.0	297.7
1	Malignant neoplasms (C00-C97)	18,759	22.1	65.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,854	19.9	59.2
3	Cerebrovascular diseases (I60-I69)	5,485	6.5	19.3
4	Alzheimer's disease (G30)	4,682	5.5	16.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,201	5.0	14.8
6	Diabetes mellitus (E10-E14)	3,943	4.7	13.9
7	Chronic lower respiratory diseases (J40-J47)	2,528	3.0	8.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,916	2.3	6.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,844	2.2	6.5
10	Influenza and pneumonia (J09-J18)	1,764	2.1	6.2
11	Septicemia (A40-A41)	1,326	1.6	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,291	1.5	4.5
13	Certain conditions originating in the perinatal period (P00-P96)	1,067	1.3	3.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	858	1.0	3.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	724	0.9	2.5
...	All other causes (residual)	17,480	20.6	61.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 1-4 years				
...	All causes	356	100.0	17.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	24.4	4.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	15.2	2.7
3	Malignant neoplasms (C00-C97)	41	11.5	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	8.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.7	*
6	Septicemia (A40-A41)	9	2.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	2.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	2.0	*
9	Influenza and pneumonia (J09-J18)	6	1.7	*
10	Anemias (D50-D64)	3	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
12	Cerebrovascular diseases (I60-I69)	2	0.6	*
13	Salmonella infections (A01-A02)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	87	24.4	4.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	523	100.0	10.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	141	27.0	2.8
2	Malignant neoplasms (C00-C97)	101	19.3	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	8.8	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	5.5	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	4.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.1	*
7	Cerebrovascular diseases (I60-I69)	11	2.1	*
7	Chronic lower respiratory diseases (J40-J47)	11	2.1	*
9	Septicemia (A40-A41)	6	1.1	*
9	Influenza and pneumonia (J09-J18)	6	1.1	*
11	Anemias (D50-D64)	4	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Nutritional deficiencies (E40-E64)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
13	Peptic ulcer (K25-K28)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	116	22.2	2.3
Hispanic, female, 15-24 years				
...	All causes	1,418	100.0	30.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	630	44.4	13.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	176	12.4	3.8
3	Malignant neoplasms (C00-C97)	121	8.5	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	118	8.3	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	2.3	0.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	2.3	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	29	2.0	0.6
8	Chronic lower respiratory diseases (J40-J47)	13	0.9	*
9	Cerebrovascular diseases (I60-I69)	12	0.8	*
9	Influenza and pneumonia (J09-J18)	12	0.8	*
11	Septicemia (A40-A41)	10	0.7	*
12	Diabetes mellitus (E10-E14)	8	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.3	*
...	All other causes (residual)	204	14.4	4.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	2,136	100.0	49.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	654	30.6	15.1
2	Malignant neoplasms (C00-C97)	362	16.9	8.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	161	7.5	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	6.3	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	124	5.8	2.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	90	4.2	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	53	2.5	1.2
8	Cerebrovascular diseases (I60-I69)	43	2.0	1.0
9	Diabetes mellitus (E10-E14)	41	1.9	0.9
10	Septicemia (A40-A41)	31	1.5	0.7
11	Influenza and pneumonia (J09-J18)	24	1.1	0.6
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	1.0	0.5
13	Chronic lower respiratory diseases (J40-J47)	16	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.5	*
...	All other causes (residual)	356	16.7	8.2
Hispanic, female, 35-44 years				
...	All causes	3,325	100.0	82.6
1	Malignant neoplasms (C00-C97)	1,077	32.4	26.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	15.7	12.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	272	8.2	6.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	167	5.0	4.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	133	4.0	3.3
6	Cerebrovascular diseases (I60-I69)	117	3.5	2.9
7	Diabetes mellitus (E10-E14)	93	2.8	2.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	2.8	2.3
9	Pregnancy, childbirth and the puerperium (O00-O99)	53	1.6	1.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	1.5	1.2
11	Septicemia (A40-A41)	43	1.3	1.1
11	Influenza and pneumonia (J09-J18)	43	1.3	1.1
13	Chronic lower respiratory diseases (J40-J47)	32	1.0	0.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.9	0.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25	0.8	0.6
...	All other causes (residual)	576	17.3	14.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	6,202	100.0	187.2
1	Malignant neoplasms (C00-C97)	2,232	36.0	67.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	718	11.6	21.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	538	8.7	16.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	343	5.5	10.4
5	Diabetes mellitus (E10-E14)	312	5.0	9.4
6	Cerebrovascular diseases (I60-I69)	279	4.5	8.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	104	1.7	3.1
8	Septicemia (A40-A41)	102	1.6	3.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	102	1.6	3.1
10	Chronic lower respiratory diseases (J40-J47)	96	1.5	2.9
11	Influenza and pneumonia (J09-J18)	92	1.5	2.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	1.1	2.0
13	Viral hepatitis (B15-B19)	64	1.0	1.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	1.0	1.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	45	0.7	1.4
...	All other causes (residual)	1,048	16.9	31.6
Hispanic, female, 55-64 years				
...	All causes	10,171	100.0	441.5
1	Malignant neoplasms (C00-C97)	3,743	36.8	162.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,522	15.0	66.1
3	Diabetes mellitus (E10-E14)	698	6.9	30.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	485	4.8	21.1
5	Cerebrovascular diseases (I60-I69)	459	4.5	19.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	427	4.2	18.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	247	2.4	10.7
8	Chronic lower respiratory diseases (J40-J47)	236	2.3	10.2
9	Septicemia (A40-A41)	159	1.6	6.9
10	Influenza and pneumonia (J09-J18)	158	1.6	6.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	120	1.2	5.2
12	Viral hepatitis (B15-B19)	106	1.0	4.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	81	0.8	3.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.5	2.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	44	0.4	1.9
...	All other causes (residual)	1,635	16.1	71.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	13,339	100.0	1,018.2
1	Malignant neoplasms (C00-C97)	4,186	31.4	319.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,495	18.7	190.5
3	Diabetes mellitus (E10-E14)	894	6.7	68.2
4	Cerebrovascular diseases (I60-I69)	759	5.7	57.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	422	3.2	32.2
6	Chronic lower respiratory diseases (J40-J47)	414	3.1	31.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	392	2.9	29.9
8	Influenza and pneumonia (J09-J18)	283	2.1	21.6
9	Septicemia (A40-A41)	275	2.1	21.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	266	2.0	20.3
11	Alzheimer's disease (G30)	224	1.7	17.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	168	1.3	12.8
13	Parkinson's disease (G20-G21)	92	0.7	7.0
14	Viral hepatitis (B15-B19)	83	0.6	6.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.5	5.1
...	All other causes (residual)	2,319	17.4	177.0
Hispanic, female, 75-84 years				
...	All causes	19,258	100.0	2,932.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,412	22.9	671.8
2	Malignant neoplasms (C00-C97)	4,207	21.8	640.6
3	Cerebrovascular diseases (I60-I69)	1,456	7.6	221.7
4	Alzheimer's disease (G30)	1,145	5.9	174.3
5	Diabetes mellitus (E10-E14)	1,043	5.4	158.8
6	Chronic lower respiratory diseases (J40-J47)	729	3.8	111.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	500	2.6	76.1
8	Influenza and pneumonia (J09-J18)	459	2.4	69.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	394	2.0	60.0
10	Septicemia (A40-A41)	357	1.9	54.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	356	1.8	54.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	313	1.6	47.7
13	Parkinson's disease (G20-G21)	277	1.4	42.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	126	0.7	19.2
15	Pneumonitis due to solids and liquids (J69)	120	0.6	18.3
...	All other causes (residual)	3,364	17.5	512.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	25,851	100.0	9,106.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,218	27.9	2,542.8
2	Alzheimer's disease (G30)	3,261	12.6	1,148.8
3	Malignant neoplasms (C00-C97)	2,678	10.4	943.4
4	Cerebrovascular diseases (I60-I69)	2,336	9.0	822.9
5	Chronic lower respiratory diseases (J40-J47)	977	3.8	344.2
6	Diabetes mellitus (E10-E14)	854	3.3	300.8
7	Influenza and pneumonia (J09-J18)	669	2.6	235.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	592	2.3	208.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	493	1.9	173.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	469	1.8	165.2
11	Parkinson's disease (G20-G21)	328	1.3	115.5
12	Septicemia (A40-A41)	316	1.2	111.3
13	Pneumonitis due to solids and liquids (J69)	195	0.8	68.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	137	0.5	48.3
15	Nutritional deficiencies (E40-E64)	125	0.5	44.0
...	All other causes (residual)	5,203	20.1	1,832.9
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,133,463	100.0	1,059.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	500,631	23.5	248.7
2	Malignant neoplasms (C00-C97)	466,467	21.9	231.7
3	Chronic lower respiratory diseases (J40-J47)	135,268	6.3	67.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121,641	5.7	60.4
5	Cerebrovascular diseases (I60-I69)	107,491	5.0	53.4
6	Alzheimer's disease (G30)	97,779	4.6	48.6
7	Diabetes mellitus (E10-E14)	53,399	2.5	26.5
8	Influenza and pneumonia (J09-J18)	39,748	1.9	19.7
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,531	1.7	18.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,248	1.7	17.5
11	Septicemia (A40-A41)	30,009	1.4	14.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29,432	1.4	14.6
13	Parkinson's disease (G20-G21)	25,826	1.2	12.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23,237	1.1	11.5
15	Pneumonitis due to solids and liquids (J69)	16,302	0.8	8.1
...	All other causes (residual)	414,454	19.4	205.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,978	100.0	23.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	680	34.4	8.2
2	Malignant neoplasms (C00-C97)	206	10.4	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	9.7	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	6.1	1.5
5	Influenza and pneumonia (J09-J18)	47	2.4	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	2.3	0.5
7	Septicemia (A40-A41)	34	1.7	0.4
8	Cerebrovascular diseases (I60-I69)	27	1.4	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.2	0.3
10	Chronic lower respiratory diseases (J40-J47)	22	1.1	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	20	1.0	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
15	Meningitis (G00,G03)	5	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (residual)	532	26.9	6.4
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,781	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	844	30.3	3.9
2	Malignant neoplasms (C00-C97)	440	15.8	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	311	11.2	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	6.8	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	3.5	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	2.9	0.4
7	Influenza and pneumonia (J09-J18)	44	1.6	0.2
8	Septicemia (A40-A41)	34	1.2	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	1.2	0.2
10	Chronic lower respiratory diseases (J40-J47)	32	1.2	0.1
11	Cerebrovascular diseases (I60-I69)	31	1.1	0.1
12	Diabetes mellitus (E10-E14)	15	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	12	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (residual)	597	21.5	2.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	17,515	100.0	72.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,895	50.8	36.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,756	21.4	15.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	766	4.4	3.2
4	Malignant neoplasms (C00-C97)	744	4.2	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	481	2.7	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	210	1.2	0.9
7	Diabetes mellitus (E10-E14)	113	0.6	0.5
8	Influenza and pneumonia (J09-J18)	100	0.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	87	0.5	0.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	79	0.5	0.3
11	Septicemia (A40-A41)	74	0.4	0.3
12	Cerebrovascular diseases (I60-I69)	71	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.2	0.2
14	Pneumonitis due to solids and liquids (J69)	26	0.1	0.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	24	0.1	0.1
...	All other causes (residual)	2,051	11.7	8.5
Non-Hispanic, white, both sexes, 25-34 years				
...	All causes	35,222	100.0	137.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,170	48.7	67.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,280	15.0	20.6
3	Malignant neoplasms (C00-C97)	2,207	6.3	8.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,811	5.1	7.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,140	3.2	4.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	551	1.6	2.2
7	Diabetes mellitus (E10-E14)	404	1.1	1.6
8	Cerebrovascular diseases (I60-I69)	283	0.8	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	259	0.7	1.0
10	Influenza and pneumonia (J09-J18)	247	0.7	1.0
11	Septicemia (A40-A41)	233	0.7	0.9
12	Pregnancy, childbirth and the puerperium (O00-O99)	201	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	167	0.5	0.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	111	0.3	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	108	0.3	0.4
...	All other causes (residual)	5,050	14.3	19.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-44 years				
...	All causes	48,409	100.0	205.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,057	31.1	64.0
2	Malignant neoplasms (C00-C97)	6,414	13.2	27.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,991	12.4	25.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,528	11.4	23.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,791	3.7	7.6
6	Diabetes mellitus (E10-E14)	1,080	2.2	4.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,025	2.1	4.4
8	Cerebrovascular diseases (I60-I69)	869	1.8	3.7
9	Influenza and pneumonia (J09-J18)	532	1.1	2.3
10	Septicemia (A40-A41)	515	1.1	2.2
11	Chronic lower respiratory diseases (J40-J47)	429	0.9	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	311	0.6	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	299	0.6	1.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	259	0.5	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	200	0.4	0.9
...	All other causes (residual)	8,109	16.8	34.5
Non-Hispanic, white, both sexes, 45-54 years				
...	All causes	116,634	100.0	420.0
1	Malignant neoplasms (C00-C97)	28,037	24.0	101.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,218	19.0	80.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,973	14.6	61.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,251	6.2	26.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,849	5.0	21.1
6	Diabetes mellitus (E10-E14)	3,628	3.1	13.1
7	Chronic lower respiratory diseases (J40-J47)	3,448	3.0	12.4
8	Cerebrovascular diseases (I60-I69)	2,828	2.4	10.2
9	Septicemia (A40-A41)	1,505	1.3	5.4
10	Influenza and pneumonia (J09-J18)	1,354	1.2	4.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,053	0.9	3.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	967	0.8	3.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	806	0.7	2.9
14	Viral hepatitis (B15-B19)	775	0.7	2.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	650	0.6	2.3
...	All other causes (residual)	19,292	16.5	69.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-64 years				
...	All causes	262,230	100.0	880.4
1	Malignant neoplasms (C00-C97)	85,111	32.5	285.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,029	21.0	184.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,085	6.1	54.0
4	Chronic lower respiratory diseases (J40-J47)	14,831	5.7	49.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	9,800	3.7	32.9
6	Diabetes mellitus (E10-E14)	8,699	3.3	29.2
7	Cerebrovascular diseases (I60-I69)	7,501	2.9	25.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,901	2.6	23.2
9	Septicemia (A40-A41)	3,967	1.5	13.3
10	Influenza and pneumonia (J09-J18)	3,494	1.3	11.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,176	1.2	10.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,250	0.9	7.6
13	Viral hepatitis (B15-B19)	1,978	0.8	6.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,175	0.4	3.9
15	Pneumonitis due to solids and liquids (J69)	1,138	0.4	3.8
...	All other causes (residual)	41,095	15.7	138.0
Non-Hispanic, white, both sexes, 65-74 years				
...	All causes	395,154	100.0	1,800.3
1	Malignant neoplasms (C00-C97)	130,650	33.1	595.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	84,617	21.4	385.5
3	Chronic lower respiratory diseases (J40-J47)	33,694	8.5	153.5
4	Cerebrovascular diseases (I60-I69)	15,129	3.8	68.9
5	Diabetes mellitus (E10-E14)	13,361	3.4	60.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,991	2.8	50.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,749	1.7	30.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,438	1.6	29.3
9	Septicemia (A40-A41)	6,330	1.6	28.8
10	Influenza and pneumonia (J09-J18)	6,006	1.5	27.4
11	Alzheimer's disease (G30)	5,524	1.4	25.2
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,007	1.0	18.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,523	0.9	16.1
14	Parkinson's disease (G20-G21)	3,461	0.9	15.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,555	0.6	11.6
...	All other causes (residual)	62,119	15.7	283.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-84 years				
...	All causes	514,552	100.0	4,606.0
1	Malignant neoplasms (C00-C97)	125,254	24.3	1,121.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	117,680	22.9	1,053.4
3	Chronic lower respiratory diseases (J40-J47)	44,024	8.6	394.1
4	Cerebrovascular diseases (I60-I69)	28,989	5.6	259.5
5	Alzheimer's disease (G30)	25,263	4.9	226.1
6	Diabetes mellitus (E10-E14)	13,600	2.6	121.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,365	2.6	119.6
8	Parkinson's disease (G20-G21)	10,515	2.0	94.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,132	2.0	90.7
10	Influenza and pneumonia (J09-J18)	9,820	1.9	87.9
11	Septicemia (A40-A41)	8,058	1.6	72.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,320	1.0	47.6
13	Pneumonitis due to solids and liquids (J69)	4,203	0.8	37.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,968	0.8	35.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,445	0.7	30.8
...	All other causes (residual)	90,916	17.7	813.8
Non-Hispanic, white, both sexes, 85 years and over				
...	All causes	728,886	100.0	14,062.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	212,527	29.2	4,100.3
2	Malignant neoplasms (C00-C97)	87,371	12.0	1,685.7
3	Alzheimer's disease (G30)	66,012	9.1	1,273.6
4	Cerebrovascular diseases (I60-I69)	51,703	7.1	997.5
5	Chronic lower respiratory diseases (J40-J47)	38,517	5.3	743.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,979	2.9	404.8
7	Influenza and pneumonia (J09-J18)	18,043	2.5	348.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,964	1.9	269.4
9	Diabetes mellitus (E10-E14)	12,496	1.7	241.1
10	Parkinson's disease (G20-G21)	11,231	1.5	216.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,079	1.5	213.7
12	Septicemia (A40-A41)	9,190	1.3	177.3
13	Pneumonitis due to solids and liquids (J69)	8,131	1.1	156.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,535	0.6	87.5
15	Nutritional deficiencies (E40-E64)	2,816	0.4	54.3
...	All other causes (residual)	160,292	22.0	3,092.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	1,077,362	100.0	1,085.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	266,995	24.8	269.0
2	Malignant neoplasms (C00-C97)	247,202	22.9	249.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76,033	7.1	76.6
4	Chronic lower respiratory diseases (J40-J47)	62,945	5.8	63.4
5	Cerebrovascular diseases (I60-I69)	43,713	4.1	44.0
6	Diabetes mellitus (E10-E14)	30,010	2.8	30.2
7	Alzheimer's disease (G30)	29,886	2.8	30.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,110	2.6	28.3
9	Influenza and pneumonia (J09-J18)	19,033	1.8	19.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18,522	1.7	18.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,216	1.7	18.4
12	Parkinson's disease (G20-G21)	15,641	1.5	15.8
13	Septicemia (A40-A41)	14,620	1.4	14.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,906	0.9	10.0
15	Pneumonitis due to solids and liquids (J69)	8,997	0.8	9.1
...	All other causes (residual)	187,533	17.4	189.0
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,113	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	419	37.6	9.9
2	Malignant neoplasms (C00-C97)	114	10.2	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	8.5	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	5.6	1.5
5	Septicemia (A40-A41)	23	2.1	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.1	0.5
7	Influenza and pneumonia (J09-J18)	21	1.9	0.5
8	Chronic lower respiratory diseases (J40-J47)	14	1.3	*
9	Cerebrovascular diseases (I60-I69)	12	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
14	Meningococcal infection (A39)	2	0.2	*
14	Anemias (D50-D64)	2	0.2	*
14	Meningitis (G00,G03)	2	0.2	*
14	Pneumonitis due to solids and liquids (J69)	2	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.2	*
...	All other causes (residual)	288	25.9	6.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,625	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	523	32.2	4.7
2	Malignant neoplasms (C00-C97)	264	16.2	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	194	11.9	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	6.6	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	3.7	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	2.6	0.4
7	Influenza and pneumonia (J09-J18)	24	1.5	0.2
8	Cerebrovascular diseases (I60-I69)	21	1.3	0.2
9	Septicemia (A40-A41)	19	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
14	Anemias (D50-D64)	5	0.3	*
15	Diabetes mellitus (E10-E14)	4	0.2	*
...	All other causes (residual)	314	19.3	2.8
Non-Hispanic, white, male, 15-24 years				
...	All causes	12,645	100.0	101.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,557	51.9	52.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,042	24.1	24.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	549	4.3	4.4
4	Malignant neoplasms (C00-C97)	464	3.7	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	316	2.5	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	0.9	1.0
7	Diabetes mellitus (E10-E14)	65	0.5	0.5
8	Influenza and pneumonia (J09-J18)	57	0.5	0.5
9	Chronic lower respiratory diseases (J40-J47)	53	0.4	0.4
10	Cerebrovascular diseases (I60-I69)	48	0.4	0.4
11	Septicemia (A40-A41)	43	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.2	0.2
13	Legal intervention (Y35,Y89.0)	20	0.2	0.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	14	0.1	*
15	Pneumonitis due to solids and liquids (J69)	13	0.1	*
...	All other causes (residual)	1,261	10.0	10.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-34 years				
...	All causes	24,293	100.0	187.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,533	51.6	96.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,199	17.3	32.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,227	5.1	9.5
4	Malignant neoplasms (C00-C97)	1,093	4.5	8.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	812	3.3	6.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	319	1.3	2.5
7	Diabetes mellitus (E10-E14)	235	1.0	1.8
8	Cerebrovascular diseases (I60-I69)	156	0.6	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	135	0.6	1.0
10	Influenza and pneumonia (J09-J18)	129	0.5	1.0
11	Septicemia (A40-A41)	108	0.4	0.8
12	Chronic lower respiratory diseases (J40-J47)	82	0.3	0.6
13	Legal intervention (Y35,Y89.0)	80	0.3	0.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	79	0.3	0.6
15	Aortic aneurysm and dissection (I71)	58	0.2	0.4
...	All other causes (residual)	3,048	12.5	23.5
Non-Hispanic, white, male, 35-44 years				
...	All causes	30,522	100.0	257.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,533	34.5	89.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,110	13.5	34.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,098	13.4	34.6
4	Malignant neoplasms (C00-C97)	2,870	9.4	24.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,059	3.5	8.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	694	2.3	5.9
7	Diabetes mellitus (E10-E14)	666	2.2	5.6
8	Cerebrovascular diseases (I60-I69)	485	1.6	4.1
9	Influenza and pneumonia (J09-J18)	280	0.9	2.4
10	Septicemia (A40-A41)	260	0.9	2.2
11	Chronic lower respiratory diseases (J40-J47)	221	0.7	1.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	201	0.7	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	184	0.6	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	135	0.4	1.1
15	Aortic aneurysm and dissection (I71)	131	0.4	1.1
...	All other causes (residual)	4,595	15.1	38.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-54 years				
...	All causes	70,971	100.0	513.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,750	22.2	113.9
2	Malignant neoplasms (C00-C97)	13,911	19.6	100.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,284	15.9	81.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,275	7.4	38.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,681	5.2	26.6
6	Diabetes mellitus (E10-E14)	2,400	3.4	17.4
7	Cerebrovascular diseases (I60-I69)	1,566	2.2	11.3
8	Chronic lower respiratory diseases (J40-J47)	1,540	2.2	11.1
9	Septicemia (A40-A41)	805	1.1	5.8
10	Influenza and pneumonia (J09-J18)	702	1.0	5.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	670	0.9	4.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	634	0.9	4.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	550	0.8	4.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	543	0.8	3.9
15	Viral hepatitis (B15-B19)	509	0.7	3.7
...	All other causes (residual)	11,151	15.7	80.6
Non-Hispanic, white, male, 55-64 years				
...	All causes	160,050	100.0	1,100.6
1	Malignant neoplasms (C00-C97)	46,980	29.4	323.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,770	24.2	266.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,948	6.8	75.3
4	Chronic lower respiratory diseases (J40-J47)	7,515	4.7	51.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,622	4.1	45.5
6	Diabetes mellitus (E10-E14)	5,539	3.5	38.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,088	3.2	35.0
8	Cerebrovascular diseases (I60-I69)	4,367	2.7	30.0
9	Septicemia (A40-A41)	2,185	1.4	15.0
10	Influenza and pneumonia (J09-J18)	1,926	1.2	13.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,848	1.2	12.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,459	0.9	10.0
13	Viral hepatitis (B15-B19)	1,381	0.9	9.5
14	Aortic aneurysm and dissection (I71)	777	0.5	5.3
15	Pneumonitis due to solids and liquids (J69)	697	0.4	4.8
...	All other causes (residual)	23,948	15.0	164.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-74 years				
...	All causes	227,014	100.0	2,175.8
1	Malignant neoplasms (C00-C97)	72,516	31.9	695.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,151	24.3	528.6
3	Chronic lower respiratory diseases (J40-J47)	17,047	7.5	163.4
4	Diabetes mellitus (E10-E14)	8,032	3.5	77.0
5	Cerebrovascular diseases (I60-I69)	8,025	3.5	76.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,056	3.1	67.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,317	1.9	41.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,523	1.6	33.8
9	Influenza and pneumonia (J09-J18)	3,338	1.5	32.0
10	Septicemia (A40-A41)	3,330	1.5	31.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,156	1.4	30.2
12	Parkinson's disease (G20-G21)	2,327	1.0	22.3
13	Alzheimer's disease (G30)	2,286	1.0	21.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,953	0.9	18.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,510	0.7	14.5
...	All other causes (residual)	33,447	14.7	320.6
Non-Hispanic, white, male, 75-84 years				
...	All causes	265,420	100.0	5,382.6
1	Malignant neoplasms (C00-C97)	67,030	25.3	1,359.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	66,153	24.9	1,341.5
3	Chronic lower respiratory diseases (J40-J47)	20,760	7.8	421.0
4	Cerebrovascular diseases (I60-I69)	13,231	5.0	268.3
5	Alzheimer's disease (G30)	9,654	3.6	195.8
6	Diabetes mellitus (E10-E14)	7,531	2.8	152.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,307	2.8	148.2
8	Parkinson's disease (G20-G21)	6,785	2.6	137.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,514	2.1	111.8
10	Influenza and pneumonia (J09-J18)	5,110	1.9	103.6
11	Septicemia (A40-A41)	3,960	1.5	80.3
12	Pneumonitis due to solids and liquids (J69)	2,443	0.9	49.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,390	0.9	48.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,283	0.9	46.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,049	0.8	41.6
...	All other causes (residual)	43,220	16.3	876.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85 years and over				
...	All causes	278,054	100.0	15,274.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	85,388	30.7	4,690.7
2	Malignant neoplasms (C00-C97)	41,944	15.1	2,304.1
3	Alzheimer's disease (G30)	17,539	6.3	963.5
4	Cerebrovascular diseases (I60-I69)	15,764	5.7	866.0
5	Chronic lower respiratory diseases (J40-J47)	15,683	5.6	861.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,513	3.1	467.7
7	Influenza and pneumonia (J09-J18)	7,402	2.7	406.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,429	2.3	353.2
9	Parkinson's disease (G20-G21)	6,128	2.2	336.6
10	Diabetes mellitus (E10-E14)	5,537	2.0	304.2
11	Pneumonitis due to solids and liquids (J69)	4,175	1.5	229.3
12	Septicemia (A40-A41)	3,846	1.4	211.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,408	1.2	187.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,102	0.8	115.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	997	0.4	54.8
...	All other causes (residual)	53,199	19.1	2,922.4
Non-Hispanic, white, female, all ages				
...	All causes	1,056,101	100.0	1,034.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	233,636	22.1	228.9
2	Malignant neoplasms (C00-C97)	219,265	20.8	214.8
3	Chronic lower respiratory diseases (J40-J47)	72,323	6.8	70.8
4	Alzheimer's disease (G30)	67,893	6.4	66.5
5	Cerebrovascular diseases (I60-I69)	63,778	6.0	62.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45,608	4.3	44.7
7	Diabetes mellitus (E10-E14)	23,389	2.2	22.9
8	Influenza and pneumonia (J09-J18)	20,715	2.0	20.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,032	1.6	16.7
10	Septicemia (A40-A41)	15,389	1.5	15.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,331	1.3	13.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,910	1.0	10.7
13	Parkinson's disease (G20-G21)	10,185	1.0	10.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,421	0.8	8.2
15	Pneumonitis due to solids and liquids (J69)	7,305	0.7	7.2
...	All other causes (residual)	226,921	21.5	222.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	865	100.0	21.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	261	30.2	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	11.1	2.4
3	Malignant neoplasms (C00-C97)	92	10.6	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	6.8	1.5
5	Influenza and pneumonia (J09-J18)	26	3.0	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	2.5	0.5
7	Cerebrovascular diseases (I60-I69)	15	1.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.6	*
9	Septicemia (A40-A41)	11	1.3	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	8	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
14	Meningitis (G00,G03)	3	0.3	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (residual)	238	27.5	5.9
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,156	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	27.8	3.0
2	Malignant neoplasms (C00-C97)	176	15.2	1.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	117	10.1	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	7.0	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	3.5	0.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	3.2	0.3
7	Influenza and pneumonia (J09-J18)	20	1.7	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
9	Septicemia (A40-A41)	15	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
11	Diabetes mellitus (E10-E14)	11	1.0	*
12	Cerebrovascular diseases (I60-I69)	10	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (residual)	281	24.3	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,870	100.0	41.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,338	48.0	19.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	714	14.7	6.1
3	Malignant neoplasms (C00-C97)	280	5.7	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	217	4.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	165	3.4	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.9	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	79	1.6	0.7
8	Diabetes mellitus (E10-E14)	48	1.0	0.4
9	Influenza and pneumonia (J09-J18)	43	0.9	0.4
10	Chronic lower respiratory diseases (J40-J47)	34	0.7	0.3
11	Septicemia (A40-A41)	31	0.6	0.3
12	Cerebrovascular diseases (I60-I69)	23	0.5	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.3	*
15	Pneumonitis due to solids and liquids (J69)	13	0.3	*
...	All other causes (residual)	766	15.7	6.5
Non-Hispanic, white, female, 25-34 years				
...	All causes	10,929	100.0	86.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,637	42.4	36.7
2	Malignant neoplasms (C00-C97)	1,114	10.2	8.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,081	9.9	8.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	584	5.3	4.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	328	3.0	2.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	232	2.1	1.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	201	1.8	1.6
8	Diabetes mellitus (E10-E14)	169	1.5	1.3
9	Cerebrovascular diseases (I60-I69)	127	1.2	1.0
10	Septicemia (A40-A41)	125	1.1	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	1.1	1.0
12	Influenza and pneumonia (J09-J18)	118	1.1	0.9
13	Chronic lower respiratory diseases (J40-J47)	85	0.8	0.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	0.5	0.4
15	Pneumonitis due to solids and liquids (J69)	37	0.3	0.3
...	All other causes (residual)	1,914	17.5	15.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-44 years				
...	All causes	17,887	100.0	152.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,524	25.3	38.7
2	Malignant neoplasms (C00-C97)	3,544	19.8	30.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,893	10.6	16.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,418	7.9	12.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	732	4.1	6.3
6	Diabetes mellitus (E10-E14)	414	2.3	3.5
7	Cerebrovascular diseases (I60-I69)	384	2.1	3.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	331	1.9	2.8
9	Septicemia (A40-A41)	255	1.4	2.2
10	Influenza and pneumonia (J09-J18)	252	1.4	2.2
11	Chronic lower respiratory diseases (J40-J47)	208	1.2	1.8
12	Pregnancy, childbirth and the puerperium (O00-O99)	168	0.9	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	164	0.9	1.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	0.7	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	83	0.5	0.7
...	All other causes (residual)	3,390	19.0	29.0
Non-Hispanic, white, female, 45-54 years				
...	All causes	45,663	100.0	327.5
1	Malignant neoplasms (C00-C97)	14,126	30.9	101.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,468	14.2	46.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,689	12.5	40.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,168	4.7	15.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,976	4.3	14.2
6	Chronic lower respiratory diseases (J40-J47)	1,908	4.2	13.7
7	Cerebrovascular diseases (I60-I69)	1,262	2.8	9.1
8	Diabetes mellitus (E10-E14)	1,228	2.7	8.8
9	Septicemia (A40-A41)	700	1.5	5.0
10	Influenza and pneumonia (J09-J18)	652	1.4	4.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	419	0.9	3.0
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	297	0.7	2.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	270	0.6	1.9
14	Viral hepatitis (B15-B19)	266	0.6	1.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	263	0.6	1.9
...	All other causes (residual)	7,971	17.5	57.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-64 years				
...	All causes	102,180	100.0	670.2
1	Malignant neoplasms (C00-C97)	38,131	37.3	250.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,259	15.9	106.6
3	Chronic lower respiratory diseases (J40-J47)	7,316	7.2	48.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,137	5.0	33.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,178	3.1	20.8
6	Diabetes mellitus (E10-E14)	3,160	3.1	20.7
7	Cerebrovascular diseases (I60-I69)	3,134	3.1	20.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,813	1.8	11.9
9	Septicemia (A40-A41)	1,782	1.7	11.7
10	Influenza and pneumonia (J09-J18)	1,568	1.5	10.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,328	1.3	8.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	791	0.8	5.2
13	Viral hepatitis (B15-B19)	597	0.6	3.9
14	Alzheimer's disease (G30)	530	0.5	3.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	514	0.5	3.4
...	All other causes (residual)	16,942	16.6	111.1
Non-Hispanic, white, female, 65-74 years				
...	All causes	168,140	100.0	1,460.2
1	Malignant neoplasms (C00-C97)	58,134	34.6	504.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,466	17.5	255.9
3	Chronic lower respiratory diseases (J40-J47)	16,647	9.9	144.6
4	Cerebrovascular diseases (I60-I69)	7,104	4.2	61.7
5	Diabetes mellitus (E10-E14)	5,329	3.2	46.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,935	2.3	34.2
7	Alzheimer's disease (G30)	3,238	1.9	28.1
8	Septicemia (A40-A41)	3,000	1.8	26.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,915	1.7	25.3
10	Influenza and pneumonia (J09-J18)	2,668	1.6	23.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,432	1.4	21.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,570	0.9	13.6
13	Parkinson's disease (G20-G21)	1,134	0.7	9.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,045	0.6	9.1
15	Pneumonitis due to solids and liquids (J69)	864	0.5	7.5
...	All other causes (residual)	28,659	17.0	248.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-84 years				
...	All causes	249,132	100.0	3,992.4
1	Malignant neoplasms (C00-C97)	58,224	23.4	933.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,527	20.7	825.7
3	Chronic lower respiratory diseases (J40-J47)	23,264	9.3	372.8
4	Cerebrovascular diseases (I60-I69)	15,758	6.3	252.5
5	Alzheimer's disease (G30)	15,609	6.3	250.1
6	Diabetes mellitus (E10-E14)	6,069	2.4	97.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,058	2.4	97.1
8	Influenza and pneumonia (J09-J18)	4,710	1.9	75.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,618	1.9	74.0
10	Septicemia (A40-A41)	4,098	1.6	65.7
11	Parkinson's disease (G20-G21)	3,730	1.5	59.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,930	1.2	47.0
13	Pneumonitis due to solids and liquids (J69)	1,760	0.7	28.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,685	0.7	27.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,537	0.6	24.6
...	All other causes (residual)	47,555	19.1	762.1
Non-Hispanic, white, female, 85 years and over				
...	All causes	450,832	100.0	13,406.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	127,139	28.2	3,780.8
2	Alzheimer's disease (G30)	48,473	10.8	1,441.4
3	Malignant neoplasms (C00-C97)	45,427	10.1	1,350.9
4	Cerebrovascular diseases (I60-I69)	35,939	8.0	1,068.7
5	Chronic lower respiratory diseases (J40-J47)	22,834	5.1	679.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,466	2.8	370.7
7	Influenza and pneumonia (J09-J18)	10,641	2.4	316.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,671	1.7	228.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,535	1.7	224.1
10	Diabetes mellitus (E10-E14)	6,959	1.5	206.9
11	Septicemia (A40-A41)	5,344	1.2	158.9
12	Parkinson's disease (G20-G21)	5,103	1.1	151.7
13	Pneumonitis due to solids and liquids (J69)	3,956	0.9	117.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,433	0.5	72.4
15	Nutritional deficiencies (E40-E64)	1,921	0.4	57.1
...	All other causes (residual)	106,991	23.7	3,181.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	326,810	100.0	775.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,608	23.4	181.8
2	Malignant neoplasms (C00-C97)	69,725	21.3	165.5
3	Cerebrovascular diseases (I60-I69)	18,189	5.6	43.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,089	5.5	42.9
5	Diabetes mellitus (E10-E14)	14,053	4.3	33.3
6	Chronic lower respiratory diseases (J40-J47)	10,838	3.3	25.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9,860	3.0	23.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,200	2.8	21.8
9	Alzheimer's disease (G30)	8,577	2.6	20.4
10	Septicemia (A40-A41)	6,684	2.0	15.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,924	1.8	14.1
12	Influenza and pneumonia (J09-J18)	5,468	1.7	13.0
13	Certain conditions originating in the perinatal period (P00-P96)	3,775	1.2	9.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3,152	1.0	7.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3,149	1.0	7.5
...	All other causes (residual)	63,519	19.4	150.7
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,016	100.0	41.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	277	27.3	11.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	13.9	5.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	10.9	4.5
4	Malignant neoplasms (C00-C97)	58	5.7	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.3	1.4
5	Influenza and pneumonia (J09-J18)	34	3.3	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	26	2.6	1.1
8	Chronic lower respiratory diseases (J40-J47)	19	1.9	*
9	Cerebrovascular diseases (I60-I69)	15	1.5	*
10	Septicemia (A40-A41)	14	1.4	*
11	Anemias (D50-D64)	13	1.3	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
14	Meningitis (G00,G03)	6	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
...	All other causes (residual)	248	24.4	10.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,233	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	388	31.5	6.3
2	Malignant neoplasms (C00-C97)	144	11.7	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	118	9.6	1.9
4	Chronic lower respiratory diseases (J40-J47)	88	7.1	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	5.0	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	4.1	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	3.6	0.7
8	Influenza and pneumonia (J09-J18)	21	1.7	0.3
9	Cerebrovascular diseases (I60-I69)	20	1.6	0.3
10	Septicemia (A40-A41)	12	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.9	*
12	Anemias (D50-D64)	10	0.8	*
13	Diabetes mellitus (E10-E14)	8	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
15	Meningitis (G00,G03)	5	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
...	All other causes (residual)	239	19.4	3.9
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	7,703	100.0	115.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,281	42.6	49.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,869	24.3	27.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	656	8.5	9.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	300	3.9	4.5
5	Malignant neoplasms (C00-C97)	230	3.0	3.4
6	Chronic lower respiratory diseases (J40-J47)	78	1.0	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	0.9	1.1
8	Pregnancy, childbirth and the puerperium (O00-O99)	67	0.9	1.0
9	Diabetes mellitus (E10-E14)	64	0.8	1.0
10	Anemias (D50-D64)	59	0.8	0.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	52	0.7	0.8
12	Influenza and pneumonia (J09-J18)	50	0.6	0.7
13	Septicemia (A40-A41)	43	0.6	0.6
14	Cerebrovascular diseases (I60-I69)	33	0.4	0.5
15	Legal intervention (Y35,Y89.0)	32	0.4	0.5
...	All other causes (residual)	817	10.6	12.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-34 years				
...	All causes	11,512	100.0	183.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,101	26.9	49.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,751	23.9	43.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,017	8.8	16.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	729	6.3	11.6
5	Malignant neoplasms (C00-C97)	640	5.6	10.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	338	2.9	5.4
7	Diabetes mellitus (E10-E14)	249	2.2	4.0
8	Pregnancy, childbirth and the puerperium (O00-O99)	154	1.3	2.4
9	Cerebrovascular diseases (I60-I69)	139	1.2	2.2
10	Chronic lower respiratory diseases (J40-J47)	134	1.2	2.1
11	Anemias (D50-D64)	131	1.1	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	108	0.9	1.7
13	Influenza and pneumonia (J09-J18)	107	0.9	1.7
14	Septicemia (A40-A41)	98	0.9	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	89	0.8	1.4
...	All other causes (residual)	1,727	15.0	27.5
Non-Hispanic, black, both sexes, 35-44 years				
...	All causes	15,195	100.0	286.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,864	18.8	54.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,569	16.9	48.5
3	Malignant neoplasms (C00-C97)	1,874	12.3	35.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,643	10.8	31.0
5	Diabetes mellitus (E10-E14)	581	3.8	11.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	541	3.6	10.2
7	Cerebrovascular diseases (I60-I69)	502	3.3	9.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	452	3.0	8.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	300	2.0	5.7
10	Septicemia (A40-A41)	243	1.6	4.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	205	1.3	3.9
12	Influenza and pneumonia (J09-J18)	184	1.2	3.5
13	Chronic lower respiratory diseases (J40-J47)	176	1.2	3.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	166	1.1	3.1
15	Anemias (D50-D64)	138	0.9	2.6
...	All other causes (residual)	2,757	18.1	52.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-54 years				
...	All causes	31,638	100.0	588.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,697	24.3	143.2
2	Malignant neoplasms (C00-C97)	6,823	21.6	126.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,165	10.0	58.9
4	Diabetes mellitus (E10-E14)	1,501	4.7	27.9
5	Cerebrovascular diseases (I60-I69)	1,447	4.6	26.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	867	2.7	16.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	806	2.5	15.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	721	2.3	13.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	643	2.0	12.0
10	Septicemia (A40-A41)	591	1.9	11.0
11	Chronic lower respiratory diseases (J40-J47)	582	1.8	10.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	559	1.8	10.4
13	Influenza and pneumonia (J09-J18)	419	1.3	7.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	339	1.1	6.3
15	Aortic aneurysm and dissection (I71)	166	0.5	3.1
...	All other causes (residual)	5,312	16.8	98.8
Non-Hispanic, black, both sexes, 55-64 years				
...	All causes	62,821	100.0	1,311.2
1	Malignant neoplasms (C00-C97)	18,210	29.0	380.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,534	24.7	324.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,229	5.1	67.4
4	Diabetes mellitus (E10-E14)	3,061	4.9	63.9
5	Cerebrovascular diseases (I60-I69)	2,956	4.7	61.7
6	Chronic lower respiratory diseases (J40-J47)	2,079	3.3	43.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,658	2.6	34.6
8	Septicemia (A40-A41)	1,378	2.2	28.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,331	2.1	27.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,029	1.6	21.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	920	1.5	19.2
12	Influenza and pneumonia (J09-J18)	883	1.4	18.4
13	Viral hepatitis (B15-B19)	481	0.8	10.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	470	0.7	9.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	258	0.4	5.4
...	All other causes (residual)	9,344	14.9	195.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-74 years				
...	All causes	67,628	100.0	2,459.5
1	Malignant neoplasms (C00-C97)	19,726	29.2	717.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,010	25.2	618.6
3	Cerebrovascular diseases (I60-I69)	3,891	5.8	141.5
4	Diabetes mellitus (E10-E14)	3,609	5.3	131.3
5	Chronic lower respiratory diseases (J40-J47)	2,983	4.4	108.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,217	3.3	80.6
7	Septicemia (A40-A41)	1,591	2.4	57.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,565	2.3	56.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,292	1.9	47.0
10	Influenza and pneumonia (J09-J18)	1,127	1.7	41.0
11	Alzheimer's disease (G30)	690	1.0	25.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	644	1.0	23.4
13	Pneumonitis due to solids and liquids (J69)	369	0.5	13.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	347	0.5	12.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	290	0.4	10.5
...	All other causes (residual)	10,277	15.2	373.8
Non-Hispanic, black, both sexes, 75-84 years				
...	All causes	63,074	100.0	5,123.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,067	25.5	1,305.1
2	Malignant neoplasms (C00-C97)	14,407	22.8	1,170.3
3	Cerebrovascular diseases (I60-I69)	4,500	7.1	365.5
4	Diabetes mellitus (E10-E14)	2,974	4.7	241.6
5	Chronic lower respiratory diseases (J40-J47)	2,815	4.5	228.7
6	Alzheimer's disease (G30)	2,702	4.3	219.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,313	3.7	187.9
8	Septicemia (A40-A41)	1,531	2.4	124.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,331	2.1	108.1
10	Influenza and pneumonia (J09-J18)	1,240	2.0	100.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	996	1.6	80.9
12	Parkinson's disease (G20-G21)	579	0.9	47.0
13	Pneumonitis due to solids and liquids (J69)	491	0.8	39.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	330	0.5	26.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	208	0.3	16.9
...	All other causes (residual)	10,590	16.8	860.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85 years and over				
...	All causes	58,116	100.0	12,171.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,944	27.4	3,339.1
2	Malignant neoplasms (C00-C97)	7,599	13.1	1,591.4
3	Alzheimer's disease (G30)	5,070	8.7	1,061.8
4	Cerebrovascular diseases (I60-I69)	4,648	8.0	973.4
5	Diabetes mellitus (E10-E14)	2,003	3.4	419.5
6	Chronic lower respiratory diseases (J40-J47)	1,872	3.2	392.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,837	3.2	384.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,480	2.5	309.9
9	Influenza and pneumonia (J09-J18)	1,340	2.3	280.6
10	Septicemia (A40-A41)	1,114	1.9	233.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	863	1.5	180.7
12	Pneumonitis due to solids and liquids (J69)	552	0.9	115.6
13	Parkinson's disease (G20-G21)	443	0.8	92.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	300	0.5	62.8
15	Nutritional deficiencies (E40-E64)	297	0.5	62.2
...	All other causes (residual)	12,754	21.9	2,671.0
Non-Hispanic, black, male, all ages				
...	All causes	168,750	100.0	836.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,044	23.7	198.4
2	Malignant neoplasms (C00-C97)	35,215	20.9	174.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,453	7.4	61.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,532	5.1	42.3
5	Cerebrovascular diseases (I60-I69)	8,115	4.8	40.2
6	Diabetes mellitus (E10-E14)	6,976	4.1	34.6
7	Chronic lower respiratory diseases (J40-J47)	5,519	3.3	27.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,348	2.6	21.5
9	Septicemia (A40-A41)	3,074	1.8	15.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,805	1.7	13.9
11	Influenza and pneumonia (J09-J18)	2,705	1.6	13.4
12	Alzheimer's disease (G30)	2,451	1.5	12.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,137	1.3	10.6
14	Certain conditions originating in the perinatal period (P00-P96)	2,101	1.2	10.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2,072	1.2	10.3
...	All other causes (residual)	30,203	17.9	149.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	575	100.0	46.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	29.0	13.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	12.7	5.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	9.4	4.3
4	Malignant neoplasms (C00-C97)	32	5.6	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.8	1.8
6	Influenza and pneumonia (J09-J18)	18	3.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	17	3.0	*
8	Chronic lower respiratory diseases (J40-J47)	14	2.4	*
9	Cerebrovascular diseases (I60-I69)	7	1.2	*
9	Acute bronchitis and bronchiolitis (J20-J21)	7	1.2	*
11	Septicemia (A40-A41)	6	1.0	*
12	Anemias (D50-D64)	5	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
13	Meningitis (G00,G03)	4	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	*
...	All other causes (residual)	141	24.5	11.3
Non-Hispanic, black, male, 5-14 years				
...	All causes	701	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	33.2	7.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	10.1	2.3
3	Malignant neoplasms (C00-C97)	66	9.4	2.1
4	Chronic lower respiratory diseases (J40-J47)	58	8.3	1.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	4.7	1.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	4.1	0.9
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.3	0.7
8	Influenza and pneumonia (J09-J18)	14	2.0	*
9	Cerebrovascular diseases (I60-I69)	12	1.7	*
10	Anemias (D50-D64)	8	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.0	*
12	Diabetes mellitus (E10-E14)	5	0.7	*
13	Septicemia (A40-A41)	3	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
15	Meningitis (G00,G03)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	132	18.8	4.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,909	100.0	173.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,955	50.0	86.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,339	22.7	39.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	524	8.9	15.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	3.3	5.7
5	Malignant neoplasms (C00-C97)	137	2.3	4.0
6	Chronic lower respiratory diseases (J40-J47)	48	0.8	1.4
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	0.8	1.4
8	Anemias (D50-D64)	35	0.6	1.0
9	Diabetes mellitus (E10-E14)	33	0.6	1.0
10	Legal intervention (Y35,Y89.0)	31	0.5	0.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.5	0.8
12	Influenza and pneumonia (J09-J18)	24	0.4	0.7
13	Septicemia (A40-A41)	21	0.4	0.6
14	Cerebrovascular diseases (I60-I69)	17	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
...	All other causes (residual)	466	7.9	13.7
Non-Hispanic, black, male, 25-34 years				
...	All causes	8,106	100.0	263.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,762	34.1	89.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,047	25.3	66.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	633	7.8	20.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	603	7.4	19.6
5	Malignant neoplasms (C00-C97)	274	3.4	8.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	243	3.0	7.9
7	Diabetes mellitus (E10-E14)	138	1.7	4.5
8	Cerebrovascular diseases (I60-I69)	84	1.0	2.7
9	Chronic lower respiratory diseases (J40-J47)	82	1.0	2.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	0.8	2.0
11	Septicemia (A40-A41)	54	0.7	1.8
11	Anemias (D50-D64)	54	0.7	1.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	0.6	1.7
14	Influenza and pneumonia (J09-J18)	49	0.6	1.6
15	Legal intervention (Y35,Y89.0)	43	0.5	1.4
...	All other causes (residual)	926	11.4	30.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-44 years				
...	All causes	9,316	100.0	372.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,848	19.8	73.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,839	19.7	73.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,445	15.5	57.7
4	Malignant neoplasms (C00-C97)	709	7.6	28.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	352	3.8	14.1
6	Diabetes mellitus (E10-E14)	322	3.5	12.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	307	3.3	12.3
8	Cerebrovascular diseases (I60-I69)	281	3.0	11.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	157	1.7	6.3
10	Septicemia (A40-A41)	120	1.3	4.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	103	1.1	4.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	102	1.1	4.1
13	Influenza and pneumonia (J09-J18)	96	1.0	3.8
14	Chronic lower respiratory diseases (J40-J47)	95	1.0	3.8
15	Aortic aneurysm and dissection (I71)	74	0.8	3.0
...	All other causes (residual)	1,466	15.7	58.6
Non-Hispanic, black, male, 45-54 years				
...	All causes	18,033	100.0	715.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,886	27.1	193.9
2	Malignant neoplasms (C00-C97)	3,001	16.6	119.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,205	12.2	87.5
4	Diabetes mellitus (E10-E14)	874	4.8	34.7
5	Cerebrovascular diseases (I60-I69)	781	4.3	31.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	659	3.7	26.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	536	3.0	21.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	406	2.3	16.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	387	2.1	15.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	322	1.8	12.8
11	Septicemia (A40-A41)	291	1.6	11.5
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	267	1.5	10.6
13	Chronic lower respiratory diseases (J40-J47)	254	1.4	10.1
14	Influenza and pneumonia (J09-J18)	229	1.3	9.1
15	Aortic aneurysm and dissection (I71)	117	0.6	4.6
...	All other causes (residual)	2,818	15.6	111.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-64 years				
...	All causes	36,501	100.0	1,667.8
1	Malignant neoplasms (C00-C97)	9,759	26.7	445.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,738	26.7	444.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,352	6.4	107.5
4	Diabetes mellitus (E10-E14)	1,694	4.6	77.4
5	Cerebrovascular diseases (I60-I69)	1,675	4.6	76.5
6	Chronic lower respiratory diseases (J40-J47)	1,043	2.9	47.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	914	2.5	41.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	889	2.4	40.6
9	Septicemia (A40-A41)	682	1.9	31.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	651	1.8	29.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	641	1.8	29.3
12	Influenza and pneumonia (J09-J18)	492	1.3	22.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	373	1.0	17.0
14	Viral hepatitis (B15-B19)	324	0.9	14.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	202	0.6	9.2
...	All other causes (residual)	5,072	13.9	231.8
Non-Hispanic, black, male, 65-74 years				
...	All causes	37,301	100.0	3,156.2
1	Malignant neoplasms (C00-C97)	10,681	28.6	903.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,821	26.3	831.0
3	Cerebrovascular diseases (I60-I69)	2,057	5.5	174.1
4	Diabetes mellitus (E10-E14)	1,916	5.1	162.1
5	Chronic lower respiratory diseases (J40-J47)	1,649	4.4	139.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,137	3.0	96.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,102	3.0	93.2
8	Septicemia (A40-A41)	809	2.2	68.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	694	1.9	58.7
10	Influenza and pneumonia (J09-J18)	622	1.7	52.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	447	1.2	37.8
12	Alzheimer's disease (G30)	303	0.8	25.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	253	0.7	21.4
14	Pneumonitis due to solids and liquids (J69)	214	0.6	18.1
15	Parkinson's disease (G20-G21)	175	0.5	14.8
...	All other causes (residual)	5,421	14.5	458.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-84 years				
...	All causes	29,653	100.0	6,277.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,673	25.9	1,624.4
2	Malignant neoplasms (C00-C97)	7,208	24.3	1,526.0
3	Cerebrovascular diseases (I60-I69)	1,950	6.6	412.8
4	Chronic lower respiratory diseases (J40-J47)	1,443	4.9	305.5
5	Diabetes mellitus (E10-E14)	1,352	4.6	286.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,005	3.4	212.8
7	Alzheimer's disease (G30)	964	3.3	204.1
8	Septicemia (A40-A41)	680	2.3	144.0
9	Influenza and pneumonia (J09-J18)	611	2.1	129.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	596	2.0	126.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	577	1.9	122.2
12	Parkinson's disease (G20-G21)	359	1.2	76.0
13	Pneumonitis due to solids and liquids (J69)	259	0.9	54.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	167	0.6	35.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	127	0.4	26.9
...	All other causes (residual)	4,682	15.8	991.2
Non-Hispanic, black, male, 85 years and over				
...	All causes	18,901	100.0	13,320.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,160	27.3	3,636.4
2	Malignant neoplasms (C00-C97)	3,338	17.7	2,352.4
3	Cerebrovascular diseases (I60-I69)	1,229	6.5	866.1
4	Alzheimer's disease (G30)	1,129	6.0	795.6
5	Chronic lower respiratory diseases (J40-J47)	824	4.4	580.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	647	3.4	456.0
7	Diabetes mellitus (E10-E14)	640	3.4	451.0
8	Influenza and pneumonia (J09-J18)	515	2.7	362.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	434	2.3	305.9
10	Septicemia (A40-A41)	370	2.0	260.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	342	1.8	241.0
12	Pneumonitis due to solids and liquids (J69)	227	1.2	160.0
13	Parkinson's disease (G20-G21)	211	1.1	148.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	100	0.5	70.5
15	Nutritional deficiencies (E40-E64)	99	0.5	69.8
...	All other causes (residual)	3,636	19.2	2,562.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	158,060	100.0	719.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,564	23.1	166.5
2	Malignant neoplasms (C00-C97)	34,510	21.8	157.1
3	Cerebrovascular diseases (I60-I69)	10,074	6.4	45.9
4	Diabetes mellitus (E10-E14)	7,077	4.5	32.2
5	Alzheimer's disease (G30)	6,126	3.9	27.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,636	3.6	25.7
7	Chronic lower respiratory diseases (J40-J47)	5,319	3.4	24.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,852	3.1	22.1
9	Septicemia (A40-A41)	3,610	2.3	16.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,119	2.0	14.2
11	Influenza and pneumonia (J09-J18)	2,763	1.7	12.6
12	Certain conditions originating in the perinatal period (P00-P96)	1,674	1.1	7.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,328	0.8	6.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,151	0.7	5.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,077	0.7	4.9
...	All other causes (residual)	33,180	21.0	151.1
Non-Hispanic, black, female, 1-4 years				
...	All causes	441	100.0	36.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110	24.9	9.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	15.4	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	12.9	4.7
4	Malignant neoplasms (C00-C97)	26	5.9	2.2
5	Influenza and pneumonia (J09-J18)	16	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.7	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	2.0	*
8	Septicemia (A40-A41)	8	1.8	*
8	Anemias (D50-D64)	8	1.8	*
8	Cerebrovascular diseases (I60-I69)	8	1.8	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Meningitis (G00,G03)	2	0.5	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	102	23.1	8.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	532	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	29.1	5.1
2	Malignant neoplasms (C00-C97)	78	14.7	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	8.8	1.5
4	Chronic lower respiratory diseases (J40-J47)	30	5.6	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	5.5	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.3	0.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	2.8	*
8	Septicemia (A40-A41)	9	1.7	*
9	Cerebrovascular diseases (I60-I69)	8	1.5	*
10	Influenza and pneumonia (J09-J18)	7	1.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Meningitis (G00,G03)	3	0.6	*
13	Pneumonitis due to solids and liquids (J69)	3	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
...	All other causes (residual)	105	19.7	3.5
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,794	100.0	54.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	530	29.5	16.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	326	18.2	9.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	132	7.4	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	5.9	3.2
5	Malignant neoplasms (C00-C97)	93	5.2	2.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	67	3.7	2.0
7	Diabetes mellitus (E10-E14)	31	1.7	0.9
8	Chronic lower respiratory diseases (J40-J47)	30	1.7	0.9
9	Influenza and pneumonia (J09-J18)	26	1.4	0.8
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	1.4	0.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	24	1.3	0.7
11	Anemias (D50-D64)	24	1.3	0.7
13	Septicemia (A40-A41)	22	1.2	0.7
14	Cerebrovascular diseases (I60-I69)	16	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
...	All other causes (residual)	327	18.2	9.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-34 years				
...	All causes	3,406	100.0	106.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	704	20.7	22.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	384	11.3	12.0
3	Malignant neoplasms (C00-C97)	366	10.7	11.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	10.0	10.6
5	Pregnancy, childbirth and the puerperium (O00-O99)	154	4.5	4.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	3.7	3.9
7	Diabetes mellitus (E10-E14)	111	3.3	3.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	95	2.8	3.0
9	Anemias (D50-D64)	77	2.3	2.4
10	Influenza and pneumonia (J09-J18)	58	1.7	1.8
11	Cerebrovascular diseases (I60-I69)	55	1.6	1.7
12	Chronic lower respiratory diseases (J40-J47)	52	1.5	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.4	1.4
14	Septicemia (A40-A41)	44	1.3	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.1	1.2
...	All other causes (residual)	758	22.3	23.6
Non-Hispanic, black, female, 35-44 years				
...	All causes	5,879	100.0	210.1
1	Malignant neoplasms (C00-C97)	1,165	19.8	41.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,016	17.3	36.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	730	12.4	26.1
4	Diabetes mellitus (E10-E14)	259	4.4	9.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	234	4.0	8.4
6	Cerebrovascular diseases (I60-I69)	221	3.8	7.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	198	3.4	7.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	2.4	5.1
9	Septicemia (A40-A41)	123	2.1	4.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	108	1.8	3.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	103	1.8	3.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	1.7	3.6
13	Influenza and pneumonia (J09-J18)	88	1.5	3.1
14	Chronic lower respiratory diseases (J40-J47)	81	1.4	2.9
15	Anemias (D50-D64)	73	1.2	2.6
...	All other causes (residual)	1,237	21.0	44.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-54 years				
...	All causes	13,605	100.0	476.5
1	Malignant neoplasms (C00-C97)	3,822	28.1	133.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,811	20.7	98.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	960	7.1	33.6
4	Cerebrovascular diseases (I60-I69)	666	4.9	23.3
5	Diabetes mellitus (E10-E14)	627	4.6	22.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	331	2.4	11.6
7	Chronic lower respiratory diseases (J40-J47)	328	2.4	11.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	315	2.3	11.0
9	Septicemia (A40-A41)	300	2.2	10.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	256	1.9	9.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	237	1.7	8.3
12	Influenza and pneumonia (J09-J18)	190	1.4	6.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	1.1	5.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	0.5	2.5
15	Anemias (D50-D64)	64	0.5	2.2
...	All other causes (residual)	2,479	18.2	86.8
Non-Hispanic, black, female, 55-64 years				
...	All causes	26,320	100.0	1,011.3
1	Malignant neoplasms (C00-C97)	8,451	32.1	324.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,796	22.0	222.7
3	Diabetes mellitus (E10-E14)	1,367	5.2	52.5
4	Cerebrovascular diseases (I60-I69)	1,281	4.9	49.2
5	Chronic lower respiratory diseases (J40-J47)	1,036	3.9	39.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	877	3.3	33.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	744	2.8	28.6
8	Septicemia (A40-A41)	696	2.6	26.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	442	1.7	17.0
10	Influenza and pneumonia (J09-J18)	391	1.5	15.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	388	1.5	14.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	269	1.0	10.3
13	Viral hepatitis (B15-B19)	157	0.6	6.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	100	0.4	3.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	0.4	3.7
...	All other causes (residual)	4,228	16.1	162.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-74 years				
...	All causes	30,327	100.0	1,934.3
1	Malignant neoplasms (C00-C97)	9,045	29.8	576.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,189	23.7	458.5
3	Cerebrovascular diseases (I60-I69)	1,834	6.0	117.0
4	Diabetes mellitus (E10-E14)	1,693	5.6	108.0
5	Chronic lower respiratory diseases (J40-J47)	1,334	4.4	85.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,080	3.6	68.9
7	Septicemia (A40-A41)	782	2.6	49.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	598	2.0	38.1
9	Influenza and pneumonia (J09-J18)	505	1.7	32.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	463	1.5	29.5
11	Alzheimer's disease (G30)	387	1.3	24.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	197	0.6	12.6
13	Pneumonitis due to solids and liquids (J69)	155	0.5	9.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	133	0.4	8.5
15	Parkinson's disease (G20-G21)	98	0.3	6.3
...	All other causes (residual)	4,834	15.9	308.3
Non-Hispanic, black, female, 75-84 years				
...	All causes	33,421	100.0	4,404.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,394	25.1	1,106.3
2	Malignant neoplasms (C00-C97)	7,199	21.5	948.8
3	Cerebrovascular diseases (I60-I69)	2,550	7.6	336.1
4	Alzheimer's disease (G30)	1,738	5.2	229.1
5	Diabetes mellitus (E10-E14)	1,622	4.9	213.8
6	Chronic lower respiratory diseases (J40-J47)	1,372	4.1	180.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,308	3.9	172.4
8	Septicemia (A40-A41)	851	2.5	112.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	754	2.3	99.4
10	Influenza and pneumonia (J09-J18)	629	1.9	82.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	400	1.2	52.7
12	Pneumonitis due to solids and liquids (J69)	232	0.7	30.6
13	Parkinson's disease (G20-G21)	220	0.7	29.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	163	0.5	21.5
15	Nutritional deficiencies (E40-E64)	109	0.3	14.4
...	All other causes (residual)	5,880	17.6	775.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85 years and over				
...	All causes	39,215	100.0	11,685.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,784	27.5	3,213.4
2	Malignant neoplasms (C00-C97)	4,261	10.9	1,269.7
3	Alzheimer's disease (G30)	3,941	10.0	1,174.3
4	Cerebrovascular diseases (I60-I69)	3,419	8.7	1,018.8
5	Diabetes mellitus (E10-E14)	1,363	3.5	406.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,190	3.0	354.6
7	Chronic lower respiratory diseases (J40-J47)	1,048	2.7	312.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,046	2.7	311.7
9	Influenza and pneumonia (J09-J18)	825	2.1	245.8
10	Septicemia (A40-A41)	744	1.9	221.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	1.3	155.2
12	Pneumonitis due to solids and liquids (J69)	325	0.8	96.8
13	Parkinson's disease (G20-G21)	232	0.6	69.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	200	0.5	59.6
15	Nutritional deficiencies (E40-E64)	198	0.5	59.0
...	All other causes (residual)	9,118	23.3	2,716.9
Non-Hispanic, American Indian or Alaska Native, both sexes, all ages				
...	All causes	18,595	100.0	685.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,332	17.9	122.9
2	Malignant neoplasms (C00-C97)	3,225	17.3	119.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,100	11.3	77.5
4	Diabetes mellitus (E10-E14)	1,078	5.8	39.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,010	5.4	37.3
6	Chronic lower respiratory diseases (J40-J47)	853	4.6	31.5
7	Cerebrovascular diseases (I60-I69)	632	3.4	23.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	580	3.1	21.4
9	Alzheimer's disease (G30)	377	2.0	13.9
10	Influenza and pneumonia (J09-J18)	360	1.9	13.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	1.8	12.7
12	Septicemia (A40-A41)	295	1.6	10.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	279	1.5	10.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	193	1.0	7.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	0.6	4.4
...	All other causes (residual)	3,818	20.5	140.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 1-4 years				
...	All causes	82	100.0	51.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	32.9	16.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	12.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	9.8	*
4	Malignant neoplasms (C00-C97)	3	3.7	*
4	Influenza and pneumonia (J09-J18)	3	3.7	*
6	Meningococcal infection (A39)	2	2.4	*
6	Septicemia (A40-A41)	2	2.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.4	*
9	Diabetes mellitus (E10-E14)	1	1.2	*
9	Cerebrovascular diseases (I60-I69)	1	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.2	*
...	All other causes (residual)	22	26.8	13.8
Non-Hispanic, American Indian or Alaska Native, both sexes, 5-14 years				
...	All causes	92	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	33.7	7.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	14.1	*
3	Malignant neoplasms (C00-C97)	8	8.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	4.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	3.3	*
6	Septicemia (A40-A41)	2	2.2	*
6	Meningitis (G00,G03)	2	2.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.2	*
9	Meningococcal infection (A39)	1	1.1	*
9	Diabetes mellitus (E10-E14)	1	1.1	*
9	Cerebrovascular diseases (I60-I69)	1	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.1	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.1	*
...	All other causes (residual)	21	22.8	5.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 15-24 years				
...	All causes	591	100.0	136.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	270	45.7	62.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	26.4	36.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	9.5	12.9
4	Malignant neoplasms (C00-C97)	10	1.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	1.2	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.0	*
7	Diabetes mellitus (E10-E14)	4	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.5	*
9	Legal intervention (Y35,Y89.0)	3	0.5	*
12	Cerebrovascular diseases (I60-I69)	2	0.3	*
12	Influenza and pneumonia (J09-J18)	2	0.3	*
14	Septicemia (A40-A41)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	61	10.3	14.1
Non-Hispanic, American Indian or Alaska Native, both sexes, 25-34 years				
...	All causes	1,103	100.0	275.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	400	36.3	99.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	15.4	42.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	99	9.0	24.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	7.2	19.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	5.5	15.2
6	Malignant neoplasms (C00-C97)	31	2.8	7.7
7	Diabetes mellitus (E10-E14)	25	2.3	6.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.1	*
9	Cerebrovascular diseases (I60-I69)	10	0.9	*
9	Legal intervention (Y35,Y89.0)	10	0.9	*
11	Influenza and pneumonia (J09-J18)	8	0.7	*
12	Septicemia (A40-A41)	6	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	6	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.3	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.3	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.3	*
...	All other causes (residual)	171	15.5	42.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 35-44 years				
...	All causes	1,332	100.0	407.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	363	27.3	111.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	171	12.8	52.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	160	12.0	48.9
4	Malignant neoplasms (C00-C97)	102	7.7	31.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	102	7.7	31.2
6	Diabetes mellitus (E10-E14)	68	5.1	20.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	4.2	17.1
8	Cerebrovascular diseases (I60-I69)	24	1.8	7.3
9	Influenza and pneumonia (J09-J18)	20	1.5	6.1
10	Septicemia (A40-A41)	16	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.9	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.8	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.6	*
13	Pneumonitis due to solids and liquids (J69)	8	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.5	*
...	All other causes (residual)	206	15.5	63.0
Non-Hispanic, American Indian or Alaska Native, both sexes, 45-54 years				
...	All causes	2,292	100.0	676.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	17.1	115.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	344	15.0	101.5
3	Malignant neoplasms (C00-C97)	323	14.1	95.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	287	12.5	84.7
5	Diabetes mellitus (E10-E14)	132	5.8	39.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	2.7	18.6
7	Cerebrovascular diseases (I60-I69)	49	2.1	14.5
8	Chronic lower respiratory diseases (J40-J47)	45	2.0	13.3
9	Influenza and pneumonia (J09-J18)	42	1.8	12.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	1.7	11.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	1.7	11.2
12	Septicemia (A40-A41)	34	1.5	10.0
13	Viral hepatitis (B15-B19)	24	1.0	7.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	0.8	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.5	*
...	All other causes (residual)	451	19.7	133.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 55-64 years				
...	All causes	3,433	100.0	1,087.9
1	Malignant neoplasms (C00-C97)	826	24.1	261.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	622	18.1	197.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	301	8.8	95.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	265	7.7	84.0
5	Diabetes mellitus (E10-E14)	238	6.9	75.4
6	Chronic lower respiratory diseases (J40-J47)	150	4.4	47.5
7	Cerebrovascular diseases (I60-I69)	88	2.6	27.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	1.7	18.7
9	Septicemia (A40-A41)	55	1.6	17.4
9	Influenza and pneumonia (J09-J18)	55	1.6	17.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	1.5	16.5
12	Viral hepatitis (B15-B19)	45	1.3	14.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	0.8	8.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	0.7	7.3
15	Pneumonitis due to solids and liquids (J69)	15	0.4	*
...	All other causes (residual)	611	17.8	193.6
Non-Hispanic, American Indian or Alaska Native, both sexes, 65-74 years				
...	All causes	3,637	100.0	1,975.1
1	Malignant neoplasms (C00-C97)	937	25.8	508.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	771	21.2	418.7
3	Diabetes mellitus (E10-E14)	280	7.7	152.1
4	Chronic lower respiratory diseases (J40-J47)	275	7.6	149.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	3.5	70.1
6	Cerebrovascular diseases (I60-I69)	119	3.3	64.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	112	3.1	60.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	92	2.5	50.0
9	Septicemia (A40-A41)	80	2.2	43.4
10	Influenza and pneumonia (J09-J18)	64	1.8	34.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	1.2	23.4
12	Alzheimer's disease (G30)	40	1.1	21.7
13	Pneumonitis due to solids and liquids (J69)	21	0.6	11.4
14	Parkinson's disease (G20-G21)	19	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.5	*
...	All other causes (residual)	638	17.5	346.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 75-84 years				
...	All causes	3,332	100.0	4,416.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	759	22.8	1,006.1
2	Malignant neoplasms (C00-C97)	704	21.1	933.2
3	Chronic lower respiratory diseases (J40-J47)	242	7.3	320.8
4	Diabetes mellitus (E10-E14)	222	6.7	294.3
5	Cerebrovascular diseases (I60-I69)	178	5.3	235.9
6	Alzheimer's disease (G30)	121	3.6	160.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	2.9	127.3
8	Influenza and pneumonia (J09-J18)	81	2.4	107.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	2.1	92.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	60	1.8	79.5
11	Septicemia (A40-A41)	58	1.7	76.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	51	1.5	67.6
13	Parkinson's disease (G20-G21)	41	1.2	54.3
14	Pneumonitis due to solids and liquids (J69)	29	0.9	38.4
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	23	0.7	30.5
...	All other causes (residual)	597	17.9	791.3
Non-Hispanic, American Indian or Alaska Native, both sexes, 85 years and over				
...	All causes	2,389	100.0	9,445.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	601	25.2	2,376.2
2	Malignant neoplasms (C00-C97)	280	11.7	1,107.0
3	Alzheimer's disease (G30)	205	8.6	810.5
4	Cerebrovascular diseases (I60-I69)	160	6.7	632.6
5	Chronic lower respiratory diseases (J40-J47)	125	5.2	494.2
6	Diabetes mellitus (E10-E14)	107	4.5	423.0
7	Influenza and pneumonia (J09-J18)	83	3.5	328.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	2.9	276.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	2.7	253.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	39	1.6	154.2
11	Septicemia (A40-A41)	38	1.6	150.2
12	Pneumonitis due to solids and liquids (J69)	27	1.1	106.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.7	*
13	Nutritional deficiencies (E40-E64)	16	0.7	*
13	Parkinson's disease (G20-G21)	16	0.7	*
...	All other causes (residual)	542	22.7	2,142.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, all ages				
...	All causes	10,280	100.0	772.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,951	19.0	146.7
2	Malignant neoplasms (C00-C97)	1,723	16.8	129.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,404	13.7	105.5
4	Diabetes mellitus (E10-E14)	574	5.6	43.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	554	5.4	41.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	441	4.3	33.2
7	Chronic lower respiratory diseases (J40-J47)	430	4.2	32.3
8	Cerebrovascular diseases (I60-I69)	288	2.8	21.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	209	2.0	15.7
10	Influenza and pneumonia (J09-J18)	176	1.7	13.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	1.6	12.5
12	Septicemia (A40-A41)	128	1.2	9.6
13	Alzheimer's disease (G30)	121	1.2	9.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	0.8	6.1
15	Certain conditions originating in the perinatal period (P00-P96)	72	0.7	5.4
...	All other causes (residual)	1,962	19.1	147.5
Non-Hispanic, American Indian or Alaska Native, male, 1-4 years				
...	All causes	38	100.0	46.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	36.8	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	10.5	*
3	Malignant neoplasms (C00-C97)	2	5.3	*
3	Influenza and pneumonia (J09-J18)	2	5.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	5.3	*
6	Meningococcal infection (A39)	1	2.6	*
6	Septicemia (A40-A41)	1	2.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.6	*
6	Cerebrovascular diseases (I60-I69)	1	2.6	*
...	All other causes (residual)	10	26.3	*
Non-Hispanic, American Indian or Alaska Native, male, 5-14 years				
...	All causes	49	100.0	23.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	42.9	10.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	14.3	*
3	Malignant neoplasms (C00-C97)	3	6.1	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.1	*
5	Diabetes mellitus (E10-E14)	1	2.0	*
5	Meningitis (G00,G03)	1	2.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.0	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.0	*
...	All other causes (residual)	12	24.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 15-24 years				
...	All causes	404	100.0	183.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	46.0	84.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	114	28.2	51.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	10.4	19.0
4	Malignant neoplasms (C00-C97)	6	1.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	1.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.0	*
7	Diabetes mellitus (E10-E14)	3	0.7	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
8	Legal intervention (Y35,Y89.0)	2	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
11	Cerebrovascular diseases (I60-I69)	1	0.2	*
11	Influenza and pneumonia (J09-J18)	1	0.2	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	33	8.2	15.0
Non-Hispanic, American Indian or Alaska Native, male, 25-34 years				
...	All causes	729	100.0	362.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	264	36.2	131.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	133	18.2	66.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	8.8	31.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	59	8.1	29.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	5.2	18.9
6	Diabetes mellitus (E10-E14)	17	2.3	*
7	Malignant neoplasms (C00-C97)	13	1.8	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.2	*
8	Legal intervention (Y35,Y89.0)	9	1.2	*
10	Cerebrovascular diseases (I60-I69)	8	1.1	*
11	Influenza and pneumonia (J09-J18)	4	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Septicemia (A40-A41)	2	0.3	*
14	Viral hepatitis (B15-B19)	2	0.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
...	All other causes (residual)	99	13.6	49.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 35-44 years				
...	All causes	825	100.0	514.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	29.2	150.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	13.0	66.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	96	11.6	59.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	9.1	46.8
5	Malignant neoplasms (C00-C97)	44	5.3	27.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	5.0	25.6
7	Diabetes mellitus (E10-E14)	39	4.7	24.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.2	*
9	Cerebrovascular diseases (I60-I69)	9	1.1	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	1.0	*
10	Influenza and pneumonia (J09-J18)	8	1.0	*
12	Septicemia (A40-A41)	6	0.7	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
14	Pneumonitis due to solids and liquids (J69)	3	0.4	*
14	Legal intervention (Y35,Y89.0)	3	0.4	*
...	All other causes (residual)	126	15.3	78.6
Non-Hispanic, American Indian or Alaska Native, male, 45-54 years				
...	All causes	1,363	100.0	831.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	268	19.7	163.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	242	17.8	147.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	156	11.4	95.1
4	Malignant neoplasms (C00-C97)	145	10.6	88.4
5	Diabetes mellitus (E10-E14)	79	5.8	48.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	3.7	31.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	2.3	19.5
8	Influenza and pneumonia (J09-J18)	25	1.8	15.2
9	Cerebrovascular diseases (I60-I69)	23	1.7	14.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.5	12.8
11	Septicemia (A40-A41)	19	1.4	*
12	Chronic lower respiratory diseases (J40-J47)	18	1.3	*
13	Viral hepatitis (B15-B19)	11	0.8	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.7	*
15	Aortic aneurysm and dissection (I71)	9	0.7	*
...	All other causes (residual)	254	18.6	154.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 55-64 years				
...	All causes	2,065	100.0	1,398.5
1	Malignant neoplasms (C00-C97)	475	23.0	321.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	393	19.0	266.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	222	10.8	150.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	152	7.4	102.9
5	Diabetes mellitus (E10-E14)	141	6.8	95.5
6	Chronic lower respiratory diseases (J40-J47)	76	3.7	51.5
7	Cerebrovascular diseases (I60-I69)	50	2.4	33.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	2.1	29.8
9	Influenza and pneumonia (J09-J18)	42	2.0	28.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.3	18.3
11	Viral hepatitis (B15-B19)	26	1.3	17.6
12	Septicemia (A40-A41)	23	1.1	15.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.9	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.4	*
...	All other causes (residual)	352	17.0	238.4
Non-Hispanic, American Indian or Alaska Native, male, 65-74 years				
...	All causes	2,047	100.0	2,384.9
1	Malignant neoplasms (C00-C97)	540	26.4	629.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	484	23.6	563.9
3	Diabetes mellitus (E10-E14)	161	7.9	187.6
4	Chronic lower respiratory diseases (J40-J47)	135	6.6	157.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	4.3	101.4
6	Cerebrovascular diseases (I60-I69)	65	3.2	75.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	60	2.9	69.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	2.3	55.9
9	Influenza and pneumonia (J09-J18)	37	1.8	43.1
10	Septicemia (A40-A41)	31	1.5	36.1
11	Parkinson's disease (G20-G21)	16	0.8	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.8	*
13	Alzheimer's disease (G30)	15	0.7	*
14	Pneumonitis due to solids and liquids (J69)	13	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
...	All other causes (residual)	330	16.1	384.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 75-84 years				
...	All causes	1,627	100.0	4,984.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	409	25.1	1,253.0
2	Malignant neoplasms (C00-C97)	363	22.3	1,112.1
3	Chronic lower respiratory diseases (J40-J47)	131	8.1	401.3
4	Diabetes mellitus (E10-E14)	92	5.7	281.8
5	Cerebrovascular diseases (I60-I69)	77	4.7	235.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	3.6	180.7
7	Alzheimer's disease (G30)	42	2.6	128.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.9	95.0
9	Influenza and pneumonia (J09-J18)	29	1.8	88.8
10	Septicemia (A40-A41)	28	1.7	85.8
10	Parkinson's disease (G20-G21)	28	1.7	85.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.5	76.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	1.2	*
14	Pneumonitis due to solids and liquids (J69)	16	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
15	Aortic aneurysm and dissection (I71)	7	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	7	0.4	*
...	All other causes (residual)	250	15.4	765.9
Non-Hispanic, American Indian or Alaska Native, male, 85 years and over				
...	All causes	950	100.0	10,474.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	268	28.2	2,954.8
2	Malignant neoplasms (C00-C97)	132	13.9	1,455.3
3	Chronic lower respiratory diseases (J40-J47)	61	6.4	672.5
4	Alzheimer's disease (G30)	57	6.0	628.4
5	Cerebrovascular diseases (I60-I69)	54	5.7	595.4
6	Diabetes mellitus (E10-E14)	41	4.3	452.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	2.9	308.7
8	Influenza and pneumonia (J09-J18)	27	2.8	297.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	2.7	286.7
10	Septicemia (A40-A41)	17	1.8	*
11	Pneumonitis due to solids and liquids (J69)	12	1.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.9	*
12	Parkinson's disease (G20-G21)	9	0.9	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.9	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	7	0.7	*
...	All other causes (residual)	193	20.3	2,127.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, all ages				
...	All causes	8,315	100.0	602.2
1	Malignant neoplasms (C00-C97)	1,502	18.1	108.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,381	16.6	100.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	696	8.4	50.4
4	Diabetes mellitus (E10-E14)	504	6.1	36.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	456	5.5	33.0
6	Chronic lower respiratory diseases (J40-J47)	423	5.1	30.6
7	Cerebrovascular diseases (I60-I69)	344	4.1	24.9
8	Alzheimer's disease (G30)	256	3.1	18.5
9	Influenza and pneumonia (J09-J18)	184	2.2	13.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	2.1	12.8
11	Septicemia (A40-A41)	167	2.0	12.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	139	1.7	10.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	112	1.3	8.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	0.8	5.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	0.8	4.7
...	All other causes (residual)	1,839	22.1	133.2
Non-Hispanic, American Indian or Alaska Native, female, 1-4 years				
...	All causes	44	100.0	55.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	29.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	18.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	9.1	*
4	Meningococcal infection (A39)	1	2.3	*
4	Septicemia (A40-A41)	1	2.3	*
4	Malignant neoplasms (C00-C97)	1	2.3	*
4	Diabetes mellitus (E10-E14)	1	2.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.3	*
4	Influenza and pneumonia (J09-J18)	1	2.3	*
4	Certain conditions originating in the perinatal period (P00-P96)	1	2.3	*
...	All other causes (residual)	12	27.3	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 5-14 years				
...	All causes	43	100.0	21.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	23.3	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	14.0	*
3	Malignant neoplasms (C00-C97)	5	11.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	9.3	*
5	Septicemia (A40-A41)	2	4.7	*
6	Meningococcal infection (A39)	1	2.3	*
6	Meningitis (G00,G03)	1	2.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.3	*
6	Cerebrovascular diseases (I60-I69)	1	2.3	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.3	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	2.3	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.3	*
...	All other causes (residual)	9	20.9	*
Non-Hispanic, American Indian or Alaska Native, female, 15-24 years				
...	All causes	187	100.0	87.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	44.9	39.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	22.5	19.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	7.5	*
4	Malignant neoplasms (C00-C97)	4	2.1	*
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4	2.1	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.6	*
7	Septicemia (A40-A41)	1	0.5	*
7	Diabetes mellitus (E10-E14)	1	0.5	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	0.5	*
7	Cerebrovascular diseases (I60-I69)	1	0.5	*
7	Influenza and pneumonia (J09-J18)	1	0.5	*
7	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
7	Legal intervention (Y35,Y89.0)	1	0.5	*
...	All other causes (residual)	28	15.0	13.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 25-34 years				
...	All causes	374	100.0	187.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	36.4	68.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	40	10.7	20.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	9.9	18.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	6.1	11.5
5	Malignant neoplasms (C00-C97)	18	4.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	4.0	*
7	Diabetes mellitus (E10-E14)	8	2.1	*
8	Septicemia (A40-A41)	4	1.1	*
8	Influenza and pneumonia (J09-J18)	4	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.8	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.8	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
14	Anemias (D50-D64)	2	0.5	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
14	Cerebrovascular diseases (I60-I69)	2	0.5	*
...	All other causes (residual)	69	18.4	34.6
Non-Hispanic, American Indian or Alaska Native, female, 35-44 years				
...	All causes	507	100.0	304.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	24.1	73.3
2	Chronic liver disease and cirrhosis (K70,K73-K74)	75	14.8	45.0
3	Malignant neoplasms (C00-C97)	58	11.4	34.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	53	10.5	31.8
5	Diabetes mellitus (E10-E14)	29	5.7	17.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	5.3	16.2
7	Cerebrovascular diseases (I60-I69)	15	3.0	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	3.0	*
9	Influenza and pneumonia (J09-J18)	12	2.4	*
10	Septicemia (A40-A41)	10	2.0	*
11	Pneumonitis due to solids and liquids (J69)	5	1.0	*
12	Viral hepatitis (B15-B19)	4	0.8	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.4	*
...	All other causes (residual)	66	13.0	39.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 45-54 years				
...	All causes	929	100.0	531.1
1	Malignant neoplasms (C00-C97)	178	19.2	101.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	131	14.1	74.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	13.2	70.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	102	11.0	58.3
5	Diabetes mellitus (E10-E14)	53	5.7	30.3
6	Chronic lower respiratory diseases (J40-J47)	27	2.9	15.4
7	Cerebrovascular diseases (I60-I69)	26	2.8	14.9
8	Influenza and pneumonia (J09-J18)	17	1.8	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.8	*
10	Septicemia (A40-A41)	15	1.6	*
11	Viral hepatitis (B15-B19)	13	1.4	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	1.3	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.0	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	0.6	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	6	0.6	*
...	All other causes (residual)	194	20.9	110.9
Non-Hispanic, American Indian or Alaska Native, female, 55-64 years				
...	All causes	1,368	100.0	814.8
1	Malignant neoplasms (C00-C97)	351	25.7	209.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	229	16.7	136.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	113	8.3	67.3
4	Diabetes mellitus (E10-E14)	97	7.1	57.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	5.8	47.1
6	Chronic lower respiratory diseases (J40-J47)	74	5.4	44.1
7	Cerebrovascular diseases (I60-I69)	38	2.8	22.6
8	Septicemia (A40-A41)	32	2.3	19.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	2.3	19.1
10	Viral hepatitis (B15-B19)	19	1.4	*
11	Influenza and pneumonia (J09-J18)	13	1.0	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.7	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	0.6	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	0.6	*
15	Pneumonitis due to solids and liquids (J69)	7	0.5	*
...	All other causes (residual)	258	18.9	153.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 65-74 years				
...	All causes	1,590	100.0	1,617.3
1	Malignant neoplasms (C00-C97)	397	25.0	403.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	287	18.1	291.9
3	Chronic lower respiratory diseases (J40-J47)	140	8.8	142.4
4	Diabetes mellitus (E10-E14)	119	7.5	121.0
5	Cerebrovascular diseases (I60-I69)	54	3.4	54.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	52	3.3	52.9
7	Septicemia (A40-A41)	49	3.1	49.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	2.8	44.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	2.6	42.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	1.7	27.5
10	Influenza and pneumonia (J09-J18)	27	1.7	27.5
12	Alzheimer's disease (G30)	25	1.6	25.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.5	*
13	Pneumonitis due to solids and liquids (J69)	8	0.5	*
15	Viral hepatitis (B15-B19)	6	0.4	*
...	All other causes (residual)	305	19.2	310.2
Non-Hispanic, American Indian or Alaska Native, female, 75-84 years				
...	All causes	1,705	100.0	3,983.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	350	20.5	817.8
2	Malignant neoplasms (C00-C97)	341	20.0	796.7
3	Diabetes mellitus (E10-E14)	130	7.6	303.7
4	Chronic lower respiratory diseases (J40-J47)	111	6.5	259.4
5	Cerebrovascular diseases (I60-I69)	101	5.9	236.0
6	Alzheimer's disease (G30)	79	4.6	184.6
7	Influenza and pneumonia (J09-J18)	52	3.0	121.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	2.3	91.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	2.2	86.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	35	2.1	81.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	1.9	74.8
12	Septicemia (A40-A41)	30	1.8	70.1
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	16	0.9	*
14	Parkinson's disease (G20-G21)	13	0.8	*
14	Pneumonitis due to solids and liquids (J69)	13	0.8	*
...	All other causes (residual)	326	19.1	761.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 85 years and over				
...	All causes	1,439	100.0	8,870.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	23.1	2,052.6
2	Malignant neoplasms (C00-C97)	148	10.3	912.3
2	Alzheimer's disease (G30)	148	10.3	912.3
4	Cerebrovascular diseases (I60-I69)	106	7.4	653.4
5	Diabetes mellitus (E10-E14)	66	4.6	406.8
6	Chronic lower respiratory diseases (J40-J47)	64	4.4	394.5
7	Influenza and pneumonia (J09-J18)	56	3.9	345.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	2.9	258.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	2.6	234.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	2.1	184.9
11	Septicemia (A40-A41)	21	1.5	129.4
12	Pneumonitis due to solids and liquids (J69)	15	1.0	*
13	Nutritional deficiencies (E40-E64)	10	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
14	Parkinson's disease (G20-G21)	7	0.5	*
...	All other causes (residual)	348	24.2	2,145.1
Non-Hispanic, Asian or Pacific Islander, both sexes, all ages				
...	All causes	68,235	100.0	350.3
1	Malignant neoplasms (C00-C97)	17,718	26.0	91.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,519	21.3	74.5
3	Cerebrovascular diseases (I60-I69)	5,217	7.6	26.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,101	4.5	15.9
5	Diabetes mellitus (E10-E14)	2,697	4.0	13.8
6	Alzheimer's disease (G30)	2,363	3.5	12.1
7	Influenza and pneumonia (J09-J18)	2,132	3.1	10.9
8	Chronic lower respiratory diseases (J40-J47)	1,914	2.8	9.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,363	2.0	7.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,348	2.0	6.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,280	1.9	6.6
12	Septicemia (A40-A41)	786	1.2	4.0
13	Parkinson's disease (G20-G21)	749	1.1	3.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	632	0.9	3.2
15	Certain conditions originating in the perinatal period (P00-P96)	527	0.8	2.7
...	All other causes (residual)	11,889	17.4	61.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	160	100.0	17.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	26.9	4.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	13.1	2.2
3	Malignant neoplasms (C00-C97)	20	12.5	2.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	6.3	*
5	Septicemia (A40-A41)	5	3.1	*
6	Cerebrovascular diseases (I60-I69)	4	2.5	*
7	Influenza and pneumonia (J09-J18)	3	1.9	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.3	*
9	Diabetes mellitus (E10-E14)	2	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	2	1.3	*
12	Meningitis (G00,G03)	1	0.6	*
12	Pneumonitis due to solids and liquids (J69)	1	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (residual)	40	25.0	4.3
Non-Hispanic, Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	218	100.0	9.2
1	Malignant neoplasms (C00-C97)	60	27.5	2.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	17.0	1.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	6.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	6.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	5.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	4.1	*
7	Influenza and pneumonia (J09-J18)	7	3.2	*
8	Cerebrovascular diseases (I60-I69)	6	2.8	*
9	Septicemia (A40-A41)	5	2.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	4	1.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	*
12	Scarlet fever and erysipelas (A38,A46)	1	0.5	*
12	Diabetes mellitus (E10-E14)	1	0.5	*
12	Meningitis (G00,G03)	1	0.5	*
12	Diseases of appendix (K35-K38)	1	0.5	*
...	All other causes (residual)	45	20.6	1.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	971	100.0	37.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	31.6	11.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	305	31.4	11.7
3	Malignant neoplasms (C00-C97)	73	7.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	7.0	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	4.2	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.4	*
7	Cerebrovascular diseases (I60-I69)	8	0.8	*
8	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	6	0.6	*
10	Septicemia (A40-A41)	3	0.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.3	*
10	Diabetes mellitus (E10-E14)	3	0.3	*
10	Aortic aneurysm and dissection (I71)	3	0.3	*
10	Influenza and pneumonia (J09-J18)	3	0.3	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.2	*
...	All other causes (residual)	125	12.9	4.8
Non-Hispanic, Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	1,401	100.0	42.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	372	26.6	11.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	264	18.8	7.9
3	Malignant neoplasms (C00-C97)	199	14.2	6.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	7.6	3.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	5.5	2.3
6	Cerebrovascular diseases (I60-I69)	35	2.5	1.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	23	1.6	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	23	1.6	0.7
9	Septicemia (A40-A41)	13	0.9	*
9	Influenza and pneumonia (J09-J18)	13	0.9	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	0.9	*
12	Diabetes mellitus (E10-E14)	11	0.8	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	10	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
...	All other causes (residual)	225	16.1	6.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	2,303	100.0	75.2
1	Malignant neoplasms (C00-C97)	693	30.1	22.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	355	15.4	11.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	13.2	10.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	228	9.9	7.4
5	Cerebrovascular diseases (I60-I69)	123	5.3	4.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	2.6	1.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	57	2.5	1.9
8	Diabetes mellitus (E10-E14)	44	1.9	1.4
9	Influenza and pneumonia (J09-J18)	23	1.0	0.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	0.8	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	19	0.8	*
12	Septicemia (A40-A41)	17	0.7	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.7	*
15	Aortic aneurysm and dissection (I71)	14	0.6	*
...	All other causes (residual)	314	13.6	10.2
Non-Hispanic, Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	4,366	100.0	166.7
1	Malignant neoplasms (C00-C97)	1,697	38.9	64.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	722	16.5	27.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	316	7.2	12.1
4	Cerebrovascular diseases (I60-I69)	286	6.6	10.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	204	4.7	7.8
6	Diabetes mellitus (E10-E14)	161	3.7	6.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	110	2.5	4.2
8	Influenza and pneumonia (J09-J18)	65	1.5	2.5
9	Septicemia (A40-A41)	49	1.1	1.9
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	1.1	1.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	1.1	1.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.1	1.8
13	Viral hepatitis (B15-B19)	29	0.7	1.1
14	Chronic lower respiratory diseases (J40-J47)	28	0.6	1.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.6	1.0
...	All other causes (residual)	535	12.3	20.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	8,508	100.0	405.1
1	Malignant neoplasms (C00-C97)	3,576	42.0	170.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,626	19.1	77.4
3	Cerebrovascular diseases (I60-I69)	514	6.0	24.5
4	Diabetes mellitus (E10-E14)	397	4.7	18.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	4.0	16.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	2.0	8.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	139	1.6	6.6
8	Influenza and pneumonia (J09-J18)	138	1.6	6.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	1.6	6.3
10	Chronic lower respiratory diseases (J40-J47)	128	1.5	6.1
11	Septicemia (A40-A41)	94	1.1	4.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	91	1.1	4.3
13	Viral hepatitis (B15-B19)	71	0.8	3.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	0.7	2.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.5	2.1
...	All other causes (residual)	989	11.6	47.1
Non-Hispanic, Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	12,025	100.0	893.1
1	Malignant neoplasms (C00-C97)	4,355	36.2	323.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,501	20.8	185.8
3	Cerebrovascular diseases (I60-I69)	851	7.1	63.2
4	Diabetes mellitus (E10-E14)	603	5.0	44.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	385	3.2	28.6
6	Chronic lower respiratory diseases (J40-J47)	320	2.7	23.8
7	Influenza and pneumonia (J09-J18)	280	2.3	20.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	254	2.1	18.9
9	Septicemia (A40-A41)	175	1.5	13.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	156	1.3	11.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	135	1.1	10.0
12	Parkinson's disease (G20-G21)	115	1.0	8.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	97	0.8	7.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	83	0.7	6.2
15	Aortic aneurysm and dissection (I71)	72	0.6	5.3
...	All other causes (residual)	1,643	13.7	122.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	16,274	100.0	2,566.8
1	Malignant neoplasms (C00-C97)	4,318	26.5	681.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,601	22.1	568.0
3	Cerebrovascular diseases (I60-I69)	1,394	8.6	219.9
4	Diabetes mellitus (E10-E14)	755	4.6	119.1
5	Chronic lower respiratory diseases (J40-J47)	610	3.7	96.2
6	Alzheimer's disease (G30)	524	3.2	82.6
7	Influenza and pneumonia (J09-J18)	522	3.2	82.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	461	2.8	72.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	426	2.6	67.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	339	2.1	53.5
11	Parkinson's disease (G20-G21)	327	2.0	51.6
12	Septicemia (A40-A41)	204	1.3	32.2
13	Pneumonitis due to solids and liquids (J69)	131	0.8	20.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	123	0.8	19.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	113	0.7	17.8
...	All other causes (residual)	2,426	14.9	382.6
Non-Hispanic, Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	21,077	100.0	8,481.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,531	26.2	2,225.6
2	Malignant neoplasms (C00-C97)	2,725	12.9	1,096.5
3	Cerebrovascular diseases (I60-I69)	1,989	9.4	800.4
4	Alzheimer's disease (G30)	1,752	8.3	705.0
5	Influenza and pneumonia (J09-J18)	1,075	5.1	432.6
6	Chronic lower respiratory diseases (J40-J47)	799	3.8	321.5
7	Diabetes mellitus (E10-E14)	720	3.4	289.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	626	3.0	251.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	511	2.4	205.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	460	2.2	185.1
11	Parkinson's disease (G20-G21)	285	1.4	114.7
12	Pneumonitis due to solids and liquids (J69)	251	1.2	101.0
13	Septicemia (A40-A41)	214	1.0	86.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	141	0.7	56.7
15	Nutritional deficiencies (E40-E64)	89	0.4	35.8
...	All other causes (residual)	3,909	18.5	1,572.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, all ages				
...	All causes	34,892	100.0	374.9
1	Malignant neoplasms (C00-C97)	8,978	25.7	96.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,840	22.5	84.2
3	Cerebrovascular diseases (I60-I69)	2,268	6.5	24.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,929	5.5	20.7
5	Diabetes mellitus (E10-E14)	1,414	4.1	15.2
6	Chronic lower respiratory diseases (J40-J47)	1,134	3.3	12.2
7	Influenza and pneumonia (J09-J18)	1,120	3.2	12.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	977	2.8	10.5
9	Alzheimer's disease (G30)	709	2.0	7.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	665	1.9	7.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	545	1.6	5.9
12	Parkinson's disease (G20-G21)	440	1.3	4.7
13	Septicemia (A40-A41)	393	1.1	4.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	388	1.1	4.2
15	Certain conditions originating in the perinatal period (P00-P96)	306	0.9	3.3
...	All other causes (residual)	5,786	16.6	62.2
Non-Hispanic, Asian or Pacific Islander, male, 1-4 years				
...	All causes	80	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	36.3	6.0
2	Malignant neoplasms (C00-C97)	9	11.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	11.3	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.5	*
4	Cerebrovascular diseases (I60-I69)	2	2.5	*
4	Influenza and pneumonia (J09-J18)	2	2.5	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.5	*
9	Septicemia (A40-A41)	1	1.3	*
9	Diabetes mellitus (E10-E14)	1	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.3	*
9	Pneumonitis due to solids and liquids (J69)	1	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.3	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.3	*
...	All other causes (residual)	17	21.3	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 5-14 years				
...	All causes	118	100.0	9.9
1	Malignant neoplasms (C00-C97)	32	27.1	2.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	18.6	1.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	9.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	8.5	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	3.4	*
7	Septicemia (A40-A41)	3	2.5	*
7	Cerebrovascular diseases (I60-I69)	3	2.5	*
7	Influenza and pneumonia (J09-J18)	3	2.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	2	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
11	Diabetes mellitus (E10-E14)	1	0.8	*
...	All other causes (residual)	21	17.8	1.8
Non-Hispanic, Asian or Pacific Islander, male, 15-24 years				
...	All causes	696	100.0	52.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	34.6	18.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	223	32.0	16.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	7.3	3.9
4	Malignant neoplasms (C00-C97)	44	6.3	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.7	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.1	*
7	Chronic lower respiratory diseases (J40-J47)	6	0.9	*
8	Cerebrovascular diseases (I60-I69)	4	0.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
9	Aortic aneurysm and dissection (I71)	3	0.4	*
11	Septicemia (A40-A41)	2	0.3	*
11	Diabetes mellitus (E10-E14)	2	0.3	*
13	Anemias (D50-D64)	1	0.1	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.1	*
13	Influenza and pneumonia (J09-J18)	1	0.1	*
13	Pneumonitis due to solids and liquids (J69)	1	0.1	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.1	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.1	*
13	Legal intervention (Y35,Y89.0)	1	0.1	*
...	All other causes (residual)	76	10.9	5.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 25-34 years				
...	All causes	944	100.0	58.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	274	29.0	17.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	196	20.8	12.1
3	Malignant neoplasms (C00-C97)	98	10.4	6.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	8.9	5.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	6.5	3.8
6	Cerebrovascular diseases (I60-I69)	18	1.9	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	18	1.9	*
8	Influenza and pneumonia (J09-J18)	10	1.1	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.8	*
10	Septicemia (A40-A41)	7	0.7	*
10	Diabetes mellitus (E10-E14)	7	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
15	Aortic aneurysm and dissection (I71)	4	0.4	*
15	Legal intervention (Y35,Y89.0)	4	0.4	*
...	All other causes (residual)	137	14.5	8.5
Non-Hispanic, Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,426	100.0	99.4
1	Malignant neoplasms (C00-C97)	278	19.5	19.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	276	19.4	19.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	237	16.6	16.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	172	12.1	12.0
5	Cerebrovascular diseases (I60-I69)	72	5.0	5.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	48	3.4	3.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	2.5	2.5
8	Diabetes mellitus (E10-E14)	29	2.0	2.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	16	1.1	*
9	Influenza and pneumonia (J09-J18)	16	1.1	*
11	Septicemia (A40-A41)	14	1.0	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.8	*
12	Aortic aneurysm and dissection (I71)	12	0.8	*
14	Viral hepatitis (B15-B19)	9	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.6	*
...	All other causes (residual)	183	12.8	12.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,593	100.0	212.0
1	Malignant neoplasms (C00-C97)	758	29.2	62.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	583	22.5	47.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	232	8.9	19.0
4	Cerebrovascular diseases (I60-I69)	151	5.8	12.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	136	5.2	11.1
6	Diabetes mellitus (E10-E14)	109	4.2	8.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	79	3.0	6.5
8	Influenza and pneumonia (J09-J18)	44	1.7	3.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	1.3	2.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	1.3	2.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.1	2.4
12	Septicemia (A40-A41)	26	1.0	2.1
13	Viral hepatitis (B15-B19)	24	0.9	2.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.8	1.8
15	Chronic lower respiratory diseases (J40-J47)	17	0.7	*
...	All other causes (residual)	314	12.1	25.7
Non-Hispanic, Asian or Pacific Islander, male, 55-64 years				
...	All causes	4,999	100.0	523.2
1	Malignant neoplasms (C00-C97)	1,804	36.1	188.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,183	23.7	123.8
3	Cerebrovascular diseases (I60-I69)	289	5.8	30.2
4	Diabetes mellitus (E10-E14)	258	5.2	27.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	214	4.3	22.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	122	2.4	12.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	101	2.0	10.6
8	Influenza and pneumonia (J09-J18)	82	1.6	8.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	1.6	8.2
10	Chronic lower respiratory diseases (J40-J47)	76	1.5	8.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	58	1.2	6.1
12	Septicemia (A40-A41)	53	1.1	5.5
13	Viral hepatitis (B15-B19)	52	1.0	5.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	0.7	3.7
15	Aortic aneurysm and dissection (I71)	24	0.5	2.5
...	All other causes (residual)	570	11.4	59.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 65-74 years				
...	All causes	6,793	100.0	1,134.8
1	Malignant neoplasms (C00-C97)	2,342	34.5	391.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,592	23.4	266.0
3	Cerebrovascular diseases (I60-I69)	440	6.5	73.5
4	Diabetes mellitus (E10-E14)	342	5.0	57.1
5	Chronic lower respiratory diseases (J40-J47)	202	3.0	33.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	2.9	32.6
7	Influenza and pneumonia (J09-J18)	171	2.5	28.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	1.9	22.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	1.4	16.2
10	Septicemia (A40-A41)	81	1.2	13.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	80	1.2	13.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	1.0	11.2
13	Parkinson's disease (G20-G21)	66	1.0	11.0
14	Aortic aneurysm and dissection (I71)	49	0.7	8.2
15	Viral hepatitis (B15-B19)	40	0.6	6.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.6	6.7
...	All other causes (residual)	857	12.6	143.2
Non-Hispanic, Asian or Pacific Islander, male, 75-84 years				
...	All causes	8,370	100.0	3,034.5
1	Malignant neoplasms (C00-C97)	2,306	27.6	836.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,890	22.6	685.2
3	Cerebrovascular diseases (I60-I69)	616	7.4	223.3
4	Diabetes mellitus (E10-E14)	394	4.7	142.8
5	Chronic lower respiratory diseases (J40-J47)	381	4.6	138.1
6	Influenza and pneumonia (J09-J18)	292	3.5	105.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	261	3.1	94.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	219	2.6	79.4
9	Parkinson's disease (G20-G21)	205	2.4	74.3
10	Alzheimer's disease (G30)	183	2.2	66.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	132	1.6	47.9
12	Septicemia (A40-A41)	110	1.3	39.9
13	Pneumonitis due to solids and liquids (J69)	82	1.0	29.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.8	24.3
15	Aortic aneurysm and dissection (I71)	41	0.5	14.9
...	All other causes (residual)	1,191	14.2	431.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 85 years and over				
...	All causes	8,339	100.0	9,112.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,186	26.2	2,388.8
2	Malignant neoplasms (C00-C97)	1,306	15.7	1,427.2
3	Cerebrovascular diseases (I60-I69)	667	8.0	728.9
4	Influenza and pneumonia (J09-J18)	497	6.0	543.1
5	Alzheimer's disease (G30)	492	5.9	537.6
6	Chronic lower respiratory diseases (J40-J47)	438	5.3	478.6
7	Diabetes mellitus (E10-E14)	271	3.2	296.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	210	2.5	229.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	2.5	229.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	190	2.3	207.6
11	Parkinson's disease (G20-G21)	155	1.9	169.4
12	Pneumonitis due to solids and liquids (J69)	146	1.8	159.5
13	Septicemia (A40-A41)	92	1.1	100.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.8	68.8
15	Aortic aneurysm and dissection (I71)	43	0.5	47.0
...	All other causes (residual)	1,373	16.5	1,500.4
Non-Hispanic, Asian or Pacific Islander, female, all ages				
...	All causes	33,343	100.0	327.8
1	Malignant neoplasms (C00-C97)	8,740	26.2	85.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,679	20.0	65.7
3	Cerebrovascular diseases (I60-I69)	2,949	8.8	29.0
4	Alzheimer's disease (G30)	1,654	5.0	16.3
5	Diabetes mellitus (E10-E14)	1,283	3.8	12.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,172	3.5	11.5
7	Influenza and pneumonia (J09-J18)	1,012	3.0	9.9
8	Chronic lower respiratory diseases (J40-J47)	780	2.3	7.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	735	2.2	7.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	683	2.0	6.7
11	Septicemia (A40-A41)	393	1.2	3.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	386	1.2	3.8
13	Parkinson's disease (G20-G21)	309	0.9	3.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	244	0.7	2.4
15	Certain conditions originating in the perinatal period (P00-P96)	221	0.7	2.2
...	All other causes (residual)	6,103	18.3	60.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 1-4 years				
...	All causes	80	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	17.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	15.0	*
3	Malignant neoplasms (C00-C97)	11	13.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	10.0	*
5	Septicemia (A40-A41)	4	5.0	*
6	Cerebrovascular diseases (I60-I69)	2	2.5	*
7	Diabetes mellitus (E10-E14)	1	1.3	*
7	Meningitis (G00,G03)	1	1.3	*
7	Influenza and pneumonia (J09-J18)	1	1.3	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.3	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.3	*
...	All other causes (residual)	23	28.8	5.0
Non-Hispanic, Asian or Pacific Islander, female, 5-14 years				
...	All causes	100	100.0	8.6
1	Malignant neoplasms (C00-C97)	28	28.0	2.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	15.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	7.0	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	5.0	*
5	Influenza and pneumonia (J09-J18)	4	4.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.0	*
6	Cerebrovascular diseases (I60-I69)	3	3.0	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	3.0	*
9	Septicemia (A40-A41)	2	2.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	2	2.0	*
11	Scarlet fever and erysipelas (A38,A46)	1	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
11	Meningitis (G00,G03)	1	1.0	*
11	Diseases of appendix (K35-K38)	1	1.0	*
...	All other causes (residual)	24	24.0	2.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 15-24 years				
...	All causes	275	100.0	21.4
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	29.8	6.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	24.0	5.1
3	Malignant neoplasms (C00-C97)	29	10.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	6.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	5.5	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	6	2.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	2.2	*
8	Cerebrovascular diseases (I60-I69)	4	1.5	*
9	Influenza and pneumonia (J09-J18)	2	0.7	*
10	Tuberculosis (A16-A19)	1	0.4	*
10	Septicemia (A40-A41)	1	0.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
10	Diabetes mellitus (E10-E14)	1	0.4	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
10	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.4	*
...	All other causes (residual)	40	14.5	3.1
Non-Hispanic, Asian or Pacific Islander, female, 25-34 years				
...	All causes	457	100.0	26.6
1	Malignant neoplasms (C00-C97)	101	22.1	5.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	21.4	5.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	14.9	4.0
4	Pregnancy, childbirth and the puerperium (O00-O99)	23	5.0	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.8	1.3
6	Cerebrovascular diseases (I60-I69)	17	3.7	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	3.5	*
8	Septicemia (A40-A41)	6	1.3	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	5	1.1	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.1	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.9	*
11	Diabetes mellitus (E10-E14)	4	0.9	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.9	*
14	Influenza and pneumonia (J09-J18)	3	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
...	All other causes (residual)	78	17.1	4.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 35-44 years				
...	All causes	877	100.0	53.8
1	Malignant neoplasms (C00-C97)	415	47.3	25.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	9.0	4.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	7.8	4.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	6.4	3.4
5	Cerebrovascular diseases (I60-I69)	51	5.8	3.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	2.6	1.4
7	Pregnancy, childbirth and the puerperium (O00-O99)	19	2.2	*
8	Diabetes mellitus (E10-E14)	15	1.7	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	9	1.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.9	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.8	*
11	Influenza and pneumonia (J09-J18)	7	0.8	*
13	Anemias (D50-D64)	6	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
...	All other causes (residual)	106	12.1	6.5
Non-Hispanic, Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,773	100.0	127.1
1	Malignant neoplasms (C00-C97)	939	53.0	67.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	7.8	10.0
3	Cerebrovascular diseases (I60-I69)	135	7.6	9.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	4.7	6.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	3.8	4.9
6	Diabetes mellitus (E10-E14)	52	2.9	3.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	31	1.7	2.2
8	Septicemia (A40-A41)	23	1.3	1.6
9	Influenza and pneumonia (J09-J18)	21	1.2	1.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.0	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.7	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	11	0.6	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	7	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.4	*
...	All other causes (residual)	204	11.5	14.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 55-64 years				
...	All causes	3,509	100.0	306.5
1	Malignant neoplasms (C00-C97)	1,772	50.5	154.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	443	12.6	38.7
3	Cerebrovascular diseases (I60-I69)	225	6.4	19.7
4	Diabetes mellitus (E10-E14)	139	4.0	12.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	3.6	11.1
6	Influenza and pneumonia (J09-J18)	56	1.6	4.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	1.5	4.7
8	Chronic lower respiratory diseases (J40-J47)	52	1.5	4.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	1.4	4.2
10	Septicemia (A40-A41)	41	1.2	3.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.1	3.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	0.9	2.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.7	2.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	0.7	2.0
15	Viral hepatitis (B15-B19)	19	0.5	*
...	All other causes (residual)	414	11.8	36.2
Non-Hispanic, Asian or Pacific Islander, female, 65-74 years				
...	All causes	5,232	100.0	699.6
1	Malignant neoplasms (C00-C97)	2,013	38.5	269.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	909	17.4	121.6
3	Cerebrovascular diseases (I60-I69)	411	7.9	55.0
4	Diabetes mellitus (E10-E14)	261	5.0	34.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	3.6	25.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	122	2.3	16.3
7	Chronic lower respiratory diseases (J40-J47)	118	2.3	15.8
8	Influenza and pneumonia (J09-J18)	109	2.1	14.6
9	Septicemia (A40-A41)	94	1.8	12.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	59	1.1	7.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	55	1.1	7.4
12	Parkinson's disease (G20-G21)	49	0.9	6.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.8	5.8
13	Alzheimer's disease (G30)	43	0.8	5.8
15	Viral hepatitis (B15-B19)	30	0.6	4.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	0.6	4.0
...	All other causes (residual)	696	13.3	93.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 75-84 years				
...	All causes	7,904	100.0	2,206.6
1	Malignant neoplasms (C00-C97)	2,012	25.5	561.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,711	21.6	477.7
3	Cerebrovascular diseases (I60-I69)	778	9.8	217.2
4	Diabetes mellitus (E10-E14)	361	4.6	100.8
5	Alzheimer's disease (G30)	341	4.3	95.2
6	Influenza and pneumonia (J09-J18)	230	2.9	64.2
7	Chronic lower respiratory diseases (J40-J47)	229	2.9	63.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	207	2.6	57.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	207	2.6	57.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	2.5	55.8
11	Parkinson's disease (G20-G21)	122	1.5	34.1
12	Septicemia (A40-A41)	94	1.2	26.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	75	0.9	20.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	56	0.7	15.6
15	Pneumonitis due to solids and liquids (J69)	49	0.6	13.7
...	All other causes (residual)	1,232	15.6	343.9
Non-Hispanic, Asian or Pacific Islander, female, 85 years and over				
...	All causes	12,738	100.0	8,113.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,345	26.3	2,130.5
2	Malignant neoplasms (C00-C97)	1,419	11.1	903.8
3	Cerebrovascular diseases (I60-I69)	1,322	10.4	842.0
4	Alzheimer's disease (G30)	1,260	9.9	802.5
5	Influenza and pneumonia (J09-J18)	578	4.5	368.1
6	Diabetes mellitus (E10-E14)	449	3.5	286.0
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	416	3.3	265.0
8	Chronic lower respiratory diseases (J40-J47)	361	2.8	229.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	301	2.4	191.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	270	2.1	172.0
11	Parkinson's disease (G20-G21)	130	1.0	82.8
12	Septicemia (A40-A41)	122	1.0	77.7
13	Pneumonitis due to solids and liquids (J69)	105	0.8	66.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.6	49.7
15	Nutritional deficiencies (E40-E64)	57	0.4	36.3
...	All other causes (residual)	2,525	19.8	1,608.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016