

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, all ages				
...	All causes	2,744,248	100.0	849.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	635,260	23.1	196.6
2	Malignant neoplasms (C00-C97)	598,038	21.8	185.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161,374	5.9	49.9
4	Chronic lower respiratory diseases (J40-J47)	154,596	5.6	47.8
5	Cerebrovascular diseases (I60-I69)	142,142	5.2	44.0
6	Alzheimer's disease (G30)	116,103	4.2	35.9
7	Diabetes mellitus (E10-E14)	80,058	2.9	24.8
8	Influenza and pneumonia (J09-J18)	51,537	1.9	15.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,046	1.8	15.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44,965	1.6	13.9
11	Septicemia (A40-A41)	40,613	1.5	12.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	40,545	1.5	12.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33,246	1.2	10.3
14	Parkinson's disease (G20-G21)	29,697	1.1	9.2
15	Pneumonitis due to solids and liquids (J69)	19,715	0.7	6.1
...	All other causes (residual)	546,313	19.9	169.1
All races and origins, both sexes, 1-4 years				
...	All causes	4,045	100.0	25.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,261	31.2	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	433	10.7	2.7
3	Malignant neoplasms (C00-C97)	377	9.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	8.4	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	118	2.9	0.7
6	Influenza and pneumonia (J09-J18)	103	2.5	0.6
7	Septicemia (A40-A41)	70	1.7	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	60	1.5	0.4
9	Cerebrovascular diseases (I60-I69)	55	1.4	0.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	1.2	0.3
12	Anemias (D50-D64)	20	0.5	0.1
13	Acute bronchitis and bronchiolitis (J20-J21)	18	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
15	Meningitis (G00,G03)	14	0.3	*
...	All other causes (residual)	1,062	26.3	6.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 5-14 years				
...	All causes	5,503	100.0	13.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,634	29.7	4.0
2	Malignant neoplasms (C00-C97)	880	16.0	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	443	8.1	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	349	6.3	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	286	5.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	188	3.4	0.5
7	Chronic lower respiratory diseases (J40-J47)	143	2.6	0.3
8	Cerebrovascular diseases (I60-I69)	88	1.6	0.2
9	Influenza and pneumonia (J09-J18)	87	1.6	0.2
10	Septicemia (A40-A41)	71	1.3	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	1.1	0.1
12	Diabetes mellitus (E10-E14)	26	0.5	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	26	0.5	0.1
14	Anemias (D50-D64)	24	0.4	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	23	0.4	0.1
...	All other causes (residual)	1,176	21.4	2.9
All races and origins, both sexes, 15-24 years				
...	All causes	32,575	100.0	74.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,895	42.7	31.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,723	17.6	13.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5,172	15.9	11.9
4	Malignant neoplasms (C00-C97)	1,431	4.4	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	949	2.9	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	388	1.2	0.9
7	Diabetes mellitus (E10-E14)	211	0.6	0.5
8	Chronic lower respiratory diseases (J40-J47)	206	0.6	0.5
9	Influenza and pneumonia (J09-J18)	189	0.6	0.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	184	0.6	0.4
11	Cerebrovascular diseases (I60-I69)	148	0.5	0.3
12	Septicemia (A40-A41)	143	0.4	0.3
13	Anemias (D50-D64)	78	0.2	0.2
14	Legal intervention (Y35,Y89.0)	77	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.2	0.2
...	All other causes (residual)	3,711	11.4	8.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 25-44 years				
...	All causes	135,408	100.0	159.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44,959	33.2	52.8
2	Malignant neoplasms (C00-C97)	14,694	10.9	17.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14,396	10.6	16.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,922	10.3	16.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,745	6.5	10.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,776	2.8	4.4
7	Diabetes mellitus (E10-E14)	2,841	2.1	3.3
8	Cerebrovascular diseases (I60-I69)	2,426	1.8	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,517	1.1	1.8
10	Influenza and pneumonia (J09-J18)	1,346	1.0	1.6
11	Septicemia (A40-A41)	1,316	1.0	1.5
12	Chronic lower respiratory diseases (J40-J47)	1,045	0.8	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,008	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	857	0.6	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	823	0.6	1.0
...	All other causes (residual)	21,737	16.1	25.5
All races and origins, both sexes, 45-64 years				
...	All causes	539,961	100.0	640.9
1	Malignant neoplasms (C00-C97)	157,655	29.2	187.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	112,637	20.9	133.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45,237	8.4	53.7
4	Chronic lower respiratory diseases (J40-J47)	22,117	4.1	26.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	21,812	4.0	25.9
6	Diabetes mellitus (E10-E14)	20,518	3.8	24.4
7	Cerebrovascular diseases (I60-I69)	17,663	3.3	21.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16,196	3.0	19.2
9	Septicemia (A40-A41)	8,413	1.6	10.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,790	1.4	9.2
11	Influenza and pneumonia (J09-J18)	7,167	1.3	8.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,347	1.0	6.3
13	Viral hepatitis (B15-B19)	4,158	0.8	4.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,634	0.7	4.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,559	0.7	4.2
...	All other causes (residual)	86,058	15.9	102.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 65 years and over				
...	All causes	2,003,458	100.0	4,068.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	507,118	25.3	1,029.8
2	Malignant neoplasms (C00-C97)	422,927	21.1	858.8
3	Chronic lower respiratory diseases (J40-J47)	131,002	6.5	266.0
4	Cerebrovascular diseases (I60-I69)	121,630	6.1	247.0
5	Alzheimer's disease (G30)	114,883	5.7	233.3
6	Diabetes mellitus (E10-E14)	56,452	2.8	114.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53,141	2.7	107.9
8	Influenza and pneumonia (J09-J18)	42,479	2.1	86.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41,095	2.1	83.5
10	Septicemia (A40-A41)	30,405	1.5	61.7
11	Parkinson's disease (G20-G21)	28,929	1.4	58.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27,283	1.4	55.4
13	Pneumonitis due to solids and liquids (J69)	17,281	0.9	35.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	14,903	0.7	30.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13,064	0.7	26.5
...	All other causes (residual)	380,866	19.0	773.4
All races and origins, male, all ages				
...	All causes	1,400,232	100.0	880.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	339,265	24.2	213.3
2	Malignant neoplasms (C00-C97)	314,571	22.5	197.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103,864	7.4	65.3
4	Chronic lower respiratory diseases (J40-J47)	73,045	5.2	45.9
5	Cerebrovascular diseases (I60-I69)	59,355	4.2	37.3
6	Diabetes mellitus (E10-E14)	43,763	3.1	27.5
7	Alzheimer's disease (G30)	35,372	2.5	22.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34,727	2.5	21.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	25,818	1.8	16.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,399	1.8	16.0
11	Influenza and pneumonia (J09-J18)	25,011	1.8	15.7
12	Septicemia (A40-A41)	19,678	1.4	12.4
13	Parkinson's disease (G20-G21)	17,877	1.3	11.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15,467	1.1	9.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,598	1.0	9.2
...	All other causes (residual)	252,422	18.0	158.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 1-4 years				
...	All causes	2,256	100.0	27.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	776	34.4	9.5
2	Malignant neoplasms (C00-C97)	206	9.1	2.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	206	9.1	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	176	7.8	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	2.7	0.8
6	Influenza and pneumonia (J09-J18)	53	2.3	0.6
7	Septicemia (A40-A41)	37	1.6	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	35	1.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	34	1.5	0.4
10	Cerebrovascular diseases (I60-I69)	28	1.2	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.0	0.3
12	Acute bronchitis and bronchiolitis (J20-J21)	14	0.6	*
13	Anemias (D50-D64)	7	0.3	*
13	Meningitis (G00,G03)	7	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (residual)	586	26.0	7.2
All races and origins, male, 5-14 years				
...	All causes	3,141	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	990	31.5	4.7
2	Malignant neoplasms (C00-C97)	490	15.6	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	8.7	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	187	6.0	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	167	5.3	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	3.2	0.5
7	Chronic lower respiratory diseases (J40-J47)	86	2.7	0.4
8	Cerebrovascular diseases (I60-I69)	55	1.8	0.3
9	Influenza and pneumonia (J09-J18)	49	1.6	0.2
10	Septicemia (A40-A41)	37	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.2	0.2
12	Anemias (D50-D64)	16	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	14	0.4	*
14	Diabetes mellitus (E10-E14)	12	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.4	*
...	All other causes (residual)	618	19.7	3.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 15-24 years				
...	All causes	24,013	100.0	107.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,241	42.6	45.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,575	19.1	20.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,479	18.7	20.1
4	Malignant neoplasms (C00-C97)	902	3.8	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	629	2.6	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	231	1.0	1.0
7	Chronic lower respiratory diseases (J40-J47)	127	0.5	0.6
8	Diabetes mellitus (E10-E14)	121	0.5	0.5
9	Influenza and pneumonia (J09-J18)	105	0.4	0.5
10	Cerebrovascular diseases (I60-I69)	92	0.4	0.4
11	Septicemia (A40-A41)	78	0.3	0.3
12	Legal intervention (Y35,Y89.0)	75	0.3	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.2	0.2
14	Anemias (D50-D64)	45	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.2	0.2
...	All other causes (residual)	2,229	9.3	10.0
All races and origins, male, 25-44 years				
...	All causes	89,472	100.0	209.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,718	36.6	76.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,181	12.5	26.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,451	10.6	22.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,248	8.1	17.0
5	Malignant neoplasms (C00-C97)	6,450	7.2	15.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,326	2.6	5.4
7	Diabetes mellitus (E10-E14)	1,694	1.9	4.0
8	Cerebrovascular diseases (I60-I69)	1,391	1.6	3.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,047	1.2	2.4
10	Influenza and pneumonia (J09-J18)	735	0.8	1.7
11	Septicemia (A40-A41)	670	0.7	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	561	0.6	1.3
13	Chronic lower respiratory diseases (J40-J47)	558	0.6	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	449	0.5	1.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	355	0.4	0.8
...	All other causes (residual)	12,638	14.1	29.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 45-64 years				
...	All causes	327,470	100.0	796.7
1	Malignant neoplasms (C00-C97)	83,696	25.6	203.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,012	23.8	189.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31,110	9.5	75.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	14,599	4.5	35.5
5	Diabetes mellitus (E10-E14)	12,760	3.9	31.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,943	3.6	29.1
7	Chronic lower respiratory diseases (J40-J47)	10,995	3.4	26.7
8	Cerebrovascular diseases (I60-I69)	10,138	3.1	24.7
9	Septicemia (A40-A41)	4,549	1.4	11.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,505	1.4	11.0
11	Influenza and pneumonia (J09-J18)	3,997	1.2	9.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,409	1.0	8.3
13	Viral hepatitis (B15-B19)	2,855	0.9	6.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2,719	0.8	6.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,658	0.8	6.5
...	All other causes (residual)	49,525	15.1	120.5
All races and origins, male, 65 years and over				
...	All causes	940,916	100.0	4,317.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250,840	26.7	1,151.0
2	Malignant neoplasms (C00-C97)	222,788	23.7	1,022.3
3	Chronic lower respiratory diseases (J40-J47)	61,221	6.5	280.9
4	Cerebrovascular diseases (I60-I69)	47,569	5.1	218.3
5	Alzheimer's disease (G30)	34,855	3.7	159.9
6	Diabetes mellitus (E10-E14)	29,171	3.1	133.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,301	2.9	125.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,266	2.2	93.0
9	Influenza and pneumonia (J09-J18)	19,966	2.1	91.6
10	Parkinson's disease (G20-G21)	17,373	1.8	79.7
11	Septicemia (A40-A41)	14,195	1.5	65.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,820	1.1	49.6
13	Pneumonitis due to solids and liquids (J69)	9,390	1.0	43.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	8,863	0.9	40.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,863	0.7	31.5
...	All other causes (residual)	159,435	16.9	731.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, all ages				
...	All causes	1,344,016	100.0	819.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	295,995	22.0	180.4
2	Malignant neoplasms (C00-C97)	283,467	21.1	172.8
3	Cerebrovascular diseases (I60-I69)	82,787	6.2	50.5
4	Chronic lower respiratory diseases (J40-J47)	81,551	6.1	49.7
5	Alzheimer's disease (G30)	80,731	6.0	49.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57,510	4.3	35.1
7	Diabetes mellitus (E10-E14)	36,295	2.7	22.1
8	Influenza and pneumonia (J09-J18)	26,526	2.0	16.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,647	1.8	15.0
10	Septicemia (A40-A41)	20,935	1.6	12.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,648	1.4	11.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	14,727	1.1	9.0
13	Parkinson's disease (G20-G21)	11,820	0.9	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,238	0.8	6.2
15	Pneumonitis due to solids and liquids (J69)	8,853	0.7	5.4
...	All other causes (residual)	289,286	21.5	176.3
All races and origins, female, 1-4 years				
...	All causes	1,789	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	485	27.1	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	12.7	2.9
3	Malignant neoplasms (C00-C97)	171	9.6	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	9.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	3.1	0.7
6	Influenza and pneumonia (J09-J18)	50	2.8	0.6
7	Septicemia (A40-A41)	33	1.8	0.4
8	Cerebrovascular diseases (I60-I69)	27	1.5	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.5	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	25	1.4	0.3
11	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
12	Anemias (D50-D64)	13	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.6	*
14	Meningitis (G00,G03)	7	0.4	*
15	Diabetes mellitus (E10-E14)	5	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (residual)	469	26.2	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 5-14 years				
...	All causes	2,362	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	644	27.3	3.2
2	Malignant neoplasms (C00-C97)	390	16.5	1.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	171	7.2	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	6.9	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	5.0	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	89	3.8	0.4
7	Chronic lower respiratory diseases (J40-J47)	57	2.4	0.3
8	Influenza and pneumonia (J09-J18)	38	1.6	0.2
9	Septicemia (A40-A41)	34	1.4	0.2
10	Cerebrovascular diseases (I60-I69)	33	1.4	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.9	0.1
12	Diabetes mellitus (E10-E14)	14	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	12	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.4	*
...	All other causes (residual)	556	23.5	2.8
All races and origins, female, 15-24 years				
...	All causes	8,562	100.0	40.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,654	42.7	17.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,148	13.4	5.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	693	8.1	3.3
4	Malignant neoplasms (C00-C97)	529	6.2	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	320	3.7	1.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	184	2.1	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	157	1.8	0.7
8	Diabetes mellitus (E10-E14)	90	1.1	0.4
9	Influenza and pneumonia (J09-J18)	84	1.0	0.4
10	Chronic lower respiratory diseases (J40-J47)	79	0.9	0.4
11	Septicemia (A40-A41)	65	0.8	0.3
12	Cerebrovascular diseases (I60-I69)	56	0.7	0.3
13	Anemias (D50-D64)	33	0.4	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	30	0.4	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.3	0.1
...	All other causes (residual)	1,411	16.5	6.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 25-44 years				
...	All causes	45,936	100.0	108.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,241	26.6	28.9
2	Malignant neoplasms (C00-C97)	8,244	17.9	19.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,471	9.7	10.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,215	7.0	7.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,497	3.3	3.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,450	3.2	3.4
7	Diabetes mellitus (E10-E14)	1,147	2.5	2.7
8	Cerebrovascular diseases (I60-I69)	1,035	2.3	2.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	823	1.8	1.9
10	Septicemia (A40-A41)	646	1.4	1.5
11	Influenza and pneumonia (J09-J18)	611	1.3	1.4
12	Chronic lower respiratory diseases (J40-J47)	487	1.1	1.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	470	1.0	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	447	1.0	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	408	0.9	1.0
...	All other causes (residual)	8,744	19.0	20.6
All races and origins, female, 45-64 years				
...	All causes	212,491	100.0	492.5
1	Malignant neoplasms (C00-C97)	73,959	34.8	171.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,625	16.3	80.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,127	6.6	32.7
4	Chronic lower respiratory diseases (J40-J47)	11,122	5.2	25.8
5	Diabetes mellitus (E10-E14)	7,758	3.7	18.0
6	Cerebrovascular diseases (I60-I69)	7,525	3.5	17.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	7,213	3.4	16.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,253	2.0	9.9
9	Septicemia (A40-A41)	3,864	1.8	9.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,285	1.5	7.6
11	Influenza and pneumonia (J09-J18)	3,170	1.5	7.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,938	0.9	4.5
13	Viral hepatitis (B15-B19)	1,303	0.6	3.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	949	0.4	2.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	938	0.4	2.2
...	All other causes (residual)	36,462	17.2	84.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 65 years and over				
...	All causes	1,062,542	100.0	3,870.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	256,278	24.1	933.6
2	Malignant neoplasms (C00-C97)	200,139	18.8	729.1
3	Alzheimer's disease (G30)	80,028	7.5	291.5
4	Cerebrovascular diseases (I60-I69)	74,061	7.0	269.8
5	Chronic lower respiratory diseases (J40-J47)	69,781	6.6	254.2
6	Diabetes mellitus (E10-E14)	27,281	2.6	99.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25,840	2.4	94.1
8	Influenza and pneumonia (J09-J18)	22,513	2.1	82.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,829	2.0	75.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16,463	1.5	60.0
11	Septicemia (A40-A41)	16,210	1.5	59.0
12	Parkinson's disease (G20-G21)	11,556	1.1	42.1
13	Pneumonitis due to solids and liquids (J69)	7,891	0.7	28.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,201	0.6	22.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	6,040	0.6	22.0
...	All other causes (residual)	221,431	20.8	806.6
Hispanic, both sexes, all ages				
...	All causes	188,254	100.0	327.6
1	Malignant neoplasms (C00-C97)	39,263	20.9	68.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,799	20.1	65.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,711	8.3	27.3
4	Cerebrovascular diseases (I60-I69)	10,283	5.5	17.9
5	Diabetes mellitus (E10-E14)	8,546	4.5	14.9
6	Alzheimer's disease (G30)	6,833	3.6	11.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,141	3.3	10.7
8	Chronic lower respiratory diseases (J40-J47)	5,287	2.8	9.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,775	2.0	6.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,668	1.9	6.4
11	Influenza and pneumonia (J09-J18)	3,641	1.9	6.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,187	1.7	5.5
13	Septicemia (A40-A41)	2,718	1.4	4.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,488	1.3	4.3
15	Certain conditions originating in the perinatal period (P00-P96)	2,415	1.3	4.2
...	All other causes (residual)	36,499	19.4	63.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 1-4 years				
...	All causes	799	100.0	19.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	29.2	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	12.4	2.4
3	Malignant neoplasms (C00-C97)	89	11.1	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	7.8	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	3.4	0.7
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	2.0	*
6	Influenza and pneumonia (J09-J18)	16	2.0	*
8	Septicemia (A40-A41)	14	1.8	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.5	*
10	Cerebrovascular diseases (I60-I69)	8	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	8	1.0	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
13	Anemias (D50-D64)	3	0.4	*
14	Meningitis (G00,G03)	2	0.3	*
14	Hernia (K40-K46)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	*
...	All other causes (residual)	200	25.0	4.9
Hispanic, both sexes, 5-14 years				
...	All causes	1,164	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	28.4	3.2
2	Malignant neoplasms (C00-C97)	226	19.4	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	7.5	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	5.2	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	4.7	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.3	0.4
7	Cerebrovascular diseases (I60-I69)	29	2.5	0.3
8	Chronic lower respiratory diseases (J40-J47)	22	1.9	0.2
9	Septicemia (A40-A41)	18	1.5	*
10	Influenza and pneumonia (J09-J18)	14	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.0	*
12	Anemias (D50-D64)	7	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Diseases of appendix (K35-K38)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (residual)	254	21.8	2.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-24 years				
...	All causes	5,722	100.0	59.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,527	44.2	26.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	987	17.2	10.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	841	14.7	8.8
4	Malignant neoplasms (C00-C97)	370	6.5	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	120	2.1	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	1.5	0.9
7	Cerebrovascular diseases (I60-I69)	34	0.6	0.4
7	Influenza and pneumonia (J09-J18)	34	0.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	30	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	29	0.5	0.3
11	Diabetes mellitus (E10-E14)	25	0.4	0.3
12	Septicemia (A40-A41)	22	0.4	0.2
13	Legal intervention (Y35,Y89.0)	21	0.4	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.2	*
...	All other causes (residual)	572	10.0	6.0
Hispanic, both sexes, 25-44 years				
...	All causes	18,467	100.0	106.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,804	31.4	33.5
2	Malignant neoplasms (C00-C97)	2,499	13.5	14.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,604	8.7	9.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,540	8.3	8.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,504	8.1	8.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	816	4.4	4.7
7	Cerebrovascular diseases (I60-I69)	434	2.4	2.5
8	Diabetes mellitus (E10-E14)	367	2.0	2.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	223	1.2	1.3
10	Influenza and pneumonia (J09-J18)	209	1.1	1.2
11	Septicemia (A40-A41)	171	0.9	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	143	0.8	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	136	0.7	0.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	0.6	0.7
15	Chronic lower respiratory diseases (J40-J47)	104	0.6	0.6
...	All other causes (residual)	2,800	15.2	16.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-64 years				
...	All causes	45,340	100.0	406.5
1	Malignant neoplasms (C00-C97)	12,499	27.6	112.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,222	18.1	73.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,077	9.0	36.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,276	7.2	29.4
5	Diabetes mellitus (E10-E14)	2,598	5.7	23.3
6	Cerebrovascular diseases (I60-I69)	1,926	4.2	17.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	887	2.0	8.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	877	1.9	7.9
9	Chronic lower respiratory diseases (J40-J47)	719	1.6	6.4
10	Septicemia (A40-A41)	704	1.6	6.3
11	Influenza and pneumonia (J09-J18)	681	1.5	6.1
12	Viral hepatitis (B15-B19)	606	1.3	5.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	484	1.1	4.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	473	1.0	4.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	421	0.9	3.8
...	All other causes (residual)	6,890	15.2	61.8
Hispanic, both sexes, 65 years and over				
...	All causes	111,932	100.0	2,820.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,822	24.9	701.0
2	Malignant neoplasms (C00-C97)	23,560	21.0	593.6
3	Cerebrovascular diseases (I60-I69)	7,827	7.0	197.2
4	Alzheimer's disease (G30)	6,740	6.0	169.8
5	Diabetes mellitus (E10-E14)	5,554	5.0	139.9
6	Chronic lower respiratory diseases (J40-J47)	4,402	3.9	110.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,731	2.4	68.8
8	Influenza and pneumonia (J09-J18)	2,653	2.4	66.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,566	2.3	64.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,038	1.8	51.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,920	1.7	48.4
12	Septicemia (A40-A41)	1,744	1.6	43.9
13	Parkinson's disease (G20-G21)	1,575	1.4	39.7
14	Pneumonitis due to solids and liquids (J69)	798	0.7	20.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	660	0.6	16.6
...	All other causes (residual)	19,342	17.3	487.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	103,532	100.0	356.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,945	20.2	72.2
2	Malignant neoplasms (C00-C97)	20,504	19.8	70.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,510	11.1	39.7
4	Cerebrovascular diseases (I60-I69)	4,798	4.6	16.5
5	Diabetes mellitus (E10-E14)	4,603	4.4	15.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,225	4.1	14.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,944	2.8	10.1
8	Chronic lower respiratory diseases (J40-J47)	2,759	2.7	9.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,657	2.6	9.2
10	Alzheimer's disease (G30)	2,151	2.1	7.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,931	1.9	6.7
12	Influenza and pneumonia (J09-J18)	1,877	1.8	6.5
13	Septicemia (A40-A41)	1,392	1.3	4.8
14	Certain conditions originating in the perinatal period (P00-P96)	1,348	1.3	4.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,197	1.2	4.1
...	All other causes (residual)	18,691	18.1	64.4
Hispanic, male, 1-4 years				
...	All causes	443	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	33.0	7.0
2	Malignant neoplasms (C00-C97)	48	10.8	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	10.2	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	7.4	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.2	*
6	Influenza and pneumonia (J09-J18)	10	2.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.8	*
8	Cerebrovascular diseases (I60-I69)	6	1.4	*
9	Septicemia (A40-A41)	5	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
12	Acute bronchitis and bronchiolitis (J20-J21)	3	0.7	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Hernia (K40-K46)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	108	24.4	5.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-14 years				
...	All causes	641	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	29.5	3.6
2	Malignant neoplasms (C00-C97)	125	19.5	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	6.4	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	4.8	0.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	4.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.4	0.4
7	Cerebrovascular diseases (I60-I69)	18	2.8	*
8	Septicemia (A40-A41)	12	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.7	*
11	Influenza and pneumonia (J09-J18)	8	1.2	*
12	Anemias (D50-D64)	3	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
15	Meningococcal infection (A39)	1	0.2	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (residual)	125	19.5	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,304	100.0	87.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,897	44.1	38.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	869	20.2	17.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	665	15.5	13.5
4	Malignant neoplasms (C00-C97)	249	5.8	5.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	2.0	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	1.2	1.1
7	Cerebrovascular diseases (I60-I69)	22	0.5	0.4
7	Influenza and pneumonia (J09-J18)	22	0.5	0.4
9	Legal intervention (Y35,Y89.0)	21	0.5	0.4
10	Diabetes mellitus (E10-E14)	17	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	17	0.4	*
12	Septicemia (A40-A41)	12	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.1	*
15	Pneumonitis due to solids and liquids (J69)	5	0.1	*
...	All other causes (residual)	354	8.2	7.2
Hispanic, male, 25-44 years				
...	All causes	13,006	100.0	145.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,629	35.6	51.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,312	10.1	14.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,310	10.1	14.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,108	8.5	12.4
5	Malignant neoplasms (C00-C97)	1,060	8.2	11.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	596	4.6	6.6
7	Cerebrovascular diseases (I60-I69)	274	2.1	3.1
8	Diabetes mellitus (E10-E14)	233	1.8	2.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	181	1.4	2.0
10	Influenza and pneumonia (J09-J18)	142	1.1	1.6
11	Septicemia (A40-A41)	97	0.7	1.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	0.6	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	0.5	0.8
14	Legal intervention (Y35,Y89.0)	60	0.5	0.7
15	Chronic lower respiratory diseases (J40-J47)	56	0.4	0.6
...	All other causes (residual)	1,806	13.9	20.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	28,967	100.0	523.0
1	Malignant neoplasms (C00-C97)	6,524	22.5	117.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,982	20.7	108.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,112	10.7	56.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,448	8.5	44.2
5	Diabetes mellitus (E10-E14)	1,588	5.5	28.7
6	Cerebrovascular diseases (I60-I69)	1,188	4.1	21.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	702	2.4	12.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	528	1.8	9.5
9	Septicemia (A40-A41)	443	1.5	8.0
10	Viral hepatitis (B15-B19)	436	1.5	7.9
11	Influenza and pneumonia (J09-J18)	431	1.5	7.8
12	Chronic lower respiratory diseases (J40-J47)	387	1.3	7.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	384	1.3	6.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	338	1.2	6.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	298	1.0	5.4
...	All other causes (residual)	4,178	14.4	75.4
Hispanic, male, 65 years and over				
...	All causes	53,484	100.0	3,112.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,697	25.6	797.2
2	Malignant neoplasms (C00-C97)	12,489	23.4	726.9
3	Cerebrovascular diseases (I60-I69)	3,276	6.1	190.7
4	Diabetes mellitus (E10-E14)	2,763	5.2	160.8
5	Chronic lower respiratory diseases (J40-J47)	2,282	4.3	132.8
6	Alzheimer's disease (G30)	2,110	3.9	122.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,437	2.7	83.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,316	2.5	76.6
9	Influenza and pneumonia (J09-J18)	1,242	2.3	72.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,174	2.2	68.3
11	Parkinson's disease (G20-G21)	878	1.6	51.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	847	1.6	49.3
13	Septicemia (A40-A41)	796	1.5	46.3
14	Pneumonitis due to solids and liquids (J69)	427	0.8	24.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	330	0.6	19.2
...	All other causes (residual)	8,420	15.7	490.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	84,722	100.0	297.7
1	Malignant neoplasms (C00-C97)	18,759	22.1	65.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,854	19.9	59.2
3	Cerebrovascular diseases (I60-I69)	5,485	6.5	19.3
4	Alzheimer's disease (G30)	4,682	5.5	16.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,201	5.0	14.8
6	Diabetes mellitus (E10-E14)	3,943	4.7	13.9
7	Chronic lower respiratory diseases (J40-J47)	2,528	3.0	8.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,916	2.3	6.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,844	2.2	6.5
10	Influenza and pneumonia (J09-J18)	1,764	2.1	6.2
11	Septicemia (A40-A41)	1,326	1.6	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,291	1.5	4.5
13	Certain conditions originating in the perinatal period (P00-P96)	1,067	1.3	3.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	858	1.0	3.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	724	0.9	2.5
...	All other causes (residual)	17,480	20.6	61.4
Hispanic, female, 1-4 years				
...	All causes	356	100.0	17.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	24.4	4.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	15.2	2.7
3	Malignant neoplasms (C00-C97)	41	11.5	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	8.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.7	*
6	Septicemia (A40-A41)	9	2.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	2.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	2.0	*
9	Influenza and pneumonia (J09-J18)	6	1.7	*
10	Anemias (D50-D64)	3	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
12	Cerebrovascular diseases (I60-I69)	2	0.6	*
13	Salmonella infections (A01-A02)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	87	24.4	4.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	523	100.0	10.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	141	27.0	2.8
2	Malignant neoplasms (C00-C97)	101	19.3	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	8.8	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	5.5	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	4.6	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.1	*
7	Cerebrovascular diseases (I60-I69)	11	2.1	*
7	Chronic lower respiratory diseases (J40-J47)	11	2.1	*
9	Septicemia (A40-A41)	6	1.1	*
9	Influenza and pneumonia (J09-J18)	6	1.1	*
11	Anemias (D50-D64)	4	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
13	Nutritional deficiencies (E40-E64)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
13	Peptic ulcer (K25-K28)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	116	22.2	2.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-24 years				
...	All causes	1,418	100.0	30.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	630	44.4	13.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	176	12.4	3.8
3	Malignant neoplasms (C00-C97)	121	8.5	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	118	8.3	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	2.3	0.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	2.3	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	29	2.0	0.6
8	Chronic lower respiratory diseases (J40-J47)	13	0.9	*
9	Cerebrovascular diseases (I60-I69)	12	0.8	*
9	Influenza and pneumonia (J09-J18)	12	0.8	*
11	Septicemia (A40-A41)	10	0.7	*
12	Diabetes mellitus (E10-E14)	8	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.3	*
...	All other causes (residual)	204	14.4	4.4
Hispanic, female, 25-44 years				
...	All causes	5,461	100.0	65.4
1	Malignant neoplasms (C00-C97)	1,439	26.4	17.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,175	21.5	14.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	396	7.3	4.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	294	5.4	3.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	228	4.2	2.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	220	4.0	2.6
7	Cerebrovascular diseases (I60-I69)	160	2.9	1.9
8	Pregnancy, childbirth and the puerperium (O00-O99)	143	2.6	1.7
9	Diabetes mellitus (E10-E14)	134	2.5	1.6
10	Septicemia (A40-A41)	74	1.4	0.9
11	Influenza and pneumonia (J09-J18)	67	1.2	0.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	1.1	0.7
13	Chronic lower respiratory diseases (J40-J47)	48	0.9	0.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	0.8	0.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	42	0.8	0.5
...	All other causes (residual)	934	17.1	11.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-64 years				
...	All causes	16,373	100.0	291.5
1	Malignant neoplasms (C00-C97)	5,975	36.5	106.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,240	13.7	39.9
3	Diabetes mellitus (E10-E14)	1,010	6.2	18.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	965	5.9	17.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	828	5.1	14.7
6	Cerebrovascular diseases (I60-I69)	738	4.5	13.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	349	2.1	6.2
8	Chronic lower respiratory diseases (J40-J47)	332	2.0	5.9
9	Septicemia (A40-A41)	261	1.6	4.6
10	Influenza and pneumonia (J09-J18)	250	1.5	4.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	186	1.1	3.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	185	1.1	3.3
13	Viral hepatitis (B15-B19)	170	1.0	3.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	89	0.5	1.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	0.5	1.5
...	All other causes (residual)	2,712	16.6	48.3
Hispanic, female, 65 years and over				
...	All causes	58,448	100.0	2,597.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,125	24.2	627.6
2	Malignant neoplasms (C00-C97)	11,071	18.9	491.9
3	Alzheimer's disease (G30)	4,630	7.9	205.7
4	Cerebrovascular diseases (I60-I69)	4,551	7.8	202.2
5	Diabetes mellitus (E10-E14)	2,791	4.8	124.0
6	Chronic lower respiratory diseases (J40-J47)	2,120	3.6	94.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,415	2.4	62.9
8	Influenza and pneumonia (J09-J18)	1,411	2.4	62.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,129	1.9	50.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,073	1.8	47.7
11	Septicemia (A40-A41)	948	1.6	42.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	864	1.5	38.4
13	Parkinson's disease (G20-G21)	697	1.2	31.0
14	Pneumonitis due to solids and liquids (J69)	371	0.6	16.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	330	0.6	14.7
...	All other causes (residual)	10,922	18.7	485.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,133,463	100.0	1,059.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	500,631	23.5	248.7
2	Malignant neoplasms (C00-C97)	466,467	21.9	231.7
3	Chronic lower respiratory diseases (J40-J47)	135,268	6.3	67.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121,641	5.7	60.4
5	Cerebrovascular diseases (I60-I69)	107,491	5.0	53.4
6	Alzheimer's disease (G30)	97,779	4.6	48.6
7	Diabetes mellitus (E10-E14)	53,399	2.5	26.5
8	Influenza and pneumonia (J09-J18)	39,748	1.9	19.7
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,531	1.7	18.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,248	1.7	17.5
11	Septicemia (A40-A41)	30,009	1.4	14.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29,432	1.4	14.6
13	Parkinson's disease (G20-G21)	25,826	1.2	12.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23,237	1.1	11.5
15	Pneumonitis due to solids and liquids (J69)	16,302	0.8	8.1
...	All other causes (residual)	414,454	19.4	205.9
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,978	100.0	23.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	680	34.4	8.2
2	Malignant neoplasms (C00-C97)	206	10.4	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	9.7	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	6.1	1.5
5	Influenza and pneumonia (J09-J18)	47	2.4	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	2.3	0.5
7	Septicemia (A40-A41)	34	1.7	0.4
8	Cerebrovascular diseases (I60-I69)	27	1.4	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.2	0.3
10	Chronic lower respiratory diseases (J40-J47)	22	1.1	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	20	1.0	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
15	Meningitis (G00,G03)	5	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (residual)	532	26.9	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,781	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	844	30.3	3.9
2	Malignant neoplasms (C00-C97)	440	15.8	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	311	11.2	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	6.8	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	3.5	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	2.9	0.4
7	Influenza and pneumonia (J09-J18)	44	1.6	0.2
8	Septicemia (A40-A41)	34	1.2	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	1.2	0.2
10	Chronic lower respiratory diseases (J40-J47)	32	1.2	0.1
11	Cerebrovascular diseases (I60-I69)	31	1.1	0.1
12	Diabetes mellitus (E10-E14)	15	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	12	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (residual)	597	21.5	2.7
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	17,515	100.0	72.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,895	50.8	36.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,756	21.4	15.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	766	4.4	3.2
4	Malignant neoplasms (C00-C97)	744	4.2	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	481	2.7	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	210	1.2	0.9
7	Diabetes mellitus (E10-E14)	113	0.6	0.5
8	Influenza and pneumonia (J09-J18)	100	0.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	87	0.5	0.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	79	0.5	0.3
11	Septicemia (A40-A41)	74	0.4	0.3
12	Cerebrovascular diseases (I60-I69)	71	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.2	0.2
14	Pneumonitis due to solids and liquids (J69)	26	0.1	0.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	24	0.1	0.1
...	All other causes (residual)	2,051	11.7	8.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 25-44 years				
...	All causes	83,631	100.0	170.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,227	38.5	65.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,808	12.9	22.0
3	Malignant neoplasms (C00-C97)	8,621	10.3	17.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,802	9.3	15.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,342	2.8	4.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,165	2.6	4.4
7	Diabetes mellitus (E10-E14)	1,484	1.8	3.0
8	Cerebrovascular diseases (I60-I69)	1,152	1.4	2.3
9	Influenza and pneumonia (J09-J18)	779	0.9	1.6
10	Septicemia (A40-A41)	748	0.9	1.5
11	Chronic lower respiratory diseases (J40-J47)	596	0.7	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	558	0.7	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	419	0.5	0.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	370	0.4	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	369	0.4	0.8
...	All other causes (residual)	13,191	15.8	26.9
Non-Hispanic, white, both sexes, 45-64 years				
...	All causes	378,864	100.0	658.3
1	Malignant neoplasms (C00-C97)	113,148	29.9	196.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,247	20.4	134.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33,058	8.7	57.4
4	Chronic lower respiratory diseases (J40-J47)	18,279	4.8	31.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	15,649	4.1	27.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14,152	3.7	24.6
7	Diabetes mellitus (E10-E14)	12,327	3.3	21.4
8	Cerebrovascular diseases (I60-I69)	10,329	2.7	17.9
9	Septicemia (A40-A41)	5,472	1.4	9.5
10	Influenza and pneumonia (J09-J18)	4,848	1.3	8.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,229	1.1	7.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,056	0.8	5.3
13	Viral hepatitis (B15-B19)	2,753	0.7	4.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,683	0.4	2.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,597	0.4	2.8
...	All other causes (residual)	61,037	16.1	106.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65 years and over				
...	All causes	1,638,592	100.0	4,277.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	414,824	25.3	1,083.0
2	Malignant neoplasms (C00-C97)	343,275	20.9	896.2
3	Chronic lower respiratory diseases (J40-J47)	116,235	7.1	303.5
4	Alzheimer's disease (G30)	96,799	5.9	252.7
5	Cerebrovascular diseases (I60-I69)	95,821	5.8	250.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45,335	2.8	118.4
7	Diabetes mellitus (E10-E14)	39,457	2.4	103.0
8	Influenza and pneumonia (J09-J18)	33,869	2.1	88.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30,534	1.9	79.7
10	Parkinson's disease (G20-G21)	25,207	1.5	65.8
11	Septicemia (A40-A41)	23,578	1.4	61.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19,922	1.2	52.0
13	Pneumonitis due to solids and liquids (J69)	14,516	0.9	37.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	11,413	0.7	29.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11,058	0.7	28.9
...	All other causes (residual)	316,749	19.3	826.9
Non-Hispanic, white, male, all ages				
...	All causes	1,077,362	100.0	1,085.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	266,995	24.8	269.0
2	Malignant neoplasms (C00-C97)	247,202	22.9	249.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76,033	7.1	76.6
4	Chronic lower respiratory diseases (J40-J47)	62,945	5.8	63.4
5	Cerebrovascular diseases (I60-I69)	43,713	4.1	44.0
6	Diabetes mellitus (E10-E14)	30,010	2.8	30.2
7	Alzheimer's disease (G30)	29,886	2.8	30.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,110	2.6	28.3
9	Influenza and pneumonia (J09-J18)	19,033	1.8	19.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18,522	1.7	18.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,216	1.7	18.4
12	Parkinson's disease (G20-G21)	15,641	1.5	15.8
13	Septicemia (A40-A41)	14,620	1.4	14.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,906	0.9	10.0
15	Pneumonitis due to solids and liquids (J69)	8,997	0.8	9.1
...	All other causes (residual)	187,533	17.4	189.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,113	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	419	37.6	9.9
2	Malignant neoplasms (C00-C97)	114	10.2	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	8.5	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	5.6	1.5
5	Septicemia (A40-A41)	23	2.1	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.1	0.5
7	Influenza and pneumonia (J09-J18)	21	1.9	0.5
8	Chronic lower respiratory diseases (J40-J47)	14	1.3	*
9	Cerebrovascular diseases (I60-I69)	12	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
14	Meningococcal infection (A39)	2	0.2	*
14	Anemias (D50-D64)	2	0.2	*
14	Meningitis (G00,G03)	2	0.2	*
14	Pneumonitis due to solids and liquids (J69)	2	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.2	*
...	All other causes (residual)	288	25.9	6.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,625	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	523	32.2	4.7
2	Malignant neoplasms (C00-C97)	264	16.2	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	194	11.9	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	6.6	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	3.7	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	2.6	0.4
7	Influenza and pneumonia (J09-J18)	24	1.5	0.2
8	Cerebrovascular diseases (I60-I69)	21	1.3	0.2
9	Septicemia (A40-A41)	19	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
14	Anemias (D50-D64)	5	0.3	*
15	Diabetes mellitus (E10-E14)	4	0.2	*
...	All other causes (residual)	314	19.3	2.8
Non-Hispanic, white, male, 15-24 years				
...	All causes	12,645	100.0	101.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,557	51.9	52.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,042	24.1	24.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	549	4.3	4.4
4	Malignant neoplasms (C00-C97)	464	3.7	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	316	2.5	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	0.9	1.0
7	Diabetes mellitus (E10-E14)	65	0.5	0.5
8	Influenza and pneumonia (J09-J18)	57	0.5	0.5
9	Chronic lower respiratory diseases (J40-J47)	53	0.4	0.4
10	Cerebrovascular diseases (I60-I69)	48	0.4	0.4
11	Septicemia (A40-A41)	43	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.2	0.2
13	Legal intervention (Y35,Y89.0)	20	0.2	0.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	14	0.1	*
15	Pneumonitis due to solids and liquids (J69)	13	0.1	*
...	All other causes (residual)	1,261	10.0	10.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-44 years				
...	All causes	54,815	100.0	221.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23,066	42.1	93.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,309	15.2	33.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,325	9.7	21.5
4	Malignant neoplasms (C00-C97)	3,963	7.2	16.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,506	2.7	6.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,378	2.5	5.6
7	Diabetes mellitus (E10-E14)	901	1.6	3.6
8	Cerebrovascular diseases (I60-I69)	641	1.2	2.6
9	Influenza and pneumonia (J09-J18)	409	0.7	1.6
10	Septicemia (A40-A41)	368	0.7	1.5
11	Chronic lower respiratory diseases (J40-J47)	303	0.6	1.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	280	0.5	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	270	0.5	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	239	0.4	1.0
15	Aortic aneurysm and dissection (I71)	189	0.3	0.8
...	All other causes (residual)	7,668	14.0	30.9
Non-Hispanic, white, male, 45-64 years				
...	All causes	231,021	100.0	814.4
1	Malignant neoplasms (C00-C97)	60,891	26.4	214.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,520	23.6	192.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,232	9.6	78.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,363	4.5	36.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	10,303	4.5	36.3
6	Chronic lower respiratory diseases (J40-J47)	9,055	3.9	31.9
7	Diabetes mellitus (E10-E14)	7,939	3.4	28.0
8	Cerebrovascular diseases (I60-I69)	5,933	2.6	20.9
9	Septicemia (A40-A41)	2,990	1.3	10.5
10	Influenza and pneumonia (J09-J18)	2,628	1.1	9.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,482	1.1	8.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,002	0.9	7.1
13	Viral hepatitis (B15-B19)	1,890	0.8	6.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,159	0.5	4.1
15	Aortic aneurysm and dissection (I71)	1,117	0.5	3.9
...	All other causes (residual)	35,517	15.4	125.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65 years and over				
...	All causes	770,488	100.0	4,483.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	206,692	26.8	1,202.7
2	Malignant neoplasms (C00-C97)	181,490	23.6	1,056.1
3	Chronic lower respiratory diseases (J40-J47)	53,490	6.9	311.3
4	Cerebrovascular diseases (I60-I69)	37,020	4.8	215.4
5	Alzheimer's disease (G30)	29,479	3.8	171.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,876	3.0	133.1
7	Diabetes mellitus (E10-E14)	21,100	2.7	122.8
8	Influenza and pneumonia (J09-J18)	15,850	2.1	92.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,466	2.0	90.0
10	Parkinson's disease (G20-G21)	15,240	2.0	88.7
11	Septicemia (A40-A41)	11,136	1.4	64.8
12	Pneumonitis due to solids and liquids (J69)	7,936	1.0	46.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,751	1.0	45.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	6,823	0.9	39.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,202	0.8	36.1
...	All other causes (residual)	131,937	17.1	767.7
Non-Hispanic, white, female, all ages				
...	All causes	1,056,101	100.0	1,034.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	233,636	22.1	228.9
2	Malignant neoplasms (C00-C97)	219,265	20.8	214.8
3	Chronic lower respiratory diseases (J40-J47)	72,323	6.8	70.8
4	Alzheimer's disease (G30)	67,893	6.4	66.5
5	Cerebrovascular diseases (I60-I69)	63,778	6.0	62.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45,608	4.3	44.7
7	Diabetes mellitus (E10-E14)	23,389	2.2	22.9
8	Influenza and pneumonia (J09-J18)	20,715	2.0	20.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,032	1.6	16.7
10	Septicemia (A40-A41)	15,389	1.5	15.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,331	1.3	13.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,910	1.0	10.7
13	Parkinson's disease (G20-G21)	10,185	1.0	10.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,421	0.8	8.2
15	Pneumonitis due to solids and liquids (J69)	7,305	0.7	7.2
...	All other causes (residual)	226,921	21.5	222.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	865	100.0	21.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	261	30.2	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	11.1	2.4
3	Malignant neoplasms (C00-C97)	92	10.6	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	6.8	1.5
5	Influenza and pneumonia (J09-J18)	26	3.0	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	2.5	0.5
7	Cerebrovascular diseases (I60-I69)	15	1.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.6	*
9	Septicemia (A40-A41)	11	1.3	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	8	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
14	Meningitis (G00,G03)	3	0.3	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (residual)	238	27.5	5.9
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,156	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	27.8	3.0
2	Malignant neoplasms (C00-C97)	176	15.2	1.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	117	10.1	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	7.0	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	3.5	0.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	3.2	0.3
7	Influenza and pneumonia (J09-J18)	20	1.7	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
9	Septicemia (A40-A41)	15	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
11	Diabetes mellitus (E10-E14)	11	1.0	*
12	Cerebrovascular diseases (I60-I69)	10	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (residual)	281	24.3	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,870	100.0	41.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,338	48.0	19.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	714	14.7	6.1
3	Malignant neoplasms (C00-C97)	280	5.7	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	217	4.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	165	3.4	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.9	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	79	1.6	0.7
8	Diabetes mellitus (E10-E14)	48	1.0	0.4
9	Influenza and pneumonia (J09-J18)	43	0.9	0.4
10	Chronic lower respiratory diseases (J40-J47)	34	0.7	0.3
11	Septicemia (A40-A41)	31	0.6	0.3
12	Cerebrovascular diseases (I60-I69)	23	0.5	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.3	*
15	Pneumonitis due to solids and liquids (J69)	13	0.3	*
...	All other causes (residual)	766	15.7	6.5
Non-Hispanic, white, female, 25-44 years				
...	All causes	28,816	100.0	118.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,161	31.8	37.7
2	Malignant neoplasms (C00-C97)	4,658	16.2	19.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,499	8.7	10.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,477	8.6	10.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	964	3.3	4.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	659	2.3	2.7
7	Diabetes mellitus (E10-E14)	583	2.0	2.4
8	Cerebrovascular diseases (I60-I69)	511	1.8	2.1
9	Septicemia (A40-A41)	380	1.3	1.6
10	Influenza and pneumonia (J09-J18)	370	1.3	1.5
11	Pregnancy, childbirth and the puerperium (O00-O99)	369	1.3	1.5
12	Chronic lower respiratory diseases (J40-J47)	293	1.0	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	288	1.0	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	180	0.6	0.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	102	0.4	0.4
...	All other causes (residual)	5,322	18.5	21.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-64 years				
...	All causes	147,843	100.0	506.5
1	Malignant neoplasms (C00-C97)	52,257	35.3	179.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,727	15.4	77.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,826	7.3	37.1
4	Chronic lower respiratory diseases (J40-J47)	9,224	6.2	31.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,346	3.6	18.3
6	Cerebrovascular diseases (I60-I69)	4,396	3.0	15.1
7	Diabetes mellitus (E10-E14)	4,388	3.0	15.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,789	2.6	13.0
9	Septicemia (A40-A41)	2,482	1.7	8.5
10	Influenza and pneumonia (J09-J18)	2,220	1.5	7.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,747	1.2	6.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,054	0.7	3.6
13	Viral hepatitis (B15-B19)	863	0.6	3.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	749	0.5	2.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	692	0.5	2.4
...	All other causes (residual)	25,083	17.0	85.9
Non-Hispanic, white, female, 65 years and over				
...	All causes	868,104	100.0	4,110.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	208,132	24.0	985.6
2	Malignant neoplasms (C00-C97)	161,785	18.6	766.1
3	Alzheimer's disease (G30)	67,320	7.8	318.8
4	Chronic lower respiratory diseases (J40-J47)	62,745	7.2	297.1
5	Cerebrovascular diseases (I60-I69)	58,801	6.8	278.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,459	2.6	106.3
7	Diabetes mellitus (E10-E14)	18,357	2.1	86.9
8	Influenza and pneumonia (J09-J18)	18,019	2.1	85.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,068	1.7	71.4
10	Septicemia (A40-A41)	12,442	1.4	58.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,171	1.4	57.6
12	Parkinson's disease (G20-G21)	9,967	1.1	47.2
13	Pneumonitis due to solids and liquids (J69)	6,580	0.8	31.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,163	0.6	24.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,590	0.5	21.7
...	All other causes (residual)	184,505	21.3	873.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	326,810	100.0	775.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,608	23.4	181.8
2	Malignant neoplasms (C00-C97)	69,725	21.3	165.5
3	Cerebrovascular diseases (I60-I69)	18,189	5.6	43.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,089	5.5	42.9
5	Diabetes mellitus (E10-E14)	14,053	4.3	33.3
6	Chronic lower respiratory diseases (J40-J47)	10,838	3.3	25.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9,860	3.0	23.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,200	2.8	21.8
9	Alzheimer's disease (G30)	8,577	2.6	20.4
10	Septicemia (A40-A41)	6,684	2.0	15.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,924	1.8	14.1
12	Influenza and pneumonia (J09-J18)	5,468	1.7	13.0
13	Certain conditions originating in the perinatal period (P00-P96)	3,775	1.2	9.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3,152	1.0	7.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3,149	1.0	7.5
...	All other causes (residual)	63,519	19.4	150.7
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,016	100.0	41.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	277	27.3	11.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	13.9	5.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	10.9	4.5
4	Malignant neoplasms (C00-C97)	58	5.7	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.3	1.4
5	Influenza and pneumonia (J09-J18)	34	3.3	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	26	2.6	1.1
8	Chronic lower respiratory diseases (J40-J47)	19	1.9	*
9	Cerebrovascular diseases (I60-I69)	15	1.5	*
10	Septicemia (A40-A41)	14	1.4	*
11	Anemias (D50-D64)	13	1.3	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
14	Meningitis (G00,G03)	6	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
...	All other causes (residual)	248	24.4	10.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,233	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	388	31.5	6.3
2	Malignant neoplasms (C00-C97)	144	11.7	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	118	9.6	1.9
4	Chronic lower respiratory diseases (J40-J47)	88	7.1	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	5.0	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	4.1	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	3.6	0.7
8	Influenza and pneumonia (J09-J18)	21	1.7	0.3
9	Cerebrovascular diseases (I60-I69)	20	1.6	0.3
10	Septicemia (A40-A41)	12	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.9	*
12	Anemias (D50-D64)	10	0.8	*
13	Diabetes mellitus (E10-E14)	8	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
15	Meningitis (G00,G03)	5	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
...	All other causes (residual)	239	19.4	3.9
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	7,703	100.0	115.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,281	42.6	49.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,869	24.3	27.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	656	8.5	9.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	300	3.9	4.5
5	Malignant neoplasms (C00-C97)	230	3.0	3.4
6	Chronic lower respiratory diseases (J40-J47)	78	1.0	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	0.9	1.1
8	Pregnancy, childbirth and the puerperium (O00-O99)	67	0.9	1.0
9	Diabetes mellitus (E10-E14)	64	0.8	1.0
10	Anemias (D50-D64)	59	0.8	0.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	52	0.7	0.8
12	Influenza and pneumonia (J09-J18)	50	0.6	0.7
13	Septicemia (A40-A41)	43	0.6	0.6
14	Cerebrovascular diseases (I60-I69)	33	0.4	0.5
15	Legal intervention (Y35,Y89.0)	32	0.4	0.5
...	All other causes (residual)	817	10.6	12.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-44 years				
...	All causes	26,707	100.0	230.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,320	19.9	45.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,744	17.8	40.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,881	14.5	33.5
4	Malignant neoplasms (C00-C97)	2,514	9.4	21.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,181	4.4	10.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	879	3.3	7.6
7	Diabetes mellitus (E10-E14)	830	3.1	7.2
8	Cerebrovascular diseases (I60-I69)	641	2.4	5.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	408	1.5	3.5
10	Septicemia (A40-A41)	341	1.3	2.9
11	Chronic lower respiratory diseases (J40-J47)	310	1.2	2.7
12	Influenza and pneumonia (J09-J18)	291	1.1	2.5
13	Anemias (D50-D64)	269	1.0	2.3
14	Pregnancy, childbirth and the puerperium (O00-O99)	262	1.0	2.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	260	1.0	2.2
...	All other causes (residual)	4,576	17.1	39.5
Non-Hispanic, black, both sexes, 45-64 years				
...	All causes	94,459	100.0	929.1
1	Malignant neoplasms (C00-C97)	25,033	26.5	246.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,231	24.6	228.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,394	6.8	62.9
4	Diabetes mellitus (E10-E14)	4,562	4.8	44.9
5	Cerebrovascular diseases (I60-I69)	4,403	4.7	43.3
6	Chronic lower respiratory diseases (J40-J47)	2,661	2.8	26.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,379	2.5	23.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,974	2.1	19.4
9	Septicemia (A40-A41)	1,969	2.1	19.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,787	1.9	17.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,588	1.7	15.6
12	Influenza and pneumonia (J09-J18)	1,302	1.4	12.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,276	1.4	12.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	597	0.6	5.9
15	Viral hepatitis (B15-B19)	585	0.6	5.8
...	All other causes (residual)	14,718	15.6	144.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65 years and over				
...	All causes	188,818	100.0	4,235.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,021	26.0	1,099.6
2	Malignant neoplasms (C00-C97)	41,732	22.1	936.1
3	Cerebrovascular diseases (I60-I69)	13,039	6.9	292.5
4	Diabetes mellitus (E10-E14)	8,586	4.5	192.6
5	Alzheimer's disease (G30)	8,462	4.5	189.8
6	Chronic lower respiratory diseases (J40-J47)	7,670	4.1	172.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,367	3.4	142.8
8	Septicemia (A40-A41)	4,236	2.2	95.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,103	2.2	92.0
10	Influenza and pneumonia (J09-J18)	3,707	2.0	83.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,424	1.8	76.8
12	Pneumonitis due to solids and liquids (J69)	1,412	0.7	31.7
13	Parkinson's disease (G20-G21)	1,295	0.7	29.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	920	0.5	20.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	913	0.5	20.5
...	All other causes (residual)	33,931	18.0	761.1
Non-Hispanic, black, male, all ages				
...	All causes	168,750	100.0	836.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,044	23.7	198.4
2	Malignant neoplasms (C00-C97)	35,215	20.9	174.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,453	7.4	61.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,532	5.1	42.3
5	Cerebrovascular diseases (I60-I69)	8,115	4.8	40.2
6	Diabetes mellitus (E10-E14)	6,976	4.1	34.6
7	Chronic lower respiratory diseases (J40-J47)	5,519	3.3	27.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,348	2.6	21.5
9	Septicemia (A40-A41)	3,074	1.8	15.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,805	1.7	13.9
11	Influenza and pneumonia (J09-J18)	2,705	1.6	13.4
12	Alzheimer's disease (G30)	2,451	1.5	12.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,137	1.3	10.6
14	Certain conditions originating in the perinatal period (P00-P96)	2,101	1.2	10.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2,072	1.2	10.3
...	All other causes (residual)	30,203	17.9	149.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	575	100.0	46.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	29.0	13.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	12.7	5.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	9.4	4.3
4	Malignant neoplasms (C00-C97)	32	5.6	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.8	1.8
6	Influenza and pneumonia (J09-J18)	18	3.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	17	3.0	*
8	Chronic lower respiratory diseases (J40-J47)	14	2.4	*
9	Cerebrovascular diseases (I60-I69)	7	1.2	*
9	Acute bronchitis and bronchiolitis (J20-J21)	7	1.2	*
11	Septicemia (A40-A41)	6	1.0	*
12	Anemias (D50-D64)	5	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
13	Meningitis (G00,G03)	4	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	*
...	All other causes (residual)	141	24.5	11.3
Non-Hispanic, black, male, 5-14 years				
...	All causes	701	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	33.2	7.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	10.1	2.3
3	Malignant neoplasms (C00-C97)	66	9.4	2.1
4	Chronic lower respiratory diseases (J40-J47)	58	8.3	1.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	4.7	1.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	4.1	0.9
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.3	0.7
8	Influenza and pneumonia (J09-J18)	14	2.0	*
9	Cerebrovascular diseases (I60-I69)	12	1.7	*
10	Anemias (D50-D64)	8	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.0	*
12	Diabetes mellitus (E10-E14)	5	0.7	*
13	Septicemia (A40-A41)	3	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
15	Meningitis (G00,G03)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	132	18.8	4.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,909	100.0	173.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,955	50.0	86.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,339	22.7	39.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	524	8.9	15.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	3.3	5.7
5	Malignant neoplasms (C00-C97)	137	2.3	4.0
6	Chronic lower respiratory diseases (J40-J47)	48	0.8	1.4
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	0.8	1.4
8	Anemias (D50-D64)	35	0.6	1.0
9	Diabetes mellitus (E10-E14)	33	0.6	1.0
10	Legal intervention (Y35,Y89.0)	31	0.5	0.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.5	0.8
12	Influenza and pneumonia (J09-J18)	24	0.4	0.7
13	Septicemia (A40-A41)	21	0.4	0.6
14	Cerebrovascular diseases (I60-I69)	17	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
...	All other causes (residual)	466	7.9	13.7
Non-Hispanic, black, male, 25-44 years				
...	All causes	17,422	100.0	312.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,207	24.1	75.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,886	22.3	69.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,481	14.2	44.4
4	Malignant neoplasms (C00-C97)	983	5.6	17.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	955	5.5	17.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	550	3.2	9.9
7	Diabetes mellitus (E10-E14)	460	2.6	8.2
8	Cerebrovascular diseases (I60-I69)	365	2.1	6.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	219	1.3	3.9
10	Chronic lower respiratory diseases (J40-J47)	177	1.0	3.2
11	Septicemia (A40-A41)	174	1.0	3.1
12	Influenza and pneumonia (J09-J18)	145	0.8	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	130	0.7	2.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	129	0.7	2.3
15	Anemias (D50-D64)	119	0.7	2.1
...	All other causes (residual)	2,442	14.0	43.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-64 years				
...	All causes	54,534	100.0	1,158.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,624	26.8	310.6
2	Malignant neoplasms (C00-C97)	12,760	23.4	271.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,557	8.4	96.8
4	Diabetes mellitus (E10-E14)	2,568	4.7	54.5
5	Cerebrovascular diseases (I60-I69)	2,456	4.5	52.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,320	2.4	28.0
7	Chronic lower respiratory diseases (J40-J47)	1,297	2.4	27.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,276	2.3	27.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,187	2.2	25.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,032	1.9	21.9
11	Septicemia (A40-A41)	973	1.8	20.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	963	1.8	20.5
13	Influenza and pneumonia (J09-J18)	721	1.3	15.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	469	0.9	10.0
15	Viral hepatitis (B15-B19)	386	0.7	8.2
...	All other causes (residual)	7,945	14.6	168.7
Non-Hispanic, black, male, 65 years and over				
...	All causes	85,855	100.0	4,780.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,654	26.4	1,261.3
2	Malignant neoplasms (C00-C97)	21,227	24.7	1,181.8
3	Cerebrovascular diseases (I60-I69)	5,236	6.1	291.5
4	Chronic lower respiratory diseases (J40-J47)	3,916	4.6	218.0
5	Diabetes mellitus (E10-E14)	3,908	4.6	217.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,789	3.2	155.3
7	Alzheimer's disease (G30)	2,396	2.8	133.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,040	2.4	113.6
9	Septicemia (A40-A41)	1,859	2.2	103.5
10	Influenza and pneumonia (J09-J18)	1,748	2.0	97.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,705	2.0	94.9
12	Parkinson's disease (G20-G21)	745	0.9	41.5
13	Pneumonitis due to solids and liquids (J69)	700	0.8	39.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	595	0.7	33.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	424	0.5	23.6
...	All other causes (residual)	13,913	16.2	774.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	158,060	100.0	719.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,564	23.1	166.5
2	Malignant neoplasms (C00-C97)	34,510	21.8	157.1
3	Cerebrovascular diseases (I60-I69)	10,074	6.4	45.9
4	Diabetes mellitus (E10-E14)	7,077	4.5	32.2
5	Alzheimer's disease (G30)	6,126	3.9	27.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,636	3.6	25.7
7	Chronic lower respiratory diseases (J40-J47)	5,319	3.4	24.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,852	3.1	22.1
9	Septicemia (A40-A41)	3,610	2.3	16.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,119	2.0	14.2
11	Influenza and pneumonia (J09-J18)	2,763	1.7	12.6
12	Certain conditions originating in the perinatal period (P00-P96)	1,674	1.1	7.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,328	0.8	6.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,151	0.7	5.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,077	0.7	4.9
...	All other causes (residual)	33,180	21.0	151.1
Non-Hispanic, black, female, 1-4 years				
...	All causes	441	100.0	36.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110	24.9	9.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	15.4	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	12.9	4.7
4	Malignant neoplasms (C00-C97)	26	5.9	2.2
5	Influenza and pneumonia (J09-J18)	16	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.7	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	2.0	*
8	Septicemia (A40-A41)	8	1.8	*
8	Anemias (D50-D64)	8	1.8	*
8	Cerebrovascular diseases (I60-I69)	8	1.8	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Meningitis (G00,G03)	2	0.5	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	102	23.1	8.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	532	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	29.1	5.1
2	Malignant neoplasms (C00-C97)	78	14.7	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	8.8	1.5
4	Chronic lower respiratory diseases (J40-J47)	30	5.6	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	5.5	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.3	0.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	2.8	*
8	Septicemia (A40-A41)	9	1.7	*
9	Cerebrovascular diseases (I60-I69)	8	1.5	*
10	Influenza and pneumonia (J09-J18)	7	1.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Meningitis (G00,G03)	3	0.6	*
13	Pneumonitis due to solids and liquids (J69)	3	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
...	All other causes (residual)	105	19.7	3.5
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,794	100.0	54.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	530	29.5	16.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	326	18.2	9.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	132	7.4	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	5.9	3.2
5	Malignant neoplasms (C00-C97)	93	5.2	2.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	67	3.7	2.0
7	Diabetes mellitus (E10-E14)	31	1.7	0.9
8	Chronic lower respiratory diseases (J40-J47)	30	1.7	0.9
9	Influenza and pneumonia (J09-J18)	26	1.4	0.8
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	1.4	0.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	24	1.3	0.7
11	Anemias (D50-D64)	24	1.3	0.7
13	Septicemia (A40-A41)	22	1.2	0.7
14	Cerebrovascular diseases (I60-I69)	16	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
...	All other causes (residual)	327	18.2	9.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-44 years				
...	All causes	9,285	100.0	154.6
1	Malignant neoplasms (C00-C97)	1,531	16.5	25.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,434	15.4	23.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,400	15.1	23.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	537	5.8	8.9
5	Diabetes mellitus (E10-E14)	370	4.0	6.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	329	3.5	5.5
7	Cerebrovascular diseases (I60-I69)	276	3.0	4.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	262	2.8	4.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	226	2.4	3.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	189	2.0	3.1
11	Septicemia (A40-A41)	167	1.8	2.8
12	Anemias (D50-D64)	150	1.6	2.5
13	Influenza and pneumonia (J09-J18)	146	1.6	2.4
14	Chronic lower respiratory diseases (J40-J47)	133	1.4	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	131	1.4	2.2
...	All other causes (residual)	2,004	21.6	33.4
Non-Hispanic, black, female, 45-64 years				
...	All causes	39,925	100.0	731.6
1	Malignant neoplasms (C00-C97)	12,273	30.7	224.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,607	21.6	157.7
3	Diabetes mellitus (E10-E14)	1,994	5.0	36.5
4	Cerebrovascular diseases (I60-I69)	1,947	4.9	35.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,837	4.6	33.7
6	Chronic lower respiratory diseases (J40-J47)	1,364	3.4	25.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,059	2.7	19.4
8	Septicemia (A40-A41)	996	2.5	18.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	698	1.7	12.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	625	1.6	11.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	600	1.5	11.0
12	Influenza and pneumonia (J09-J18)	581	1.5	10.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	244	0.6	4.5
14	Viral hepatitis (B15-B19)	199	0.5	3.6
15	Anemias (D50-D64)	154	0.4	2.8
...	All other causes (residual)	6,747	16.9	123.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65 years and over				
...	All causes	102,963	100.0	3,867.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,367	25.6	990.4
2	Malignant neoplasms (C00-C97)	20,505	19.9	770.2
3	Cerebrovascular diseases (I60-I69)	7,803	7.6	293.1
4	Alzheimer's disease (G30)	6,066	5.9	227.9
5	Diabetes mellitus (E10-E14)	4,678	4.5	175.7
6	Chronic lower respiratory diseases (J40-J47)	3,754	3.6	141.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,578	3.5	134.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,398	2.3	90.1
9	Septicemia (A40-A41)	2,377	2.3	89.3
10	Influenza and pneumonia (J09-J18)	1,959	1.9	73.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,384	1.3	52.0
12	Pneumonitis due to solids and liquids (J69)	712	0.7	26.7
13	Parkinson's disease (G20-G21)	550	0.5	20.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	496	0.5	18.6
15	Nutritional deficiencies (E40-E64)	361	0.4	13.6
...	All other causes (residual)	19,975	19.4	750.3
Non-Hispanic, American Indian or Alaska Native, both sexes, all ages				
...	All causes	18,595	100.0	685.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,332	17.9	122.9
2	Malignant neoplasms (C00-C97)	3,225	17.3	119.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,100	11.3	77.5
4	Diabetes mellitus (E10-E14)	1,078	5.8	39.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,010	5.4	37.3
6	Chronic lower respiratory diseases (J40-J47)	853	4.6	31.5
7	Cerebrovascular diseases (I60-I69)	632	3.4	23.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	580	3.1	21.4
9	Alzheimer's disease (G30)	377	2.0	13.9
10	Influenza and pneumonia (J09-J18)	360	1.9	13.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	1.8	12.7
12	Septicemia (A40-A41)	295	1.6	10.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	279	1.5	10.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	193	1.0	7.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	0.6	4.4
...	All other causes (residual)	3,818	20.5	140.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 1-4 years				
...	All causes	82	100.0	51.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	32.9	16.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	12.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	9.8	*
4	Malignant neoplasms (C00-C97)	3	3.7	*
4	Influenza and pneumonia (J09-J18)	3	3.7	*
6	Meningococcal infection (A39)	2	2.4	*
6	Septicemia (A40-A41)	2	2.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.4	*
9	Diabetes mellitus (E10-E14)	1	1.2	*
9	Cerebrovascular diseases (I60-I69)	1	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.2	*
...	All other causes (residual)	22	26.8	13.8
Non-Hispanic, American Indian or Alaska Native, both sexes, 5-14 years				
...	All causes	92	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	33.7	7.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	14.1	*
3	Malignant neoplasms (C00-C97)	8	8.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	4.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	3.3	*
6	Septicemia (A40-A41)	2	2.2	*
6	Meningitis (G00,G03)	2	2.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.2	*
9	Meningococcal infection (A39)	1	1.1	*
9	Diabetes mellitus (E10-E14)	1	1.1	*
9	Cerebrovascular diseases (I60-I69)	1	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.1	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.1	*
...	All other causes (residual)	21	22.8	5.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 15-24 years				
...	All causes	591	100.0	136.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	270	45.7	62.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	26.4	36.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	9.5	12.9
4	Malignant neoplasms (C00-C97)	10	1.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	1.2	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.0	*
7	Diabetes mellitus (E10-E14)	4	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.5	*
9	Legal intervention (Y35,Y89.0)	3	0.5	*
12	Cerebrovascular diseases (I60-I69)	2	0.3	*
12	Influenza and pneumonia (J09-J18)	2	0.3	*
14	Septicemia (A40-A41)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	61	10.3	14.1
Non-Hispanic, American Indian or Alaska Native, both sexes, 25-44 years				
...	All causes	2,435	100.0	334.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	763	31.3	104.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	272	11.2	37.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	270	11.1	37.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	221	9.1	30.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	5.5	18.5
6	Malignant neoplasms (C00-C97)	133	5.5	18.3
7	Diabetes mellitus (E10-E14)	93	3.8	12.8
8	Cerebrovascular diseases (I60-I69)	34	1.4	4.7
9	Influenza and pneumonia (J09-J18)	28	1.1	3.8
10	Septicemia (A40-A41)	22	0.9	3.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.7	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.5	*
13	Legal intervention (Y35,Y89.0)	13	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	11	0.5	*
...	All other causes (residual)	380	15.6	52.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 45-64 years				
...	All causes	5,725	100.0	874.8
1	Malignant neoplasms (C00-C97)	1,149	20.1	175.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	966	16.9	147.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	692	12.1	105.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	552	9.6	84.3
5	Diabetes mellitus (E10-E14)	370	6.5	56.5
6	Chronic lower respiratory diseases (J40-J47)	195	3.4	29.8
7	Cerebrovascular diseases (I60-I69)	137	2.4	20.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	2.0	17.6
9	Influenza and pneumonia (J09-J18)	97	1.7	14.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	97	1.7	14.8
11	Septicemia (A40-A41)	89	1.6	13.6
12	Viral hepatitis (B15-B19)	69	1.2	10.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	1.1	9.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	47	0.8	7.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.4	3.5
...	All other causes (residual)	1,066	18.6	162.9
Non-Hispanic, American Indian or Alaska Native, both sexes, 65 years and over				
...	All causes	9,358	100.0	3,284.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,131	22.8	748.0
2	Malignant neoplasms (C00-C97)	1,921	20.5	674.3
3	Chronic lower respiratory diseases (J40-J47)	642	6.9	225.4
4	Diabetes mellitus (E10-E14)	609	6.5	213.8
5	Cerebrovascular diseases (I60-I69)	457	4.9	160.4
6	Alzheimer's disease (G30)	366	3.9	128.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	3.2	103.6
8	Influenza and pneumonia (J09-J18)	228	2.4	80.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	226	2.4	79.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	182	1.9	63.9
11	Septicemia (A40-A41)	176	1.9	61.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	133	1.4	46.7
13	Pneumonitis due to solids and liquids (J69)	77	0.8	27.0
14	Parkinson's disease (G20-G21)	76	0.8	26.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.5	16.8
...	All other causes (residual)	1,791	19.1	628.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, all ages				
...	All causes	10,280	100.0	772.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,951	19.0	146.7
2	Malignant neoplasms (C00-C97)	1,723	16.8	129.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,404	13.7	105.5
4	Diabetes mellitus (E10-E14)	574	5.6	43.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	554	5.4	41.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	441	4.3	33.2
7	Chronic lower respiratory diseases (J40-J47)	430	4.2	32.3
8	Cerebrovascular diseases (I60-I69)	288	2.8	21.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	209	2.0	15.7
10	Influenza and pneumonia (J09-J18)	176	1.7	13.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	1.6	12.5
12	Septicemia (A40-A41)	128	1.2	9.6
13	Alzheimer's disease (G30)	121	1.2	9.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	0.8	6.1
15	Certain conditions originating in the perinatal period (P00-P96)	72	0.7	5.4
...	All other causes (residual)	1,962	19.1	147.5
Non-Hispanic, American Indian or Alaska Native, male, 1-4 years				
...	All causes	38	100.0	46.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	36.8	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	10.5	*
3	Malignant neoplasms (C00-C97)	2	5.3	*
3	Influenza and pneumonia (J09-J18)	2	5.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	5.3	*
6	Meningococcal infection (A39)	1	2.6	*
6	Septicemia (A40-A41)	1	2.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.6	*
6	Cerebrovascular diseases (I60-I69)	1	2.6	*
...	All other causes (residual)	10	26.3	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 5-14 years				
...	All causes	49	100.0	23.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	42.9	10.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	14.3	*
3	Malignant neoplasms (C00-C97)	3	6.1	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.1	*
5	Diabetes mellitus (E10-E14)	1	2.0	*
5	Meningitis (G00,G03)	1	2.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.0	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.0	*
...	All other causes (residual)	12	24.5	*
Non-Hispanic, American Indian or Alaska Native, male, 15-24 years				
...	All causes	404	100.0	183.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	46.0	84.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	114	28.2	51.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	10.4	19.0
4	Malignant neoplasms (C00-C97)	6	1.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	1.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.0	*
7	Diabetes mellitus (E10-E14)	3	0.7	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
8	Legal intervention (Y35,Y89.0)	2	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
11	Cerebrovascular diseases (I60-I69)	1	0.2	*
11	Influenza and pneumonia (J09-J18)	1	0.2	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	33	8.2	15.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 25-44 years				
...	All causes	1,554	100.0	429.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	505	32.5	139.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	208	13.4	57.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	155	10.0	42.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	9.3	40.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	6.8	29.0
6	Malignant neoplasms (C00-C97)	57	3.7	15.8
7	Diabetes mellitus (E10-E14)	56	3.6	15.5
8	Cerebrovascular diseases (I60-I69)	17	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.8	*
10	Influenza and pneumonia (J09-J18)	12	0.8	*
10	Legal intervention (Y35,Y89.0)	12	0.8	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.6	*
14	Septicemia (A40-A41)	8	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.5	*
...	All other causes (residual)	233	15.0	64.4
Non-Hispanic, American Indian or Alaska Native, male, 45-64 years				
...	All causes	3,428	100.0	1,100.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	635	18.5	203.8
2	Malignant neoplasms (C00-C97)	620	18.1	199.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	490	14.3	157.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	308	9.0	98.8
5	Diabetes mellitus (E10-E14)	220	6.4	70.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	95	2.8	30.5
7	Chronic lower respiratory diseases (J40-J47)	94	2.7	30.2
8	Cerebrovascular diseases (I60-I69)	73	2.1	23.4
9	Influenza and pneumonia (J09-J18)	67	2.0	21.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.4	15.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	1.4	15.1
12	Septicemia (A40-A41)	42	1.2	13.5
13	Viral hepatitis (B15-B19)	37	1.1	11.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	0.8	9.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.5	*
...	All other causes (residual)	608	17.7	195.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 65 years and over				
...	All causes	4,624	100.0	3,625.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,161	25.1	910.3
2	Malignant neoplasms (C00-C97)	1,035	22.4	811.5
3	Chronic lower respiratory diseases (J40-J47)	327	7.1	256.4
4	Diabetes mellitus (E10-E14)	294	6.4	230.5
5	Cerebrovascular diseases (I60-I69)	196	4.2	153.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	3.8	136.4
7	Alzheimer's disease (G30)	114	2.5	89.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	2.3	82.3
9	Influenza and pneumonia (J09-J18)	93	2.0	72.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	89	1.9	69.8
11	Septicemia (A40-A41)	76	1.6	59.6
12	Parkinson's disease (G20-G21)	53	1.1	41.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.0	34.5
14	Pneumonitis due to solids and liquids (J69)	41	0.9	32.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.5	19.6
...	All other causes (residual)	797	17.2	624.9
Non-Hispanic, American Indian or Alaska Native, female, all ages				
...	All causes	8,315	100.0	602.2
1	Malignant neoplasms (C00-C97)	1,502	18.1	108.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,381	16.6	100.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	696	8.4	50.4
4	Diabetes mellitus (E10-E14)	504	6.1	36.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	456	5.5	33.0
6	Chronic lower respiratory diseases (J40-J47)	423	5.1	30.6
7	Cerebrovascular diseases (I60-I69)	344	4.1	24.9
8	Alzheimer's disease (G30)	256	3.1	18.5
9	Influenza and pneumonia (J09-J18)	184	2.2	13.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	2.1	12.8
11	Septicemia (A40-A41)	167	2.0	12.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	139	1.7	10.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	112	1.3	8.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	0.8	5.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	0.8	4.7
...	All other causes (residual)	1,839	22.1	133.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 1-4 years				
...	All causes	44	100.0	55.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	29.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	18.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	9.1	*
4	Meningococcal infection (A39)	1	2.3	*
4	Septicemia (A40-A41)	1	2.3	*
4	Malignant neoplasms (C00-C97)	1	2.3	*
4	Diabetes mellitus (E10-E14)	1	2.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.3	*
4	Influenza and pneumonia (J09-J18)	1	2.3	*
4	Certain conditions originating in the perinatal period (P00-P96)	1	2.3	*
...	All other causes (residual)	12	27.3	*
Non-Hispanic, American Indian or Alaska Native, female, 5-14 years				
...	All causes	43	100.0	21.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	23.3	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	14.0	*
3	Malignant neoplasms (C00-C97)	5	11.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	9.3	*
5	Septicemia (A40-A41)	2	4.7	*
6	Meningococcal infection (A39)	1	2.3	*
6	Meningitis (G00,G03)	1	2.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.3	*
6	Cerebrovascular diseases (I60-I69)	1	2.3	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.3	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	2.3	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.3	*
...	All other causes (residual)	9	20.9	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 15-24 years				
...	All causes	187	100.0	87.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	44.9	39.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	22.5	19.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	7.5	*
4	Malignant neoplasms (C00-C97)	4	2.1	*
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4	2.1	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.6	*
7	Septicemia (A40-A41)	1	0.5	*
7	Diabetes mellitus (E10-E14)	1	0.5	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	0.5	*
7	Cerebrovascular diseases (I60-I69)	1	0.5	*
7	Influenza and pneumonia (J09-J18)	1	0.5	*
7	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
7	Legal intervention (Y35,Y89.0)	1	0.5	*
...	All other causes (residual)	28	15.0	13.1
Non-Hispanic, American Indian or Alaska Native, female, 25-44 years				
...	All causes	881	100.0	240.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	29.3	70.5
2	Chronic liver disease and cirrhosis (K70,K73-K74)	115	13.1	31.4
3	Malignant neoplasms (C00-C97)	76	8.6	20.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	8.6	20.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	7.3	17.5
6	Diabetes mellitus (E10-E14)	37	4.2	10.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	3.4	8.2
8	Cerebrovascular diseases (I60-I69)	17	1.9	*
9	Influenza and pneumonia (J09-J18)	16	1.8	*
10	Septicemia (A40-A41)	14	1.6	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
12	Pneumonitis due to solids and liquids (J69)	5	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	5	0.6	*
...	All other causes (residual)	151	17.1	41.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 45-64 years				
...	All causes	2,297	100.0	670.0
1	Malignant neoplasms (C00-C97)	529	23.0	154.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	331	14.4	96.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	244	10.6	71.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	8.8	58.9
5	Diabetes mellitus (E10-E14)	150	6.5	43.8
6	Chronic lower respiratory diseases (J40-J47)	101	4.4	29.5
7	Cerebrovascular diseases (I60-I69)	64	2.8	18.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	2.1	14.3
9	Septicemia (A40-A41)	47	2.0	13.7
10	Viral hepatitis (B15-B19)	32	1.4	9.3
11	Influenza and pneumonia (J09-J18)	30	1.3	8.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	0.9	5.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	0.8	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	10	0.4	*
...	All other causes (residual)	445	19.4	129.8
Non-Hispanic, American Indian or Alaska Native, female, 65 years and over				
...	All causes	4,734	100.0	3,008.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	970	20.5	616.5
2	Malignant neoplasms (C00-C97)	886	18.7	563.1
3	Diabetes mellitus (E10-E14)	315	6.7	200.2
3	Chronic lower respiratory diseases (J40-J47)	315	6.7	200.2
5	Cerebrovascular diseases (I60-I69)	261	5.5	165.9
6	Alzheimer's disease (G30)	252	5.3	160.2
7	Influenza and pneumonia (J09-J18)	135	2.9	85.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	121	2.6	76.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	2.6	76.9
10	Septicemia (A40-A41)	100	2.1	63.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	93	2.0	59.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.9	56.6
13	Pneumonitis due to solids and liquids (J69)	36	0.8	22.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.5	14.6
14	Parkinson's disease (G20-G21)	23	0.5	14.6
...	All other causes (residual)	994	21.0	631.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, all ages				
...	All causes	68,235	100.0	350.3
1	Malignant neoplasms (C00-C97)	17,718	26.0	91.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,519	21.3	74.5
3	Cerebrovascular diseases (I60-I69)	5,217	7.6	26.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,101	4.5	15.9
5	Diabetes mellitus (E10-E14)	2,697	4.0	13.8
6	Alzheimer's disease (G30)	2,363	3.5	12.1
7	Influenza and pneumonia (J09-J18)	2,132	3.1	10.9
8	Chronic lower respiratory diseases (J40-J47)	1,914	2.8	9.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,363	2.0	7.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,348	2.0	6.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,280	1.9	6.6
12	Septicemia (A40-A41)	786	1.2	4.0
13	Parkinson's disease (G20-G21)	749	1.1	3.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	632	0.9	3.2
15	Certain conditions originating in the perinatal period (P00-P96)	527	0.8	2.7
...	All other causes (residual)	11,889	17.4	61.0
Non-Hispanic, Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	160	100.0	17.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	26.9	4.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	13.1	2.2
3	Malignant neoplasms (C00-C97)	20	12.5	2.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	6.3	*
5	Septicemia (A40-A41)	5	3.1	*
6	Cerebrovascular diseases (I60-I69)	4	2.5	*
7	Influenza and pneumonia (J09-J18)	3	1.9	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.3	*
9	Diabetes mellitus (E10-E14)	2	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	2	1.3	*
12	Meningitis (G00,G03)	1	0.6	*
12	Pneumonitis due to solids and liquids (J69)	1	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (residual)	40	25.0	4.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	218	100.0	9.2
1	Malignant neoplasms (C00-C97)	60	27.5	2.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	17.0	1.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	6.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	6.0	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	5.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	4.1	*
7	Influenza and pneumonia (J09-J18)	7	3.2	*
8	Cerebrovascular diseases (I60-I69)	6	2.8	*
9	Septicemia (A40-A41)	5	2.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	4	1.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	*
12	Scarlet fever and erysipelas (A38,A46)	1	0.5	*
12	Diabetes mellitus (E10-E14)	1	0.5	*
12	Meningitis (G00,G03)	1	0.5	*
12	Diseases of appendix (K35-K38)	1	0.5	*
...	All other causes (residual)	45	20.6	1.9
Non-Hispanic, Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	971	100.0	37.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	31.6	11.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	305	31.4	11.7
3	Malignant neoplasms (C00-C97)	73	7.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	7.0	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	4.2	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.4	*
7	Cerebrovascular diseases (I60-I69)	8	0.8	*
8	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	6	0.6	*
10	Septicemia (A40-A41)	3	0.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.3	*
10	Diabetes mellitus (E10-E14)	3	0.3	*
10	Aortic aneurysm and dissection (I71)	3	0.3	*
10	Influenza and pneumonia (J09-J18)	3	0.3	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.2	*
...	All other causes (residual)	125	12.9	4.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 25-44 years				
...	All causes	3,704	100.0	57.9
1	Malignant neoplasms (C00-C97)	892	24.1	13.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	677	18.3	10.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	492	13.3	7.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	461	12.4	7.2
5	Cerebrovascular diseases (I60-I69)	158	4.3	2.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	3.7	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	80	2.2	1.3
8	Diabetes mellitus (E10-E14)	55	1.5	0.9
9	Pregnancy, childbirth and the puerperium (O00-O99)	42	1.1	0.7
10	Influenza and pneumonia (J09-J18)	36	1.0	0.6
11	Septicemia (A40-A41)	30	0.8	0.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	27	0.7	0.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.6	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.6	0.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.5	0.3
15	Aortic aneurysm and dissection (I71)	20	0.5	0.3
15	Chronic lower respiratory diseases (J40-J47)	20	0.5	0.3
...	All other causes (residual)	511	13.8	8.0
Non-Hispanic, Asian or Pacific Islander, both sexes, 45-64 years				
...	All causes	12,874	100.0	272.8
1	Malignant neoplasms (C00-C97)	5,273	41.0	111.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,348	18.2	49.8
3	Cerebrovascular diseases (I60-I69)	800	6.2	17.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	657	5.1	13.9
5	Diabetes mellitus (E10-E14)	558	4.3	11.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	374	2.9	7.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	249	1.9	5.3
8	Influenza and pneumonia (J09-J18)	203	1.6	4.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	178	1.4	3.8
10	Chronic lower respiratory diseases (J40-J47)	156	1.2	3.3
11	Septicemia (A40-A41)	143	1.1	3.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	1.1	2.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	0.8	2.2
14	Viral hepatitis (B15-B19)	100	0.8	2.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.5	1.4
...	All other causes (residual)	1,527	11.9	32.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 65 years and over				
...	All causes	49,376	100.0	2,215.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,633	23.6	521.9
2	Malignant neoplasms (C00-C97)	11,398	23.1	511.4
3	Cerebrovascular diseases (I60-I69)	4,234	8.6	190.0
4	Alzheimer's disease (G30)	2,346	4.8	105.3
5	Diabetes mellitus (E10-E14)	2,078	4.2	93.2
6	Influenza and pneumonia (J09-J18)	1,877	3.8	84.2
7	Chronic lower respiratory diseases (J40-J47)	1,729	3.5	77.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,357	2.7	60.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,140	2.3	51.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,121	2.3	50.3
11	Parkinson's disease (G20-G21)	727	1.5	32.6
12	Septicemia (A40-A41)	593	1.2	26.6
13	Pneumonitis due to solids and liquids (J69)	432	0.9	19.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	347	0.7	15.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	302	0.6	13.5
...	All other causes (residual)	8,062	16.3	361.7
Non-Hispanic, Asian or Pacific Islander, male, all ages				
...	All causes	34,892	100.0	374.9
1	Malignant neoplasms (C00-C97)	8,978	25.7	96.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,840	22.5	84.2
3	Cerebrovascular diseases (I60-I69)	2,268	6.5	24.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,929	5.5	20.7
5	Diabetes mellitus (E10-E14)	1,414	4.1	15.2
6	Chronic lower respiratory diseases (J40-J47)	1,134	3.3	12.2
7	Influenza and pneumonia (J09-J18)	1,120	3.2	12.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	977	2.8	10.5
9	Alzheimer's disease (G30)	709	2.0	7.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	665	1.9	7.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	545	1.6	5.9
12	Parkinson's disease (G20-G21)	440	1.3	4.7
13	Septicemia (A40-A41)	393	1.1	4.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	388	1.1	4.2
15	Certain conditions originating in the perinatal period (P00-P96)	306	0.9	3.3
...	All other causes (residual)	5,786	16.6	62.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 1-4 years				
...	All causes	80	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	36.3	6.0
2	Malignant neoplasms (C00-C97)	9	11.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	11.3	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.5	*
4	Cerebrovascular diseases (I60-I69)	2	2.5	*
4	Influenza and pneumonia (J09-J18)	2	2.5	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.5	*
9	Septicemia (A40-A41)	1	1.3	*
9	Diabetes mellitus (E10-E14)	1	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.3	*
9	Pneumonitis due to solids and liquids (J69)	1	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.3	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.3	*
...	All other causes (residual)	17	21.3	*
Non-Hispanic, Asian or Pacific Islander, male, 5-14 years				
...	All causes	118	100.0	9.9
1	Malignant neoplasms (C00-C97)	32	27.1	2.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	18.6	1.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	9.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	8.5	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	3.4	*
7	Septicemia (A40-A41)	3	2.5	*
7	Cerebrovascular diseases (I60-I69)	3	2.5	*
7	Influenza and pneumonia (J09-J18)	3	2.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	2	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
11	Diabetes mellitus (E10-E14)	1	0.8	*
...	All other causes (residual)	21	17.8	1.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 15-24 years				
...	All causes	696	100.0	52.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	34.6	18.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	223	32.0	16.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	7.3	3.9
4	Malignant neoplasms (C00-C97)	44	6.3	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.7	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.1	*
7	Chronic lower respiratory diseases (J40-J47)	6	0.9	*
8	Cerebrovascular diseases (I60-I69)	4	0.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
9	Aortic aneurysm and dissection (I71)	3	0.4	*
11	Septicemia (A40-A41)	2	0.3	*
11	Diabetes mellitus (E10-E14)	2	0.3	*
13	Anemias (D50-D64)	1	0.1	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.1	*
13	Influenza and pneumonia (J09-J18)	1	0.1	*
13	Pneumonitis due to solids and liquids (J69)	1	0.1	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.1	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.1	*
13	Legal intervention (Y35,Y89.0)	1	0.1	*
...	All other causes (residual)	76	10.9	5.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 25-44 years				
...	All causes	2,370	100.0	77.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	511	21.6	16.8
2	Malignant neoplasms (C00-C97)	376	15.9	12.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	368	15.5	12.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	360	15.2	11.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	4.1	3.2
6	Cerebrovascular diseases (I60-I69)	90	3.8	3.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	66	2.8	2.2
8	Diabetes mellitus (E10-E14)	36	1.5	1.2
9	Influenza and pneumonia (J09-J18)	26	1.1	0.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.9	0.7
11	Septicemia (A40-A41)	21	0.9	0.7
12	Aortic aneurysm and dissection (I71)	16	0.7	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.7	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	13	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.5	*
...	All other causes (residual)	326	13.8	10.7
Non-Hispanic, Asian or Pacific Islander, male, 45-64 years				
...	All causes	7,592	100.0	348.5
1	Malignant neoplasms (C00-C97)	2,562	33.7	117.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,766	23.3	81.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	446	5.9	20.5
4	Cerebrovascular diseases (I60-I69)	440	5.8	20.2
5	Diabetes mellitus (E10-E14)	367	4.8	16.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	258	3.4	11.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	180	2.4	8.3
8	Influenza and pneumonia (J09-J18)	126	1.7	5.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	1.4	4.9
10	Chronic lower respiratory diseases (J40-J47)	93	1.2	4.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	92	1.2	4.2
12	Septicemia (A40-A41)	79	1.0	3.6
13	Viral hepatitis (B15-B19)	76	1.0	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	0.9	3.2
15	Aortic aneurysm and dissection (I71)	40	0.5	1.8
...	All other causes (residual)	890	11.7	40.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 65 years and over				
...	All causes	23,502	100.0	2,433.1
1	Malignant neoplasms (C00-C97)	5,954	25.3	616.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,668	24.1	586.8
3	Cerebrovascular diseases (I60-I69)	1,723	7.3	178.4
4	Chronic lower respiratory diseases (J40-J47)	1,021	4.3	105.7
5	Diabetes mellitus (E10-E14)	1,007	4.3	104.3
6	Influenza and pneumonia (J09-J18)	960	4.1	99.4
7	Alzheimer's disease (G30)	702	3.0	72.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	666	2.8	68.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	541	2.3	56.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	439	1.9	45.4
11	Parkinson's disease (G20-G21)	426	1.8	44.1
12	Septicemia (A40-A41)	283	1.2	29.3
13	Pneumonitis due to solids and liquids (J69)	258	1.1	26.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	170	0.7	17.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	142	0.6	14.7
...	All other causes (residual)	3,542	15.1	366.7
Non-Hispanic, Asian or Pacific Islander, female, all ages				
...	All causes	33,343	100.0	327.8
1	Malignant neoplasms (C00-C97)	8,740	26.2	85.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,679	20.0	65.7
3	Cerebrovascular diseases (I60-I69)	2,949	8.8	29.0
4	Alzheimer's disease (G30)	1,654	5.0	16.3
5	Diabetes mellitus (E10-E14)	1,283	3.8	12.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,172	3.5	11.5
7	Influenza and pneumonia (J09-J18)	1,012	3.0	9.9
8	Chronic lower respiratory diseases (J40-J47)	780	2.3	7.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	735	2.2	7.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	683	2.0	6.7
11	Septicemia (A40-A41)	393	1.2	3.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	386	1.2	3.8
13	Parkinson's disease (G20-G21)	309	0.9	3.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	244	0.7	2.4
15	Certain conditions originating in the perinatal period (P00-P96)	221	0.7	2.2
...	All other causes (residual)	6,103	18.3	60.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 1-4 years				
...	All causes	80	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	17.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	15.0	*
3	Malignant neoplasms (C00-C97)	11	13.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	10.0	*
5	Septicemia (A40-A41)	4	5.0	*
6	Cerebrovascular diseases (I60-I69)	2	2.5	*
7	Diabetes mellitus (E10-E14)	1	1.3	*
7	Meningitis (G00,G03)	1	1.3	*
7	Influenza and pneumonia (J09-J18)	1	1.3	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.3	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.3	*
...	All other causes (residual)	23	28.8	5.0
Non-Hispanic, Asian or Pacific Islander, female, 5-14 years				
...	All causes	100	100.0	8.6
1	Malignant neoplasms (C00-C97)	28	28.0	2.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	15.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	7.0	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	5.0	*
5	Influenza and pneumonia (J09-J18)	4	4.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.0	*
6	Cerebrovascular diseases (I60-I69)	3	3.0	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	3.0	*
9	Septicemia (A40-A41)	2	2.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	2	2.0	*
11	Scarlet fever and erysipelas (A38,A46)	1	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
11	Meningitis (G00,G03)	1	1.0	*
11	Diseases of appendix (K35-K38)	1	1.0	*
...	All other causes (residual)	24	24.0	2.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 15-24 years				
...	All causes	275	100.0	21.4
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	29.8	6.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	24.0	5.1
3	Malignant neoplasms (C00-C97)	29	10.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	6.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	5.5	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	6	2.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	2.2	*
8	Cerebrovascular diseases (I60-I69)	4	1.5	*
9	Influenza and pneumonia (J09-J18)	2	0.7	*
10	Tuberculosis (A16-A19)	1	0.4	*
10	Septicemia (A40-A41)	1	0.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
10	Diabetes mellitus (E10-E14)	1	0.4	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
10	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.4	*
...	All other causes (residual)	40	14.5	3.1
Non-Hispanic, Asian or Pacific Islander, female, 25-44 years				
...	All causes	1,334	100.0	39.8
1	Malignant neoplasms (C00-C97)	516	38.7	15.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	12.4	5.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	124	9.3	3.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	7.6	3.0
5	Cerebrovascular diseases (I60-I69)	68	5.1	2.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	42	3.1	1.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	2.9	1.2
8	Diabetes mellitus (E10-E14)	19	1.4	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.0	*
10	Influenza and pneumonia (J09-J18)	10	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.7	*
12	Septicemia (A40-A41)	9	0.7	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.6	*
14	Anemias (D50-D64)	7	0.5	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	7	0.5	*
...	All other causes (residual)	187	14.0	5.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 45-64 years				
...	All causes	5,282	100.0	207.9
1	Malignant neoplasms (C00-C97)	2,711	51.3	106.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	582	11.0	22.9
3	Cerebrovascular diseases (I60-I69)	360	6.8	14.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	211	4.0	8.3
5	Diabetes mellitus (E10-E14)	191	3.6	7.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	2.2	4.6
7	Influenza and pneumonia (J09-J18)	77	1.5	3.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	1.3	2.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	69	1.3	2.7
10	Septicemia (A40-A41)	64	1.2	2.5
11	Chronic lower respiratory diseases (J40-J47)	63	1.2	2.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	0.9	1.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.7	1.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	0.7	1.4
15	Viral hepatitis (B15-B19)	24	0.5	0.9
...	All other causes (residual)	627	11.9	24.7
Non-Hispanic, Asian or Pacific Islander, female, 65 years and over				
...	All causes	25,874	100.0	2,048.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,965	23.1	472.3
2	Malignant neoplasms (C00-C97)	5,444	21.0	431.0
3	Cerebrovascular diseases (I60-I69)	2,511	9.7	198.8
4	Alzheimer's disease (G30)	1,644	6.4	130.2
5	Diabetes mellitus (E10-E14)	1,071	4.1	84.8
6	Influenza and pneumonia (J09-J18)	917	3.5	72.6
7	Chronic lower respiratory diseases (J40-J47)	708	2.7	56.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	691	2.7	54.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	682	2.6	54.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	599	2.3	47.4
11	Septicemia (A40-A41)	310	1.2	24.5
12	Parkinson's disease (G20-G21)	301	1.2	23.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	177	0.7	14.0
14	Pneumonitis due to solids and liquids (J69)	174	0.7	13.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	160	0.6	12.7
...	All other causes (residual)	4,520	17.5	357.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016