

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|-----------|-------------------------|---------|
| United States | | | | |
| ... | All causes | 2,712,630 | 100.0 | 844.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 633,842 | 23.4 | 197.2 |
| 2 | Malignant neoplasms (C00-C97) | 595,930 | 22.0 | 185.4 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 155,041 | 5.7 | 48.2 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 146,571 | 5.4 | 45.6 |
| 5 | Cerebrovascular diseases (I60-I69) | 140,323 | 5.2 | 43.7 |
| 6 | Alzheimer's disease (G30) | 110,561 | 4.1 | 34.4 |
| 7 | Diabetes mellitus (E10-E14) | 79,535 | 2.9 | 24.7 |
| 8 | Influenza and pneumonia (J09-J18) | 57,062 | 2.1 | 17.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 49,959 | 1.8 | 15.5 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 44,193 | 1.6 | 13.7 |
| 11 | Septicemia (A40-A41) | 40,773 | 1.5 | 12.7 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 40,326 | 1.5 | 12.5 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 32,200 | 1.2 | 10.0 |
| 14 | Parkinson's disease (G20-G21) | 27,972 | 1.0 | 8.7 |
| 15 | Pneumonitis due to solids and liquids (J69) | 19,803 | 0.7 | 6.2 |
| ... | All other causes (residual) | 538,539 | 19.9 | 167.6 |
| Alabama | | | | |
| ... | All causes | 51,909 | 100.0 | 1,068.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 12,981 | 25.0 | 267.2 |
| 2 | Malignant neoplasms (C00-C97) | 10,354 | 19.9 | 213.1 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 3,279 | 6.3 | 67.5 |
| 4 | Cerebrovascular diseases (I60-I69) | 2,937 | 5.7 | 60.4 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,552 | 4.9 | 52.5 |
| 6 | Alzheimer's disease (G30) | 2,282 | 4.4 | 47.0 |
| 7 | Diabetes mellitus (E10-E14) | 1,255 | 2.4 | 25.8 |
| 8 | Influenza and pneumonia (J09-J18) | 1,097 | 2.1 | 22.6 |
| 9 | Septicemia (A40-A41) | 1,046 | 2.0 | 21.5 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,040 | 2.0 | 21.4 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 750 | 1.4 | 15.4 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 716 | 1.4 | 14.7 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 519 | 1.0 | 10.7 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 473 | 0.9 | 9.7 |
| 15 | Parkinson's disease (G20-G21) | 461 | 0.9 | 9.5 |
| ... | All other causes (residual) | 10,167 | 19.6 | 209.2 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Alaska | | | | |
| ... | All causes | 4,316 | 100.0 | 584.5 |
| 1 | Malignant neoplasms (C00-C97) | 978 | 22.7 | 132.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 846 | 19.6 | 114.6 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 388 | 9.0 | 52.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 204 | 4.7 | 27.6 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 201 | 4.7 | 27.2 |
| 6 | Cerebrovascular diseases (I60-I69) | 182 | 4.2 | 24.6 |
| 7 | Diabetes mellitus (E10-E14) | 142 | 3.3 | 19.2 |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 114 | 2.6 | 15.4 |
| 9 | Alzheimer's disease (G30) | 68 | 1.6 | 9.2 |
| 10 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 62 | 1.4 | 8.4 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 50 | 1.2 | 6.8 |
| 12 | Influenza and pneumonia (J09-J18) | 41 | 0.9 | 5.6 |
| 13 | Parkinson's disease (G20-G21) | 38 | 0.9 | 5.1 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 38 | 0.9 | 5.1 |
| 15 | Septicemia (A40-A41) | 37 | 0.9 | 5.0 |
| ... | All other causes (residual) | 927 | 21.5 | 125.5 |
| Arizona | | | | |
| ... | All causes | 54,299 | 100.0 | 795.2 |
| 1 | Malignant neoplasms (C00-C97) | 11,776 | 21.7 | 172.5 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,458 | 21.1 | 167.8 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 3,681 | 6.8 | 53.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,539 | 6.5 | 51.8 |
| 5 | Alzheimer's disease (G30) | 2,943 | 5.4 | 43.1 |
| 6 | Cerebrovascular diseases (I60-I69) | 2,522 | 4.6 | 36.9 |
| 7 | Diabetes mellitus (E10-E14) | 2,081 | 3.8 | 30.5 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,276 | 2.3 | 18.7 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,149 | 2.1 | 16.8 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 963 | 1.8 | 14.1 |
| 11 | Influenza and pneumonia (J09-J18) | 775 | 1.4 | 11.4 |
| 12 | Parkinson's disease (G20-G21) | 720 | 1.3 | 10.5 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 458 | 0.8 | 6.7 |
| 14 | Septicemia (A40-A41) | 444 | 0.8 | 6.5 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 364 | 0.7 | 5.3 |
| ... | All other causes (residual) | 10,150 | 18.7 | 148.7 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|---------|
| Arkansas | | | | |
| ... | All causes | 31,617 | 100.0 | 1,061.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,938 | 25.1 | 266.5 |
| 2 | Malignant neoplasms (C00-C97) | 6,727 | 21.3 | 225.9 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 2,270 | 7.2 | 76.2 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,653 | 5.2 | 55.5 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,538 | 4.9 | 51.6 |
| 6 | Alzheimer's disease (G30) | 1,457 | 4.6 | 48.9 |
| 7 | Diabetes mellitus (E10-E14) | 886 | 2.8 | 29.7 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 707 | 2.2 | 23.7 |
| 9 | Influenza and pneumonia (J09-J18) | 700 | 2.2 | 23.5 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 577 | 1.8 | 19.4 |
| 11 | Septicemia (A40-A41) | 525 | 1.7 | 17.6 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 421 | 1.3 | 14.1 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 329 | 1.0 | 11.0 |
| 14 | Parkinson's disease (G20-G21) | 248 | 0.8 | 8.3 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 217 | 0.7 | 7.3 |
| ... | All other causes (residual) | 5,424 | 17.2 | 182.1 |
| California | | | | |
| ... | All causes | 259,206 | 100.0 | 662.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 61,289 | 23.6 | 156.6 |
| 2 | Malignant neoplasms (C00-C97) | 59,629 | 23.0 | 152.3 |
| 3 | Alzheimer's disease (G30) | 15,065 | 5.8 | 38.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 15,065 | 5.8 | 38.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 13,621 | 5.3 | 34.8 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 12,544 | 4.8 | 32.0 |
| 7 | Diabetes mellitus (E10-E14) | 8,845 | 3.4 | 22.6 |
| 8 | Influenza and pneumonia (J09-J18) | 6,188 | 2.4 | 15.8 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 5,425 | 2.1 | 13.9 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 5,118 | 2.0 | 13.1 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 4,167 | 1.6 | 10.6 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,557 | 1.4 | 9.1 |
| 13 | Parkinson's disease (G20-G21) | 2,896 | 1.1 | 7.4 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,987 | 0.8 | 5.1 |
| 15 | Pneumonitis due to solids and liquids (J69) | 1,752 | 0.7 | 4.5 |
| ... | All other causes (residual) | 42,058 | 16.2 | 107.4 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Colorado | | | | |
| ... | All causes | 36,349 | 100.0 | 666.2 |
| 1 | Malignant neoplasms (C00-C97) | 7,604 | 20.9 | 139.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,009 | 19.3 | 128.5 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,725 | 7.5 | 49.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,577 | 7.1 | 47.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 1,856 | 5.1 | 34.0 |
| 6 | Alzheimer's disease (G30) | 1,612 | 4.4 | 29.5 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,093 | 3.0 | 20.0 |
| 8 | Diabetes mellitus (E10-E14) | 887 | 2.4 | 16.3 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 735 | 2.0 | 13.5 |
| 10 | Influenza and pneumonia (J09-J18) | 659 | 1.8 | 12.1 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 458 | 1.3 | 8.4 |
| 12 | Septicemia (A40-A41) | 456 | 1.3 | 8.4 |
| 13 | Parkinson's disease (G20-G21) | 452 | 1.2 | 8.3 |
| 14 | Pneumonitis due to solids and liquids (J69) | 346 | 1.0 | 6.3 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 273 | 0.8 | 5.0 |
| ... | All other causes (residual) | 7,607 | 20.9 | 139.4 |
| Connecticut | | | | |
| ... | All causes | 30,535 | 100.0 | 850.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,205 | 23.6 | 200.6 |
| 2 | Malignant neoplasms (C00-C97) | 6,666 | 21.8 | 185.6 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,799 | 5.9 | 50.1 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,388 | 4.5 | 38.7 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,369 | 4.5 | 38.1 |
| 6 | Alzheimer's disease (G30) | 966 | 3.2 | 26.9 |
| 7 | Influenza and pneumonia (J09-J18) | 667 | 2.2 | 18.6 |
| 8 | Diabetes mellitus (E10-E14) | 653 | 2.1 | 18.2 |
| 9 | Septicemia (A40-A41) | 646 | 2.1 | 18.0 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 584 | 1.9 | 16.3 |
| 11 | Pneumonitis due to solids and liquids (J69) | 386 | 1.3 | 10.7 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 384 | 1.3 | 10.7 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 365 | 1.2 | 10.2 |
| 14 | Parkinson's disease (G20-G21) | 334 | 1.1 | 9.3 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 317 | 1.0 | 8.8 |
| ... | All other causes (residual) | 6,806 | 22.3 | 189.5 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|----------------------|--|--------|-------------------------|-------|
| Delaware | | | | |
| ... | All causes | 8,582 | 100.0 | 907.3 |
| 1 | Malignant neoplasms (C00-C97) | 2,010 | 23.4 | 212.5 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,940 | 22.6 | 205.1 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 510 | 5.9 | 53.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 466 | 5.4 | 49.3 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 449 | 5.2 | 47.5 |
| 6 | Alzheimer's disease (G30) | 264 | 3.1 | 27.9 |
| 7 | Diabetes mellitus (E10-E14) | 215 | 2.5 | 22.7 |
| 8 | Influenza and pneumonia (J09-J18) | 189 | 2.2 | 20.0 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 174 | 2.0 | 18.4 |
| 10 | Septicemia (A40-A41) | 145 | 1.7 | 15.3 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 122 | 1.4 | 12.9 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 111 | 1.3 | 11.7 |
| 13 | Parkinson's disease (G20-G21) | 84 | 1.0 | 8.9 |
| 14 | Pneumonitis due to solids and liquids (J69) | 73 | 0.9 | 7.7 |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 67 | 0.8 | 7.1 |
| ... | All other causes (residual) | 1,763 | 20.5 | 186.4 |
| District of Columbia | | | | |
| ... | All causes | 4,871 | 100.0 | 724.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,217 | 25.0 | 181.0 |
| 2 | Malignant neoplasms (C00-C97) | 1,072 | 22.0 | 159.5 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 265 | 5.4 | 39.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 242 | 5.0 | 36.0 |
| 5 | Diabetes mellitus (E10-E14) | 162 | 3.3 | 24.1 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 147 | 3.0 | 21.9 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 136 | 2.8 | 20.2 |
| 8 | Alzheimer's disease (G30) | 129 | 2.6 | 19.2 |
| 9 | Influenza and pneumonia (J09-J18) | 104 | 2.1 | 15.5 |
| 10 | Septicemia (A40-A41) | 88 | 1.8 | 13.1 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 87 | 1.8 | 12.9 |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 70 | 1.4 | 10.4 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 65 | 1.3 | 9.7 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 61 | 1.3 | 9.1 |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 46 | 0.9 | 6.8 |
| ... | All other causes (residual) | 980 | 20.1 | 145.8 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|-------|
| Florida | | | | |
| ... | All causes | 191,737 | 100.0 | 945.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 45,441 | 23.7 | 224.2 |
| 2 | Malignant neoplasms (C00-C97) | 44,027 | 23.0 | 217.2 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 11,705 | 6.1 | 57.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 11,433 | 6.0 | 56.4 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 10,578 | 5.5 | 52.2 |
| 6 | Alzheimer's disease (G30) | 7,031 | 3.7 | 34.7 |
| 7 | Diabetes mellitus (E10-E14) | 5,403 | 2.8 | 26.7 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 3,205 | 1.7 | 15.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,203 | 1.7 | 15.8 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,084 | 1.6 | 15.2 |
| 11 | Influenza and pneumonia (J09-J18) | 2,676 | 1.4 | 13.2 |
| 12 | Septicemia (A40-A41) | 2,368 | 1.2 | 11.7 |
| 13 | Parkinson's disease (G20-G21) | 2,228 | 1.2 | 11.0 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 2,187 | 1.1 | 10.8 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,479 | 0.8 | 7.3 |
| ... | All other causes (residual) | 35,689 | 18.6 | 176.1 |
| Georgia | | | | |
| ... | All causes | 79,942 | 100.0 | 782.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 17,769 | 22.2 | 174.0 |
| 2 | Malignant neoplasms (C00-C97) | 16,945 | 21.2 | 165.9 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 4,607 | 5.8 | 45.1 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,344 | 5.4 | 42.5 |
| 5 | Cerebrovascular diseases (I60-I69) | 4,335 | 5.4 | 42.4 |
| 6 | Alzheimer's disease (G30) | 3,714 | 4.6 | 36.4 |
| 7 | Diabetes mellitus (E10-E14) | 2,210 | 2.8 | 21.6 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,881 | 2.4 | 18.4 |
| 9 | Septicemia (A40-A41) | 1,526 | 1.9 | 14.9 |
| 10 | Influenza and pneumonia (J09-J18) | 1,467 | 1.8 | 14.4 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,317 | 1.6 | 12.9 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,076 | 1.3 | 10.5 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,003 | 1.3 | 9.8 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 738 | 0.9 | 7.2 |
| 15 | Parkinson's disease (G20-G21) | 733 | 0.9 | 7.2 |
| ... | All other causes (residual) | 16,277 | 20.4 | 159.3 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Hawaii | | | | |
| ... | All causes | 11,053 | 100.0 | 772.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,605 | 23.6 | 182.0 |
| 2 | Malignant neoplasms (C00-C97) | 2,462 | 22.3 | 172.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 735 | 6.6 | 51.3 |
| 4 | Influenza and pneumonia (J09-J18) | 557 | 5.0 | 38.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 536 | 4.8 | 37.4 |
| 6 | Alzheimer's disease (G30) | 422 | 3.8 | 29.5 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 332 | 3.0 | 23.2 |
| 8 | Diabetes mellitus (E10-E14) | 263 | 2.4 | 18.4 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 202 | 1.8 | 14.1 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 201 | 1.8 | 14.0 |
| 11 | Parkinson's disease (G20-G21) | 141 | 1.3 | 9.8 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 138 | 1.2 | 9.6 |
| 13 | Septicemia (A40-A41) | 123 | 1.1 | 8.6 |
| 14 | Pneumonitis due to solids and liquids (J69) | 108 | 1.0 | 7.5 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 86 | 0.8 | 6.0 |
| ... | All other causes (residual) | 2,142 | 19.4 | 149.6 |
| Idaho | | | | |
| ... | All causes | 13,026 | 100.0 | 787.1 |
| 1 | Malignant neoplasms (C00-C97) | 2,849 | 21.9 | 172.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,825 | 21.7 | 170.7 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 843 | 6.5 | 50.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 746 | 5.7 | 45.1 |
| 5 | Cerebrovascular diseases (I60-I69) | 641 | 4.9 | 38.7 |
| 6 | Alzheimer's disease (G30) | 552 | 4.2 | 33.4 |
| 7 | Diabetes mellitus (E10-E14) | 403 | 3.1 | 24.4 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 359 | 2.8 | 21.7 |
| 9 | Influenza and pneumonia (J09-J18) | 218 | 1.7 | 13.2 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 218 | 1.7 | 13.2 |
| 11 | Parkinson's disease (G20-G21) | 169 | 1.3 | 10.2 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 154 | 1.2 | 9.3 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 101 | 0.8 | 6.1 |
| 14 | Pneumonitis due to solids and liquids (J69) | 100 | 0.8 | 6.0 |
| 15 | Septicemia (A40-A41) | 84 | 0.6 | 5.1 |
| ... | All other causes (residual) | 2,764 | 21.2 | 167.0 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|-------|
| Illinois | | | | |
| ... | All causes | 106,872 | 100.0 | 831.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 25,652 | 24.0 | 199.5 |
| 2 | Malignant neoplasms (C00-C97) | 24,713 | 23.1 | 192.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 5,709 | 5.3 | 44.4 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 5,544 | 5.2 | 43.1 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,850 | 4.5 | 37.7 |
| 6 | Alzheimer's disease (G30) | 3,686 | 3.4 | 28.7 |
| 7 | Diabetes mellitus (E10-E14) | 2,817 | 2.6 | 21.9 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,543 | 2.4 | 19.8 |
| 9 | Influenza and pneumonia (J09-J18) | 2,343 | 2.2 | 18.2 |
| 10 | Septicemia (A40-A41) | 1,819 | 1.7 | 14.1 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,363 | 1.3 | 10.6 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,289 | 1.2 | 10.0 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,137 | 1.1 | 8.8 |
| 14 | Parkinson's disease (G20-G21) | 1,079 | 1.0 | 8.4 |
| 15 | Pneumonitis due to solids and liquids (J69) | 960 | 0.9 | 7.5 |
| ... | All other causes (residual) | 21,368 | 20.0 | 166.2 |
| Indiana | | | | |
| ... | All causes | 62,713 | 100.0 | 947.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 13,948 | 22.2 | 210.7 |
| 2 | Malignant neoplasms (C00-C97) | 13,511 | 21.5 | 204.1 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 4,212 | 6.7 | 63.6 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,258 | 5.2 | 49.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 2,959 | 4.7 | 44.7 |
| 6 | Alzheimer's disease (G30) | 2,513 | 4.0 | 38.0 |
| 7 | Diabetes mellitus (E10-E14) | 2,030 | 3.2 | 30.7 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,450 | 2.3 | 21.9 |
| 9 | Septicemia (A40-A41) | 1,139 | 1.8 | 17.2 |
| 10 | Influenza and pneumonia (J09-J18) | 1,044 | 1.7 | 15.8 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 960 | 1.5 | 14.5 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 859 | 1.4 | 13.0 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 694 | 1.1 | 10.5 |
| 14 | Parkinson's disease (G20-G21) | 638 | 1.0 | 9.6 |
| 15 | Pneumonitis due to solids and liquids (J69) | 465 | 0.7 | 7.0 |
| ... | All other causes (residual) | 13,033 | 20.8 | 196.9 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Iowa | | | | |
| ... | All causes | 29,600 | 100.0 | 947.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,813 | 23.0 | 218.1 |
| 2 | Malignant neoplasms (C00-C97) | 6,513 | 22.0 | 208.5 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 2,011 | 6.8 | 64.4 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,537 | 5.2 | 49.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 1,418 | 4.8 | 45.4 |
| 6 | Alzheimer's disease (G30) | 1,339 | 4.5 | 42.9 |
| 7 | Diabetes mellitus (E10-E14) | 1,077 | 3.6 | 34.5 |
| 8 | Influenza and pneumonia (J09-J18) | 618 | 2.1 | 19.8 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 433 | 1.5 | 13.9 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 390 | 1.3 | 12.5 |
| 11 | Parkinson's disease (G20-G21) | 341 | 1.2 | 10.9 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 336 | 1.1 | 10.8 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 328 | 1.1 | 10.5 |
| 14 | Septicemia (A40-A41) | 252 | 0.9 | 8.1 |
| 15 | Pneumonitis due to solids and liquids (J69) | 222 | 0.8 | 7.1 |
| ... | All other causes (residual) | 5,972 | 20.2 | 191.2 |
| Kansas | | | | |
| ... | All causes | 26,664 | 100.0 | 915.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,624 | 21.1 | 193.2 |
| 2 | Malignant neoplasms (C00-C97) | 5,604 | 21.0 | 192.5 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 1,704 | 6.4 | 58.5 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,475 | 5.5 | 50.7 |
| 5 | Cerebrovascular diseases (I60-I69) | 1,364 | 5.1 | 46.8 |
| 6 | Alzheimer's disease (G30) | 865 | 3.2 | 29.7 |
| 7 | Diabetes mellitus (E10-E14) | 684 | 2.6 | 23.5 |
| 8 | Influenza and pneumonia (J09-J18) | 682 | 2.6 | 23.4 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 582 | 2.2 | 20.0 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 477 | 1.8 | 16.4 |
| 11 | Atherosclerosis (I70) | 451 | 1.7 | 15.5 |
| 12 | Septicemia (A40-A41) | 381 | 1.4 | 13.1 |
| 13 | Parkinson's disease (G20-G21) | 326 | 1.2 | 11.2 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 313 | 1.2 | 10.7 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 245 | 0.9 | 8.4 |
| ... | All other causes (residual) | 5,887 | 22.1 | 202.2 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|---------|
| Kentucky | | | | |
| ... | All causes | 46,564 | 100.0 | 1,052.3 |
| 1 | Malignant neoplasms (C00-C97) | 10,312 | 22.1 | 233.0 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,077 | 21.6 | 227.7 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 3,331 | 7.2 | 75.3 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,962 | 6.4 | 66.9 |
| 5 | Cerebrovascular diseases (I60-I69) | 2,050 | 4.4 | 46.3 |
| 6 | Alzheimer's disease (G30) | 1,694 | 3.6 | 38.3 |
| 7 | Diabetes mellitus (E10-E14) | 1,458 | 3.1 | 32.9 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 998 | 2.1 | 22.6 |
| 9 | Influenza and pneumonia (J09-J18) | 967 | 2.1 | 21.9 |
| 10 | Septicemia (A40-A41) | 873 | 1.9 | 19.7 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 776 | 1.7 | 17.5 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 657 | 1.4 | 14.8 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 441 | 0.9 | 10.0 |
| 14 | Parkinson's disease (G20-G21) | 395 | 0.8 | 8.9 |
| 15 | Pneumonitis due to solids and liquids (J69) | 329 | 0.7 | 7.4 |
| ... | All other causes (residual) | 9,244 | 19.9 | 208.9 |
| Louisiana | | | | |
| ... | All causes | 43,716 | 100.0 | 936.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,665 | 24.4 | 228.3 |
| 2 | Malignant neoplasms (C00-C97) | 9,397 | 21.5 | 201.2 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,578 | 5.9 | 55.2 |
| 4 | Cerebrovascular diseases (I60-I69) | 2,280 | 5.2 | 48.8 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 2,178 | 5.0 | 46.6 |
| 6 | Alzheimer's disease (G30) | 2,018 | 4.6 | 43.2 |
| 7 | Diabetes mellitus (E10-E14) | 1,199 | 2.7 | 25.7 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,160 | 2.7 | 24.8 |
| 9 | Septicemia (A40-A41) | 991 | 2.3 | 21.2 |
| 10 | Influenza and pneumonia (J09-J18) | 753 | 1.7 | 16.1 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 722 | 1.7 | 15.5 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 569 | 1.3 | 12.2 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 553 | 1.3 | 11.8 |
| 14 | Parkinson's disease (G20-G21) | 371 | 0.8 | 7.9 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 369 | 0.8 | 7.9 |
| ... | All other causes (residual) | 7,913 | 18.1 | 169.4 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|---------|
| Maine | | | | |
| ... | All causes | 14,479 | 100.0 | 1,089.2 |
| 1 | Malignant neoplasms (C00-C97) | 3,398 | 23.5 | 255.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,009 | 20.8 | 226.4 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 1,025 | 7.1 | 77.1 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 802 | 5.5 | 60.3 |
| 5 | Cerebrovascular diseases (I60-I69) | 616 | 4.3 | 46.3 |
| 6 | Alzheimer's disease (G30) | 544 | 3.8 | 40.9 |
| 7 | Diabetes mellitus (E10-E14) | 407 | 2.8 | 30.6 |
| 8 | Influenza and pneumonia (J09-J18) | 333 | 2.3 | 25.1 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 235 | 1.6 | 17.7 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 211 | 1.5 | 15.9 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 195 | 1.3 | 14.7 |
| 12 | Parkinson's disease (G20-G21) | 147 | 1.0 | 11.1 |
| 13 | Septicemia (A40-A41) | 145 | 1.0 | 10.9 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 105 | 0.7 | 7.9 |
| 15 | Pneumonitis due to solids and liquids (J69) | 102 | 0.7 | 7.7 |
| ... | All other causes (residual) | 3,205 | 22.1 | 241.1 |
| Maryland | | | | |
| ... | All causes | 47,247 | 100.0 | 786.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,481 | 24.3 | 191.1 |
| 2 | Malignant neoplasms (C00-C97) | 10,568 | 22.4 | 175.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,540 | 5.4 | 42.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,040 | 4.3 | 34.0 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,903 | 4.0 | 31.7 |
| 6 | Diabetes mellitus (E10-E14) | 1,243 | 2.6 | 20.7 |
| 7 | Influenza and pneumonia (J09-J18) | 1,176 | 2.5 | 19.6 |
| 8 | Alzheimer's disease (G30) | 1,095 | 2.3 | 18.2 |
| 9 | Septicemia (A40-A41) | 913 | 1.9 | 15.2 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 834 | 1.8 | 13.9 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 596 | 1.3 | 9.9 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 553 | 1.2 | 9.2 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 521 | 1.1 | 8.7 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 471 | 1.0 | 7.8 |
| 15 | Parkinson's disease (G20-G21) | 461 | 1.0 | 7.7 |
| ... | All other causes (residual) | 10,852 | 23.0 | 180.7 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Massachusetts | | | | |
| ... | All causes | 57,806 | 100.0 | 850.8 |
| 1 | Malignant neoplasms (C00-C97) | 12,750 | 22.1 | 187.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 12,130 | 21.0 | 178.5 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,229 | 5.6 | 47.5 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,784 | 4.8 | 41.0 |
| 5 | Cerebrovascular diseases (I60-I69) | 2,475 | 4.3 | 36.4 |
| 6 | Alzheimer's disease (G30) | 1,815 | 3.1 | 26.7 |
| 7 | Influenza and pneumonia (J09-J18) | 1,512 | 2.6 | 22.3 |
| 8 | Diabetes mellitus (E10-E14) | 1,398 | 2.4 | 20.6 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,228 | 2.1 | 18.1 |
| 10 | Septicemia (A40-A41) | 881 | 1.5 | 13.0 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 661 | 1.1 | 9.7 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 642 | 1.1 | 9.4 |
| 13 | Parkinson's disease (G20-G21) | 640 | 1.1 | 9.4 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 603 | 1.0 | 8.9 |
| 15 | Pneumonitis due to solids and liquids (J69) | 560 | 1.0 | 8.2 |
| ... | All other causes (residual) | 14,498 | 25.1 | 213.4 |
| Michigan | | | | |
| ... | All causes | 95,140 | 100.0 | 958.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 24,794 | 26.1 | 249.9 |
| 2 | Malignant neoplasms (C00-C97) | 20,732 | 21.8 | 208.9 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 5,848 | 6.1 | 58.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 4,666 | 4.9 | 47.0 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,647 | 4.9 | 46.8 |
| 6 | Alzheimer's disease (G30) | 3,771 | 4.0 | 38.0 |
| 7 | Diabetes mellitus (E10-E14) | 2,751 | 2.9 | 27.7 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,921 | 2.0 | 19.4 |
| 9 | Influenza and pneumonia (J09-J18) | 1,894 | 2.0 | 19.1 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,410 | 1.5 | 14.2 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,287 | 1.4 | 13.0 |
| 12 | Septicemia (A40-A41) | 1,268 | 1.3 | 12.8 |
| 13 | Parkinson's disease (G20-G21) | 960 | 1.0 | 9.7 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 922 | 1.0 | 9.3 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 597 | 0.6 | 6.0 |
| ... | All other causes (residual) | 17,672 | 18.6 | 178.1 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|---------|
| Minnesota | | | | |
| ... | All causes | 42,800 | 100.0 | 779.7 |
| 1 | Malignant neoplasms (C00-C97) | 9,925 | 23.2 | 180.8 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,844 | 18.3 | 142.9 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,574 | 6.0 | 46.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,351 | 5.5 | 42.8 |
| 5 | Cerebrovascular diseases (I60-I69) | 2,238 | 5.2 | 40.8 |
| 6 | Alzheimer's disease (G30) | 1,789 | 4.2 | 32.6 |
| 7 | Diabetes mellitus (E10-E14) | 1,221 | 2.9 | 22.2 |
| 8 | Influenza and pneumonia (J09-J18) | 738 | 1.7 | 13.4 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 730 | 1.7 | 13.3 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 639 | 1.5 | 11.6 |
| 11 | Parkinson's disease (G20-G21) | 620 | 1.4 | 11.3 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 542 | 1.3 | 9.9 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 518 | 1.2 | 9.4 |
| 14 | Septicemia (A40-A41) | 439 | 1.0 | 8.0 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 316 | 0.7 | 5.8 |
| ... | All other causes (residual) | 10,316 | 24.1 | 187.9 |
| Mississippi | | | | |
| ... | All causes | 31,783 | 100.0 | 1,062.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,969 | 25.1 | 266.3 |
| 2 | Malignant neoplasms (C00-C97) | 6,485 | 20.4 | 216.7 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 1,924 | 6.1 | 64.3 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,814 | 5.7 | 60.6 |
| 5 | Cerebrovascular diseases (I60-I69) | 1,734 | 5.5 | 57.9 |
| 6 | Alzheimer's disease (G30) | 1,402 | 4.4 | 46.9 |
| 7 | Diabetes mellitus (E10-E14) | 1,092 | 3.4 | 36.5 |
| 8 | Influenza and pneumonia (J09-J18) | 791 | 2.5 | 26.4 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 731 | 2.3 | 24.4 |
| 10 | Septicemia (A40-A41) | 676 | 2.1 | 22.6 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 508 | 1.6 | 17.0 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 431 | 1.4 | 14.4 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 382 | 1.2 | 12.8 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 325 | 1.0 | 10.9 |
| 15 | Parkinson's disease (G20-G21) | 247 | 0.8 | 8.3 |
| ... | All other causes (residual) | 5,272 | 16.6 | 176.2 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Missouri | | | | |
| ... | All causes | 59,871 | 100.0 | 984.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 14,808 | 24.7 | 243.4 |
| 2 | Malignant neoplasms (C00-C97) | 12,965 | 21.7 | 213.1 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 3,935 | 6.6 | 64.7 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,309 | 5.5 | 54.4 |
| 5 | Cerebrovascular diseases (I60-I69) | 3,037 | 5.1 | 49.9 |
| 6 | Alzheimer's disease (G30) | 2,173 | 3.6 | 35.7 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,483 | 2.5 | 24.4 |
| 8 | Diabetes mellitus (E10-E14) | 1,468 | 2.5 | 24.1 |
| 9 | Influenza and pneumonia (J09-J18) | 1,337 | 2.2 | 22.0 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,052 | 1.8 | 17.3 |
| 11 | Septicemia (A40-A41) | 856 | 1.4 | 14.1 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 678 | 1.1 | 11.1 |
| 13 | Parkinson's disease (G20-G21) | 560 | 0.9 | 9.2 |
| 14 | Pneumonitis due to solids and liquids (J69) | 552 | 0.9 | 9.1 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 547 | 0.9 | 9.0 |
| ... | All other causes (residual) | 11,111 | 18.6 | 182.6 |
| Montana | | | | |
| ... | All causes | 9,942 | 100.0 | 962.5 |
| 1 | Malignant neoplasms (C00-C97) | 2,130 | 21.4 | 206.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,104 | 21.2 | 203.7 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 679 | 6.8 | 65.7 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 637 | 6.4 | 61.7 |
| 5 | Cerebrovascular diseases (I60-I69) | 458 | 4.6 | 44.3 |
| 6 | Diabetes mellitus (E10-E14) | 321 | 3.2 | 31.1 |
| 7 | Alzheimer's disease (G30) | 277 | 2.8 | 26.8 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 272 | 2.7 | 26.3 |
| 9 | Influenza and pneumonia (J09-J18) | 183 | 1.8 | 17.7 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 179 | 1.8 | 17.3 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 121 | 1.2 | 11.7 |
| 12 | Parkinson's disease (G20-G21) | 114 | 1.1 | 11.0 |
| 13 | Septicemia (A40-A41) | 112 | 1.1 | 10.8 |
| 14 | Pneumonitis due to solids and liquids (J69) | 67 | 0.7 | 6.5 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 66 | 0.7 | 6.4 |
| ... | All other causes (residual) | 2,222 | 22.3 | 215.1 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Nebraska | | | | |
| ... | All causes | 16,740 | 100.0 | 882.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,591 | 21.5 | 189.4 |
| 2 | Malignant neoplasms (C00-C97) | 3,514 | 21.0 | 185.3 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 1,174 | 7.0 | 61.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 799 | 4.8 | 42.1 |
| 5 | Cerebrovascular diseases (I60-I69) | 776 | 4.6 | 40.9 |
| 6 | Alzheimer's disease (G30) | 598 | 3.6 | 31.5 |
| 7 | Diabetes mellitus (E10-E14) | 553 | 3.3 | 29.2 |
| 8 | Influenza and pneumonia (J09-J18) | 397 | 2.4 | 20.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 264 | 1.6 | 13.9 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 247 | 1.5 | 13.0 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 223 | 1.3 | 11.8 |
| 12 | Parkinson's disease (G20-G21) | 220 | 1.3 | 11.6 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 165 | 1.0 | 8.7 |
| 14 | Septicemia (A40-A41) | 158 | 0.9 | 8.3 |
| 15 | Pneumonitis due to solids and liquids (J69) | 146 | 0.9 | 7.7 |
| ... | All other causes (residual) | 3,915 | 23.4 | 206.5 |
| Nevada | | | | |
| ... | All causes | 22,879 | 100.0 | 791.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,114 | 26.7 | 211.5 |
| 2 | Malignant neoplasms (C00-C97) | 5,015 | 21.9 | 173.5 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 1,617 | 7.1 | 55.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,340 | 5.9 | 46.4 |
| 5 | Cerebrovascular diseases (I60-I69) | 1,078 | 4.7 | 37.3 |
| 6 | Alzheimer's disease (G30) | 874 | 3.8 | 30.2 |
| 7 | Influenza and pneumonia (J09-J18) | 630 | 2.8 | 21.8 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 558 | 2.4 | 19.3 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 475 | 2.1 | 16.4 |
| 10 | Diabetes mellitus (E10-E14) | 420 | 1.8 | 14.5 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 316 | 1.4 | 10.9 |
| 12 | Parkinson's disease (G20-G21) | 213 | 0.9 | 7.4 |
| 13 | Septicemia (A40-A41) | 205 | 0.9 | 7.1 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 194 | 0.8 | 6.7 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 191 | 0.8 | 6.6 |
| ... | All other causes (residual) | 3,639 | 15.9 | 125.9 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| New Hampshire | | | | |
| ... | All causes | 11,984 | 100.0 | 900.6 |
| 1 | Malignant neoplasms (C00-C97) | 2,773 | 23.1 | 208.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,571 | 21.5 | 193.2 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 815 | 6.8 | 61.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 705 | 5.9 | 53.0 |
| 5 | Cerebrovascular diseases (I60-I69) | 457 | 3.8 | 34.3 |
| 6 | Alzheimer's disease (G30) | 432 | 3.6 | 32.5 |
| 7 | Diabetes mellitus (E10-E14) | 308 | 2.6 | 23.1 |
| 8 | Influenza and pneumonia (J09-J18) | 275 | 2.3 | 20.7 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 228 | 1.9 | 17.1 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 194 | 1.6 | 14.6 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 161 | 1.3 | 12.1 |
| 12 | Parkinson's disease (G20-G21) | 143 | 1.2 | 10.7 |
| 13 | Septicemia (A40-A41) | 119 | 1.0 | 8.9 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 89 | 0.7 | 6.7 |
| 15 | Pneumonitis due to solids and liquids (J69) | 75 | 0.6 | 5.6 |
| ... | All other causes (residual) | 2,639 | 22.0 | 198.3 |
| New Jersey | | | | |
| ... | All causes | 72,271 | 100.0 | 806.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 18,647 | 25.8 | 208.2 |
| 2 | Malignant neoplasms (C00-C97) | 16,270 | 22.5 | 181.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,413 | 4.7 | 38.1 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,218 | 4.5 | 35.9 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 3,202 | 4.4 | 35.7 |
| 6 | Alzheimer's disease (G30) | 2,260 | 3.1 | 25.2 |
| 7 | Septicemia (A40-A41) | 1,957 | 2.7 | 21.8 |
| 8 | Diabetes mellitus (E10-E14) | 1,933 | 2.7 | 21.6 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,582 | 2.2 | 17.7 |
| 10 | Influenza and pneumonia (J09-J18) | 1,402 | 1.9 | 15.7 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 789 | 1.1 | 8.8 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 782 | 1.1 | 8.7 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 733 | 1.0 | 8.2 |
| 14 | Pneumonitis due to solids and liquids (J69) | 725 | 1.0 | 8.1 |
| 15 | Parkinson's disease (G20-G21) | 716 | 1.0 | 8.0 |
| ... | All other causes (residual) | 14,642 | 20.3 | 163.5 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|-------|
| New Mexico | | | | |
| ... | All causes | 17,685 | 100.0 | 848.2 |
| 1 | Malignant neoplasms (C00-C97) | 3,591 | 20.3 | 172.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,508 | 19.8 | 168.2 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,430 | 8.1 | 68.6 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,109 | 6.3 | 53.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 786 | 4.4 | 37.7 |
| 6 | Diabetes mellitus (E10-E14) | 608 | 3.4 | 29.2 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 566 | 3.2 | 27.1 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 500 | 2.8 | 24.0 |
| 9 | Alzheimer's disease (G30) | 483 | 2.7 | 23.2 |
| 10 | Influenza and pneumonia (J09-J18) | 322 | 1.8 | 15.4 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 316 | 1.8 | 15.2 |
| 12 | Septicemia (A40-A41) | 232 | 1.3 | 11.1 |
| 13 | Parkinson's disease (G20-G21) | 172 | 1.0 | 8.2 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 157 | 0.9 | 7.5 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 154 | 0.9 | 7.4 |
| ... | All other causes (residual) | 3,751 | 21.2 | 179.9 |
| New York | | | | |
| ... | All causes | 153,628 | 100.0 | 776.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 44,450 | 28.9 | 224.5 |
| 2 | Malignant neoplasms (C00-C97) | 35,089 | 22.8 | 177.3 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 7,109 | 4.6 | 35.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 6,515 | 4.2 | 32.9 |
| 5 | Cerebrovascular diseases (I60-I69) | 6,292 | 4.1 | 31.8 |
| 6 | Influenza and pneumonia (J09-J18) | 4,881 | 3.2 | 24.7 |
| 7 | Diabetes mellitus (E10-E14) | 4,045 | 2.6 | 20.4 |
| 8 | Alzheimer's disease (G30) | 3,174 | 2.1 | 16.0 |
| 9 | Septicemia (A40-A41) | 2,601 | 1.7 | 13.1 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 2,287 | 1.5 | 11.6 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,245 | 1.5 | 11.3 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,652 | 1.1 | 8.3 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,571 | 1.0 | 7.9 |
| 14 | Parkinson's disease (G20-G21) | 1,247 | 0.8 | 6.3 |
| 15 | Pneumonitis due to solids and liquids (J69) | 1,075 | 0.7 | 5.4 |
| ... | All other causes (residual) | 29,395 | 19.1 | 148.5 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| North Carolina | | | | |
| ... | All causes | 89,133 | 100.0 | 887.5 |
| 1 | Malignant neoplasms (C00-C97) | 19,322 | 21.7 | 192.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 18,474 | 20.7 | 184.0 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 5,221 | 5.9 | 52.0 |
| 4 | Cerebrovascular diseases (I60-I69) | 5,033 | 5.6 | 50.1 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,991 | 5.6 | 49.7 |
| 6 | Alzheimer's disease (G30) | 3,803 | 4.3 | 37.9 |
| 7 | Diabetes mellitus (E10-E14) | 2,746 | 3.1 | 27.3 |
| 8 | Influenza and pneumonia (J09-J18) | 2,115 | 2.4 | 21.1 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,821 | 2.0 | 18.1 |
| 10 | Septicemia (A40-A41) | 1,557 | 1.7 | 15.5 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,406 | 1.6 | 14.0 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,254 | 1.4 | 12.5 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 944 | 1.1 | 9.4 |
| 14 | Parkinson's disease (G20-G21) | 897 | 1.0 | 8.9 |
| 15 | Pneumonitis due to solids and liquids (J69) | 723 | 0.8 | 7.2 |
| ... | All other causes (residual) | 18,826 | 21.1 | 187.5 |
| North Dakota | | | | |
| ... | All causes | 6,223 | 100.0 | 822.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,323 | 21.3 | 174.8 |
| 2 | Malignant neoplasms (C00-C97) | 1,320 | 21.2 | 174.4 |
| 3 | Alzheimer's disease (G30) | 376 | 6.0 | 49.7 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 368 | 5.9 | 48.6 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 351 | 5.6 | 46.4 |
| 6 | Cerebrovascular diseases (I60-I69) | 308 | 4.9 | 40.7 |
| 7 | Diabetes mellitus (E10-E14) | 195 | 3.1 | 25.8 |
| 8 | Influenza and pneumonia (J09-J18) | 172 | 2.8 | 22.7 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 124 | 2.0 | 16.4 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 118 | 1.9 | 15.6 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 86 | 1.4 | 11.4 |
| 12 | Septicemia (A40-A41) | 81 | 1.3 | 10.7 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 79 | 1.3 | 10.4 |
| 14 | Pneumonitis due to solids and liquids (J69) | 77 | 1.2 | 10.2 |
| 15 | Parkinson's disease (G20-G21) | 65 | 1.0 | 8.6 |
| ... | All other causes (residual) | 1,180 | 19.0 | 155.9 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|---------|
| Ohio | | | | |
| ... | All causes | 118,188 | 100.0 | 1,017.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 28,069 | 23.7 | 241.7 |
| 2 | Malignant neoplasms (C00-C97) | 25,396 | 21.5 | 218.7 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 7,211 | 6.1 | 62.1 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 6,756 | 5.7 | 58.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 5,945 | 5.0 | 51.2 |
| 6 | Alzheimer's disease (G30) | 4,643 | 3.9 | 40.0 |
| 7 | Diabetes mellitus (E10-E14) | 3,645 | 3.1 | 31.4 |
| 8 | Influenza and pneumonia (J09-J18) | 2,445 | 2.1 | 21.1 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,100 | 1.8 | 18.1 |
| 10 | Septicemia (A40-A41) | 1,957 | 1.7 | 16.9 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,650 | 1.4 | 14.2 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,506 | 1.3 | 13.0 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,395 | 1.2 | 12.0 |
| 14 | Parkinson's disease (G20-G21) | 1,139 | 1.0 | 9.8 |
| 15 | Pneumonitis due to solids and liquids (J69) | 876 | 0.7 | 7.5 |
| ... | All other causes (residual) | 23,455 | 19.8 | 202.0 |
| Oklahoma | | | | |
| ... | All causes | 39,422 | 100.0 | 1,007.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,310 | 26.2 | 263.6 |
| 2 | Malignant neoplasms (C00-C97) | 8,280 | 21.0 | 211.7 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 2,924 | 7.4 | 74.8 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,422 | 6.1 | 61.9 |
| 5 | Cerebrovascular diseases (I60-I69) | 1,881 | 4.8 | 48.1 |
| 6 | Alzheimer's disease (G30) | 1,498 | 3.8 | 38.3 |
| 7 | Diabetes mellitus (E10-E14) | 1,442 | 3.7 | 36.9 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 790 | 2.0 | 20.2 |
| 9 | Influenza and pneumonia (J09-J18) | 717 | 1.8 | 18.3 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 616 | 1.6 | 15.7 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 597 | 1.5 | 15.3 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 502 | 1.3 | 12.8 |
| 13 | Septicemia (A40-A41) | 419 | 1.1 | 10.7 |
| 14 | Parkinson's disease (G20-G21) | 334 | 0.8 | 8.5 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 324 | 0.8 | 8.3 |
| ... | All other causes (residual) | 6,366 | 16.1 | 162.8 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|---------|
| Oregon | | | | |
| ... | All causes | 35,705 | 100.0 | 886.2 |
| 1 | Malignant neoplasms (C00-C97) | 8,093 | 22.7 | 200.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,859 | 19.2 | 170.2 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 2,119 | 5.9 | 52.6 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,999 | 5.6 | 49.6 |
| 5 | Cerebrovascular diseases (I60-I69) | 1,873 | 5.2 | 46.5 |
| 6 | Alzheimer's disease (G30) | 1,652 | 4.6 | 41.0 |
| 7 | Diabetes mellitus (E10-E14) | 1,148 | 3.2 | 28.5 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 762 | 2.1 | 18.9 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 667 | 1.9 | 16.6 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 565 | 1.6 | 14.0 |
| 11 | Influenza and pneumonia (J09-J18) | 453 | 1.3 | 11.2 |
| 12 | Parkinson's disease (G20-G21) | 429 | 1.2 | 10.6 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 409 | 1.1 | 10.2 |
| 14 | Septicemia (A40-A41) | 227 | 0.6 | 5.6 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 222 | 0.6 | 5.5 |
| ... | All other causes (residual) | 8,228 | 23.0 | 204.2 |
| Pennsylvania | | | | |
| ... | All causes | 132,598 | 100.0 | 1,035.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 32,042 | 24.2 | 250.3 |
| 2 | Malignant neoplasms (C00-C97) | 28,697 | 21.6 | 224.2 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 7,324 | 5.5 | 57.2 |
| 4 | Cerebrovascular diseases (I60-I69) | 6,987 | 5.3 | 54.6 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 6,664 | 5.0 | 52.1 |
| 6 | Alzheimer's disease (G30) | 4,012 | 3.0 | 31.3 |
| 7 | Diabetes mellitus (E10-E14) | 3,777 | 2.8 | 29.5 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,021 | 2.3 | 23.6 |
| 9 | Influenza and pneumonia (J09-J18) | 2,899 | 2.2 | 22.6 |
| 10 | Septicemia (A40-A41) | 2,343 | 1.8 | 18.3 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,894 | 1.4 | 14.8 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,438 | 1.1 | 11.2 |
| 13 | Parkinson's disease (G20-G21) | 1,348 | 1.0 | 10.5 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,108 | 0.8 | 8.7 |
| 15 | Pneumonitis due to solids and liquids (J69) | 1,087 | 0.8 | 8.5 |
| ... | All other causes (residual) | 27,957 | 21.1 | 218.4 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Rhode Island | | | | |
| ... | All causes | 10,163 | 100.0 | 962.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,371 | 23.3 | 224.5 |
| 2 | Malignant neoplasms (C00-C97) | 2,226 | 21.9 | 210.7 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 649 | 6.4 | 61.4 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 510 | 5.0 | 48.3 |
| 5 | Alzheimer's disease (G30) | 453 | 4.5 | 42.9 |
| 6 | Cerebrovascular diseases (I60-I69) | 394 | 3.9 | 37.3 |
| 7 | Diabetes mellitus (E10-E14) | 276 | 2.7 | 26.1 |
| 8 | Influenza and pneumonia (J09-J18) | 246 | 2.4 | 23.3 |
| 9 | Septicemia (A40-A41) | 143 | 1.4 | 13.5 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 142 | 1.4 | 13.4 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 139 | 1.4 | 13.2 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 127 | 1.2 | 12.0 |
| 13 | Parkinson's disease (G20-G21) | 117 | 1.2 | 11.1 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 108 | 1.1 | 10.2 |
| 15 | Pneumonitis due to solids and liquids (J69) | 88 | 0.9 | 8.3 |
| ... | All other causes (residual) | 2,174 | 21.4 | 205.8 |
| South Carolina | | | | |
| ... | All causes | 47,198 | 100.0 | 964.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,092 | 21.4 | 206.1 |
| 2 | Malignant neoplasms (C00-C97) | 9,950 | 21.1 | 203.2 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 2,908 | 6.2 | 59.4 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,737 | 5.8 | 55.9 |
| 5 | Cerebrovascular diseases (I60-I69) | 2,600 | 5.5 | 53.1 |
| 6 | Alzheimer's disease (G30) | 2,453 | 5.2 | 50.1 |
| 7 | Diabetes mellitus (E10-E14) | 1,347 | 2.9 | 27.5 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 886 | 1.9 | 18.1 |
| 9 | Septicemia (A40-A41) | 854 | 1.8 | 17.4 |
| 10 | Influenza and pneumonia (J09-J18) | 848 | 1.8 | 17.3 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 742 | 1.6 | 15.2 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 716 | 1.5 | 14.6 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 482 | 1.0 | 9.8 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 448 | 0.9 | 9.2 |
| 15 | Parkinson's disease (G20-G21) | 420 | 0.9 | 8.6 |
| ... | All other causes (residual) | 9,715 | 20.6 | 198.4 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|---------|
| South Dakota | | | | |
| ... | All causes | 7,731 | 100.0 | 900.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,711 | 22.1 | 199.3 |
| 2 | Malignant neoplasms (C00-C97) | 1,640 | 21.2 | 191.0 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 502 | 6.5 | 58.5 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 469 | 6.1 | 54.6 |
| 5 | Alzheimer's disease (G30) | 421 | 5.4 | 49.0 |
| 6 | Cerebrovascular diseases (I60-I69) | 383 | 5.0 | 44.6 |
| 7 | Diabetes mellitus (E10-E14) | 282 | 3.6 | 32.8 |
| 8 | Influenza and pneumonia (J09-J18) | 213 | 2.8 | 24.8 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 173 | 2.2 | 20.2 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 139 | 1.8 | 16.2 |
| 11 | Septicemia (A40-A41) | 120 | 1.6 | 14.0 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 104 | 1.3 | 12.1 |
| 13 | Parkinson's disease (G20-G21) | 80 | 1.0 | 9.3 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 79 | 1.0 | 9.2 |
| 15 | Pneumonitis due to solids and liquids (J69) | 58 | 0.8 | 6.8 |
| ... | All other causes (residual) | 1,357 | 17.6 | 158.1 |
| Tennessee | | | | |
| ... | All causes | 66,570 | 100.0 | 1,008.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 15,730 | 23.6 | 238.3 |
| 2 | Malignant neoplasms (C00-C97) | 14,214 | 21.4 | 215.4 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 4,239 | 6.4 | 64.2 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,873 | 5.8 | 58.7 |
| 5 | Cerebrovascular diseases (I60-I69) | 3,447 | 5.2 | 52.2 |
| 6 | Alzheimer's disease (G30) | 3,122 | 4.7 | 47.3 |
| 7 | Diabetes mellitus (E10-E14) | 1,798 | 2.7 | 27.2 |
| 8 | Influenza and pneumonia (J09-J18) | 1,723 | 2.6 | 26.1 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,090 | 1.6 | 16.5 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,068 | 1.6 | 16.2 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 957 | 1.4 | 14.5 |
| 12 | Septicemia (A40-A41) | 924 | 1.4 | 14.0 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 649 | 1.0 | 9.8 |
| 14 | Parkinson's disease (G20-G21) | 604 | 0.9 | 9.2 |
| 15 | Pneumonitis due to solids and liquids (J69) | 556 | 0.8 | 8.4 |
| ... | All other causes (residual) | 12,576 | 18.9 | 190.5 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|---------|-------------------------|-------|
| Texas | | | | |
| ... | All causes | 189,654 | 100.0 | 690.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 43,298 | 22.8 | 157.6 |
| 2 | Malignant neoplasms (C00-C97) | 39,121 | 20.6 | 142.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 10,485 | 5.5 | 38.2 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 10,231 | 5.4 | 37.2 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 9,976 | 5.3 | 36.3 |
| 6 | Alzheimer's disease (G30) | 8,903 | 4.7 | 32.4 |
| 7 | Diabetes mellitus (E10-E14) | 5,521 | 2.9 | 20.1 |
| 8 | Septicemia (A40-A41) | 4,381 | 2.3 | 15.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4,052 | 2.1 | 14.8 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,844 | 2.0 | 14.0 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 3,403 | 1.8 | 12.4 |
| 12 | Influenza and pneumonia (J09-J18) | 3,214 | 1.7 | 11.7 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 2,089 | 1.1 | 7.6 |
| 14 | Parkinson's disease (G20-G21) | 1,983 | 1.0 | 7.2 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,538 | 0.8 | 5.6 |
| ... | All other causes (residual) | 37,615 | 19.8 | 136.9 |
| Utah | | | | |
| ... | All causes | 17,334 | 100.0 | 578.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,598 | 20.8 | 120.1 |
| 2 | Malignant neoplasms (C00-C97) | 3,091 | 17.8 | 103.2 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,223 | 7.1 | 40.8 |
| 4 | Alzheimer's disease (G30) | 906 | 5.2 | 30.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 888 | 5.1 | 29.6 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 811 | 4.7 | 27.1 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 630 | 3.6 | 21.0 |
| 8 | Diabetes mellitus (E10-E14) | 604 | 3.5 | 20.2 |
| 9 | Influenza and pneumonia (J09-J18) | 372 | 2.1 | 12.4 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 363 | 2.1 | 12.1 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 244 | 1.4 | 8.1 |
| 12 | Parkinson's disease (G20-G21) | 229 | 1.3 | 7.6 |
| 13 | Septicemia (A40-A41) | 166 | 1.0 | 5.5 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 136 | 0.8 | 4.5 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 112 | 0.6 | 3.7 |
| ... | All other causes (residual) | 3,961 | 22.9 | 132.2 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Vermont | | | | |
| ... | All causes | 5,919 | 100.0 | 945.5 |
| 1 | Malignant neoplasms (C00-C97) | 1,399 | 23.6 | 223.5 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,311 | 22.1 | 209.4 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 357 | 6.0 | 57.0 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 346 | 5.8 | 55.3 |
| 5 | Cerebrovascular diseases (I60-I69) | 307 | 5.2 | 49.0 |
| 6 | Alzheimer's disease (G30) | 298 | 5.0 | 47.6 |
| 7 | Diabetes mellitus (E10-E14) | 159 | 2.7 | 25.4 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 103 | 1.7 | 16.5 |
| 9 | Influenza and pneumonia (J09-J18) | 85 | 1.4 | 13.6 |
| 10 | Parkinson's disease (G20-G21) | 83 | 1.4 | 13.3 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 71 | 1.2 | 11.3 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 57 | 1.0 | 9.1 |
| 13 | Septicemia (A40-A41) | 50 | 0.8 | 8.0 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 39 | 0.7 | 6.2 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 27 | 0.5 | 4.3 |
| ... | All other causes (residual) | 1,227 | 20.7 | 196.0 |
| Virginia | | | | |
| ... | All causes | 65,577 | 100.0 | 782.3 |
| 1 | Malignant neoplasms (C00-C97) | 14,947 | 22.8 | 178.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 14,077 | 21.5 | 167.9 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,429 | 5.2 | 40.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 3,393 | 5.2 | 40.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 3,370 | 5.1 | 40.2 |
| 6 | Alzheimer's disease (G30) | 2,248 | 3.4 | 26.8 |
| 7 | Diabetes mellitus (E10-E14) | 2,044 | 3.1 | 24.4 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,470 | 2.2 | 17.5 |
| 9 | Influenza and pneumonia (J09-J18) | 1,416 | 2.2 | 16.9 |
| 10 | Septicemia (A40-A41) | 1,187 | 1.8 | 14.2 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,118 | 1.7 | 13.3 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 943 | 1.4 | 11.2 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 684 | 1.0 | 8.2 |
| 14 | Parkinson's disease (G20-G21) | 638 | 1.0 | 7.6 |
| 15 | Pneumonitis due to solids and liquids (J69) | 590 | 0.9 | 7.0 |
| ... | All other causes (residual) | 14,023 | 21.4 | 167.3 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|---------|
| Washington | | | | |
| ... | All causes | 54,595 | 100.0 | 761.4 |
| 1 | Malignant neoplasms (C00-C97) | 12,687 | 23.2 | 176.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,025 | 20.2 | 153.8 |
| 3 | Alzheimer's disease (G30) | 3,490 | 6.4 | 48.7 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,192 | 5.8 | 44.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 3,154 | 5.8 | 44.0 |
| 6 | Cerebrovascular diseases (I60-I69) | 2,703 | 5.0 | 37.7 |
| 7 | Diabetes mellitus (E10-E14) | 1,811 | 3.3 | 25.3 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,137 | 2.1 | 15.9 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,024 | 1.9 | 14.3 |
| 10 | Influenza and pneumonia (J09-J18) | 851 | 1.6 | 11.9 |
| 11 | Parkinson's disease (G20-G21) | 654 | 1.2 | 9.1 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 594 | 1.1 | 8.3 |
| 13 | Pneumonitis due to solids and liquids (J69) | 590 | 1.1 | 8.2 |
| 14 | Septicemia (A40-A41) | 501 | 0.9 | 7.0 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 474 | 0.9 | 6.6 |
| ... | All other causes (residual) | 10,708 | 19.6 | 149.3 |
| West Virginia | | | | |
| ... | All causes | 22,752 | 100.0 | 1,233.8 |
| 1 | Malignant neoplasms (C00-C97) | 4,839 | 21.3 | 262.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,727 | 20.8 | 256.3 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 1,628 | 7.2 | 88.3 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,516 | 6.7 | 82.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 1,079 | 4.7 | 58.5 |
| 6 | Diabetes mellitus (E10-E14) | 784 | 3.4 | 42.5 |
| 7 | Alzheimer's disease (G30) | 738 | 3.2 | 40.0 |
| 8 | Influenza and pneumonia (J09-J18) | 526 | 2.3 | 28.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 510 | 2.2 | 27.7 |
| 10 | Septicemia (A40-A41) | 348 | 1.5 | 18.9 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 340 | 1.5 | 18.4 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 338 | 1.5 | 18.3 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 297 | 1.3 | 16.1 |
| 14 | Parkinson's disease (G20-G21) | 163 | 0.7 | 8.8 |
| 15 | Pneumonitis due to solids and liquids (J69) | 159 | 0.7 | 8.6 |
| ... | All other causes (residual) | 4,760 | 20.9 | 258.1 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death: United States and each State, 2015

[Rates per 100,000 population]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>) and State | Number | Percent of total deaths | Rate |
|-------------------|--|--------|-------------------------|-------|
| Wisconsin | | | | |
| ... | All causes | 51,264 | 100.0 | 888.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,473 | 22.4 | 198.8 |
| 2 | Malignant neoplasms (C00-C97) | 11,423 | 22.3 | 197.9 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,206 | 6.3 | 55.6 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 2,846 | 5.6 | 49.3 |
| 5 | Cerebrovascular diseases (I60-I69) | 2,618 | 5.1 | 45.4 |
| 6 | Alzheimer's disease (G30) | 2,087 | 4.1 | 36.2 |
| 7 | Diabetes mellitus (E10-E14) | 1,382 | 2.7 | 23.9 |
| 8 | Influenza and pneumonia (J09-J18) | 1,052 | 2.1 | 18.2 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 996 | 1.9 | 17.3 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 877 | 1.7 | 15.2 |
| 11 | Parkinson's disease (G20-G21) | 600 | 1.2 | 10.4 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 591 | 1.2 | 10.2 |
| 13 | Septicemia (A40-A41) | 517 | 1.0 | 9.0 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 485 | 0.9 | 8.4 |
| 15 | Pneumonitis due to solids and liquids (J69) | 407 | 0.8 | 7.1 |
| ... | All other causes (residual) | 10,704 | 20.9 | 185.5 |
| Wyoming | | | | |
| ... | All causes | 4,778 | 100.0 | 815.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,030 | 21.6 | 175.7 |
| 2 | Malignant neoplasms (C00-C97) | 931 | 19.5 | 158.8 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 400 | 8.4 | 68.2 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 368 | 7.7 | 62.8 |
| 5 | Cerebrovascular diseases (I60-I69) | 198 | 4.1 | 33.8 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 157 | 3.3 | 26.8 |
| 7 | Alzheimer's disease (G30) | 151 | 3.2 | 25.8 |
| 8 | Diabetes mellitus (E10-E14) | 136 | 2.8 | 23.2 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 118 | 2.5 | 20.1 |
| 10 | Influenza and pneumonia (J09-J18) | 99 | 2.1 | 16.9 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 86 | 1.8 | 14.7 |
| 12 | Septicemia (A40-A41) | 46 | 1.0 | 7.8 |
| 13 | Parkinson's disease (G20-G21) | 42 | 0.9 | 7.2 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 41 | 0.9 | 7.0 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 34 | 0.7 | 5.8 |
| ... | All other causes (residual) | 941 | 19.7 | 160.6 |

... Category not applicable

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.