

Person with disabilities experiencing problems accessing
NCHS birth and death tables should contact
nchsed@cdc.gov, or call 301-458-4688

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Arkansas												
All races, both sexes.....	2,918	8	1	1	5	4	7	6	5	12	24	47
Male.....	1,487	5	-	-	4	1	4	5	1	9	11	25
Female.....	1,431	3	1	1	1	3	3	1	4	3	13	22
White, both sexes.....	2,692	5	1	-	4	2	6	5	3	11	20	40
Male.....	1,374	2	-	-	3	-	3	4	-	9	10	22
Female.....	1,318	3	1	-	1	2	3	1	3	2	10	18
Black, both sexes.....	218	3	-	1	1	2	1	1	2	1	4	6
Male.....	110	3	-	-	1	1	1	1	1	-	1	2
Female.....	108	-	-	1	-	1	-	-	1	1	3	4
American Indian 1/, both sexes.....	3	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	22,257	55	29	19	14	22	24	33	35	64	127	263
Male.....	10,562	30	14	10	12	15	14	19	19	30	74	142
Female.....	11,695	25	15	9	2	7	10	14	16	34	53	121
White, both sexes.....	19,222	43	20	13	9	19	19	24	25	42	96	195
Male.....	8,943	25	7	6	7	14	11	14	14	21	59	106
Female.....	10,279	18	13	7	2	5	8	10	11	21	37	89
Black, both sexes.....	1,269	7	5	4	3	1	3	8	5	11	19	47
Male.....	632	3	3	2	3	-	2	4	3	5	7	24
Female.....	637	4	2	2	-	1	1	4	2	6	12	23
American Indian 1/, both sexes.....	116	1	-	-	-	-	-	-	-	3	3	3
Male.....	59	-	-	-	-	-	-	-	-	1	2	2
Female.....	57	1	-	-	-	-	-	-	-	2	1	1
Asian or Pacific Islander 2/, both sexes.....	1,650	4	4	2	2	2	2	1	5	8	9	18
Male.....	928	2	4	2	2	1	1	1	2	3	6	10
Female.....	722	2	-	-	-	1	1	-	3	5	3	8

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Iowa												
All races, both sexes.....	3,010	1	1	2	1	2	4	5	4	5	18	25
Male.....	1,484	1	1	1	1	2	3	4	3	1	8	12
Female.....	1,526	-	-	1	-	-	1	1	1	4	10	13
White, both sexes.....	2,980	1	-	2	1	2	4	4	4	5	16	23
Male.....	1,463	1	-	1	1	2	3	3	3	1	7	10
Female.....	1,517	-	-	1	-	-	1	1	1	4	9	13
Black, both sexes.....	29	-	1	-	-	-	-	1	-	-	2	2
Male.....	20	-	1	-	-	-	-	1	-	-	1	2
Female.....	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes.....	2,645	2	2	3	-	2	3	2	6	3	14	26
Male.....	1,291	2	1	2	-	-	2	-	2	2	12	6
Female.....	1,354	-	1	1	-	2	1	2	4	1	2	20
White, both sexes.....	2,536	1	2	2	-	1	3	2	5	3	12	21
Male.....	1,229	1	1	1	-	-	2	-	2	2	10	2
Female.....	1,307	-	1	1	-	1	1	2	3	1	2	19
Black, both sexes.....	69	1	-	1	-	-	-	-	1	-	2	4
Male.....	41	1	-	1	-	-	-	-	-	-	2	3
Female.....	28	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes.....	32	-	-	-	-	-	-	-	-	-	-	1
Male.....	17	-	-	-	-	-	-	-	-	-	-	1
Female.....	15	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	8	-	-	-	-	1	-	-	-	-	-	-
Male.....	4	-	-	-	-	-	-	-	-	-	-	-
Female.....	4	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
Massachusetts												
All races, both sexes.....	5,139	6	3	-	-	1	8	2	3	15	27	47
Male.....	2,304	4	-	-	-	-	6	1	3	10	14	27
Female.....	2,835	2	3	-	-	1	2	1	-	5	13	20
White, both sexes.....	4,971	5	3	-	-	1	6	2	3	12	25	39
Male.....	2,221	3	-	-	-	-	5	1	3	8	12	24
Female.....	2,750	2	3	-	-	1	1	1	-	4	13	15
Black, both sexes.....	115	1	-	-	-	-	2	-	-	3	1	8
Male.....	54	1	-	-	-	-	1	-	-	2	1	3
Female.....	61	-	-	-	-	-	1	-	-	1	-	5
American Indian 1/, both sexes.....	4	-	-	-	-	-	-	-	-	-	1	-
Male.....	3	-	-	-	-	-	-	-	-	-	1	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	49	-	-	-	-	-	-	-	-	-	-	-
Male.....	26	-	-	-	-	-	-	-	-	-	-	-
Female.....	23	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes.....	7,707	14	9	5	7	9	8	14	11	31	47	93
Male.....	3,751	6	6	4	3	5	3	7	4	18	21	56
Female.....	3,956	8	3	1	4	4	5	7	7	13	26	37
White, both sexes.....	6,892	7	6	2	3	5	3	8	6	13	36	63
Male.....	3,342	3	4	2	1	4	2	5	3	7	19	41
Female.....	3,550	4	2	-	2	1	1	3	3	6	17	22
Black, both sexes.....	748	6	2	3	4	4	5	6	5	15	11	30
Male.....	375	3	1	2	2	1	1	2	1	10	2	15
Female.....	373	3	1	1	2	3	4	4	4	5	9	15
American Indian 1/, both sexes.....	36	-	-	-	-	-	-	-	-	2	-	-
Male.....	16	-	-	-	-	-	-	-	-	1	-	-
Female.....	20	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	31	1	1	-	-	-	-	-	-	1	-	-
Male.....	18	-	1	-	-	-	-	-	-	-	-	-
Female.....	13	1	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
X. Diseases of the respiratory system (J00-J98,U04)												
New Mexico												
All races, both sexes.....	1,489	-	6	-	1	1	1	4	6	11	14	21
Male.....	750	-	4	-	1	-	-	3	2	5	8	13
Female.....	739	-	2	-	-	1	1	1	4	6	6	8
White, both sexes.....	1,386	-	4	-	1	1	-	4	5	6	11	18
Male.....	701	-	3	-	1	-	-	3	2	2	6	11
Female.....	685	-	1	-	-	1	-	1	3	4	5	7
Black, both sexes.....	12	-	1	-	-	-	-	-	-	1	-	-
Male.....	8	-	-	-	-	-	-	-	-	-	-	-
Female.....	4	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	87	-	1	-	-	-	1	-	1	4	3	3
Male.....	40	-	1	-	-	-	-	-	-	3	2	2
Female.....	47	-	-	-	-	-	1	-	1	1	1	1
Asian or Pacific Islander 2/, both sexes.....	4	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	13,246	42	19	12	9	8	24	27	30	48	88	186
Male.....	6,057	24	9	8	5	5	14	22	18	24	54	104
Female.....	7,189	18	10	4	4	3	10	5	12	24	34	82
White, both sexes.....	11,591	23	10	9	3	3	19	18	21	28	65	132
Male.....	5,269	15	4	7	2	2	11	15	15	12	39	77
Female.....	6,322	8	6	2	1	1	8	3	6	16	26	55
Black, both sexes.....	1,368	16	7	2	6	4	3	8	9	19	19	45
Male.....	631	8	3	-	3	3	2	6	3	11	13	25
Female.....	737	8	4	2	3	1	1	2	6	8	6	20
American Indian 1/, both sexes.....	22	-	-	-	-	-	-	-	-	-	1	2
Male.....	12	-	-	-	-	-	-	-	-	-	1	1
Female.....	10	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	265	3	2	1	-	1	2	1	-	1	3	7
Male.....	145	1	2	1	-	-	1	1	-	1	1	1
Female.....	120	2	-	-	-	1	1	-	-	-	2	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

United States

All races, both sexes.....	4,985	7,732	12,197	17,207	24,486	34,903	43,139	39,657	24,280	9,014	1,902	10
Male.....	2,751	4,307	6,624	9,170	12,903	17,908	21,183	17,882	9,273	2,752	441	10
Female.....	2,234	3,425	5,573	8,037	11,583	16,995	21,956	21,775	15,007	6,262	1,461	-
White, both sexes.....	4,001	6,447	10,732	15,354	22,108	31,963	39,899	36,860	22,600	8,311	1,665	9
Male.....	2,232	3,607	5,746	8,131	11,572	16,302	19,505	16,583	8,597	2,523	379	9
Female.....	1,769	2,840	4,986	7,223	10,536	15,661	20,394	20,277	14,003	5,788	1,286	-
Black, both sexes.....	872	1,122	1,243	1,530	1,916	2,255	2,402	2,014	1,167	500	178	-
Male.....	454	610	742	853	1,064	1,226	1,223	879	438	141	37	-
Female.....	418	512	501	677	852	1,029	1,179	1,135	729	359	141	-
American Indian 1/, both sexes.....	57	74	101	143	143	164	155	125	61	30	7	-
Male.....	33	37	63	71	83	72	75	51	25	11	2	-
Female.....	24	37	38	72	60	92	80	74	36	19	5	-
Asian or Pacific Islander 2/, both sexes.....	55	89	121	180	319	521	683	658	452	173	52	1
Male.....	32	53	73	115	184	308	380	369	213	77	23	1
Female.....	23	36	48	65	135	213	303	289	239	96	29	-

Alabama

All races, both sexes.....	116	160	331	420	550	697	814	646	358	118	22	-
Male.....	72	86	190	240	306	360	386	304	125	41	4	-
Female.....	44	74	141	180	244	337	428	342	233	77	18	-
White, both sexes.....	89	129	276	360	467	615	731	567	309	99	14	-
Male.....	57	69	156	207	261	316	339	265	108	33	2	-
Female.....	32	60	120	153	206	299	392	302	201	66	12	-
Black, both sexes.....	27	30	53	60	83	80	82	77	48	17	8	-
Male.....	15	16	33	33	45	42	46	39	17	8	2	-
Female.....	12	14	20	27	38	38	36	38	31	9	6	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	1	1	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	2	-	-	1	1	2	-	1	-	-
Male.....	-	1	1	-	-	1	1	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	2	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Alaska												
All races, both sexes.....	13	8	25	34	42	63	45	26	17	5	-	-
Male.....	5	2	17	17	24	28	25	13	4	1	-	-
Female.....	8	6	8	17	18	35	20	13	13	4	-	-
White, both sexes.....	9	6	19	23	31	49	37	19	15	5	-	-
Male.....	3	2	13	13	16	22	19	9	4	1	-	-
Female.....	6	4	6	10	15	27	18	10	11	4	-	-
Black, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	3	1	6	9	11	13	8	4	2	-	-	-
Male.....	1	-	4	4	8	5	6	1	-	-	-	-
Female.....	2	1	2	5	3	8	2	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	2	-	1	-	3	-	-	-	-
Male.....	1	-	-	-	-	1	-	3	-	-	-	-
Female.....	-	-	-	2	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes.....	96	142	250	336	529	696	816	717	382	157	23	-
Male.....	49	76	123	189	281	364	402	360	160	67	7	-
Female.....	47	66	127	147	248	332	414	357	222	90	16	-
White, both sexes.....	85	136	236	320	505	660	790	690	365	147	19	-
Male.....	44	73	117	180	270	347	395	345	155	62	6	-
Female.....	41	63	119	140	235	313	395	345	210	85	13	-
Black, both sexes.....	6	3	4	3	6	12	9	8	1	2	1	-
Male.....	2	1	1	2	1	5	1	5	-	1	-	-
Female.....	4	2	3	1	5	7	8	3	1	1	1	-
American Indian 1/, both sexes.....	4	3	8	12	12	21	14	15	14	8	3	-
Male.....	2	2	4	6	7	9	5	7	5	4	1	-
Female.....	2	1	4	6	5	12	9	8	9	4	2	-
Asian or Pacific Islander 2/, both sexes.....	1	-	2	1	6	3	3	4	2	-	-	-
Male.....	1	-	1	1	3	3	1	3	-	-	-	-
Female.....	-	-	1	-	3	-	2	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Arkansas

All races, both sexes.....	83	108	188	264	317	478	520	450	258	110	22	-
Male.....	49	58	116	146	172	275	259	204	101	38	4	-
Female.....	34	50	72	118	145	203	261	246	157	72	18	-
White, both sexes.....	56	95	174	243	293	455	493	425	240	101	20	-
Male.....	34	53	106	135	159	263	245	192	95	35	4	-
Female.....	22	42	68	108	134	192	248	233	145	66	16	-
Black, both sexes.....	27	13	14	19	23	20	27	25	17	9	2	-
Male.....	15	5	10	11	12	11	14	12	6	3	-	-
Female.....	12	8	4	8	11	9	13	13	11	6	2	-
American Indian 1/, both sexes.....	-	-	-	2	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	3	-	-	1	-	-	-
Male.....	-	-	-	-	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	2	-	-	1	-	-	-

California

All races, both sexes.....	445	670	1,025	1,482	2,238	3,304	4,324	4,194	2,619	1,017	253	1
Male.....	227	402	518	790	1,139	1,682	2,054	1,925	1,045	333	67	1
Female.....	218	268	507	692	1,099	1,622	2,270	2,269	1,574	684	186	-
White, both sexes.....	355	540	871	1,265	1,924	2,875	3,811	3,692	2,301	878	204	1
Male.....	182	328	430	661	964	1,436	1,777	1,647	898	285	50	1
Female.....	173	212	441	604	960	1,439	2,034	2,045	1,403	593	154	-
Black, both sexes.....	58	93	101	120	160	166	174	139	80	44	21	-
Male.....	30	52	58	62	86	89	87	63	36	9	4	-
Female.....	28	41	43	58	74	77	87	76	44	35	17	-
American Indian 1/, both sexes.....	7	10	5	11	19	14	13	19	6	2	-	-
Male.....	2	6	2	7	10	5	8	10	4	-	-	-
Female.....	5	4	3	4	9	9	5	9	2	2	-	-
Asian or Pacific Islander 2/, both sexes.....	25	27	48	86	135	249	326	344	232	93	28	-
Male.....	13	16	28	60	79	152	182	205	107	39	13	-
Female.....	12	11	20	26	56	97	144	139	125	54	15	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Colorado

All races, both sexes.....	88	125	187	239	338	500	583	588	349	132	33	-
Male.....	53	78	104	132	181	269	305	277	154	41	8	-
Female.....	35	47	83	107	157	231	278	311	195	91	25	-
White, both sexes.....	80	120	177	229	325	491	569	577	337	130	33	-
Male.....	48	75	101	125	173	266	296	275	149	41	8	-
Female.....	32	45	76	104	152	225	273	302	188	89	25	-
Black, both sexes.....	7	2	9	6	9	3	6	4	10	1	-	-
Male.....	4	2	3	5	5	1	5	1	4	-	-	-
Female.....	3	-	6	1	4	2	1	3	6	1	-	-
American Indian 1/, both sexes.....	-	3	1	3	1	4	2	5	-	-	-	-
Male.....	-	1	-	1	1	1	2	1	-	-	-	-
Female.....	-	2	1	2	-	3	-	4	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	1	3	2	6	2	2	1	-	-
Male.....	1	-	-	1	2	1	2	-	1	-	-	-
Female.....	-	-	-	-	1	1	4	2	1	1	-	-

Connecticut

All races, both sexes.....	34	71	127	167	221	377	582	620	452	176	36	-
Male.....	24	36	65	82	110	205	280	299	163	64	5	-
Female.....	10	35	62	85	111	172	302	321	289	112	31	-
White, both sexes.....	28	59	114	156	207	358	555	603	445	170	34	-
Male.....	20	28	60	75	104	194	270	289	162	60	5	-
Female.....	8	31	54	81	103	164	285	314	283	110	29	-
Black, both sexes.....	6	10	13	11	13	15	25	17	6	4	2	-
Male.....	4	6	5	7	5	9	10	10	1	2	-	-
Female.....	2	4	8	4	8	6	15	7	5	2	2	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	1	4	2	-	1	2	-	-
Male.....	-	1	-	-	1	2	-	-	-	2	-	-
Female.....	-	-	-	-	-	2	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Delaware

All races, both sexes.....	15	18	36	41	78	119	122	110	53	15	3	-
Male.....	5	11	24	23	38	51	61	41	16	5	1	-
Female.....	10	7	12	18	40	68	61	69	37	10	2	-
White, both sexes.....	11	15	32	31	68	107	110	103	50	15	3	-
Male.....	3	10	22	18	32	45	54	37	14	5	1	-
Female.....	8	5	10	13	36	62	56	66	36	10	2	-
Black, both sexes.....	4	3	4	9	10	12	12	7	3	-	-	-
Male.....	2	1	2	4	6	6	7	4	2	-	-	-
Female.....	2	2	2	5	4	6	5	3	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Dist. of Columbia

All races, both sexes.....	15	14	16	18	21	38	42	37	37	16	2	-
Male.....	9	9	11	12	16	20	21	14	15	2	1	-
Female.....	6	5	5	6	5	18	21	23	22	14	1	-
White, both sexes.....	2	2	4	4	5	9	17	11	21	6	-	-
Male.....	2	1	4	4	3	4	8	7	7	1	-	-
Female.....	-	1	-	-	2	5	9	4	14	5	-	-
Black, both sexes.....	13	12	12	14	16	28	25	26	14	10	2	-
Male.....	7	8	7	8	13	15	13	7	7	1	1	-
Female.....	6	4	5	6	3	13	12	19	7	9	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	2	-	-	-
Male.....	-	-	-	-	-	1	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Florida

All races, both sexes.....	287	457	783	1,086	1,608	2,270	2,783	2,495	1,472	478	81	-
Male.....	164	267	440	584	857	1,182	1,378	1,113	556	171	22	-
Female.....	123	190	343	502	751	1,088	1,405	1,382	916	307	59	-
White, both sexes.....	239	393	697	988	1,498	2,152	2,660	2,396	1,402	456	71	-
Male.....	147	231	384	530	794	1,113	1,308	1,072	534	159	20	-
Female.....	92	162	313	458	704	1,039	1,352	1,324	868	297	51	-
Black, both sexes.....	45	60	79	93	102	109	113	92	64	20	10	-
Male.....	15	33	51	54	61	65	64	38	21	11	2	-
Female.....	30	27	28	39	41	44	49	54	43	9	8	-
American Indian 1/, both sexes.....	2	-	3	1	2	1	4	-	-	-	-	-
Male.....	2	-	2	-	-	-	2	-	-	-	-	-
Female.....	-	-	1	1	2	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	4	4	4	6	8	6	7	6	2	-	-
Male.....	-	3	3	-	2	4	4	3	1	1	-	-
Female.....	1	1	1	4	4	4	2	4	5	1	-	-

Georgia

All races, both sexes.....	168	284	409	553	787	911	1,164	946	554	238	34	-
Male.....	96	157	231	301	412	443	537	404	184	68	9	-
Female.....	72	127	178	252	375	468	627	542	370	170	25	-
White, both sexes.....	119	207	325	456	656	796	1,019	816	480	201	23	-
Male.....	67	117	181	242	336	385	463	351	162	56	4	-
Female.....	52	90	144	214	320	411	556	465	318	145	19	-
Black, both sexes.....	48	77	84	92	123	108	134	126	72	36	11	-
Male.....	28	40	50	57	70	53	66	52	20	11	5	-
Female.....	20	37	34	35	53	55	68	74	52	25	6	-
American Indian 1/, both sexes.....	-	-	-	1	2	2	1	-	-	-	-	-
Male.....	-	-	-	-	1	2	-	-	-	-	-	-
Female.....	-	-	-	1	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	4	6	5	10	4	2	1	-	-
Male.....	1	-	-	2	5	3	8	1	2	1	-	-
Female.....	-	-	-	2	1	2	2	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Hawaii

All races, both sexes.....	12	25	32	33	63	106	162	155	105	48	12	1
Male.....	8	16	19	17	36	61	94	79	48	24	8	1
Female.....	4	9	13	16	27	45	68	76	57	24	4	-
White, both sexes.....	6	12	14	15	24	33	43	48	30	18	2	1
Male.....	4	8	7	6	13	19	28	23	12	8	1	1
Female.....	2	4	7	9	11	14	15	25	18	10	1	-
Black, both sexes.....	-	-	-	-	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	6	13	18	18	39	72	119	106	75	30	10	-
Male.....	4	8	12	11	23	42	66	56	36	16	7	-
Female.....	2	5	6	7	16	30	53	50	39	14	3	-

Idaho

All races, both sexes.....	25	39	55	81	123	165	217	185	126	45	8	-
Male.....	19	16	29	53	66	81	101	95	45	18	2	-
Female.....	6	23	26	28	57	84	116	90	81	27	6	-
White, both sexes.....	24	37	53	81	122	162	214	183	126	45	8	-
Male.....	18	16	29	53	66	79	99	94	45	18	2	-
Female.....	6	21	24	28	56	83	115	89	81	27	6	-
Black, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	2	1	-	1	3	1	-	-	-	-	-
Male.....	1	-	-	-	-	2	1	-	-	-	-	-
Female.....	-	2	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	-	2	2	-	-	-	-
Male.....	-	-	-	-	-	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Illinois

All races, both sexes.....	220	296	460	639	900	1,434	1,778	1,666	1,099	409	93	-
Male.....	120	162	266	346	457	735	854	720	389	117	17	-
Female.....	100	134	194	293	443	699	924	946	710	292	76	-
White, both sexes.....	164	218	377	536	781	1,273	1,612	1,538	1,012	390	84	-
Male.....	91	121	215	286	397	645	770	670	355	112	13	-
Female.....	73	97	162	250	384	628	842	868	657	278	71	-
Black, both sexes.....	54	72	79	97	113	143	144	113	70	16	7	-
Male.....	28	37	47	56	57	77	70	43	29	3	3	-
Female.....	26	35	32	41	56	66	74	70	41	13	4	-
American Indian 1/, both sexes.....	-	-	1	3	1	-	2	-	1	-	-	-
Male.....	-	-	1	2	-	-	1	-	-	-	-	-
Female.....	-	-	-	1	1	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	6	3	3	5	18	20	15	16	3	2	-
Male.....	1	4	3	2	3	13	13	7	5	2	1	-
Female.....	1	2	-	1	2	5	7	8	11	1	1	-

Indiana

All races, both sexes.....	127	218	312	459	620	818	1,029	927	500	182	55	-
Male.....	70	105	172	239	326	416	517	433	187	40	10	-
Female.....	57	113	140	220	294	402	512	494	313	142	45	-
White, both sexes.....	117	195	304	444	593	779	989	893	478	177	54	-
Male.....	65	93	168	231	317	402	491	418	180	38	10	-
Female.....	52	102	136	213	276	377	498	475	298	139	44	-
Black, both sexes.....	10	23	8	14	26	38	40	33	22	5	1	-
Male.....	5	12	4	8	9	14	26	15	7	2	-	-
Female.....	5	11	4	6	17	24	14	18	15	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Iowa												
All races, both sexes.....	42	81	137	173	307	432	578	588	414	157	33	-
Male.....	27	43	74	83	175	231	321	279	166	41	7	-
Female.....	15	38	63	90	132	201	257	309	248	116	26	-
White, both sexes.....	42	78	134	172	301	425	575	588	413	157	33	-
Male.....	27	42	71	82	171	227	318	279	166	41	7	-
Female.....	15	36	63	90	130	198	257	309	247	116	26	-
Black, both sexes.....	-	3	3	1	5	7	3	-	1	-	-	-
Male.....	-	1	3	1	3	4	3	-	-	-	-	-
Female.....	-	2	-	-	2	3	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes.....	45	82	119	172	263	401	551	463	330	134	22	-
Male.....	24	37	70	92	149	209	285	236	125	31	4	-
Female.....	21	45	49	80	114	192	266	227	205	103	18	-
White, both sexes.....	41	73	111	165	254	381	531	449	325	132	22	-
Male.....	22	34	64	86	144	197	275	229	122	31	4	-
Female.....	19	39	47	79	110	184	256	220	203	101	18	-
Black, both sexes.....	3	5	6	4	3	12	12	9	4	2	-	-
Male.....	1	2	5	4	2	8	6	4	2	-	-	-
Female.....	2	3	1	-	1	4	6	5	2	2	-	-
American Indian 1/, both sexes.....	-	4	2	3	3	7	6	5	1	-	-	-
Male.....	-	1	1	2	2	3	3	3	1	-	-	-
Female.....	-	3	1	1	1	4	3	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	3	1	2	-	-	-	-	-
Male.....	1	-	-	-	1	1	1	-	-	-	-	-
Female.....	-	-	-	-	2	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Kentucky

All races, both sexes.....	122	192	269	418	555	691	823	647	394	151	31	-
Male.....	71	109	142	209	311	343	407	272	164	40	4	-
Female.....	51	83	127	209	244	348	416	375	230	111	27	-
White, both sexes.....	112	178	255	396	538	665	782	627	382	147	28	-
Male.....	67	103	131	197	300	327	387	261	161	39	4	-
Female.....	45	75	124	199	238	338	395	366	221	108	24	-
Black, both sexes.....	10	13	14	22	17	24	40	19	12	4	3	-
Male.....	4	5	11	12	11	15	19	11	3	1	-	-
Female.....	6	8	3	10	6	9	21	8	9	3	3	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	1	1	-	-	-	-
Male.....	-	1	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Louisiana

All races, both sexes.....	81	137	191	327	375	526	580	516	281	111	16	-
Male.....	44	78	106	190	195	256	271	191	98	30	5	-
Female.....	37	59	85	137	180	270	309	325	183	81	11	-
White, both sexes.....	58	96	140	264	310	451	508	435	226	89	10	-
Male.....	30	51	78	151	156	216	234	158	79	23	2	-
Female.....	28	45	62	113	154	235	274	277	147	66	8	-
Black, both sexes.....	23	41	48	62	64	75	71	80	55	22	6	-
Male.....	14	27	26	39	39	40	37	32	19	7	3	-
Female.....	9	14	22	23	25	35	34	48	36	15	3	-
American Indian 1/, both sexes.....	-	-	1	-	1	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	1	-	-	1	1	-	-	-	-
Male.....	-	-	1	-	-	-	-	1	-	-	-	-
Female.....	-	-	1	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Maine

All races, both sexes.....	20	40	77	100	147	182	230	246	148	48	12	-
Male.....	5	26	33	61	77	92	115	108	50	15	2	-
Female.....	15	14	44	39	70	90	115	138	98	33	10	-
White, both sexes.....	19	40	76	95	146	182	230	246	147	48	12	-
Male.....	5	26	33	59	77	92	115	108	50	15	2	-
Female.....	14	14	43	36	69	90	115	138	97	33	10	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	1	4	1	-	-	-	1	-	-	-
Male.....	-	-	-	2	-	-	-	-	-	-	-	-
Female.....	1	-	1	2	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	76	115	181	225	359	542	730	714	459	153	24	-
Male.....	49	54	101	114	202	265	326	307	182	48	6	-
Female.....	27	61	80	111	157	277	404	407	277	105	18	-
White, both sexes.....	52	78	151	177	289	440	627	627	414	140	20	-
Male.....	35	37	79	89	165	221	277	271	166	44	6	-
Female.....	17	41	72	88	124	219	350	356	248	96	14	-
Black, both sexes.....	21	35	30	46	65	97	92	81	42	12	4	-
Male.....	12	15	22	23	33	41	41	34	14	4	-	-
Female.....	9	20	8	23	32	56	51	47	28	8	4	-
American Indian 1/, both sexes.....	-	-	-	1	1	1	-	-	-	1	-	-
Male.....	-	-	-	1	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	3	2	-	1	4	4	11	6	3	-	-	-
Male.....	2	2	-	1	3	3	8	2	2	-	-	-
Female.....	1	-	-	-	1	1	3	4	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Massachusetts

All races, both sexes.....	58	151	200	305	469	696	1,008	1,045	759	276	60	-
Male.....	22	82	104	161	219	355	482	465	269	71	9	-
Female.....	36	69	96	144	250	341	526	580	490	205	51	-
White, both sexes.....	52	143	190	295	453	668	986	1,021	740	270	57	-
Male.....	22	80	97	153	211	339	473	453	258	70	9	-
Female.....	30	63	93	142	242	329	513	568	482	200	48	-
Black, both sexes.....	6	6	8	9	12	20	14	15	6	3	1	-
Male.....	-	2	5	7	5	11	5	8	3	-	-	-
Female.....	6	4	3	2	7	9	9	7	3	3	1	-
American Indian 1/, both sexes.....	-	-	-	1	1	-	-	1	-	-	-	-
Male.....	-	-	-	1	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	2	-	3	8	8	8	13	3	2	-
Male.....	-	-	2	-	2	5	4	4	8	1	-	-
Female.....	-	2	-	-	1	3	4	4	5	2	2	-

Michigan

All races, both sexes.....	189	272	404	576	811	1,239	1,496	1,299	827	288	58	-
Male.....	104	158	216	292	439	627	716	602	358	90	16	-
Female.....	85	114	188	284	372	612	780	697	469	198	42	-
White, both sexes.....	139	215	359	493	725	1,140	1,371	1,211	765	272	50	-
Male.....	73	132	192	249	383	576	653	564	327	87	15	-
Female.....	66	83	167	244	342	564	718	647	438	185	35	-
Black, both sexes.....	49	53	44	72	76	89	114	82	55	15	8	-
Male.....	30	25	23	37	49	47	58	34	28	3	1	-
Female.....	19	28	21	35	27	42	56	48	27	12	7	-
American Indian 1/, both sexes.....	1	3	-	7	6	5	5	4	2	1	-	-
Male.....	1	1	-	3	4	2	2	2	-	-	-	-
Female.....	-	2	-	4	2	3	3	2	2	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	4	4	5	6	2	5	-	-	-
Male.....	-	-	1	3	3	2	3	2	3	-	-	-
Female.....	-	1	-	1	1	3	3	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Minnesota

All races, both sexes.....	47	77	127	203	286	488	619	580	397	148	38	-
Male.....	24	49	69	109	166	251	320	284	165	57	11	-
Female.....	23	28	58	94	120	237	299	296	232	91	27	-
White, both sexes.....	38	65	120	192	277	480	601	573	391	148	36	-
Male.....	17	43	65	102	162	248	309	280	161	57	11	-
Female.....	21	22	55	90	115	232	292	293	230	91	25	-
Black, both sexes.....	3	8	3	3	3	4	8	3	3	-	1	-
Male.....	2	4	1	2	2	1	6	2	1	-	-	-
Female.....	1	4	2	1	1	3	2	1	2	-	1	-
American Indian 1/, both sexes.....	5	3	4	6	4	1	6	2	1	-	-	-
Male.....	4	1	3	4	2	1	3	1	1	-	-	-
Female.....	1	2	1	2	2	-	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	2	2	3	4	2	2	-	1	-
Male.....	1	1	-	1	-	1	2	1	2	-	-	-
Female.....	-	-	-	1	2	2	2	1	-	-	1	-

Mississippi

All races, both sexes.....	74	114	143	241	310	420	420	357	228	90	25	-
Male.....	41	67	81	133	188	225	223	160	81	28	2	-
Female.....	33	47	62	108	122	195	197	197	147	62	23	-
White, both sexes.....	45	84	107	186	254	346	336	287	188	65	18	-
Male.....	22	49	58	96	154	177	173	127	60	22	1	-
Female.....	23	35	49	90	100	169	163	160	128	43	17	-
Black, both sexes.....	29	30	32	55	53	74	82	69	39	24	7	-
Male.....	19	18	20	37	33	48	49	32	20	6	1	-
Female.....	10	12	12	18	20	26	33	37	19	18	6	-
American Indian 1/, both sexes.....	-	-	3	-	-	-	-	-	-	1	-	-
Male.....	-	-	3	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	3	-	2	1	1	-	-	-
Male.....	-	-	-	-	1	-	1	1	1	-	-	-
Female.....	-	-	1	-	2	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Missouri

All races, both sexes.....	118	189	314	429	612	810	972	955	618	237	56	1
Male.....	64	113	163	237	305	426	488	436	247	68	12	1
Female.....	54	76	151	192	307	384	484	519	371	169	44	-
White, both sexes.....	98	159	287	397	576	765	919	922	595	228	52	1
Male.....	51	97	147	220	284	401	466	425	242	66	10	1
Female.....	47	62	140	177	292	364	453	497	353	162	42	-
Black, both sexes.....	20	29	27	31	32	42	45	29	20	9	4	-
Male.....	13	15	16	16	20	23	19	10	5	2	2	-
Female.....	7	14	11	15	12	19	26	19	15	7	2	-
American Indian 1/, both sexes.....	-	1	-	1	2	1	3	-	-	-	-	-
Male.....	-	1	-	1	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	2	5	4	3	-	-	-
Male.....	-	-	-	-	-	1	3	1	-	-	-	-
Female.....	-	-	-	-	2	1	2	3	3	-	-	-

Montana

All races, both sexes.....	19	23	43	83	118	170	185	174	100	39	15	-
Male.....	9	13	24	44	64	80	100	81	34	11	3	-
Female.....	10	10	19	39	54	90	85	93	66	28	12	-
White, both sexes.....	17	21	36	79	112	161	176	170	97	39	15	-
Male.....	8	12	21	42	61	73	96	78	33	11	3	-
Female.....	9	9	15	37	51	88	80	92	64	28	12	-
Black, both sexes.....	-	-	1	-	-	1	1	-	-	-	-	-
Male.....	-	-	1	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	2	2	6	4	6	8	7	4	3	-	-	-
Male.....	1	1	2	2	3	6	4	3	1	-	-	-
Female.....	1	1	4	2	3	2	3	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Nebraska

All races, both sexes.....	40	31	63	112	141	262	296	298	196	84	18	1
Male.....	23	22	31	60	75	149	153	137	79	27	5	1
Female.....	17	9	32	52	66	113	143	161	117	57	13	-
White, both sexes.....	37	28	59	107	135	254	292	295	195	82	18	1
Male.....	22	19	28	58	71	145	151	136	79	26	5	1
Female.....	15	9	31	49	64	109	141	159	116	56	13	-
Black, both sexes.....	1	3	4	4	6	4	2	3	1	2	-	-
Male.....	-	3	3	2	4	3	1	1	-	1	-	-
Female.....	1	-	1	2	2	1	1	2	1	1	-	-
American Indian 1/, both sexes.....	2	-	-	1	-	2	2	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	-	-	1	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Nevada

All races, both sexes.....	46	60	108	157	218	314	338	254	146	33	6	-
Male.....	26	38	52	87	128	159	154	105	51	15	2	-
Female.....	20	22	56	70	90	155	184	149	95	18	4	-
White, both sexes.....	43	56	103	144	205	294	325	244	137	31	4	-
Male.....	25	37	50	79	116	146	148	102	47	14	2	-
Female.....	18	19	53	65	89	148	177	142	90	17	2	-
Black, both sexes.....	2	1	4	9	7	8	10	6	4	1	2	-
Male.....	-	-	2	6	7	6	5	2	2	1	-	-
Female.....	2	1	2	3	-	2	5	4	2	-	2	-
American Indian 1/, both sexes.....	1	-	-	1	1	4	-	-	1	-	-	-
Male.....	1	-	-	1	1	2	-	-	-	-	-	-
Female.....	-	-	-	-	-	2	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	3	1	3	5	8	3	4	4	1	-	-
Male.....	-	1	-	1	4	5	1	1	2	-	-	-
Female.....	-	2	1	2	1	3	2	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

New Hampshire

All races, both sexes.....	18	26	65	72	110	166	189	174	111	44	10	-
Male.....	9	13	32	42	46	86	82	77	40	11	1	-
Female.....	9	13	33	30	64	80	107	97	71	33	9	-
White, both sexes.....	18	26	65	72	110	166	189	174	110	44	10	-
Male.....	9	13	32	42	46	86	82	77	40	11	1	-
Female.....	9	13	33	30	64	80	107	97	70	33	9	-
Black, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	87	160	242	384	520	847	1,177	1,144	722	264	40	-
Male.....	51	82	129	193	270	430	555	491	267	76	4	-
Female.....	36	78	113	191	250	417	622	653	455	188	36	-
White, both sexes.....	57	122	210	330	454	783	1,093	1,077	676	238	37	-
Male.....	39	67	108	159	239	388	515	459	251	72	4	-
Female.....	18	55	102	171	215	395	578	618	425	166	33	-
Black, both sexes.....	28	36	28	49	58	59	72	54	38	24	3	-
Male.....	12	14	17	31	25	38	35	26	12	4	-	-
Female.....	16	22	11	18	33	21	37	28	26	20	3	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	1	-	-	-	-	-
Male.....	-	-	-	-	1	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	2	4	5	7	4	11	13	8	2	-	-
Male.....	-	1	4	3	5	4	4	6	4	-	-	-
Female.....	2	1	-	2	2	-	7	7	4	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

New Mexico

All races, both sexes.....	32	52	72	114	160	260	291	244	128	59	12	-
Male.....	16	31	44	63	87	131	149	113	52	24	4	-
Female.....	16	21	28	51	73	129	142	131	76	35	8	-
White, both sexes.....	29	48	64	107	152	248	279	227	122	51	9	-
Male.....	15	29	38	58	81	128	143	108	49	21	3	-
Female.....	14	19	26	49	71	120	136	119	73	30	6	-
Black, both sexes.....	1	1	1	-	2	3	1	-	1	-	-	-
Male.....	-	1	1	-	2	2	1	-	1	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	2	7	7	6	9	10	16	5	8	3	-
Male.....	1	1	5	5	4	1	4	5	2	3	1	-
Female.....	-	1	2	2	2	8	6	11	3	5	2	-
Asian or Pacific Islander 2/, both sexes.	1	1	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	1	-	-	-	-	-	1	-	-	-	-

New York

All races, both sexes.....	283	406	609	819	1,223	1,916	2,528	2,474	1,660	681	154	-
Male.....	171	222	332	408	609	932	1,202	1,075	602	184	33	-
Female.....	112	184	277	411	614	984	1,326	1,399	1,058	497	121	-
White, both sexes.....	206	330	494	699	1,048	1,699	2,278	2,225	1,534	618	129	-
Male.....	129	186	268	347	522	823	1,085	965	553	162	30	-
Female.....	77	144	226	352	526	876	1,193	1,260	981	456	99	-
Black, both sexes.....	73	70	105	103	141	182	208	197	84	46	21	-
Male.....	40	33	56	51	64	93	92	80	27	15	3	-
Female.....	33	37	49	52	77	89	116	117	57	31	18	-
American Indian 1/, both sexes.....	1	1	3	2	3	4	3	1	1	-	-	-
Male.....	1	1	3	1	2	1	1	-	-	-	-	-
Female.....	-	-	-	1	1	3	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.	3	5	7	15	31	31	39	51	41	17	4	-
Male.....	1	2	5	9	21	15	24	30	22	7	-	-
Female.....	2	3	2	6	10	16	15	21	19	10	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Ohio												
All races, both sexes.....	242	358	543	861	1,123	1,657	2,065	1,710	967	355	69	-
Male.....	116	185	290	443	585	809	1,016	733	336	110	19	-
Female.....	126	173	253	418	538	848	1,049	977	631	245	50	-
White, both sexes.....	206	307	490	776	1,030	1,531	1,940	1,609	916	335	61	-
Male.....	100	160	253	398	527	747	952	691	321	104	18	-
Female.....	106	147	237	378	503	784	988	918	595	231	43	-
Black, both sexes.....	35	48	52	82	92	122	119	97	48	20	8	-
Male.....	16	24	37	43	57	59	60	40	14	6	1	-
Female.....	19	24	15	39	35	63	59	57	34	14	7	-
American Indian 1/, both sexes.....	-	1	-	1	-	-	2	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	1	-	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	2	1	2	1	4	4	3	3	-	-	-
Male.....	-	1	-	2	1	3	3	2	1	-	-	-
Female.....	1	1	1	-	-	1	1	1	2	-	-	-
Oklahoma												
All races, both sexes.....	101	174	259	342	480	649	639	547	302	114	25	-
Male.....	47	100	129	186	249	336	327	265	115	35	4	-
Female.....	54	74	130	156	231	313	312	282	187	79	21	-
White, both sexes.....	78	153	230	307	436	603	599	515	285	104	25	-
Male.....	35	89	109	166	227	315	305	253	107	29	4	-
Female.....	43	64	121	141	209	288	294	262	178	75	21	-
Black, both sexes.....	10	6	15	13	18	14	14	9	7	5	-	-
Male.....	4	3	12	9	10	6	9	3	4	2	-	-
Female.....	6	3	3	4	8	8	5	6	3	3	-	-
American Indian 1/, both sexes.....	12	15	13	22	26	32	22	23	10	5	-	-
Male.....	7	8	8	11	12	15	11	9	4	4	-	-
Female.....	5	7	5	11	14	17	11	14	6	1	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	1	-	-	-	4	-	-	-	-	-
Male.....	1	-	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	1	-	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
X. Diseases of the respiratory system (J00-J98,U04)												
Oregon												
All races, both sexes.....	67	105	185	243	319	451	535	538	319	96	17	-
Male.....	44	64	97	124	167	244	255	245	141	31	3	-
Female.....	23	41	88	119	152	207	280	293	178	65	14	-
White, both sexes.....	64	99	174	234	310	441	517	527	313	93	17	-
Male.....	44	61	91	121	160	239	248	239	136	30	3	-
Female.....	20	38	83	113	150	202	269	288	177	63	14	-
Black, both sexes.....	1	2	2	1	4	3	1	3	2	2	-	-
Male.....	-	-	2	-	3	1	-	1	2	1	-	-
Female.....	1	2	-	1	1	2	1	2	-	1	-	-
American Indian 1/, both sexes.....	2	3	8	8	2	3	9	1	2	-	-	-
Male.....	-	2	4	3	1	2	4	-	2	-	-	-
Female.....	2	1	4	5	1	1	5	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	-	3	4	8	7	2	1	-	-
Male.....	-	1	-	-	3	2	3	5	1	-	-	-
Female.....	-	-	1	-	-	2	5	2	1	1	-	-
Pennsylvania												
All races, both sexes.....	203	335	543	722	1,126	1,755	2,349	2,263	1,364	471	90	2
Male.....	103	196	297	401	575	911	1,162	1,037	507	150	25	2
Female.....	100	139	246	321	551	844	1,187	1,226	857	321	65	-
White, both sexes.....	159	276	491	651	1,024	1,630	2,236	2,162	1,303	448	85	1
Male.....	84	159	269	361	526	846	1,094	1,000	485	146	24	1
Female.....	75	117	222	290	498	784	1,142	1,162	818	302	61	-
Black, both sexes.....	44	58	48	69	98	117	104	89	55	20	4	-
Male.....	19	37	25	38	48	62	62	30	20	2	1	-
Female.....	25	21	23	31	50	55	42	59	35	18	3	-
American Indian 1/, both sexes.....	-	1	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	3	2	4	8	9	12	6	3	1	1
Male.....	-	-	2	2	1	3	6	7	2	2	-	1
Female.....	-	-	1	-	3	5	3	5	4	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Rhode Island

All races, both sexes.....	13	26	42	33	83	120	167	175	109	54	7	-
Male.....	6	14	23	14	48	59	90	82	38	11	-	-
Female.....	7	12	19	19	35	61	77	93	71	43	7	-
White, both sexes.....	12	25	41	32	81	115	164	173	108	53	7	-
Male.....	6	13	22	14	46	54	88	82	38	11	-	-
Female.....	6	12	19	18	35	61	76	91	70	42	7	-
Black, both sexes.....	1	1	1	-	2	5	1	1	1	-	-	-
Male.....	-	1	1	-	2	5	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	1	1	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	2	-	-	1	-	-
Male.....	-	-	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-

South Carolina

All races, both sexes.....	100	151	253	334	452	570	637	497	266	86	23	1
Male.....	52	91	151	182	242	304	319	213	85	31	3	1
Female.....	48	60	102	152	210	266	318	284	181	55	20	-
White, both sexes.....	65	111	206	283	393	497	555	428	228	65	17	1
Male.....	31	65	118	155	210	267	282	176	73	22	2	1
Female.....	34	46	88	128	183	230	273	252	155	43	15	-
Black, both sexes.....	35	39	47	49	59	73	80	69	38	21	6	-
Male.....	21	26	33	27	32	37	36	37	12	9	1	-
Female.....	14	13	14	22	27	36	44	32	26	12	5	-
American Indian 1/, both sexes.....	-	1	-	2	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

South Dakota

All races, both sexes.....	11	12	34	56	78	111	131	132	99	51	13	-
Male.....	7	8	19	28	44	61	76	69	46	17	4	-
Female.....	4	4	15	28	34	50	55	63	53	34	9	-
White, both sexes.....	8	11	32	52	74	107	124	128	96	50	13	-
Male.....	6	8	18	27	41	59	73	68	45	17	4	-
Female.....	2	3	14	25	33	48	51	60	51	33	9	-
Black, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	3	1	2	4	4	4	6	4	3	1	-	-
Male.....	1	-	1	1	3	2	2	1	1	-	-	-
Female.....	2	1	1	3	1	2	4	3	2	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	158	258	390	504	664	866	1,014	912	521	172	40	1
Male.....	88	135	216	277	336	444	507	378	195	43	12	1
Female.....	70	123	174	227	328	422	507	534	326	129	28	-
White, both sexes.....	125	229	350	457	606	803	937	853	479	161	33	1
Male.....	74	116	192	251	303	410	470	356	177	39	10	1
Female.....	51	113	158	206	303	393	467	497	302	122	23	-
Black, both sexes.....	33	28	38	43	56	59	77	57	40	11	7	-
Male.....	14	18	22	23	32	33	37	21	17	4	2	-
Female.....	19	10	16	20	24	26	40	36	23	7	5	-
American Indian 1/, both sexes.....	-	1	1	1	-	-	-	-	-	-	-	-
Male.....	-	1	1	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	3	2	4	-	2	2	-	-	-
Male.....	-	-	1	2	1	1	-	1	1	-	-	-
Female.....	-	-	-	1	1	3	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Texas

All races, both sexes.....	375	553	880	1,201	1,687	2,277	2,635	2,286	1,281	511	103	2
Male.....	217	308	464	645	898	1,196	1,320	1,062	515	148	25	2
Female.....	158	245	416	556	789	1,081	1,315	1,224	766	363	78	-
White, both sexes.....	313	471	796	1,104	1,533	2,082	2,456	2,128	1,205	466	90	2
Male.....	176	261	418	595	808	1,082	1,228	982	483	134	21	2
Female.....	137	210	378	509	725	1,000	1,228	1,146	722	332	69	-
Black, both sexes.....	59	74	72	85	136	164	149	141	66	40	11	-
Male.....	39	43	41	42	83	103	81	73	26	13	3	-
Female.....	20	31	31	43	53	61	68	68	40	27	8	-
American Indian 1/, both sexes.....	2	3	-	3	2	6	8	2	-	-	-	-
Male.....	1	1	-	1	2	-	2	1	-	-	-	-
Female.....	1	2	-	2	-	6	6	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	5	12	9	16	25	22	15	10	5	2	-
Male.....	1	3	5	7	5	11	9	6	6	1	1	-
Female.....	-	2	7	2	11	14	13	9	4	4	1	-

Utah

All races, both sexes.....	27	63	66	93	146	171	232	194	117	36	4	-
Male.....	18	30	43	55	90	95	126	94	61	10	1	-
Female.....	9	33	23	38	56	76	106	100	56	26	3	-
White, both sexes.....	27	62	62	88	143	168	226	188	116	35	4	-
Male.....	18	29	41	52	88	92	120	91	60	9	1	-
Female.....	9	33	21	36	55	76	106	97	56	26	3	-
Black, both sexes.....	-	-	-	1	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	1	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	3	3	-	1	1	1	-	-	-	-
Male.....	-	1	2	2	-	1	1	1	-	-	-	-
Female.....	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	1	3	2	4	4	1	1	-	-
Male.....	-	-	-	1	2	2	4	1	1	1	-	-
Female.....	-	-	1	-	1	-	-	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Vermont

All races, both sexes.....	8	12	24	35	57	67	87	98	52	19	7	-
Male.....	4	4	15	14	29	30	50	45	19	5	2	-
Female.....	4	8	9	21	28	37	37	53	33	14	5	-
White, both sexes.....	8	12	24	35	57	66	87	98	52	19	7	-
Male.....	4	4	15	14	29	29	50	45	19	5	2	-
Female.....	4	8	9	21	28	37	37	53	33	14	5	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	110	166	291	441	574	882	1,010	971	564	188	48	-
Male.....	70	88	153	230	305	456	481	424	201	53	9	-
Female.....	40	78	138	211	269	426	529	547	363	135	39	-
White, both sexes.....	80	118	231	371	491	769	894	886	499	161	38	-
Male.....	52	62	121	191	257	385	429	378	177	47	8	-
Female.....	28	56	110	180	234	384	465	508	322	114	30	-
Black, both sexes.....	30	46	57	66	78	99	106	77	60	25	9	-
Male.....	18	25	32	37	46	62	47	39	22	5	-	-
Female.....	12	21	25	29	32	37	59	38	38	20	9	-
American Indian 1/, both sexes.....	-	-	1	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	2	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	2	4	5	12	10	8	5	2	1	-
Male.....	-	1	-	2	2	7	5	7	2	1	1	-
Female.....	-	1	2	2	3	5	5	1	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Washington

All races, both sexes.....	98	161	239	359	537	687	779	824	459	178	33	-
Male.....	52	79	135	189	283	353	368	386	181	55	10	-
Female.....	46	82	104	170	254	334	411	438	278	123	23	-
White, both sexes.....	90	142	221	338	513	653	746	793	454	174	31	-
Male.....	46	66	123	177	268	329	352	374	180	52	9	-
Female.....	44	76	98	161	245	324	394	419	274	122	22	-
Black, both sexes.....	3	10	8	10	5	9	6	3	2	-	1	-
Male.....	1	6	4	5	2	6	3	-	1	-	1	-
Female.....	2	4	4	5	3	3	3	3	1	-	-	-
American Indian 1/, both sexes.....	2	4	8	7	8	5	4	6	2	-	-	-
Male.....	2	4	6	4	6	4	2	2	-	-	-	-
Female.....	-	-	2	3	2	1	2	4	2	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	5	2	4	11	20	23	22	1	4	1	-
Male.....	3	3	2	3	7	14	11	10	-	3	-	-
Female.....	-	2	-	1	4	6	12	12	1	1	1	-

West Virginia

All races, both sexes.....	54	93	139	201	271	356	416	345	207	54	13	-
Male.....	32	58	81	106	136	193	226	175	92	16	2	-
Female.....	22	35	58	95	135	163	190	170	115	38	11	-
White, both sexes.....	52	86	137	200	266	351	411	343	203	51	13	-
Male.....	30	53	79	106	134	190	223	173	89	16	2	-
Female.....	22	33	58	94	132	161	188	170	114	35	11	-
Black, both sexes.....	2	7	2	1	4	5	5	2	4	3	-	-
Male.....	2	5	2	-	2	3	3	2	3	-	-	-
Female.....	-	2	-	1	2	2	2	-	1	3	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

X. Diseases of the respiratory system (J00-J98,U04)

Wisconsin

All races, both sexes.....	75	119	190	279	441	623	829	837	550	207	35	-
Male.....	51	68	114	145	238	326	432	359	217	66	10	-
Female.....	24	51	76	134	203	297	397	478	333	141	25	-
White, both sexes.....	64	110	171	261	422	601	807	816	539	205	34	-
Male.....	48	63	100	138	224	318	419	347	211	66	10	-
Female.....	16	47	71	123	198	283	388	469	328	139	24	-
Black, both sexes.....	8	6	11	15	14	15	14	14	6	1	1	-
Male.....	1	4	9	6	12	6	8	7	3	-	-	-
Female.....	7	2	2	9	2	9	6	7	3	1	1	-
American Indian 1/, both sexes.....	3	2	6	3	3	4	5	5	2	1	-	-
Male.....	2	-	3	1	2	1	2	3	1	-	-	-
Female.....	1	2	3	2	1	3	3	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	2	-	2	3	3	2	3	-	-	-
Male.....	-	1	2	-	-	1	3	2	2	-	-	-
Female.....	-	-	-	-	2	2	-	-	1	-	-	-

Wyoming

All races, both sexes.....	13	13	26	44	55	68	88	94	46	24	3	-
Male.....	5	7	14	25	31	37	38	35	16	11	2	-
Female.....	8	6	12	19	24	31	50	59	30	13	1	-
White, both sexes.....	12	13	26	43	53	67	85	94	46	23	3	-
Male.....	4	7	14	24	29	37	37	35	16	10	2	-
Female.....	8	6	12	19	24	30	48	59	30	13	1	-
Black, both sexes.....	-	-	-	-	1	-	1	-	-	1	-	-
Male.....	-	-	-	-	1	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	1	1	1	1	-	-	-	-	-
Male.....	1	-	-	1	1	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute upper respiratory infections (J00-J06)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute upper respiratory infections (J00-J06)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute upper respiratory infections (J00-J06)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute upper respiratory infections (J00-J06)

New Mexico

All races, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	11	3	-	-	-	1	-	2	-	1	-	-
Male.....	5	2	-	-	-	-	-	2	-	-	-	-
Female.....	6	1	-	-	-	1	-	-	-	1	-	-
White, both sexes.....	9	2	-	-	-	-	-	2	-	1	-	-
Male.....	4	1	-	-	-	-	-	2	-	-	-	-
Female.....	5	1	-	-	-	-	-	-	-	1	-	-
Black, both sexes.....	1	1	-	-	-	-	-	-	-	-	-	-
Male.....	1	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute upper respiratory infections (J00-J06)

Colorado

All races, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Connecticut

All races, both sexes.....	-	-	-	-	-	1	-	2	1	-	-	-
Male.....	-	-	-	-	-	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute upper respiratory infections (J00-J06)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute upper respiratory infections (J00-J06)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute pharyngitis and tonsillitis (J02-J03)

Alaska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute pharyngitis and tonsillitis (J02-J03)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute pharyngitis and tonsillitis (J02-J03)

Kentucky

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute pharyngitis and tonsillitis (J02-J03)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute pharyngitis and tonsillitis (J02-J03)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute pharyngitis and tonsillitis (J02-J03)

Missouri

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute pharyngitis and tonsillitis (J02-J03)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute pharyngitis and tonsillitis (J02-J03)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute pharyngitis and tonsillitis (J02-J03)

Ohio

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute pharyngitis and tonsillitis (J02-J03)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute pharyngitis and tonsillitis (J02-J03)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute pharyngitis and tonsillitis (J02-J03)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

Alaska

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

Massachusetts

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Michigan

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

Minnesota

Table with 12 columns for age groups and rows for Minnesota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 12 columns for age groups and rows for Mississippi categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

Missouri

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

Nebraska

Table with 12 columns for age groups and rows for race and sex categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 12 columns for age groups and rows for race and sex categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

North Carolina

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute laryngitis and tracheitis (J04)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute laryngitis and tracheitis (J04)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute laryngitis and tracheitis (J04)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Acute laryngitis and tracheitis (J04)

Colorado

Table with 13 columns (Age groups) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Age groups) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Acute laryngitis and tracheitis (J04)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute laryngitis and tracheitis (J04)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute laryngitis and tracheitis (J04)

Massachusetts

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by Male and Female.

Michigan

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by Male and Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute laryngitis and tracheitis (J04)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute laryngitis and tracheitis (J04)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute laryngitis and tracheitis (J04)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute laryngitis and tracheitis (J04)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

New Mexico

All races, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	6	1	-	-	-	1	-	-	-	-	-	-
Male.....	2	1	-	-	-	-	-	-	-	-	-	-
Female.....	4	-	-	-	-	1	-	-	-	-	-	-
White, both sexes.....	4	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	1	1	-	-	-	-	-	-	-	-	-	-
Male.....	1	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

Colorado

All races, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Connecticut

All races, both sexes.....	-	-	-	-	-	1	-	2	1	-	-	-
Male.....	-	-	-	-	-	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

Delaware

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

Hawaii

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Hawaii. All values are dashes (-).

Idaho

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Idaho. Values are 1 for 'All races, both sexes' and 'White, both sexes' in the 50-54 age group, and dashes (-) for all other cells.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

Oregon

Table with 13 columns (Age groups) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Age groups) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)

Vermont

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of the respiratory system (J09-J98,U04)

Arkansas

All races, both sexes.....	2,917	8	1	1	5	4	7	6	5	12	24	47
Male.....	1,487	5	-	-	4	1	4	5	1	9	11	25
Female.....	1,430	3	1	1	1	3	3	1	4	3	13	22
White, both sexes.....	2,691	5	1	-	4	2	6	5	3	11	20	40
Male.....	1,374	2	-	-	3	-	3	4	-	9	10	22
Female.....	1,317	3	1	-	1	2	3	1	3	2	10	18
Black, both sexes.....	218	3	-	1	1	2	1	1	2	1	4	6
Male.....	110	3	-	-	1	1	1	1	1	-	1	2
Female.....	108	-	-	1	-	1	-	-	1	1	3	4
American Indian 1/, both sexes.....	3	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	22,237	52	27	17	14	20	22	33	35	63	126	262
Male.....	10,551	29	13	9	12	13	12	19	19	30	74	141
Female.....	11,686	23	14	8	2	7	10	14	16	33	52	121
White, both sexes.....	19,203	40	18	11	9	17	17	24	25	41	96	194
Male.....	8,932	24	6	5	7	12	9	14	14	21	59	105
Female.....	10,271	16	12	6	2	5	8	10	11	20	37	89
Black, both sexes.....	1,269	7	5	4	3	1	3	8	5	11	19	47
Male.....	632	3	3	2	3	-	2	4	3	5	7	24
Female.....	637	4	2	2	-	1	1	4	2	6	12	23
American Indian 1/, both sexes.....	116	1	-	-	-	-	-	-	-	3	3	3
Male.....	59	-	-	-	-	-	-	-	-	1	2	2
Female.....	57	1	-	-	-	-	-	-	-	2	1	1
Asian or Pacific Islander 2/, both sexes.....	1,649	4	4	2	2	2	2	1	5	8	8	18
Male.....	928	2	4	2	2	1	1	1	2	3	6	10
Female.....	721	2	-	-	-	1	1	-	3	5	2	8

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of the respiratory system (J09-J98,U04)

Iowa

All races, both sexes.....	3,007	1	1	2	1	2	4	5	4	5	18	25
Male.....	1,482	1	1	1	1	2	3	4	3	1	8	12
Female.....	1,525	-	-	1	-	-	1	1	1	4	10	13
White, both sexes.....	2,977	1	-	2	1	2	4	4	4	5	16	23
Male.....	1,461	1	-	1	1	2	3	3	3	1	7	10
Female.....	1,516	-	-	1	-	-	1	1	1	4	9	13
Black, both sexes.....	29	-	1	-	-	-	-	1	-	-	2	2
Male.....	20	-	1	-	-	-	-	1	-	-	1	2
Female.....	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Kansas

All races, both sexes.....	2,645	2	2	3	-	2	3	2	6	3	14	26
Male.....	1,291	2	1	2	-	-	2	-	2	2	12	6
Female.....	1,354	-	1	1	-	2	1	2	4	1	2	20
White, both sexes.....	2,536	1	2	2	-	1	3	2	5	3	12	21
Male.....	1,229	1	1	1	-	-	2	-	2	2	10	2
Female.....	1,307	-	1	1	-	1	1	2	3	1	2	19
Black, both sexes.....	69	1	-	1	-	-	-	-	1	-	2	4
Male.....	41	1	-	1	-	-	-	-	-	-	2	3
Female.....	28	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes.....	32	-	-	-	-	-	-	-	-	-	-	1
Male.....	17	-	-	-	-	-	-	-	-	-	-	1
Female.....	15	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	8	-	-	-	-	1	-	-	-	-	-	-
Male.....	4	-	-	-	-	-	-	-	-	-	-	-
Female.....	4	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of the respiratory system (J09-J98,U04)

Massachusetts

All races, both sexes.....	5,134	6	3	-	-	1	7	2	3	15	27	47
Male.....	2,303	4	-	-	-	-	6	1	3	10	14	27
Female.....	2,831	2	3	-	-	1	1	1	-	5	13	20
White, both sexes.....	4,967	5	3	-	-	1	6	2	3	12	25	39
Male.....	2,220	3	-	-	-	-	5	1	3	8	12	24
Female.....	2,747	2	3	-	-	1	1	1	-	4	13	15
Black, both sexes.....	114	1	-	-	-	-	1	-	-	3	1	8
Male.....	54	1	-	-	-	-	1	-	-	2	1	3
Female.....	60	-	-	-	-	-	-	-	-	1	-	5
American Indian 1/, both sexes.....	4	-	-	-	-	-	-	-	-	-	1	-
Male.....	3	-	-	-	-	-	-	-	-	-	1	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	49	-	-	-	-	-	-	-	-	-	-	-
Male.....	26	-	-	-	-	-	-	-	-	-	-	-
Female.....	23	-	-	-	-	-	-	-	-	-	-	-

Michigan

All races, both sexes.....	7,705	13	9	5	7	9	8	14	11	31	47	92
Male.....	3,750	6	6	4	3	5	3	7	4	18	21	55
Female.....	3,955	7	3	1	4	4	5	7	7	13	26	37
White, both sexes.....	6,890	6	6	2	3	5	3	8	6	13	36	62
Male.....	3,341	3	4	2	1	4	2	5	3	7	19	40
Female.....	3,549	3	2	-	2	1	1	3	3	6	17	22
Black, both sexes.....	748	6	2	3	4	4	5	6	5	15	11	30
Male.....	375	3	1	2	2	1	1	2	1	10	2	15
Female.....	373	3	1	1	2	3	4	4	4	5	9	15
American Indian 1/, both sexes.....	36	-	-	-	-	-	-	-	-	2	-	-
Male.....	16	-	-	-	-	-	-	-	-	1	-	-
Female.....	20	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	31	1	1	-	-	-	-	-	-	1	-	-
Male.....	18	-	1	-	-	-	-	-	-	-	-	-
Female.....	13	1	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of the respiratory system (J09-J98,U04)												
New Mexico												
All races, both sexes.....	1,488	-	6	-	1	1	1	4	6	11	14	21
Male.....	750	-	4	-	1	-	-	3	2	5	8	13
Female.....	738	-	2	-	-	1	1	1	4	6	6	8
White, both sexes.....	1,386	-	4	-	1	1	-	4	5	6	11	18
Male.....	701	-	3	-	1	-	-	3	2	2	6	11
Female.....	685	-	1	-	-	1	-	1	3	4	5	7
Black, both sexes.....	12	-	1	-	-	-	-	-	-	1	-	-
Male.....	8	-	-	-	-	-	-	-	-	-	-	-
Female.....	4	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	86	-	1	-	-	-	1	-	1	4	3	3
Male.....	40	-	1	-	-	-	-	-	-	3	2	2
Female.....	46	-	-	-	-	-	1	-	1	1	1	1
Asian or Pacific Islander 2/, both sexes.....	4	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	13,235	39	19	12	9	7	24	25	30	47	88	186
Male.....	6,052	22	9	8	5	5	14	20	18	24	54	104
Female.....	7,183	17	10	4	4	2	10	5	12	23	34	82
White, both sexes.....	11,582	21	10	9	3	3	19	16	21	27	65	132
Male.....	5,265	14	4	7	2	2	11	13	15	12	39	77
Female.....	6,317	7	6	2	1	1	8	3	6	15	26	55
Black, both sexes.....	1,367	15	7	2	6	4	3	8	9	19	19	45
Male.....	630	7	3	-	3	3	2	6	3	11	13	25
Female.....	737	8	4	2	3	1	1	2	6	8	6	20
American Indian 1/, both sexes.....	22	-	-	-	-	-	-	-	-	-	1	2
Male.....	12	-	-	-	-	-	-	-	-	-	1	1
Female.....	10	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	264	3	2	1	-	-	2	1	-	1	3	7
Male.....	145	1	2	1	-	-	1	1	-	1	1	1
Female.....	119	2	-	-	-	-	1	-	-	-	2	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

United States

All races, both sexes.....	4,978	7,726	12,193	17,199	24,479	34,900	43,129	39,645	24,268	9,006	1,897	10
Male.....	2,744	4,303	6,621	9,166	12,900	17,907	21,181	17,878	9,269	2,750	439	10
Female.....	2,234	3,423	5,572	8,033	11,579	16,993	21,948	21,767	14,999	6,256	1,458	-
White, both sexes.....	3,994	6,441	10,729	15,349	22,101	31,961	39,890	36,849	22,589	8,305	1,660	9
Male.....	2,225	3,603	5,743	8,128	11,569	16,301	19,503	16,579	8,594	2,522	377	9
Female.....	1,769	2,838	4,986	7,221	10,532	15,660	20,387	20,270	13,995	5,783	1,283	-
Black, both sexes.....	872	1,122	1,242	1,528	1,916	2,255	2,402	2,013	1,166	499	178	-
Male.....	454	610	742	853	1,064	1,226	1,223	879	437	141	37	-
Female.....	418	512	500	675	852	1,029	1,179	1,134	729	358	141	-
American Indian 1/, both sexes.....	57	74	101	142	143	164	154	125	61	30	7	-
Male.....	33	37	63	70	83	72	75	51	25	11	2	-
Female.....	24	37	38	72	60	92	79	74	36	19	5	-
Asian or Pacific Islander 2/, both sexes.....	55	89	121	180	319	520	683	658	452	172	52	1
Male.....	32	53	73	115	184	308	380	369	213	76	23	1
Female.....	23	36	48	65	135	212	303	289	239	96	29	-

Alabama

All races, both sexes.....	116	160	331	420	550	697	814	646	357	118	22	-
Male.....	72	86	190	240	306	360	386	304	125	41	4	-
Female.....	44	74	141	180	244	337	428	342	232	77	18	-
White, both sexes.....	89	129	276	360	467	615	731	567	308	99	14	-
Male.....	57	69	156	207	261	316	339	265	108	33	2	-
Female.....	32	60	120	153	206	299	392	302	200	66	12	-
Black, both sexes.....	27	30	53	60	83	80	82	77	48	17	8	-
Male.....	15	16	33	33	45	42	46	39	17	8	2	-
Female.....	12	14	20	27	38	38	36	38	31	9	6	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	1	1	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	2	-	-	1	1	2	-	1	-	-
Male.....	-	1	1	-	-	1	1	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	2	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Alaska

All races, both sexes.....	13	8	25	34	42	63	45	26	17	5	-	-
Male.....	5	2	17	17	24	28	25	13	4	1	-	-
Female.....	8	6	8	17	18	35	20	13	13	4	-	-
White, both sexes.....	9	6	19	23	31	49	37	19	15	5	-	-
Male.....	3	2	13	13	16	22	19	9	4	1	-	-
Female.....	6	4	6	10	15	27	18	10	11	4	-	-
Black, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	3	1	6	9	11	13	8	4	2	-	-	-
Male.....	1	-	4	4	8	5	6	1	-	-	-	-
Female.....	2	1	2	5	3	8	2	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	2	-	1	-	3	-	-	-	-
Male.....	1	-	-	-	-	1	-	3	-	-	-	-
Female.....	-	-	-	2	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	95	142	250	336	528	696	816	717	382	156	23	-
Male.....	48	76	123	189	280	364	402	360	160	67	7	-
Female.....	47	66	127	147	248	332	414	357	222	89	16	-
White, both sexes.....	84	136	236	320	504	660	790	690	365	146	19	-
Male.....	43	73	117	180	269	347	395	345	155	62	6	-
Female.....	41	63	119	140	235	313	395	345	210	84	13	-
Black, both sexes.....	6	3	4	3	6	12	9	8	1	2	1	-
Male.....	2	1	1	2	1	5	1	5	-	1	-	-
Female.....	4	2	3	1	5	7	8	3	1	1	1	-
American Indian 1/, both sexes.....	4	3	8	12	12	21	14	15	14	8	3	-
Male.....	2	2	4	6	7	9	5	7	5	4	1	-
Female.....	2	1	4	6	5	12	9	8	9	4	2	-
Asian or Pacific Islander 2/, both sexes.....	1	-	2	1	6	3	3	4	2	-	-	-
Male.....	1	-	1	1	3	3	1	3	-	-	-	-
Female.....	-	-	1	-	3	-	2	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Arkansas

All races, both sexes.....	83	108	188	264	317	478	520	450	257	110	22	-
Male.....	49	58	116	146	172	275	259	204	101	38	4	-
Female.....	34	50	72	118	145	203	261	246	156	72	18	-
White, both sexes.....	56	95	174	243	293	455	493	425	239	101	20	-
Male.....	34	53	106	135	159	263	245	192	95	35	4	-
Female.....	22	42	68	108	134	192	248	233	144	66	16	-
Black, both sexes.....	27	13	14	19	23	20	27	25	17	9	2	-
Male.....	15	5	10	11	12	11	14	12	6	3	-	-
Female.....	12	8	4	8	11	9	13	13	11	6	2	-
American Indian 1/, both sexes.....	-	-	-	2	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	3	-	-	1	-	-	-
Male.....	-	-	-	-	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	2	-	-	1	-	-	-

California

All races, both sexes.....	445	669	1,024	1,481	2,238	3,304	4,323	4,193	2,619	1,016	253	1
Male.....	227	401	517	789	1,139	1,682	2,054	1,925	1,045	333	67	1
Female.....	218	268	507	692	1,099	1,622	2,269	2,268	1,574	683	186	-
White, both sexes.....	355	539	870	1,264	1,924	2,875	3,810	3,691	2,301	877	204	1
Male.....	182	327	429	660	964	1,436	1,777	1,647	898	285	50	1
Female.....	173	212	441	604	960	1,439	2,033	2,044	1,403	592	154	-
Black, both sexes.....	58	93	101	120	160	166	174	139	80	44	21	-
Male.....	30	52	58	62	86	89	87	63	36	9	4	-
Female.....	28	41	43	58	74	77	87	76	44	35	17	-
American Indian 1/, both sexes.....	7	10	5	11	19	14	13	19	6	2	-	-
Male.....	2	6	2	7	10	5	8	10	4	-	-	-
Female.....	5	4	3	4	9	9	5	9	2	2	-	-
Asian or Pacific Islander 2/, both sexes.....	25	27	48	86	135	249	326	344	232	93	28	-
Male.....	13	16	28	60	79	152	182	205	107	39	13	-
Female.....	12	11	20	26	56	97	144	139	125	54	15	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Colorado

All races, both sexes.....	88	125	187	239	337	500	583	588	349	132	33	-
Male.....	53	78	104	132	181	269	305	277	154	41	8	-
Female.....	35	47	83	107	156	231	278	311	195	91	25	-
White, both sexes.....	80	120	177	229	324	491	569	577	337	130	33	-
Male.....	48	75	101	125	173	266	296	275	149	41	8	-
Female.....	32	45	76	104	151	225	273	302	188	89	25	-
Black, both sexes.....	7	2	9	6	9	3	6	4	10	1	-	-
Male.....	4	2	3	5	5	1	5	1	4	-	-	-
Female.....	3	-	6	1	4	2	1	3	6	1	-	-
American Indian 1/, both sexes.....	-	3	1	3	1	4	2	5	-	-	-	-
Male.....	-	1	-	1	1	1	2	1	-	-	-	-
Female.....	-	2	1	2	-	3	-	4	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	1	3	2	6	2	2	1	-	-
Male.....	1	-	-	1	2	1	2	-	1	-	-	-
Female.....	-	-	-	-	1	1	4	2	1	1	-	-

Connecticut

All races, both sexes.....	34	71	127	167	221	376	582	618	451	176	36	-
Male.....	24	36	65	82	110	205	280	298	162	64	5	-
Female.....	10	35	62	85	111	171	302	320	289	112	31	-
White, both sexes.....	28	59	114	156	207	358	555	602	444	170	34	-
Male.....	20	28	60	75	104	194	270	288	161	60	5	-
Female.....	8	31	54	81	103	164	285	314	283	110	29	-
Black, both sexes.....	6	10	13	11	13	15	25	16	6	4	2	-
Male.....	4	6	5	7	5	9	10	10	1	2	-	-
Female.....	2	4	8	4	8	6	15	6	5	2	2	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	1	3	2	-	1	2	-	-
Male.....	-	1	-	-	1	2	-	-	-	2	-	-
Female.....	-	-	-	-	-	1	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Delaware

All races, both sexes.....	15	18	36	41	78	119	122	110	53	15	3	-
Male.....	5	11	24	23	38	51	61	41	16	5	1	-
Female.....	10	7	12	18	40	68	61	69	37	10	2	-
White, both sexes.....	11	15	32	31	68	107	110	103	50	15	3	-
Male.....	3	10	22	18	32	45	54	37	14	5	1	-
Female.....	8	5	10	13	36	62	56	66	36	10	2	-
Black, both sexes.....	4	3	4	9	10	12	12	7	3	-	-	-
Male.....	2	1	2	4	6	6	7	4	2	-	-	-
Female.....	2	2	2	5	4	6	5	3	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Dist. of Columbia

All races, both sexes.....	15	14	16	18	21	38	42	37	37	16	2	-
Male.....	9	9	11	12	16	20	21	14	15	2	1	-
Female.....	6	5	5	6	5	18	21	23	22	14	1	-
White, both sexes.....	2	2	4	4	5	9	17	11	21	6	-	-
Male.....	2	1	4	4	3	4	8	7	7	1	-	-
Female.....	-	1	-	-	2	5	9	4	14	5	-	-
Black, both sexes.....	13	12	12	14	16	28	25	26	14	10	2	-
Male.....	7	8	7	8	13	15	13	7	7	1	1	-
Female.....	6	4	5	6	3	13	12	19	7	9	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	2	-	-	-
Male.....	-	-	-	-	-	1	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Florida

All races, both sexes.....	287	457	783	1,085	1,608	2,270	2,783	2,495	1,471	478	81	-
Male.....	164	267	440	583	857	1,182	1,378	1,113	555	171	22	-
Female.....	123	190	343	502	751	1,088	1,405	1,382	916	307	59	-
White, both sexes.....	239	393	697	987	1,498	2,152	2,660	2,396	1,401	456	71	-
Male.....	147	231	384	529	794	1,113	1,308	1,072	533	159	20	-
Female.....	92	162	313	458	704	1,039	1,352	1,324	868	297	51	-
Black, both sexes.....	45	60	79	93	102	109	113	92	64	20	10	-
Male.....	15	33	51	54	61	65	64	38	21	11	2	-
Female.....	30	27	28	39	41	44	49	54	43	9	8	-
American Indian 1/, both sexes.....	2	-	3	1	2	1	4	-	-	-	-	-
Male.....	2	-	2	-	-	-	2	-	-	-	-	-
Female.....	-	-	1	1	2	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	4	4	4	6	8	6	7	6	2	-	-
Male.....	-	3	3	-	2	4	4	3	1	1	-	-
Female.....	1	1	1	4	4	4	2	4	5	1	-	-

Georgia

All races, both sexes.....	168	284	409	553	787	911	1,164	946	554	238	34	-
Male.....	96	157	231	301	412	443	537	404	184	68	9	-
Female.....	72	127	178	252	375	468	627	542	370	170	25	-
White, both sexes.....	119	207	325	456	656	796	1,019	816	480	201	23	-
Male.....	67	117	181	242	336	385	463	351	162	56	4	-
Female.....	52	90	144	214	320	411	556	465	318	145	19	-
Black, both sexes.....	48	77	84	92	123	108	134	126	72	36	11	-
Male.....	28	40	50	57	70	53	66	52	20	11	5	-
Female.....	20	37	34	35	53	55	68	74	52	25	6	-
American Indian 1/, both sexes.....	-	-	-	1	2	2	1	-	-	-	-	-
Male.....	-	-	-	-	1	2	-	-	-	-	-	-
Female.....	-	-	-	1	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	4	6	5	10	4	2	1	-	-
Male.....	1	-	-	2	5	3	8	1	2	1	-	-
Female.....	-	-	-	2	1	2	2	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Hawaii

All races, both sexes.....	12	25	32	33	63	106	162	155	105	48	12	1
Male.....	8	16	19	17	36	61	94	79	48	24	8	1
Female.....	4	9	13	16	27	45	68	76	57	24	4	-
White, both sexes.....	6	12	14	15	24	33	43	48	30	18	2	1
Male.....	4	8	7	6	13	19	28	23	12	8	1	1
Female.....	2	4	7	9	11	14	15	25	18	10	1	-
Black, both sexes.....	-	-	-	-	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	6	13	18	18	39	72	119	106	75	30	10	-
Male.....	4	8	12	11	23	42	66	56	36	16	7	-
Female.....	2	5	6	7	16	30	53	50	39	14	3	-

Idaho

All races, both sexes.....	24	39	55	81	123	165	217	185	126	45	8	-
Male.....	18	16	29	53	66	81	101	95	45	18	2	-
Female.....	6	23	26	28	57	84	116	90	81	27	6	-
White, both sexes.....	23	37	53	81	122	162	214	183	126	45	8	-
Male.....	17	16	29	53	66	79	99	94	45	18	2	-
Female.....	6	21	24	28	56	83	115	89	81	27	6	-
Black, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	2	1	-	1	3	1	-	-	-	-	-
Male.....	1	-	-	-	-	2	1	-	-	-	-	-
Female.....	-	2	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	-	2	2	-	-	-	-
Male.....	-	-	-	-	-	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Illinois

All races, both sexes.....	219	296	460	638	900	1,434	1,778	1,666	1,099	409	92	-
Male.....	119	162	266	345	457	735	854	720	389	117	17	-
Female.....	100	134	194	293	443	699	924	946	710	292	75	-
White, both sexes.....	163	218	377	536	781	1,273	1,612	1,538	1,012	390	83	-
Male.....	90	121	215	286	397	645	770	670	355	112	13	-
Female.....	73	97	162	250	384	628	842	868	657	278	70	-
Black, both sexes.....	54	72	79	97	113	143	144	113	70	16	7	-
Male.....	28	37	47	56	57	77	70	43	29	3	3	-
Female.....	26	35	32	41	56	66	74	70	41	13	4	-
American Indian 1/, both sexes.....	-	-	1	2	1	-	2	-	1	-	-	-
Male.....	-	-	1	1	-	-	1	-	-	-	-	-
Female.....	-	-	-	1	1	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	6	3	3	5	18	20	15	16	3	2	-
Male.....	1	4	3	2	3	13	13	7	5	2	1	-
Female.....	1	2	-	1	2	5	7	8	11	1	1	-

Indiana

All races, both sexes.....	127	218	312	459	620	818	1,028	927	500	182	55	-
Male.....	70	105	172	239	326	416	517	433	187	40	10	-
Female.....	57	113	140	220	294	402	511	494	313	142	45	-
White, both sexes.....	117	195	304	444	593	779	988	893	478	177	54	-
Male.....	65	93	168	231	317	402	491	418	180	38	10	-
Female.....	52	102	136	213	276	377	497	475	298	139	44	-
Black, both sexes.....	10	23	8	14	26	38	40	33	22	5	1	-
Male.....	5	12	4	8	9	14	26	15	7	2	-	-
Female.....	5	11	4	6	17	24	14	18	15	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Iowa

All races, both sexes.....	42	80	137	173	307	432	578	587	414	156	33	-
Male.....	27	42	74	83	175	231	321	278	166	41	7	-
Female.....	15	38	63	90	132	201	257	309	248	115	26	-
White, both sexes.....	42	77	134	172	301	425	575	587	413	156	33	-
Male.....	27	41	71	82	171	227	318	278	166	41	7	-
Female.....	15	36	63	90	130	198	257	309	247	115	26	-
Black, both sexes.....	-	3	3	1	5	7	3	-	1	-	-	-
Male.....	-	1	3	1	3	4	3	-	-	-	-	-
Female.....	-	2	-	-	2	3	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Kansas

All races, both sexes.....	45	82	119	172	263	401	551	463	330	134	22	-
Male.....	24	37	70	92	149	209	285	236	125	31	4	-
Female.....	21	45	49	80	114	192	266	227	205	103	18	-
White, both sexes.....	41	73	111	165	254	381	531	449	325	132	22	-
Male.....	22	34	64	86	144	197	275	229	122	31	4	-
Female.....	19	39	47	79	110	184	256	220	203	101	18	-
Black, both sexes.....	3	5	6	4	3	12	12	9	4	2	-	-
Male.....	1	2	5	4	2	8	6	4	2	-	-	-
Female.....	2	3	1	-	1	4	6	5	2	2	-	-
American Indian 1/, both sexes.....	-	4	2	3	3	7	6	5	1	-	-	-
Male.....	-	1	1	2	2	3	3	3	1	-	-	-
Female.....	-	3	1	1	1	4	3	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	3	1	2	-	-	-	-	-
Male.....	1	-	-	-	1	1	1	-	-	-	-	-
Female.....	-	-	-	-	2	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Kentucky

All races, both sexes.....	122	192	269	418	555	691	822	647	394	151	31	-
Male.....	71	109	142	209	311	343	407	272	164	40	4	-
Female.....	51	83	127	209	244	348	415	375	230	111	27	-
White, both sexes.....	112	178	255	396	538	665	781	627	382	147	28	-
Male.....	67	103	131	197	300	327	387	261	161	39	4	-
Female.....	45	75	124	199	238	338	394	366	221	108	24	-
Black, both sexes.....	10	13	14	22	17	24	40	19	12	4	3	-
Male.....	4	5	11	12	11	15	19	11	3	1	-	-
Female.....	6	8	3	10	6	9	21	8	9	3	3	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	1	1	-	-	-	-
Male.....	-	1	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Louisiana

All races, both sexes.....	81	136	191	327	375	526	580	516	281	111	16	-
Male.....	44	78	106	190	195	256	271	191	98	30	5	-
Female.....	37	58	85	137	180	270	309	325	183	81	11	-
White, both sexes.....	58	95	140	264	310	451	508	435	226	89	10	-
Male.....	30	51	78	151	156	216	234	158	79	23	2	-
Female.....	28	44	62	113	154	235	274	277	147	66	8	-
Black, both sexes.....	23	41	48	62	64	75	71	80	55	22	6	-
Male.....	14	27	26	39	39	40	37	32	19	7	3	-
Female.....	9	14	22	23	25	35	34	48	36	15	3	-
American Indian 1/, both sexes.....	-	-	1	-	1	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	1	-	-	1	1	-	-	-	-
Male.....	-	-	1	-	-	-	-	1	-	-	-	-
Female.....	-	-	1	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Maine

All races, both sexes.....	20	40	77	100	147	182	230	246	147	48	12	-
Male.....	5	26	33	61	77	92	115	108	50	15	2	-
Female.....	15	14	44	39	70	90	115	138	97	33	10	-
White, both sexes.....	19	40	76	95	146	182	230	246	146	48	12	-
Male.....	5	26	33	59	77	92	115	108	50	15	2	-
Female.....	14	14	43	36	69	90	115	138	96	33	10	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	1	4	1	-	-	-	1	-	-	-
Male.....	-	-	-	2	-	-	-	-	-	-	-	-
Female.....	1	-	1	2	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	76	115	180	225	359	542	729	714	458	153	24	-
Male.....	49	54	100	114	202	265	326	307	181	48	6	-
Female.....	27	61	80	111	157	277	403	407	277	105	18	-
White, both sexes.....	52	78	150	177	289	440	626	627	414	140	20	-
Male.....	35	37	78	89	165	221	277	271	166	44	6	-
Female.....	17	41	72	88	124	219	349	356	248	96	14	-
Black, both sexes.....	21	35	30	46	65	97	92	81	41	12	4	-
Male.....	12	15	22	23	33	41	41	34	13	4	-	-
Female.....	9	20	8	23	32	56	51	47	28	8	4	-
American Indian 1/, both sexes.....	-	-	-	1	1	1	-	-	-	1	-	-
Male.....	-	-	-	1	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	3	2	-	1	4	4	11	6	3	-	-	-
Male.....	2	2	-	1	3	3	8	2	2	-	-	-
Female.....	1	-	-	-	1	1	3	4	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Massachusetts

All races, both sexes.....	58	149	200	305	469	696	1,008	1,044	759	276	59	-
Male.....	22	81	104	161	219	355	482	465	269	71	9	-
Female.....	36	68	96	144	250	341	526	579	490	205	50	-
White, both sexes.....	52	141	190	295	453	668	986	1,020	740	270	56	-
Male.....	22	79	97	153	211	339	473	453	258	70	9	-
Female.....	30	62	93	142	242	329	513	567	482	200	47	-
Black, both sexes.....	6	6	8	9	12	20	14	15	6	3	1	-
Male.....	-	2	5	7	5	11	5	8	3	-	-	-
Female.....	6	4	3	2	7	9	9	7	3	3	1	-
American Indian 1/, both sexes.....	-	-	-	1	1	-	-	1	-	-	-	-
Male.....	-	-	-	1	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	2	-	3	8	8	8	13	3	2	-
Male.....	-	-	2	-	2	5	4	4	8	1	-	-
Female.....	-	2	-	-	1	3	4	4	5	2	2	-

Michigan

All races, both sexes.....	189	272	404	576	811	1,239	1,496	1,299	827	288	58	-
Male.....	104	158	216	292	439	627	716	602	358	90	16	-
Female.....	85	114	188	284	372	612	780	697	469	198	42	-
White, both sexes.....	139	215	359	493	725	1,140	1,371	1,211	765	272	50	-
Male.....	73	132	192	249	383	576	653	564	327	87	15	-
Female.....	66	83	167	244	342	564	718	647	438	185	35	-
Black, both sexes.....	49	53	44	72	76	89	114	82	55	15	8	-
Male.....	30	25	23	37	49	47	58	34	28	3	1	-
Female.....	19	28	21	35	27	42	56	48	27	12	7	-
American Indian 1/, both sexes.....	1	3	-	7	6	5	5	4	2	1	-	-
Male.....	1	1	-	3	4	2	2	2	-	-	-	-
Female.....	-	2	-	4	2	3	3	2	2	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	4	4	5	6	2	5	-	-	-
Male.....	-	-	1	3	3	2	3	2	3	-	-	-
Female.....	-	1	-	1	1	3	3	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Minnesota

All races, both sexes.....	47	77	127	203	286	488	619	579	397	148	38	-
Male.....	24	49	69	109	166	251	320	284	165	57	11	-
Female.....	23	28	58	94	120	237	299	295	232	91	27	-
White, both sexes.....	38	65	120	192	277	480	601	572	391	148	36	-
Male.....	17	43	65	102	162	248	309	280	161	57	11	-
Female.....	21	22	55	90	115	232	292	292	230	91	25	-
Black, both sexes.....	3	8	3	3	3	4	8	3	3	-	1	-
Male.....	2	4	1	2	2	1	6	2	1	-	-	-
Female.....	1	4	2	1	1	3	2	1	2	-	1	-
American Indian 1/, both sexes.....	5	3	4	6	4	1	6	2	1	-	-	-
Male.....	4	1	3	4	2	1	3	1	1	-	-	-
Female.....	1	2	1	2	2	-	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	2	2	3	4	2	2	-	1	-
Male.....	1	1	-	1	-	1	2	1	2	-	-	-
Female.....	-	-	-	1	2	2	2	1	-	-	1	-

Mississippi

All races, both sexes.....	74	114	143	241	310	420	420	357	228	90	25	-
Male.....	41	67	81	133	188	225	223	160	81	28	2	-
Female.....	33	47	62	108	122	195	197	197	147	62	23	-
White, both sexes.....	45	84	107	186	254	346	336	287	188	65	18	-
Male.....	22	49	58	96	154	177	173	127	60	22	1	-
Female.....	23	35	49	90	100	169	163	160	128	43	17	-
Black, both sexes.....	29	30	32	55	53	74	82	69	39	24	7	-
Male.....	19	18	20	37	33	48	49	32	20	6	1	-
Female.....	10	12	12	18	20	26	33	37	19	18	6	-
American Indian 1/, both sexes.....	-	-	3	-	-	-	-	-	-	1	-	-
Male.....	-	-	3	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	3	-	2	1	1	-	-	-
Male.....	-	-	-	-	1	-	1	1	1	-	-	-
Female.....	-	-	1	-	2	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Missouri

All races, both sexes.....	118	189	314	428	612	810	972	954	618	237	56	1
Male.....	64	113	163	237	305	426	488	436	247	68	12	1
Female.....	54	76	151	191	307	384	484	518	371	169	44	-
White, both sexes.....	98	159	287	396	576	765	919	921	595	228	52	1
Male.....	51	97	147	220	284	401	466	425	242	66	10	1
Female.....	47	62	140	176	292	364	453	496	353	162	42	-
Black, both sexes.....	20	29	27	31	32	42	45	29	20	9	4	-
Male.....	13	15	16	16	20	23	19	10	5	2	2	-
Female.....	7	14	11	15	12	19	26	19	15	7	2	-
American Indian 1/, both sexes.....	-	1	-	1	2	1	3	-	-	-	-	-
Male.....	-	1	-	1	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	2	5	4	3	-	-	-
Male.....	-	-	-	-	-	1	3	1	-	-	-	-
Female.....	-	-	-	-	2	1	2	3	3	-	-	-

Montana

All races, both sexes.....	19	23	43	83	118	170	185	174	100	39	15	-
Male.....	9	13	24	44	64	80	100	81	34	11	3	-
Female.....	10	10	19	39	54	90	85	93	66	28	12	-
White, both sexes.....	17	21	36	79	112	161	176	170	97	39	15	-
Male.....	8	12	21	42	61	73	96	78	33	11	3	-
Female.....	9	9	15	37	51	88	80	92	64	28	12	-
Black, both sexes.....	-	-	1	-	-	1	1	-	-	-	-	-
Male.....	-	-	1	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	2	2	6	4	6	8	7	4	3	-	-	-
Male.....	1	1	2	2	3	6	4	3	1	-	-	-
Female.....	1	1	4	2	3	2	3	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Nebraska

All races, both sexes.....	40	31	63	112	141	262	295	298	196	84	18	1
Male.....	23	22	31	60	75	149	152	137	79	27	5	1
Female.....	17	9	32	52	66	113	143	161	117	57	13	-
White, both sexes.....	37	28	59	107	135	254	291	295	195	82	18	1
Male.....	22	19	28	58	71	145	150	136	79	26	5	1
Female.....	15	9	31	49	64	109	141	159	116	56	13	-
Black, both sexes.....	1	3	4	4	6	4	2	3	1	2	-	-
Male.....	-	3	3	2	4	3	1	1	-	1	-	-
Female.....	1	-	1	2	2	1	1	2	1	1	-	-
American Indian 1/, both sexes.....	2	-	-	1	-	2	2	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	-	-	1	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Nevada

All races, both sexes.....	46	60	108	157	218	314	338	254	146	33	6	-
Male.....	26	38	52	87	128	159	154	105	51	15	2	-
Female.....	20	22	56	70	90	155	184	149	95	18	4	-
White, both sexes.....	43	56	103	144	205	294	325	244	137	31	4	-
Male.....	25	37	50	79	116	146	148	102	47	14	2	-
Female.....	18	19	53	65	89	148	177	142	90	17	2	-
Black, both sexes.....	2	1	4	9	7	8	10	6	4	1	2	-
Male.....	-	-	2	6	7	6	5	2	2	1	-	-
Female.....	2	1	2	3	-	2	5	4	2	-	2	-
American Indian 1/, both sexes.....	1	-	-	1	1	4	-	-	1	-	-	-
Male.....	1	-	-	1	1	2	-	-	-	-	-	-
Female.....	-	-	-	-	-	2	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	3	1	3	5	8	3	4	4	1	-	-
Male.....	-	1	-	1	4	5	1	1	2	-	-	-
Female.....	-	2	1	2	1	3	2	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

New Hampshire

All races, both sexes.....	18	26	65	72	110	166	189	174	110	44	9	-
Male.....	9	13	32	42	46	86	82	77	39	11	1	-
Female.....	9	13	33	30	64	80	107	97	71	33	8	-
White, both sexes.....	18	26	65	72	110	166	189	174	109	44	9	-
Male.....	9	13	32	42	46	86	82	77	39	11	1	-
Female.....	9	13	33	30	64	80	107	97	70	33	8	-
Black, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	87	160	242	384	520	847	1,177	1,143	721	264	40	-
Male.....	51	82	129	193	270	430	555	491	267	76	4	-
Female.....	36	78	113	191	250	417	622	652	454	188	36	-
White, both sexes.....	57	122	210	330	454	783	1,093	1,076	675	238	37	-
Male.....	39	67	108	159	239	388	515	459	251	72	4	-
Female.....	18	55	102	171	215	395	578	617	424	166	33	-
Black, both sexes.....	28	36	28	49	58	59	72	54	38	24	3	-
Male.....	12	14	17	31	25	38	35	26	12	4	-	-
Female.....	16	22	11	18	33	21	37	28	26	20	3	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	1	-	-	-	-	-
Male.....	-	-	-	-	1	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	2	4	5	7	4	11	13	8	2	-	-
Male.....	-	1	4	3	5	4	4	6	4	-	-	-
Female.....	2	1	-	2	2	-	7	7	4	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

New Mexico

All races, both sexes.....	32	52	72	114	160	260	290	244	128	59	12	-
Male.....	16	31	44	63	87	131	149	113	52	24	4	-
Female.....	16	21	28	51	73	129	141	131	76	35	8	-
White, both sexes.....	29	48	64	107	152	248	279	227	122	51	9	-
Male.....	15	29	38	58	81	128	143	108	49	21	3	-
Female.....	14	19	26	49	71	120	136	119	73	30	6	-
Black, both sexes.....	1	1	1	-	2	3	1	-	1	-	-	-
Male.....	-	1	1	-	2	2	1	-	1	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	2	7	7	6	9	9	16	5	8	3	-
Male.....	1	1	5	5	4	1	4	5	2	3	1	-
Female.....	-	1	2	2	2	8	5	11	3	5	2	-
Asian or Pacific Islander 2/, both sexes.	1	1	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	1	-	-	-	-	-	1	-	-	-	-

New York

All races, both sexes.....	283	406	609	819	1,223	1,916	2,528	2,474	1,658	680	153	-
Male.....	171	222	332	408	609	932	1,202	1,075	602	184	32	-
Female.....	112	184	277	411	614	984	1,326	1,399	1,056	496	121	-
White, both sexes.....	206	330	494	699	1,048	1,699	2,278	2,225	1,532	617	128	-
Male.....	129	186	268	347	522	823	1,085	965	553	162	29	-
Female.....	77	144	226	352	526	876	1,193	1,260	979	455	99	-
Black, both sexes.....	73	70	105	103	141	182	208	197	84	46	21	-
Male.....	40	33	56	51	64	93	92	80	27	15	3	-
Female.....	33	37	49	52	77	89	116	117	57	31	18	-
American Indian 1/, both sexes.....	1	1	3	2	3	4	3	1	1	-	-	-
Male.....	1	1	3	1	2	1	1	-	-	-	-	-
Female.....	-	-	-	1	1	3	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.	3	5	7	15	31	31	39	51	41	17	4	-
Male.....	1	2	5	9	21	15	24	30	22	7	-	-
Female.....	2	3	2	6	10	16	15	21	19	10	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Ohio

All races, both sexes.....	242	358	542	860	1,122	1,656	2,065	1,710	966	355	68	-
Male.....	116	185	290	443	585	809	1,016	733	336	110	18	-
Female.....	126	173	252	417	537	847	1,049	977	630	245	50	-
White, both sexes.....	206	307	490	776	1,029	1,530	1,940	1,609	915	335	60	-
Male.....	100	160	253	398	527	747	952	691	321	104	17	-
Female.....	106	147	237	378	502	783	988	918	594	231	43	-
Black, both sexes.....	35	48	51	81	92	122	119	97	48	20	8	-
Male.....	16	24	37	43	57	59	60	40	14	6	1	-
Female.....	19	24	14	38	35	63	59	57	34	14	7	-
American Indian 1/, both sexes.....	-	1	-	1	-	-	2	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	1	-	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	2	1	2	1	4	4	3	3	-	-	-
Male.....	-	1	-	2	1	3	3	2	1	-	-	-
Female.....	1	1	1	-	-	1	1	1	2	-	-	-

Oklahoma

All races, both sexes.....	101	174	259	342	480	649	639	547	302	114	25	-
Male.....	47	100	129	186	249	336	327	265	115	35	4	-
Female.....	54	74	130	156	231	313	312	282	187	79	21	-
White, both sexes.....	78	153	230	307	436	603	599	515	285	104	25	-
Male.....	35	89	109	166	227	315	305	253	107	29	4	-
Female.....	43	64	121	141	209	288	294	262	178	75	21	-
Black, both sexes.....	10	6	15	13	18	14	14	9	7	5	-	-
Male.....	4	3	12	9	10	6	9	3	4	2	-	-
Female.....	6	3	3	4	8	8	5	6	3	3	-	-
American Indian 1/, both sexes.....	12	15	13	22	26	32	22	23	10	5	-	-
Male.....	7	8	8	11	12	15	11	9	4	4	-	-
Female.....	5	7	5	11	14	17	11	14	6	1	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	1	-	-	-	4	-	-	-	-	-
Male.....	1	-	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	1	-	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Oregon

All races, both sexes.....	67	105	185	243	318	451	535	537	319	96	17	-
Male.....	44	64	97	124	166	244	255	245	141	31	3	-
Female.....	23	41	88	119	152	207	280	292	178	65	14	-
White, both sexes.....	64	99	174	234	309	441	517	526	313	93	17	-
Male.....	44	61	91	121	159	239	248	239	136	30	3	-
Female.....	20	38	83	113	150	202	269	287	177	63	14	-
Black, both sexes.....	1	2	2	1	4	3	1	3	2	2	-	-
Male.....	-	-	2	-	3	1	-	1	2	1	-	-
Female.....	1	2	-	1	1	2	1	2	-	1	-	-
American Indian 1/, both sexes.....	2	3	8	8	2	3	9	1	2	-	-	-
Male.....	-	2	4	3	1	2	4	-	2	-	-	-
Female.....	2	1	4	5	1	1	5	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	-	3	4	8	7	2	1	-	-
Male.....	-	1	-	-	3	2	3	5	1	-	-	-
Female.....	-	-	1	-	-	2	5	2	1	1	-	-

Pennsylvania

All races, both sexes.....	203	335	542	722	1,126	1,755	2,349	2,263	1,364	471	90	2
Male.....	103	196	296	401	575	911	1,162	1,037	507	150	25	2
Female.....	100	139	246	321	551	844	1,187	1,226	857	321	65	-
White, both sexes.....	159	276	490	651	1,024	1,630	2,236	2,162	1,303	448	85	1
Male.....	84	159	268	361	526	846	1,094	1,000	485	146	24	1
Female.....	75	117	222	290	498	784	1,142	1,162	818	302	61	-
Black, both sexes.....	44	58	48	69	98	117	104	89	55	20	4	-
Male.....	19	37	25	38	48	62	62	30	20	2	1	-
Female.....	25	21	23	31	50	55	42	59	35	18	3	-
American Indian 1/, both sexes.....	-	1	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	3	2	4	8	9	12	6	3	1	1
Male.....	-	-	2	2	1	3	6	7	2	2	-	1
Female.....	-	-	1	-	3	5	3	5	4	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Rhode Island

All races, both sexes.....	13	26	42	33	83	120	167	175	109	54	7	-
Male.....	6	14	23	14	48	59	90	82	38	11	-	-
Female.....	7	12	19	19	35	61	77	93	71	43	7	-
White, both sexes.....	12	25	41	32	81	115	164	173	108	53	7	-
Male.....	6	13	22	14	46	54	88	82	38	11	-	-
Female.....	6	12	19	18	35	61	76	91	70	42	7	-
Black, both sexes.....	1	1	1	-	2	5	1	1	1	-	-	-
Male.....	-	1	1	-	2	5	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	1	1	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	2	-	-	1	-	-
Male.....	-	-	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-

South Carolina

All races, both sexes.....	100	151	253	334	451	570	637	497	266	86	23	1
Male.....	52	91	151	182	242	304	319	213	85	31	3	1
Female.....	48	60	102	152	209	266	318	284	181	55	20	-
White, both sexes.....	65	111	206	283	392	497	555	428	228	65	17	1
Male.....	31	65	118	155	210	267	282	176	73	22	2	1
Female.....	34	46	88	128	182	230	273	252	155	43	15	-
Black, both sexes.....	35	39	47	49	59	73	80	69	38	21	6	-
Male.....	21	26	33	27	32	37	36	37	12	9	1	-
Female.....	14	13	14	22	27	36	44	32	26	12	5	-
American Indian 1/, both sexes.....	-	1	-	2	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

South Dakota

All races, both sexes.....	11	12	34	56	78	111	131	132	99	51	13	-
Male.....	7	8	19	28	44	61	76	69	46	17	4	-
Female.....	4	4	15	28	34	50	55	63	53	34	9	-
White, both sexes.....	8	11	32	52	74	107	124	128	96	50	13	-
Male.....	6	8	18	27	41	59	73	68	45	17	4	-
Female.....	2	3	14	25	33	48	51	60	51	33	9	-
Black, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	3	1	2	4	4	4	6	4	3	1	-	-
Male.....	1	-	1	1	3	2	2	1	1	-	-	-
Female.....	2	1	1	3	1	2	4	3	2	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	157	258	390	502	663	866	1,013	912	521	172	40	1
Male.....	87	135	216	277	335	444	507	378	195	43	12	1
Female.....	70	123	174	225	328	422	506	534	326	129	28	-
White, both sexes.....	124	229	350	456	605	803	936	853	479	161	33	1
Male.....	73	116	192	251	302	410	470	356	177	39	10	1
Female.....	51	113	158	205	303	393	466	497	302	122	23	-
Black, both sexes.....	33	28	38	42	56	59	77	57	40	11	7	-
Male.....	14	18	22	23	32	33	37	21	17	4	2	-
Female.....	19	10	16	19	24	26	40	36	23	7	5	-
American Indian 1/, both sexes.....	-	1	1	1	-	-	-	-	-	-	-	-
Male.....	-	1	1	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	3	2	4	-	2	2	-	-	-
Male.....	-	-	1	2	1	1	-	1	1	-	-	-
Female.....	-	-	-	1	1	3	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Texas

All races, both sexes.....	373	553	880	1,200	1,687	2,277	2,635	2,284	1,281	509	103	2
Male.....	215	308	464	644	898	1,196	1,320	1,061	515	147	25	2
Female.....	158	245	416	556	789	1,081	1,315	1,223	766	362	78	-
White, both sexes.....	311	471	796	1,103	1,533	2,082	2,456	2,126	1,205	465	90	2
Male.....	174	261	418	594	808	1,082	1,228	981	483	134	21	2
Female.....	137	210	378	509	725	1,000	1,228	1,145	722	331	69	-
Black, both sexes.....	59	74	72	85	136	164	149	141	66	40	11	-
Male.....	39	43	41	42	83	103	81	73	26	13	3	-
Female.....	20	31	31	43	53	61	68	68	40	27	8	-
American Indian 1/, both sexes.....	2	3	-	3	2	6	8	2	-	-	-	-
Male.....	1	1	-	1	2	-	2	1	-	-	-	-
Female.....	1	2	-	2	-	6	6	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	5	12	9	16	25	22	15	10	4	2	-
Male.....	1	3	5	7	5	11	9	6	6	-	1	-
Female.....	-	2	7	2	11	14	13	9	4	4	1	-

Utah

All races, both sexes.....	27	63	66	93	146	171	232	193	117	36	4	-
Male.....	18	30	43	55	90	95	126	93	61	10	1	-
Female.....	9	33	23	38	56	76	106	100	56	26	3	-
White, both sexes.....	27	62	62	88	143	168	226	187	116	35	4	-
Male.....	18	29	41	52	88	92	120	90	60	9	1	-
Female.....	9	33	21	36	55	76	106	97	56	26	3	-
Black, both sexes.....	-	-	-	1	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	1	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	3	3	-	1	1	1	-	-	-	-
Male.....	-	1	2	2	-	1	1	1	-	-	-	-
Female.....	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	1	3	2	4	4	1	1	-	-
Male.....	-	-	-	1	2	2	4	1	1	1	-	-
Female.....	-	-	1	-	1	-	-	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Vermont

All races, both sexes.....	8	12	24	35	57	67	87	98	52	19	7	-
Male.....	4	4	15	14	29	30	50	45	19	5	2	-
Female.....	4	8	9	21	28	37	37	53	33	14	5	-
White, both sexes.....	8	12	24	35	57	66	87	98	52	19	7	-
Male.....	4	4	15	14	29	29	50	45	19	5	2	-
Female.....	4	8	9	21	28	37	37	53	33	14	5	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	110	166	291	441	574	881	1,009	971	564	188	48	-
Male.....	70	88	153	230	305	455	481	424	201	53	9	-
Female.....	40	78	138	211	269	426	528	547	363	135	39	-
White, both sexes.....	80	118	231	371	491	768	893	886	499	161	38	-
Male.....	52	62	121	191	257	384	429	378	177	47	8	-
Female.....	28	56	110	180	234	384	464	508	322	114	30	-
Black, both sexes.....	30	46	57	66	78	99	106	77	60	25	9	-
Male.....	18	25	32	37	46	62	47	39	22	5	-	-
Female.....	12	21	25	29	32	37	59	38	38	20	9	-
American Indian 1/, both sexes.....	-	-	1	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	2	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	2	4	5	12	10	8	5	2	1	-
Male.....	-	1	-	2	2	7	5	7	2	1	1	-
Female.....	-	1	2	2	3	5	5	1	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Washington

All races, both sexes.....	98	161	239	359	537	687	778	824	459	178	33	-
Male.....	52	79	135	189	283	353	367	386	181	55	10	-
Female.....	46	82	104	170	254	334	411	438	278	123	23	-
White, both sexes.....	90	142	221	338	513	653	745	793	454	174	31	-
Male.....	46	66	123	177	268	329	351	374	180	52	9	-
Female.....	44	76	98	161	245	324	394	419	274	122	22	-
Black, both sexes.....	3	10	8	10	5	9	6	3	2	-	1	-
Male.....	1	6	4	5	2	6	3	-	1	-	1	-
Female.....	2	4	4	5	3	3	3	3	1	-	-	-
American Indian 1/, both sexes.....	2	4	8	7	8	5	4	6	2	-	-	-
Male.....	2	4	6	4	6	4	2	2	-	-	-	-
Female.....	-	-	2	3	2	1	2	4	2	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	5	2	4	11	20	23	22	1	4	1	-
Male.....	3	3	2	3	7	14	11	10	-	3	-	-
Female.....	-	2	-	1	4	6	12	12	1	1	1	-

West Virginia

All races, both sexes.....	54	93	139	201	270	356	416	345	207	54	13	-
Male.....	32	58	81	106	136	193	226	175	92	16	2	-
Female.....	22	35	58	95	134	163	190	170	115	38	11	-
White, both sexes.....	52	86	137	200	265	351	411	343	203	51	13	-
Male.....	30	53	79	106	134	190	223	173	89	16	2	-
Female.....	22	33	58	94	131	161	188	170	114	35	11	-
Black, both sexes.....	2	7	2	1	4	5	5	2	4	3	-	-
Male.....	2	5	2	-	2	3	3	2	3	-	-	-
Female.....	-	2	-	1	2	2	2	-	1	3	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of the respiratory system (J09-J98,U04)

Wisconsin

All races, both sexes.....	74	118	190	279	441	623	828	837	549	207	35	-
Male.....	50	67	114	145	238	326	432	359	217	66	10	-
Female.....	24	51	76	134	203	297	396	478	332	141	25	-
White, both sexes.....	63	109	171	261	422	601	806	816	538	205	34	-
Male.....	47	62	100	138	224	318	419	347	211	66	10	-
Female.....	16	47	71	123	198	283	387	469	327	139	24	-
Black, both sexes.....	8	6	11	15	14	15	14	14	6	1	1	-
Male.....	1	4	9	6	12	6	8	7	3	-	-	-
Female.....	7	2	2	9	2	9	6	7	3	1	1	-
American Indian 1/, both sexes.....	3	2	6	3	3	4	5	5	2	1	-	-
Male.....	2	-	3	1	2	1	2	3	1	-	-	-
Female.....	1	2	3	2	1	3	3	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	2	-	2	3	3	2	3	-	-	-
Male.....	-	1	2	-	-	1	3	2	2	-	-	-
Female.....	-	-	-	-	2	2	-	-	1	-	-	-

Wyoming

All races, both sexes.....	13	13	26	44	55	68	88	94	46	23	3	-
Male.....	5	7	14	25	31	37	38	35	16	10	2	-
Female.....	8	6	12	19	24	31	50	59	30	13	1	-
White, both sexes.....	12	13	26	43	53	67	85	94	46	22	3	-
Male.....	4	7	14	24	29	37	37	35	16	9	2	-
Female.....	8	6	12	19	24	30	48	59	30	13	1	-
Black, both sexes.....	-	-	-	-	1	-	1	-	-	1	-	-
Male.....	-	-	-	-	1	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	1	1	1	1	-	-	-	-	-
Male.....	1	-	-	1	1	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Influenza (J09-J11)

Alaska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Influenza (J09-J11)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Influenza (J09-J11)

Missouri

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Influenza (J09-J11)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Influenza (J09-J11)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Arkansas												
All races, both sexes.....	727	2	1	-	-	2	3	2	3	5	3	11
Male.....	345	1	-	-	-	-	3	2	-	3	3	6
Female.....	382	1	1	-	-	2	-	-	3	2	-	5
White, both sexes.....	651	2	1	-	-	2	2	1	2	4	3	7
Male.....	305	1	-	-	-	-	2	1	-	3	3	3
Female.....	346	1	1	-	-	2	-	-	2	1	-	4
Black, both sexes.....	73	-	-	-	-	-	1	1	1	1	-	3
Male.....	39	-	-	-	-	-	1	1	-	-	-	2
Female.....	34	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	6,510	24	7	4	5	5	8	10	18	24	39	75
Male.....	2,994	9	2	3	3	3	5	6	11	13	27	41
Female.....	3,516	15	5	1	2	2	3	4	7	11	12	34
White, both sexes.....	5,322	18	5	1	4	4	7	10	13	17	30	53
Male.....	2,400	7	-	1	2	3	4	6	8	9	21	26
Female.....	2,922	11	5	-	2	1	3	4	5	8	9	27
Black, both sexes.....	454	4	-	1	1	-	-	-	2	1	6	13
Male.....	225	2	-	-	1	-	-	-	2	1	3	10
Female.....	229	2	-	1	-	-	-	-	-	-	3	3
American Indian 1/, both sexes.....	18	1	-	-	-	-	-	-	-	2	1	1
Male.....	11	-	-	-	-	-	-	-	-	1	1	1
Female.....	7	1	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	716	1	2	2	-	1	1	-	3	4	2	8
Male.....	358	-	2	2	-	-	1	-	1	2	2	4
Female.....	358	1	-	-	-	1	-	-	2	2	-	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Iowa												
All races, both sexes.....	739	1	-	-	-	1	1	2	2	1	7	8
Male.....	293	1	-	-	-	1	1	2	2	1	3	5
Female.....	446	-	-	-	-	-	-	-	-	-	4	3
White, both sexes.....	730	1	-	-	-	1	1	2	2	1	7	7
Male.....	287	1	-	-	-	1	1	2	2	1	3	4
Female.....	443	-	-	-	-	-	-	-	-	-	4	3
Black, both sexes.....	8	-	-	-	-	-	-	-	-	-	-	1
Male.....	5	-	-	-	-	-	-	-	-	-	-	1
Female.....	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes.....	662	1	1	-	-	1	2	-	3	1	4	10
Male.....	292	1	-	-	-	-	1	-	1	-	4	2
Female.....	370	-	1	-	-	1	1	-	2	1	-	8
White, both sexes.....	639	1	1	-	-	-	2	-	3	1	3	8
Male.....	280	1	-	-	-	-	1	-	1	-	3	-
Female.....	359	-	1	-	-	-	1	-	2	1	-	8
Black, both sexes.....	17	-	-	-	-	-	-	-	-	-	1	1
Male.....	8	-	-	-	-	-	-	-	-	-	1	1
Female.....	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	1
Male.....	2	-	-	-	-	-	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	-	-	-	-	1	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Massachusetts												
All races, both sexes.....	1,526	2	1	-	-	-	-	-	-	4	8	10
Male.....	660	1	-	-	-	-	-	-	-	4	3	6
Female.....	866	1	1	-	-	-	-	-	-	-	5	4
White, both sexes.....	1,471	1	1	-	-	-	-	-	-	3	8	7
Male.....	633	-	-	-	-	-	-	-	-	3	3	4
Female.....	838	1	1	-	-	-	-	-	-	-	5	3
Black, both sexes.....	33	1	-	-	-	-	-	-	-	1	-	3
Male.....	18	1	-	-	-	-	-	-	-	1	-	2
Female.....	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	21	-	-	-	-	-	-	-	-	-	-	-
Male.....	8	-	-	-	-	-	-	-	-	-	-	-
Female.....	13	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes.....	1,628	4	7	-	3	2	1	7	4	8	20	23
Male.....	755	2	5	-	1	1	1	3	-	5	10	17
Female.....	873	2	2	-	2	1	-	4	4	3	10	6
White, both sexes.....	1,393	1	6	-	2	1	1	4	2	3	16	11
Male.....	639	1	4	-	1	1	1	2	-	2	9	9
Female.....	754	-	2	-	1	-	-	2	2	1	7	2
Black, both sexes.....	224	3	-	-	1	1	-	3	2	4	4	12
Male.....	111	1	-	-	-	-	-	1	-	3	1	8
Female.....	113	2	-	-	1	1	-	2	2	1	3	4
American Indian 1/, both sexes.....	7	-	-	-	-	-	-	-	-	-	-	-
Male.....	4	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	-	1	-	-	-	-	-	-	1	-	-
Male.....	1	-	1	-	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
United States												
All races, both sexes.....	1,095	1,379	1,736	2,183	3,326	5,708	9,072	11,137	9,040	4,095	1,124	3
Male.....	651	825	1,036	1,225	1,800	2,940	4,435	4,847	3,304	1,182	256	3
Female.....	444	554	700	958	1,526	2,768	4,637	6,290	5,736	2,913	868	-
White, both sexes.....	834	1,057	1,394	1,775	2,742	4,934	8,066	10,093	8,277	3,731	980	3
Male.....	489	646	822	981	1,476	2,532	3,951	4,370	3,031	1,081	218	3
Female.....	345	411	572	794	1,266	2,402	4,115	5,723	5,246	2,650	762	-
Black, both sexes.....	229	286	283	331	467	574	736	753	500	252	105	-
Male.....	139	156	174	195	260	301	340	313	166	60	21	-
Female.....	90	130	109	136	207	273	396	440	334	192	84	-
American Indian 1/, both sexes.....	11	17	17	20	18	34	32	31	27	11	5	-
Male.....	10	11	16	10	12	13	19	10	6	5	1	-
Female.....	1	6	1	10	6	21	13	21	21	6	4	-
Asian or Pacific Islander 2/, both sexes.....	21	19	42	57	99	166	238	260	236	101	34	-
Male.....	13	12	24	39	52	94	125	154	101	36	16	-
Female.....	8	7	18	18	47	72	113	106	135	65	18	-
Alabama												
All races, both sexes.....	22	29	43	48	71	105	148	177	132	44	13	-
Male.....	15	20	24	32	45	50	71	87	49	13	3	-
Female.....	7	9	19	16	26	55	77	90	83	31	10	-
White, both sexes.....	15	22	33	38	47	84	123	144	105	36	8	-
Male.....	10	15	18	25	31	38	56	72	39	11	2	-
Female.....	5	7	15	13	16	46	67	72	66	25	6	-
Black, both sexes.....	7	6	10	10	24	21	24	33	26	7	5	-
Male.....	5	4	6	7	14	12	14	15	10	2	1	-
Female.....	2	2	4	3	10	9	10	18	16	5	4	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	-	1	-	-	1	-	-
Male.....	-	1	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Alaska												
All races, both sexes.....	3	1	4	1	3	6	8	8	8	1	-	-
Male.....	1	-	3	1	3	2	4	4	3	-	-	-
Female.....	2	1	1	-	-	4	4	4	5	1	-	-
White, both sexes.....	3	1	3	1	2	3	4	6	7	1	-	-
Male.....	1	-	2	1	2	1	1	3	3	-	-	-
Female.....	2	1	1	-	-	2	3	3	4	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	1	3	4	1	1	-	-	-
Male.....	-	-	1	-	1	1	3	-	-	-	-	-
Female.....	-	-	-	-	-	2	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes.....	30	33	32	45	71	126	158	155	120	52	14	-
Male.....	14	19	22	28	35	63	78	83	48	18	5	-
Female.....	16	14	10	17	36	63	80	72	72	34	9	-
White, both sexes.....	24	29	26	40	66	114	148	143	109	47	11	-
Male.....	11	16	18	25	33	58	75	76	45	15	4	-
Female.....	13	13	8	15	33	56	73	67	64	32	7	-
Black, both sexes.....	2	2	1	-	1	2	3	4	-	1	-	-
Male.....	-	1	-	-	-	1	-	4	-	1	-	-
Female.....	2	1	1	-	1	1	3	-	-	-	-	-
American Indian 1/, both sexes.....	3	2	4	4	2	9	5	7	11	4	3	-
Male.....	2	2	4	2	1	3	2	2	3	2	1	-
Female.....	1	-	-	2	1	6	3	5	8	2	2	-
Asian or Pacific Islander 2/, both sexes.....	1	-	1	1	2	1	2	1	-	-	-	-
Male.....	1	-	-	1	1	1	1	1	-	-	-	-
Female.....	-	-	1	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonia (J12-J18)

Arkansas

All races, both sexes.....	20	25	34	31	44	74	148	141	116	49	13	-
Male.....	12	15	19	21	24	46	65	58	47	18	2	-
Female.....	8	10	15	10	20	28	83	83	69	31	11	-
White, both sexes.....	13	19	29	26	36	66	137	134	109	46	12	-
Male.....	8	12	16	16	21	40	60	54	46	17	2	-
Female.....	5	7	13	10	15	26	77	80	63	29	10	-
Black, both sexes.....	7	6	5	5	8	7	11	7	6	3	1	-
Male.....	4	3	3	5	3	6	5	4	1	1	-	-
Female.....	3	3	2	-	5	1	6	3	5	2	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	1	-	-	-

California

All races, both sexes.....	85	127	190	256	439	709	1,175	1,407	1,144	579	179	1
Male.....	45	77	100	151	234	369	566	642	459	179	48	1
Female.....	40	50	90	105	205	340	609	765	685	400	131	-
White, both sexes.....	65	99	145	190	332	558	980	1,189	970	487	144	1
Male.....	35	63	76	108	175	287	474	515	392	151	36	1
Female.....	30	36	69	82	157	271	506	674	578	336	108	-
Black, both sexes.....	14	21	26	33	50	56	68	70	43	31	14	-
Male.....	7	10	17	20	30	28	31	38	16	7	2	-
Female.....	7	11	9	13	20	28	37	32	27	24	12	-
American Indian 1/, both sexes.....	-	-	-	2	-	3	2	6	-	-	-	-
Male.....	-	-	-	1	-	2	2	3	-	-	-	-
Female.....	-	-	-	1	-	1	-	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6	7	19	31	57	92	125	142	131	61	21	-
Male.....	3	4	7	22	29	52	59	86	51	21	10	-
Female.....	3	3	12	9	28	40	66	56	80	40	11	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Delaware												
All races, both sexes.....	2	2	4	3	13	15	12	32	22	5	2	-
Male.....	1	1	1	2	6	7	5	16	6	2	1	-
Female.....	1	1	3	1	7	8	7	16	16	3	1	-
White, both sexes.....	2	1	3	2	11	14	11	28	20	5	2	-
Male.....	1	-	1	1	5	6	4	13	5	2	1	-
Female.....	1	1	2	1	6	8	7	15	15	3	1	-
Black, both sexes.....	-	1	1	-	2	1	1	4	2	-	-	-
Male.....	-	1	-	-	1	1	1	3	1	-	-	-
Female.....	-	-	1	-	1	-	-	1	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes.....	3	4	1	4	3	5	7	18	17	11	2	-
Male.....	3	4	1	4	1	3	5	4	5	1	1	-
Female.....	-	-	-	-	2	2	2	14	12	10	1	-
White, both sexes.....	1	-	-	1	1	1	1	4	10	2	-	-
Male.....	1	-	-	1	-	1	1	2	3	-	-	-
Female.....	-	-	-	-	1	-	-	2	7	2	-	-
Black, both sexes.....	2	4	1	3	2	4	6	14	5	9	2	-
Male.....	2	4	1	3	1	2	4	2	1	1	1	-
Female.....	-	-	-	-	1	2	2	12	4	8	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	2	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia (J12-J18)

Florida

All races, both sexes.....	52	67	91	99	167	252	372	469	370	140	29	-
Male.....	34	42	52	62	89	140	195	207	129	50	7	-
Female.....	18	25	39	37	78	112	177	262	241	90	22	-
White, both sexes.....	43	51	67	80	143	230	340	438	346	133	23	-
Male.....	28	34	37	51	76	134	180	197	123	46	7	-
Female.....	15	17	30	29	67	96	160	241	223	87	16	-
Black, both sexes.....	8	16	21	17	24	22	30	30	20	6	6	-
Male.....	6	8	12	11	13	6	14	10	5	3	-	-
Female.....	2	8	9	6	11	16	16	20	15	3	6	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	1	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	2	2	-	-	1	1	4	1	-	-
Male.....	-	-	2	-	-	-	1	-	1	1	-	-
Female.....	1	-	-	2	-	-	-	1	3	-	-	-

Georgia

All races, both sexes.....	28	49	46	78	108	161	227	268	206	113	18	-
Male.....	17	28	27	40	69	71	105	100	66	33	3	-
Female.....	11	21	19	38	39	90	122	168	140	80	15	-
White, both sexes.....	14	28	33	52	84	134	189	221	173	95	12	-
Male.....	9	18	18	27	52	60	90	82	57	27	1	-
Female.....	5	10	15	25	32	74	99	139	116	68	11	-
Black, both sexes.....	13	21	13	25	24	26	33	46	32	18	6	-
Male.....	7	10	9	13	17	10	11	17	8	6	2	-
Female.....	6	11	4	12	7	16	22	29	24	12	4	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	1	-	1	5	1	1	-	-	-
Male.....	1	-	-	-	-	1	4	1	1	-	-	-
Female.....	-	-	-	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia (J12-J18)

Hawaii

All races, both sexes.....	2	3	6	3	6	27	31	52	45	13	6	-
Male.....	1	3	3	2	3	18	23	32	23	6	5	-
Female.....	1	-	3	1	3	9	8	20	22	7	1	-
White, both sexes.....	-	2	2	1	2	5	2	15	12	2	2	-
Male.....	-	2	-	-	1	4	2	8	8	1	1	-
Female.....	-	-	2	1	1	1	-	7	4	1	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	1	4	2	4	22	29	37	33	11	4	-
Male.....	1	1	3	2	2	14	21	24	15	5	4	-
Female.....	1	-	1	-	2	8	8	13	18	6	-	-

Idaho

All races, both sexes.....	2	10	9	6	3	24	32	48	50	21	4	-
Male.....	2	4	7	5	-	7	16	21	10	10	-	-
Female.....	-	6	2	1	3	17	16	27	40	11	4	-
White, both sexes.....	2	9	9	6	3	24	30	48	50	21	4	-
Male.....	2	4	7	5	-	7	15	21	10	10	-	-
Female.....	-	5	2	1	3	17	15	27	40	11	4	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonia (J12-J18)

Illinois

All races, both sexes.....	63	67	89	109	124	252	470	536	450	201	58	-
Male.....	31	45	58	60	59	125	229	214	155	53	9	-
Female.....	32	22	31	49	65	127	241	322	295	148	49	-
White, both sexes.....	50	48	64	84	97	202	400	490	405	194	52	-
Male.....	24	32	44	44	45	99	196	198	138	51	6	-
Female.....	26	16	20	40	52	103	204	292	267	143	46	-
Black, both sexes.....	11	17	24	21	26	46	61	41	37	5	4	-
Male.....	6	11	13	14	14	23	28	15	15	-	2	-
Female.....	5	6	11	7	12	23	33	26	22	5	2	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	2	1	3	1	4	9	5	7	2	2	-
Male.....	1	2	1	2	-	3	5	1	2	2	1	-
Female.....	1	-	-	1	1	1	4	4	5	-	1	-

Indiana

All races, both sexes.....	21	30	41	46	54	106	186	257	170	90	32	-
Male.....	14	16	22	26	25	51	80	110	60	21	4	-
Female.....	7	14	19	20	29	55	106	147	110	69	28	-
White, both sexes.....	19	25	38	44	47	99	175	252	162	86	32	-
Male.....	12	15	21	25	24	50	74	107	59	19	4	-
Female.....	7	10	17	19	23	49	101	145	103	67	28	-
Black, both sexes.....	2	5	3	2	7	6	11	5	8	4	-	-
Male.....	2	1	1	1	1	1	6	3	1	2	-	-
Female.....	-	4	2	1	6	5	5	2	7	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia (J12-J18)

Iowa

All races, both sexes.....	8	16	19	26	39	58	111	167	167	86	19	-
Male.....	6	6	10	12	18	26	60	64	58	15	2	-
Female.....	2	10	9	14	21	32	51	103	109	71	17	-
White, both sexes.....	8	14	18	26	37	58	109	167	166	86	19	-
Male.....	6	6	9	12	16	26	58	64	58	15	2	-
Female.....	2	8	9	14	21	32	51	103	108	71	17	-
Black, both sexes.....	-	2	1	-	1	-	2	-	1	-	-	-
Male.....	-	-	1	-	1	-	2	-	-	-	-	-
Female.....	-	2	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Kansas

All races, both sexes.....	10	13	11	18	36	60	115	142	148	70	16	-
Male.....	6	8	7	9	24	37	56	68	50	14	4	-
Female.....	4	5	4	9	12	23	59	74	98	56	12	-
White, both sexes.....	8	13	11	18	35	59	109	137	145	69	16	-
Male.....	4	8	7	9	23	36	54	66	49	14	4	-
Female.....	4	5	4	9	12	23	55	71	96	55	12	-
Black, both sexes.....	1	-	-	-	-	1	4	5	3	1	-	-
Male.....	1	-	-	-	-	1	1	2	1	-	-	-
Female.....	-	-	-	-	-	-	3	3	2	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	1	-	1	-	-	-	-	-
Male.....	1	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Maine												
All races, both sexes.....	2	5	7	9	12	23	46	53	53	16	6	-
Male.....	1	2	3	4	8	8	20	25	17	3	-	-
Female.....	1	3	4	5	4	15	26	28	36	13	6	-
White, both sexes.....	2	5	7	9	12	23	46	53	53	16	6	-
Male.....	1	2	3	4	8	8	20	25	17	3	-	-
Female.....	1	3	4	5	4	15	26	28	36	13	6	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes.....	24	26	40	35	66	101	169	201	171	73	18	-
Male.....	17	11	24	20	41	57	85	85	68	18	4	-
Female.....	7	15	16	15	25	44	84	116	103	55	14	-
White, both sexes.....	11	15	29	26	49	72	141	163	156	65	15	-
Male.....	9	7	17	17	31	43	75	70	63	16	4	-
Female.....	2	8	12	9	18	29	66	93	93	49	11	-
Black, both sexes.....	10	10	11	9	16	29	26	34	12	8	3	-
Male.....	6	3	7	3	9	14	10	14	3	2	-	-
Female.....	4	7	4	6	7	15	16	20	9	6	3	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	1	-	-	1	-	2	4	3	-	-	-
Male.....	2	1	-	-	1	-	-	1	2	-	-	-
Female.....	1	-	-	-	-	-	2	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Massachusetts												
All races, both sexes.....	19	32	29	43	89	149	269	351	345	139	36	-
Male.....	8	17	19	23	59	85	126	151	119	34	5	-
Female.....	11	15	10	20	30	64	143	200	226	105	31	-
White, both sexes.....	19	28	28	40	85	139	261	345	336	135	35	-
Male.....	8	16	18	21	55	80	124	148	114	34	5	-
Female.....	11	12	10	19	30	59	137	197	222	101	30	-
Black, both sexes.....	-	3	-	2	3	7	3	5	3	2	-	-
Male.....	-	1	-	1	3	4	-	3	2	-	-	-
Female.....	-	2	-	1	-	3	3	2	1	2	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	-	1	3	5	1	6	2	1	-
Male.....	-	-	1	-	1	1	2	-	3	-	-	-
Female.....	-	1	-	-	-	2	3	1	3	2	1	-
Michigan												
All races, both sexes.....	40	51	63	61	90	195	263	328	302	121	35	-
Male.....	21	30	41	30	46	96	133	141	127	37	8	-
Female.....	19	21	22	31	44	99	130	187	175	84	27	-
White, both sexes.....	25	33	54	40	72	174	227	298	277	115	31	-
Male.....	12	23	36	18	35	82	115	129	116	36	7	-
Female.....	13	10	18	22	37	92	112	169	161	79	24	-
Black, both sexes.....	15	17	9	21	16	19	35	29	23	6	4	-
Male.....	9	7	5	12	9	14	17	11	11	1	1	-
Female.....	6	10	4	9	7	5	18	18	12	5	3	-
American Indian 1/, both sexes.....	-	1	-	-	2	1	1	1	1	-	-	-
Male.....	-	-	-	-	2	-	1	1	-	-	-	-
Female.....	-	1	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia (J12-J18)

Minnesota

All races, both sexes.....	13	11	15	14	24	58	91	133	125	63	21	-
Male.....	8	7	11	6	13	37	46	59	60	24	6	-
Female.....	5	4	4	8	11	21	45	74	65	39	15	-
White, both sexes.....	11	10	12	13	23	55	90	130	123	63	20	-
Male.....	6	6	8	5	12	35	45	58	58	24	6	-
Female.....	5	4	4	8	11	20	45	72	65	39	14	-
Black, both sexes.....	-	1	-	-	1	1	-	2	1	-	1	-
Male.....	-	1	-	-	1	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	1	-
American Indian 1/, both sexes.....	2	-	3	1	-	1	1	1	-	-	-	-
Male.....	2	-	3	1	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	1	-	-	-
Male.....	-	-	-	-	-	1	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Mississippi

All races, both sexes.....	24	21	14	28	41	70	89	110	83	26	12	-
Male.....	15	12	9	14	20	37	43	47	22	7	1	-
Female.....	9	9	5	14	21	33	46	63	61	19	11	-
White, both sexes.....	11	11	7	18	27	53	68	79	72	18	9	-
Male.....	6	7	4	6	13	25	31	35	16	7	-	-
Female.....	5	4	3	12	14	28	37	44	56	11	9	-
Black, both sexes.....	13	10	6	10	14	17	20	30	10	7	3	-
Male.....	9	5	4	8	7	12	12	11	5	-	1	-
Female.....	4	5	2	2	7	5	8	19	5	7	2	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	-	-	-	1	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Nebraska												
All races, both sexes.....	7	3	2	8	20	39	48	78	65	32	11	-
Male.....	4	3	2	4	12	24	28	27	18	7	4	-
Female.....	3	-	-	4	8	15	20	51	47	25	7	-
White, both sexes.....	6	2	1	8	19	36	48	77	65	32	11	-
Male.....	4	2	1	4	11	22	28	27	18	7	4	-
Female.....	2	-	-	4	8	14	20	50	47	25	7	-
Black, both sexes.....	1	1	1	-	1	2	-	1	-	-	-	-
Male.....	-	1	1	-	1	2	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes.....	14	20	19	22	41	62	79	52	45	12	5	-
Male.....	9	10	12	10	29	32	32	25	17	5	2	-
Female.....	5	10	7	12	12	30	47	27	28	7	3	-
White, both sexes.....	13	17	17	21	37	57	74	48	43	11	3	-
Male.....	9	9	12	10	26	29	31	24	16	5	2	-
Female.....	4	8	5	11	11	28	43	24	27	6	1	-
Black, both sexes.....	1	1	1	1	1	1	3	2	1	-	2	-
Male.....	-	-	-	-	1	1	1	-	-	-	-	-
Female.....	1	1	1	1	-	-	2	2	1	-	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	1	-	3	3	2	2	1	1	-	-
Male.....	-	1	-	-	2	2	-	1	1	-	-	-
Female.....	-	1	1	-	1	1	2	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia (J12-J18)

New Hampshire

All races, both sexes.....	3	2	6	9	9	18	43	47	39	15	6	-
Male.....	1	2	5	2	4	7	16	24	13	2	1	-
Female.....	2	-	1	7	5	11	27	23	26	13	5	-
White, both sexes.....	3	2	6	9	9	18	43	47	38	15	6	-
Male.....	1	2	5	2	4	7	16	24	13	2	1	-
Female.....	2	-	1	7	5	11	27	23	25	13	5	-
Black, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	25	25	45	63	77	150	250	294	237	105	18	-
Male.....	15	14	29	32	39	90	127	128	82	28	2	-
Female.....	10	11	16	31	38	60	123	166	155	77	16	-
White, both sexes.....	17	15	37	50	66	138	221	278	215	94	17	-
Male.....	10	8	23	25	35	80	114	120	77	27	2	-
Female.....	7	7	14	25	31	58	107	158	138	67	15	-
Black, both sexes.....	8	10	6	13	8	10	24	14	21	9	1	-
Male.....	5	6	4	7	1	8	11	7	4	1	-	-
Female.....	3	4	2	6	7	2	13	7	17	8	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	-	3	2	5	2	1	2	-	-
Male.....	-	-	2	-	3	2	2	1	1	-	-	-
Female.....	-	-	-	-	-	-	3	1	-	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia (J12-J18)

New Mexico

All races, both sexes.....	8	8	12	15	16	39	48	56	36	25	5	-
Male.....	5	5	10	10	8	16	23	25	10	10	1	-
Female.....	3	3	2	5	8	23	25	31	26	15	4	-
White, both sexes.....	7	6	9	14	16	34	44	52	33	23	4	-
Male.....	4	4	7	9	8	16	21	24	9	9	1	-
Female.....	3	2	2	5	8	18	23	28	24	14	3	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	2	3	1	-	5	4	4	3	2	1	-
Male.....	1	1	3	1	-	-	2	1	1	1	-	-
Female.....	-	1	-	-	-	5	2	3	2	1	1	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	102	122	139	176	278	451	765	942	804	372	98	-
Male.....	72	70	82	97	153	243	369	417	283	105	19	-
Female.....	30	52	57	79	125	208	396	525	521	267	79	-
White, both sexes.....	69	87	96	125	194	349	640	796	719	326	79	-
Male.....	52	52	61	71	112	190	315	351	255	89	17	-
Female.....	17	35	35	54	82	159	325	445	464	237	62	-
Black, both sexes.....	32	31	39	43	69	89	101	114	60	33	15	-
Male.....	20	16	18	21	31	47	40	45	17	11	2	-
Female.....	12	15	21	22	38	42	61	69	43	22	13	-
American Indian 1/, both sexes.....	-	1	-	-	1	-	1	-	1	-	-	-
Male.....	-	1	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	3	4	8	14	13	23	32	24	13	4	-
Male.....	-	1	3	5	9	6	14	21	11	5	-	-
Female.....	1	2	1	3	5	7	9	11	13	8	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Oregon												
All races, both sexes.....	6	9	13	22	22	45	68	113	108	35	10	-
Male.....	4	5	6	12	12	26	34	44	52	12	3	-
Female.....	2	4	7	10	10	19	34	69	56	23	7	-
White, both sexes.....	6	8	13	21	22	45	65	110	106	33	10	-
Male.....	4	5	6	12	12	26	33	42	51	12	3	-
Female.....	2	3	7	9	10	19	32	68	55	21	7	-
Black, both sexes.....	-	-	-	-	-	-	-	-	1	1	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	1	-	1	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	2	3	1	1	-	-
Male.....	-	-	-	-	-	-	1	2	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	1	1	-	-
Pennsylvania												
All races, both sexes.....	41	64	69	96	156	274	455	595	436	193	52	1
Male.....	20	39	39	55	93	153	217	267	161	65	15	1
Female.....	21	25	30	41	63	121	238	328	275	128	37	-
White, both sexes.....	34	46	61	85	140	245	420	562	417	183	49	1
Male.....	16	26	34	50	83	139	196	255	153	64	14	1
Female.....	18	20	27	35	57	106	224	307	264	119	35	-
Black, both sexes.....	7	18	7	10	14	27	34	27	17	9	3	-
Male.....	4	13	4	4	10	13	21	8	7	-	1	-
Female.....	3	5	3	6	4	14	13	19	10	9	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	1	2	2	1	6	2	1	-	-
Male.....	-	-	1	1	-	1	-	4	1	1	-	-
Female.....	-	-	-	-	2	1	1	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia (J12-J18)

South Dakota

All races, both sexes.....	5	2	1	5	17	17	27	30	36	27	10	-
Male.....	5	1	1	3	8	6	11	12	13	7	3	-
Female.....	-	1	-	2	9	11	16	18	23	20	7	-
White, both sexes.....	4	1	1	5	15	15	26	30	34	26	10	-
Male.....	4	1	1	3	7	5	10	12	12	7	3	-
Female.....	-	-	-	2	8	10	16	18	22	19	7	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	-	-	2	2	1	-	2	1	-	-
Male.....	1	-	-	-	1	1	1	-	1	-	-	-
Female.....	-	1	-	-	1	1	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	31	48	58	64	95	179	247	296	223	89	25	-
Male.....	18	33	33	39	45	80	114	115	86	18	8	-
Female.....	13	15	25	25	50	99	133	181	137	71	17	-
White, both sexes.....	23	45	52	54	82	159	222	277	209	82	21	-
Male.....	14	30	29	32	36	70	104	109	80	16	6	-
Female.....	9	15	23	22	46	89	118	168	129	66	15	-
Black, both sexes.....	8	3	5	9	13	17	25	18	13	7	4	-
Male.....	4	3	3	6	9	9	10	6	6	2	2	-
Female.....	4	-	2	3	4	8	15	12	7	5	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	1	-	3	-	1	1	-	-	-
Male.....	-	-	1	1	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	2	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia (J12-J18)

Vermont

All races, both sexes.....	-	1	-	3	3	8	9	16	10	9	4	-
Male.....	-	-	-	3	1	5	5	8	2	5	1	-
Female.....	-	1	-	-	2	3	4	8	8	4	3	-
White, both sexes.....	-	1	-	3	3	8	9	16	10	9	4	-
Male.....	-	-	-	3	1	5	5	8	2	5	1	-
Female.....	-	1	-	-	2	3	4	8	8	4	3	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	21	25	38	68	74	137	200	282	213	86	27	-
Male.....	14	17	29	41	42	69	105	125	77	19	6	-
Female.....	7	8	9	27	32	68	95	157	136	67	21	-
White, both sexes.....	11	17	25	50	59	107	179	256	190	76	22	-
Male.....	7	11	20	29	32	52	99	109	67	19	6	-
Female.....	4	6	5	21	27	55	80	147	123	57	16	-
Black, both sexes.....	10	8	13	16	15	25	20	22	19	9	5	-
Male.....	7	6	9	11	10	14	5	13	8	-	-	-
Female.....	3	2	4	5	5	11	15	9	11	9	5	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	-	5	1	4	4	1	-	-
Male.....	-	-	-	1	-	3	1	3	2	-	-	-
Female.....	-	-	-	1	-	2	-	1	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia (J12-J18)

Wisconsin

All races, both sexes.....	18	22	19	28	56	92	172	236	208	93	19	-
Male.....	12	14	11	14	30	44	83	97	83	23	4	-
Female.....	6	8	8	14	26	48	89	139	125	70	15	-
White, both sexes.....	17	21	18	27	52	90	171	229	204	92	18	-
Male.....	11	14	10	14	29	44	82	95	80	23	4	-
Female.....	6	7	8	13	23	46	89	134	124	69	14	-
Black, both sexes.....	1	1	1	1	1	2	1	6	3	1	1	-
Male.....	1	-	1	-	1	-	1	2	2	-	-	-
Female.....	-	1	-	1	-	2	-	4	1	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	2	-	-	-	-	-	-	-

Wyoming

All races, both sexes.....	4	2	5	6	2	8	17	28	23	12	-	-
Male.....	2	1	3	3	1	3	8	10	10	5	-	-
Female.....	2	1	2	3	1	5	9	18	13	7	-	-
White, both sexes.....	3	2	5	6	1	8	16	28	23	11	-	-
Male.....	1	1	3	3	-	3	8	10	10	4	-	-
Female.....	2	1	2	3	1	5	8	18	13	7	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	1	-	-	-	-	-	-	-
Male.....	1	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Viral pneumonia, not elsewhere classified (J12)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Viral pneumonia, not elsewhere classified (J12)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Viral pneumonia, not elsewhere classified (J12)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Viral pneumonia, not elsewhere classified (J12)

Massachusetts

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.

Michigan

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Viral pneumonia, not elsewhere classified (J12)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Viral pneumonia, not elsewhere classified (J12)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Viral pneumonia, not elsewhere classified (J12)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Viral pneumonia, not elsewhere classified (J12)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Viral pneumonia, not elsewhere classified (J12)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Viral pneumonia, not elsewhere classified (J12)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Viral pneumonia, not elsewhere classified (J12)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bacterial pneumonia (J13-J15)												
Arkansas												
All races, both sexes.....	30	-	-	-	-	1	1	-	-	1	1	-
Male.....	16	-	-	-	-	-	1	-	-	1	1	-
Female.....	14	-	-	-	-	1	-	-	-	-	-	-
White, both sexes.....	29	-	-	-	-	1	-	-	-	1	1	-
Male.....	15	-	-	-	-	-	-	-	-	1	1	-
Female.....	14	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes.....	1	-	-	-	-	-	1	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	401	7	3	1	1	2	1	4	2	8	8	10
Male.....	198	2	1	1	1	-	1	3	1	3	5	7
Female.....	203	5	2	-	-	2	-	1	1	5	3	3
White, both sexes.....	318	5	2	-	1	1	1	4	-	7	7	7
Male.....	153	1	-	-	1	-	1	3	-	3	4	4
Female.....	165	4	2	-	-	1	-	1	-	4	3	3
Black, both sexes.....	37	1	-	-	-	-	-	-	-	-	-	2
Male.....	18	1	-	-	-	-	-	-	-	-	-	2
Female.....	19	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	44	1	1	1	-	1	-	-	2	1	1	1
Male.....	25	-	1	1	-	-	-	-	1	-	1	1
Female.....	19	1	-	-	-	1	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Bacterial pneumonia (J13-J15)

Iowa												
All races, both sexes.....	31	-	-	-	-	-	-	1	-	-	-	1
Male.....	14	-	-	-	-	-	-	1	-	-	-	-
Female.....	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes.....	31	-	-	-	-	-	-	1	-	-	-	1
Male.....	14	-	-	-	-	-	-	1	-	-	-	-
Female.....	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Kansas												
All races, both sexes.....	17	-	1	-	-	1	-	-	-	-	-	-
Male.....	10	-	-	-	-	-	-	-	-	-	-	-
Female.....	7	-	1	-	-	1	-	-	-	-	-	-
White, both sexes.....	16	-	1	-	-	-	-	-	-	-	-	-
Male.....	10	-	-	-	-	-	-	-	-	-	-	-
Female.....	6	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bacterial pneumonia (J13-J15)

Arkansas

All races, both sexes.....	-	2	2	3	1	4	5	7	1	1	-	-
Male.....	-	-	1	-	1	4	3	4	-	-	-	-
Female.....	-	2	1	3	-	-	2	3	1	1	-	-
White, both sexes.....	-	2	2	3	-	4	5	7	1	1	-	-
Male.....	-	-	1	-	1	4	3	4	-	-	-	-
Female.....	-	2	1	3	-	-	2	3	1	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	9	17	14	17	21	46	67	78	47	32	6	-
Male.....	3	11	8	11	11	25	35	46	13	7	3	-
Female.....	6	6	6	6	10	21	32	32	34	25	3	-
White, both sexes.....	7	8	9	9	10	37	58	68	42	29	6	-
Male.....	3	7	7	5	6	20	30	38	11	6	3	-
Female.....	4	1	2	4	4	17	28	30	31	23	3	-
Black, both sexes.....	1	8	2	5	6	3	2	4	1	2	-	-
Male.....	-	3	1	3	2	2	1	3	-	-	-	-
Female.....	1	5	1	2	4	1	1	1	1	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	3	3	5	5	6	6	4	1	-	-
Male.....	-	1	-	3	3	2	3	5	2	1	-	-
Female.....	1	-	3	-	2	3	3	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Bacterial pneumonia (J13-J15)

Massachusetts

All races, both sexes.....	-	1	2	1	6	4	8	8	5	2	1	-
Male.....	-	-	1	-	5	-	4	4	4	2	-	-
Female.....	-	1	1	1	1	4	4	4	1	-	1	-
White, both sexes.....	-	1	2	1	6	4	8	8	5	2	1	-
Male.....	-	-	1	-	5	-	4	4	4	2	-	-
Female.....	-	1	1	1	1	4	4	4	1	-	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Michigan

All races, both sexes.....	1	6	4	6	5	3	10	9	7	6	-	-
Male.....	-	4	1	3	1	2	6	3	3	2	-	-
Female.....	1	2	3	3	4	1	4	6	4	4	-	-
White, both sexes.....	1	5	2	5	5	3	10	9	6	6	-	-
Male.....	-	3	-	2	1	2	6	3	3	2	-	-
Female.....	1	2	2	3	4	1	4	6	3	4	-	-
Black, both sexes.....	-	1	2	1	-	-	-	-	-	-	-	-
Male.....	-	1	1	1	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Bacterial pneumonia (J13-J15)

New Mexico

All races, both sexes.....	3	3	2	1	2	2	1	1	-	1	-	-
Male.....	1	1	2	1	1	-	1	1	-	-	-	-
Female.....	2	2	-	-	1	2	-	-	-	1	-	-
White, both sexes.....	2	2	1	1	2	1	1	1	-	1	-	-
Male.....	-	1	1	1	1	-	1	1	-	-	-	-
Female.....	2	1	-	-	1	1	-	-	-	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	1	-	-	1	-	-	-	-	-	-
Male.....	1	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	12	22	27	18	52	45	65	75	73	15	8	-
Male.....	7	14	21	11	34	26	30	36	19	3	5	-
Female.....	5	8	6	7	18	19	35	39	54	12	3	-
White, both sexes.....	10	14	19	10	36	28	49	60	55	12	6	-
Male.....	7	9	16	6	23	17	20	27	17	3	4	-
Female.....	3	5	3	4	13	11	29	33	38	9	2	-
Black, both sexes.....	2	8	8	6	14	16	12	12	11	3	1	-
Male.....	-	5	5	4	10	8	7	7	1	-	1	-
Female.....	2	3	3	2	4	8	5	5	10	3	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	1	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	1	1	4	3	6	-	1	-
Male.....	-	-	-	1	-	1	3	2	1	-	-	-
Female.....	-	-	-	1	1	-	1	1	5	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bacterial pneumonia (J13-J15)

Oregon

All races, both sexes.....	-	3	1	2	3	3	2	10	5	2	-	-
Male.....	-	3	-	1	2	1	1	5	2	1	-	-
Female.....	-	-	1	1	1	2	1	5	3	1	-	-
White, both sexes.....	-	3	1	2	3	3	2	9	5	2	-	-
Male.....	-	3	-	1	2	1	1	5	2	1	-	-
Female.....	-	-	1	1	1	2	1	4	3	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Pennsylvania

All races, both sexes.....	1	7	3	6	5	10	9	14	7	1	-	-
Male.....	-	4	2	3	2	8	4	7	4	-	-	-
Female.....	1	3	1	3	3	2	5	7	3	1	-	-
White, both sexes.....	1	4	3	5	4	10	8	13	7	1	-	-
Male.....	-	2	2	3	2	8	4	6	4	-	-	-
Female.....	1	2	1	2	2	2	4	7	3	1	-	-
Black, both sexes.....	-	3	-	1	-	-	1	-	-	-	-	-
Male.....	-	2	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Arkansas												
All races, both sexes.....	695	2	1	-	-	1	2	2	3	4	2	11
Male.....	328	1	-	-	-	-	2	2	-	2	2	6
Female.....	367	1	1	-	-	1	-	-	3	2	-	5
White, both sexes.....	620	2	1	-	-	1	2	1	2	3	2	7
Male.....	289	1	-	-	-	-	2	1	-	2	2	3
Female.....	331	1	1	-	-	1	-	-	2	1	-	4
Black, both sexes.....	72	-	-	-	-	-	-	1	1	1	-	3
Male.....	38	-	-	-	-	-	-	1	-	-	-	2
Female.....	34	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	6,083	10	3	3	4	3	7	6	16	15	31	64
Male.....	2,788	5	-	2	2	3	4	3	10	10	22	34
Female.....	3,295	5	3	1	2	-	3	3	6	5	9	30
White, both sexes.....	4,983	6	3	1	3	3	6	6	13	10	23	45
Male.....	2,242	4	-	1	1	3	3	3	8	6	17	22
Female.....	2,741	2	3	-	2	-	3	3	5	4	6	23
Black, both sexes.....	416	3	-	1	1	-	-	-	2	1	6	11
Male.....	207	1	-	-	1	-	-	-	2	1	3	8
Female.....	209	2	-	1	-	-	-	-	-	-	3	3
American Indian 1/, both sexes.....	15	1	-	-	-	-	-	-	-	1	1	1
Male.....	9	-	-	-	-	-	-	-	-	1	1	1
Female.....	6	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	669	-	-	1	-	-	1	-	1	3	1	7
Male.....	330	-	-	1	-	-	1	-	-	2	1	3
Female.....	339	-	-	-	-	-	-	-	1	1	-	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia due to other or unspecified organisms (J16,J18)												
Massachusetts												
All races, both sexes.....	1,483	2	1	-	-	-	-	-	-	4	7	9
Male.....	639	1	-	-	-	-	-	-	-	4	3	6
Female.....	844	1	1	-	-	-	-	-	-	-	4	3
White, both sexes.....	1,428	1	1	-	-	-	-	-	-	3	7	6
Male.....	612	-	-	-	-	-	-	-	-	3	3	4
Female.....	816	1	1	-	-	-	-	-	-	-	4	2
Black, both sexes.....	33	1	-	-	-	-	-	-	-	1	-	3
Male.....	18	1	-	-	-	-	-	-	-	1	-	2
Female.....	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	21	-	-	-	-	-	-	-	-	-	-	-
Male.....	8	-	-	-	-	-	-	-	-	-	-	-
Female.....	13	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes.....	1,559	2	6	-	3	2	1	6	4	7	18	22
Male.....	722	2	4	-	1	1	1	2	-	4	8	16
Female.....	837	-	2	-	2	1	-	4	4	3	10	6
White, both sexes.....	1,332	1	5	-	2	1	1	3	2	2	14	11
Male.....	610	1	3	-	1	1	1	1	-	1	7	9
Female.....	722	-	2	-	1	-	-	2	2	1	7	2
Black, both sexes.....	217	1	-	-	1	1	-	3	2	4	4	11
Male.....	107	1	-	-	-	-	-	1	-	3	1	7
Female.....	110	-	-	-	1	1	-	2	2	1	3	4
American Indian 1/, both sexes.....	7	-	-	-	-	-	-	-	-	-	-	-
Male.....	4	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	1	-	-	-	-	-	-	1	-	-
Male.....	1	-	1	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
United States												
All races, both sexes.....	993	1,218	1,569	2,017	3,076	5,349	8,605	10,618	8,689	3,953	1,090	3
Male.....	604	728	927	1,134	1,660	2,754	4,194	4,581	3,189	1,142	246	3
Female.....	389	490	642	883	1,416	2,595	4,411	6,037	5,500	2,811	844	-
White, both sexes.....	752	938	1,260	1,639	2,537	4,624	7,659	9,628	7,968	3,599	953	3
Male.....	450	571	731	908	1,360	2,372	3,747	4,133	2,928	1,043	209	3
Female.....	302	367	529	731	1,177	2,252	3,912	5,495	5,040	2,556	744	-
Black, both sexes.....	212	248	258	309	433	540	695	713	473	243	99	-
Male.....	133	136	160	183	241	283	315	294	159	59	20	-
Female.....	79	112	98	126	192	257	380	419	314	184	79	-
American Indian 1/, both sexes.....	10	14	14	18	16	29	28	30	24	11	5	-
Male.....	9	10	13	9	10	10	16	10	5	5	1	-
Female.....	1	4	1	9	6	19	12	20	19	6	4	-
Asian or Pacific Islander 2/, both sexes.....	19	18	37	51	90	156	223	247	224	100	33	-
Male.....	12	11	23	34	49	89	116	144	97	35	16	-
Female.....	7	7	14	17	41	67	107	103	127	65	17	-
Alabama												
All races, both sexes.....	17	26	40	44	66	102	138	171	127	44	13	-
Male.....	13	18	22	28	41	48	64	85	46	13	3	-
Female.....	4	8	18	16	25	54	74	86	81	31	10	-
White, both sexes.....	12	20	31	35	42	81	114	139	100	36	8	-
Male.....	8	13	16	22	27	36	50	70	36	11	2	-
Female.....	4	7	15	13	15	45	64	69	64	25	6	-
Black, both sexes.....	5	5	9	9	24	21	23	32	26	7	5	-
Male.....	5	4	6	6	14	12	13	15	10	2	1	-
Female.....	-	1	3	3	10	9	10	17	16	5	4	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	-	1	-	-	1	-	-
Male.....	-	1	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

Alaska

All races, both sexes.....	3	1	4	1	3	6	7	8	8	1	-	-
Male.....	1	-	3	1	3	2	4	4	3	-	-	-
Female.....	2	1	1	-	-	4	3	4	5	1	-	-
White, both sexes.....	3	1	3	1	2	3	3	6	7	1	-	-
Male.....	1	-	2	1	2	1	1	3	3	-	-	-
Female.....	2	1	1	-	-	2	2	3	4	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	1	3	4	1	1	-	-	-
Male.....	-	-	1	-	1	1	3	-	-	-	-	-
Female.....	-	-	-	-	-	2	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	22	20	19	36	45	83	121	109	97	40	13	-
Male.....	14	11	15	22	22	39	56	61	42	13	5	-
Female.....	8	9	4	14	23	44	65	48	55	27	8	-
White, both sexes.....	17	17	17	33	41	75	115	101	87	36	10	-
Male.....	11	9	13	21	21	36	54	57	40	11	4	-
Female.....	6	8	4	12	20	39	61	44	47	25	6	-
Black, both sexes.....	1	1	-	-	1	1	1	2	-	-	-	-
Male.....	-	-	-	-	-	1	-	2	-	-	-	-
Female.....	1	1	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	3	2	2	3	1	6	4	6	10	4	3	-
Male.....	2	2	2	1	-	1	2	2	2	2	1	-
Female.....	1	-	-	2	1	5	2	4	8	2	2	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	2	1	1	-	-	-	-	-
Male.....	1	-	-	-	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonia due to other or unspecified organisms (J16,J18)

Arkansas

All races, both sexes.....	20	23	32	28	43	70	142	133	115	48	13	-
Male.....	12	15	18	21	23	42	61	54	47	18	2	-
Female.....	8	8	14	7	20	28	81	79	68	30	11	-
White, both sexes.....	13	17	27	23	35	62	131	126	108	45	12	-
Male.....	8	12	15	16	20	36	56	50	46	17	2	-
Female.....	5	5	12	7	15	26	75	76	62	28	10	-
Black, both sexes.....	7	6	5	5	8	7	11	7	6	3	1	-
Male.....	4	3	3	5	3	6	5	4	1	1	-	-
Female.....	3	3	2	-	5	1	6	3	5	2	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	1	-	-	-

California

All races, both sexes.....	73	110	176	237	417	662	1,103	1,328	1,096	546	172	1
Male.....	41	66	92	140	223	343	528	596	446	172	45	1
Female.....	32	44	84	97	194	319	575	732	650	374	127	-
White, both sexes.....	56	91	136	179	321	521	918	1,120	927	457	137	1
Male.....	31	56	69	103	169	267	442	477	381	145	33	1
Female.....	25	35	67	76	152	254	476	643	546	312	104	-
Black, both sexes.....	12	13	24	28	44	53	66	66	42	29	14	-
Male.....	7	7	16	17	28	26	30	35	16	7	2	-
Female.....	5	6	8	11	16	27	36	31	26	22	12	-
American Indian 1/, both sexes.....	-	-	-	2	-	2	1	6	-	-	-	-
Male.....	-	-	-	1	-	1	1	3	-	-	-	-
Female.....	-	-	-	1	-	1	-	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	6	16	28	52	86	118	136	127	60	21	-
Male.....	3	3	7	19	26	49	55	81	49	20	10	-
Female.....	2	3	9	9	26	37	63	55	78	40	11	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

Delaware

All races, both sexes.....	2	2	3	3	12	15	12	31	20	5	2	-
Male.....	1	1	1	2	5	7	5	15	6	2	1	-
Female.....	1	1	2	1	7	8	7	16	14	3	1	-
White, both sexes.....	2	1	2	2	10	14	11	27	18	5	2	-
Male.....	1	-	1	1	4	6	4	12	5	2	1	-
Female.....	1	1	1	1	6	8	7	15	13	3	1	-
Black, both sexes.....	-	1	1	-	2	1	1	4	2	-	-	-
Male.....	-	1	-	-	1	1	1	3	1	-	-	-
Female.....	-	-	1	-	1	-	-	1	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Dist. of Columbia

All races, both sexes.....	3	2	1	4	3	5	6	17	16	11	2	-
Male.....	3	2	1	4	1	3	4	4	5	1	1	-
Female.....	-	-	-	-	2	2	2	13	11	10	1	-
White, both sexes.....	1	-	-	1	1	1	1	4	10	2	-	-
Male.....	1	-	-	1	-	1	1	2	3	-	-	-
Female.....	-	-	-	-	1	-	-	2	7	2	-	-
Black, both sexes.....	2	2	1	3	2	4	5	13	4	9	2	-
Male.....	2	2	1	3	1	2	3	2	1	1	1	-
Female.....	-	-	-	-	1	2	2	11	3	8	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	2	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Florida												
All races, both sexes.....	50	58	78	90	153	232	345	434	355	134	24	-
Male.....	34	35	44	56	84	129	182	187	125	48	7	-
Female.....	16	23	34	34	69	103	163	247	230	86	17	-
White, both sexes.....	42	45	57	72	129	210	314	405	333	127	20	-
Male.....	28	28	30	46	71	123	168	177	120	44	7	-
Female.....	14	17	27	26	58	87	146	228	213	83	13	-
Black, both sexes.....	7	13	19	16	24	22	29	28	18	6	4	-
Male.....	6	7	12	10	13	6	13	10	4	3	-	-
Female.....	1	6	7	6	11	16	16	18	14	3	4	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	1	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	1	2	-	-	1	1	4	1	-	-
Male.....	-	-	1	-	-	-	1	-	1	1	-	-
Female.....	1	-	-	2	-	-	-	1	3	-	-	-
Georgia												
All races, both sexes.....	26	44	45	73	102	155	220	262	204	111	18	-
Male.....	15	25	26	38	66	67	103	99	66	33	3	-
Female.....	11	19	19	35	36	88	117	163	138	78	15	-
White, both sexes.....	13	25	32	49	79	129	182	217	171	93	12	-
Male.....	8	16	17	25	49	56	88	81	57	27	1	-
Female.....	5	9	15	24	30	73	94	136	114	66	11	-
Black, both sexes.....	12	19	13	23	23	25	33	44	32	18	6	-
Male.....	6	9	9	13	17	10	11	17	8	6	2	-
Female.....	6	10	4	10	6	15	22	27	24	12	4	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	1	-	1	5	1	1	-	-	-
Male.....	1	-	-	-	-	1	4	1	1	-	-	-
Female.....	-	-	-	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

Hawaii

All races, both sexes.....	2	3	6	3	5	26	30	50	43	13	6	-
Male.....	1	3	3	2	3	18	22	30	22	6	5	-
Female.....	1	-	3	1	2	8	8	20	21	7	1	-
White, both sexes.....	-	2	2	1	2	5	2	14	11	2	2	-
Male.....	-	2	-	-	1	4	2	7	8	1	1	-
Female.....	-	-	2	1	1	1	-	7	3	1	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	2	1	4	2	3	21	28	36	32	11	4	-
Male.....	1	1	3	2	2	14	20	23	14	5	4	-
Female.....	1	-	1	-	1	7	8	13	18	6	-	-

Idaho

All races, both sexes.....	2	9	6	5	3	22	31	46	49	20	4	-
Male.....	2	4	4	4	-	7	16	20	10	9	-	-
Female.....	-	5	2	1	3	15	15	26	39	11	4	-
White, both sexes.....	2	9	6	5	3	22	29	46	49	20	4	-
Male.....	2	4	4	4	-	7	15	20	10	9	-	-
Female.....	-	5	2	1	3	15	14	26	39	11	4	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

Illinois

All races, both sexes.....	58	62	85	106	122	245	463	527	439	200	57	-
Male.....	28	41	54	58	59	123	226	206	151	53	9	-
Female.....	30	21	31	48	63	122	237	321	288	147	48	-
White, both sexes.....	45	43	63	81	95	197	394	481	396	193	51	-
Male.....	21	28	43	42	45	97	194	190	135	51	6	-
Female.....	24	15	20	39	50	100	200	291	261	142	45	-
Black, both sexes.....	11	17	21	21	26	44	60	41	35	5	4	-
Male.....	6	11	10	14	14	23	27	15	14	-	2	-
Female.....	5	6	11	7	12	21	33	26	21	5	2	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	2	1	3	1	4	9	5	7	2	2	-
Male.....	1	2	1	2	-	3	5	1	2	2	1	-
Female.....	1	-	-	1	1	1	4	4	5	-	1	-

Indiana

All races, both sexes.....	20	30	40	42	53	104	182	255	168	88	31	-
Male.....	13	16	22	26	24	51	77	109	58	21	4	-
Female.....	7	14	18	16	29	53	105	146	110	67	27	-
White, both sexes.....	18	25	37	40	46	97	172	250	160	85	31	-
Male.....	11	15	21	25	23	50	72	106	57	19	4	-
Female.....	7	10	16	15	23	47	100	144	103	66	27	-
Black, both sexes.....	2	5	3	2	7	6	10	5	8	3	-	-
Male.....	2	1	1	1	1	1	5	3	1	2	-	-
Female.....	-	4	2	1	6	5	5	2	7	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

Iowa

All races, both sexes.....	7	13	17	22	37	54	109	161	160	84	19	-
Male.....	6	4	9	10	16	26	59	61	55	14	2	-
Female.....	1	9	8	12	21	28	50	100	105	70	17	-
White, both sexes.....	7	11	17	22	35	54	107	161	159	84	19	-
Male.....	6	4	9	10	14	26	57	61	55	14	2	-
Female.....	1	7	8	12	21	28	50	100	104	70	17	-
Black, both sexes.....	-	2	-	-	1	-	2	-	1	-	-	-
Male.....	-	-	-	-	1	-	2	-	-	-	-	-
Female.....	-	2	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Kansas

All races, both sexes.....	9	11	9	17	33	59	113	139	146	70	16	-
Male.....	5	6	5	8	22	36	56	66	49	14	4	-
Female.....	4	5	4	9	11	23	57	73	97	56	12	-
White, both sexes.....	7	11	9	17	32	58	107	134	143	69	16	-
Male.....	3	6	5	8	21	35	54	64	48	14	4	-
Female.....	4	5	4	9	11	23	53	70	95	55	12	-
Black, both sexes.....	1	-	-	-	-	1	4	5	3	1	-	-
Male.....	1	-	-	-	-	1	1	2	1	-	-	-
Female.....	-	-	-	-	-	-	3	3	2	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	1	-	1	-	-	-	-	-
Male.....	1	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

Maine

All races, both sexes.....	2	4	6	8	12	21	45	52	51	15	6	-
Male.....	1	1	3	4	8	8	19	25	17	3	-	-
Female.....	1	3	3	4	4	13	26	27	34	12	6	-
White, both sexes.....	2	4	6	8	12	21	45	52	51	15	6	-
Male.....	1	1	3	4	8	8	19	25	17	3	-	-
Female.....	1	3	3	4	4	13	26	27	34	12	6	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	24	22	38	33	63	97	164	196	166	72	18	-
Male.....	17	9	23	18	40	55	81	82	65	18	4	-
Female.....	7	13	15	15	23	42	83	114	101	54	14	-
White, both sexes.....	11	12	28	25	48	70	136	159	152	64	15	-
Male.....	9	5	16	16	31	43	71	67	61	16	4	-
Female.....	2	7	12	9	17	27	65	92	91	48	11	-
Black, both sexes.....	10	9	10	8	14	27	26	33	11	8	3	-
Male.....	6	3	7	2	8	12	10	14	2	2	-	-
Female.....	4	6	3	6	6	15	16	19	9	6	3	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	1	-	-	1	-	2	4	3	-	-	-
Male.....	2	1	-	-	1	-	-	1	2	-	-	-
Female.....	1	-	-	-	-	-	2	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonia due to other or unspecified organisms (J16,J18)

Massachusetts

All races, both sexes.....	19	30	27	42	83	144	260	343	340	137	35	-
Male.....	8	17	18	23	54	85	121	147	115	32	5	-
Female.....	11	13	9	19	29	59	139	196	225	105	30	-
White, both sexes.....	19	26	26	39	79	134	252	337	331	133	34	-
Male.....	8	16	17	21	50	80	119	144	110	32	5	-
Female.....	11	10	9	18	29	54	133	193	221	101	29	-
Black, both sexes.....	-	3	-	2	3	7	3	5	3	2	-	-
Male.....	-	1	-	1	3	4	-	3	2	-	-	-
Female.....	-	2	-	1	-	3	3	2	1	2	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	-	1	3	5	1	6	2	1	-
Male.....	-	-	1	-	1	1	2	-	3	-	-	-
Female.....	-	1	-	-	-	2	3	1	3	2	1	-

Michigan

All races, both sexes.....	39	45	58	55	85	192	253	319	292	115	35	-
Male.....	21	26	39	27	45	94	127	138	123	35	8	-
Female.....	18	19	19	28	40	98	126	181	169	80	27	-
White, both sexes.....	24	28	51	35	67	171	217	289	268	109	31	-
Male.....	12	20	35	16	34	80	109	126	112	34	7	-
Female.....	12	8	16	19	33	91	108	163	156	75	24	-
Black, both sexes.....	15	16	7	20	16	19	35	29	23	6	4	-
Male.....	9	6	4	11	9	14	17	11	11	1	1	-
Female.....	6	10	3	9	7	5	18	18	12	5	3	-
American Indian 1/, both sexes.....	-	1	-	-	2	1	1	1	1	-	-	-
Male.....	-	-	-	-	2	-	1	1	-	-	-	-
Female.....	-	1	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
Minnesota												
All races, both sexes.....	11	8	14	12	24	53	89	128	124	61	21	-
Male.....	7	6	10	6	13	33	44	58	60	23	6	-
Female.....	4	2	4	6	11	20	45	70	64	38	15	-
White, both sexes.....	9	7	11	11	23	50	88	125	122	61	20	-
Male.....	5	5	7	5	12	31	43	57	58	23	6	-
Female.....	4	2	4	6	11	19	45	68	64	38	14	-
Black, both sexes.....	-	1	-	-	1	1	-	2	1	-	1	-
Male.....	-	1	-	-	1	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	1	-
American Indian 1/, both sexes.....	2	-	3	1	-	1	1	1	-	-	-	-
Male.....	2	-	3	1	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	1	-	-	-
Male.....	-	-	-	-	-	1	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes.....	22	19	14	28	39	67	83	106	81	25	12	-
Male.....	14	12	9	14	20	35	42	46	22	7	1	-
Female.....	8	7	5	14	19	32	41	60	59	18	11	-
White, both sexes.....	9	10	7	18	25	51	62	78	70	18	9	-
Male.....	5	7	4	6	13	24	30	35	16	7	-	-
Female.....	4	3	3	12	12	27	32	43	54	11	9	-
Black, both sexes.....	13	9	6	10	14	16	20	27	10	6	3	-
Male.....	9	5	4	8	7	11	12	10	5	-	1	-
Female.....	4	4	2	2	7	5	8	17	5	6	2	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	-	-	-	1	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

Nebraska

All races, both sexes.....	7	3	2	7	17	36	47	76	65	32	11	-
Male.....	4	3	2	4	9	22	27	27	18	7	4	-
Female.....	3	-	-	3	8	14	20	49	47	25	7	-
White, both sexes.....	6	2	1	7	16	33	47	75	65	32	11	-
Male.....	4	2	1	4	8	20	27	27	18	7	4	-
Female.....	2	-	-	3	8	13	20	48	47	25	7	-
Black, both sexes.....	1	1	1	-	1	2	-	1	-	-	-	-
Male.....	-	1	1	-	1	2	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	14	20	18	21	36	59	77	51	43	11	4	-
Male.....	9	10	11	9	27	30	30	25	17	5	2	-
Female.....	5	10	7	12	9	29	47	26	26	6	2	-
White, both sexes.....	13	17	16	20	32	54	72	47	41	10	2	-
Male.....	9	9	11	9	24	27	29	24	16	5	2	-
Female.....	4	8	5	11	8	27	43	23	25	5	-	-
Black, both sexes.....	1	1	1	1	1	1	3	2	1	-	2	-
Male.....	-	-	-	-	1	1	1	-	-	-	-	-
Female.....	1	1	1	1	-	-	2	2	1	-	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	1	-	3	3	2	2	1	1	-	-
Male.....	-	1	-	-	2	2	-	1	1	-	-	-
Female.....	-	1	1	-	1	1	2	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

New Hampshire

All races, both sexes.....	2	2	5	7	8	13	41	44	37	15	6	-
Male.....	1	2	4	2	3	3	15	22	12	2	1	-
Female.....	1	-	1	5	5	10	26	22	25	13	5	-
White, both sexes.....	2	2	5	7	8	13	41	44	36	15	6	-
Male.....	1	2	4	2	3	3	15	22	12	2	1	-
Female.....	1	-	1	5	5	10	26	22	24	13	5	-
Black, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	24	25	42	62	76	146	244	285	232	103	18	-
Male.....	15	14	26	31	38	88	124	122	78	27	2	-
Female.....	9	11	16	31	38	58	120	163	154	76	16	-
White, both sexes.....	16	15	35	49	65	134	216	269	210	92	17	-
Male.....	10	8	21	24	34	78	111	114	73	26	2	-
Female.....	6	7	14	25	31	56	105	155	137	66	15	-
Black, both sexes.....	8	10	5	13	8	10	23	14	21	9	1	-
Male.....	5	6	3	7	1	8	11	7	4	1	-	-
Female.....	3	4	2	6	7	2	12	7	17	8	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	-	3	2	5	2	1	2	-	-
Male.....	-	-	2	-	3	2	2	1	1	-	-	-
Female.....	-	-	-	-	-	-	3	1	-	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

New Mexico

All races, both sexes.....	5	5	10	14	14	37	47	53	36	24	5	-
Male.....	4	4	8	9	7	16	22	24	10	10	1	-
Female.....	1	1	2	5	7	21	25	29	26	14	4	-
White, both sexes.....	5	4	8	13	14	33	43	49	33	22	4	-
Male.....	4	3	6	8	7	16	20	23	9	9	1	-
Female.....	1	1	2	5	7	17	23	26	24	13	3	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	2	1	-	4	4	4	3	2	1	-
Male.....	-	1	2	1	-	-	2	1	1	1	-	-
Female.....	-	-	-	-	-	4	2	3	2	1	1	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	90	100	111	157	226	406	698	866	730	357	90	-
Male.....	65	56	61	86	119	217	338	381	264	102	14	-
Female.....	25	44	50	71	107	189	360	485	466	255	76	-
White, both sexes.....	59	73	76	114	158	321	589	735	664	314	73	-
Male.....	45	43	45	65	89	173	294	324	238	86	13	-
Female.....	14	30	31	49	69	148	295	411	426	228	60	-
Black, both sexes.....	30	23	31	37	55	73	89	102	48	30	14	-
Male.....	20	11	13	17	21	39	33	38	16	11	1	-
Female.....	10	12	18	20	34	34	56	64	32	19	13	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	1	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	3	4	6	13	12	19	29	18	13	3	-
Male.....	-	1	3	4	9	5	11	19	10	5	-	-
Female.....	1	2	1	2	4	7	8	10	8	8	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

Oregon

All races, both sexes.....	6	6	10	20	19	42	65	103	103	33	10	-
Male.....	4	2	4	11	10	25	33	39	50	11	3	-
Female.....	2	4	6	9	9	17	32	64	53	22	7	-
White, both sexes.....	6	5	10	19	19	42	62	101	101	31	10	-
Male.....	4	2	4	11	10	25	32	37	49	11	3	-
Female.....	2	3	6	8	9	17	30	64	52	20	7	-
Black, both sexes.....	-	-	-	-	-	-	-	-	1	1	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	1	-	1	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	2	2	1	1	-	-
Male.....	-	-	-	-	-	-	1	2	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	1	1	-	-

Pennsylvania

All races, both sexes.....	39	57	66	89	151	264	445	578	429	192	52	1
Male.....	20	35	37	52	91	145	212	258	157	65	15	1
Female.....	19	22	29	37	60	119	233	320	272	127	37	-
White, both sexes.....	32	42	58	79	136	235	411	546	410	182	49	1
Male.....	16	24	32	47	81	131	191	247	149	64	14	1
Female.....	16	18	26	32	55	104	220	299	261	118	35	-
Black, both sexes.....	7	15	7	9	14	27	33	27	17	9	3	-
Male.....	4	11	4	4	10	13	21	8	7	-	1	-
Female.....	3	4	3	5	4	14	12	19	10	9	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	1	1	2	1	5	2	1	-	-
Male.....	-	-	1	1	-	1	-	3	1	1	-	-
Female.....	-	-	-	-	1	1	1	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia due to other or unspecified organisms (J16,J18)												
South Dakota												
All races, both sexes.....	4	2	-	5	16	17	25	29	34	27	10	-
Male.....	4	1	-	3	7	6	9	11	13	7	3	-
Female.....	-	1	-	2	9	11	16	18	21	20	7	-
White, both sexes.....	3	1	-	5	14	15	24	29	33	26	10	-
Male.....	3	1	-	3	6	5	8	11	12	7	3	-
Female.....	-	-	-	2	8	10	16	18	21	19	7	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	-	-	2	2	1	-	1	1	-	-
Male.....	1	-	-	-	1	1	1	-	1	-	-	-
Female.....	-	1	-	-	1	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes.....	30	47	52	59	83	167	234	280	212	83	25	-
Male.....	17	32	30	37	40	76	109	109	82	17	8	-
Female.....	13	15	22	22	43	91	125	171	130	66	17	-
White, both sexes.....	23	45	46	50	74	149	211	261	198	76	21	-
Male.....	14	30	26	30	34	67	100	103	76	15	6	-
Female.....	9	15	20	20	40	82	111	158	122	61	15	-
Black, both sexes.....	7	2	5	8	9	15	23	18	13	7	4	-
Male.....	3	2	3	6	6	8	9	6	6	2	2	-
Female.....	4	-	2	2	3	7	14	12	7	5	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	1	-	3	-	1	1	-	-	-
Male.....	-	-	1	1	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	2	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

Vermont

All races, both sexes.....	-	1	-	3	3	8	9	15	9	9	4	-
Male.....	-	-	-	3	1	5	5	7	2	5	1	-
Female.....	-	1	-	-	2	3	4	8	7	4	3	-
White, both sexes.....	-	1	-	3	3	8	9	15	9	9	4	-
Male.....	-	-	-	3	1	5	5	7	2	5	1	-
Female.....	-	1	-	-	2	3	4	8	7	4	3	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	20	23	35	62	70	132	192	273	210	86	26	-
Male.....	13	15	26	37	39	65	102	119	77	19	6	-
Female.....	7	8	9	25	31	67	90	154	133	67	20	-
White, both sexes.....	10	15	22	45	57	103	173	248	187	76	22	-
Male.....	6	9	17	25	30	49	97	104	67	19	6	-
Female.....	4	6	5	20	27	54	76	144	120	57	16	-
Black, both sexes.....	10	8	13	15	13	24	18	21	19	9	4	-
Male.....	7	6	9	11	9	13	4	12	8	-	-	-
Female.....	3	2	4	4	4	11	14	9	11	9	4	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	-	5	1	4	4	1	-	-
Male.....	-	-	-	1	-	3	1	3	2	-	-	-
Female.....	-	-	-	1	-	2	-	1	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonia due to other or unspecified organisms (J16,J18)

Wisconsin

All races, both sexes.....	17	19	19	26	52	90	163	229	203	90	18	-
Male.....	12	12	11	13	27	43	79	96	81	23	3	-
Female.....	5	7	8	13	25	47	84	133	122	67	15	-
White, both sexes.....	16	18	18	25	48	88	162	223	200	89	17	-
Male.....	11	12	10	13	26	43	78	94	78	23	3	-
Female.....	5	6	8	12	22	45	84	129	122	66	14	-
Black, both sexes.....	1	1	1	1	1	2	1	5	2	1	1	-
Male.....	1	-	1	-	1	-	1	2	2	-	-	-
Female.....	-	1	-	1	-	2	-	3	-	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	2	-	-	-	-	-	-	-

Wyoming

All races, both sexes.....	4	2	4	6	2	8	17	28	20	12	-	-
Male.....	2	1	3	3	1	3	8	10	8	5	-	-
Female.....	2	1	1	3	1	5	9	18	12	7	-	-
White, both sexes.....	3	2	4	6	1	8	16	28	20	11	-	-
Male.....	1	1	3	3	-	3	8	10	8	4	-	-
Female.....	2	1	1	3	1	5	8	18	12	7	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	1	-	-	-	-	-	-	-
Male.....	1	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other acute lower respiratory infections (J20-J22,U04)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other acute lower respiratory infections (J20-J22,U04)

Colorado

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other acute lower respiratory infections (J20-J22,U04)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other acute lower respiratory infections (J20-J22,U04)

Oregon

All races, both sexes.....	-	-	-	-	1	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	1	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	2	-	-	-	2	3	2	5	1	-	-	-
Male.....	1	-	-	-	1	1	1	3	-	-	-	-
Female.....	1	-	-	-	1	2	1	2	1	-	-	-
White, both sexes.....	2	-	-	-	2	3	2	4	1	-	-	-
Male.....	1	-	-	-	1	1	1	3	-	-	-	-
Female.....	1	-	-	-	1	2	1	1	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute bronchitis and bronchiolitis (J20-J21)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute bronchitis and bronchiolitis (J20-J21)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute bronchitis and bronchiolitis (J20-J21)

Massachusetts

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute bronchitis and bronchiolitis (J20-J21)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Acute bronchitis and bronchiolitis (J20-J21)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified acute lower respiratory infections (J22,U04)

Alaska

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified acute lower respiratory infections (J22,U04)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other and unspecified acute lower respiratory infections (J22,U04)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified acute lower respiratory infections (J22,U04)

Rhode Island

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other and unspecified acute lower respiratory infections (J22,U04)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified acute lower respiratory infections (J22,U04)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other and unspecified acute lower respiratory infections (J22,U04)

Vermont

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other and unspecified acute lower respiratory infections (J22,U04)

Washington

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Alaska

Table with 13 columns for age groups and 13 rows for race/sex categories in Alaska. All cells contain dashes (-).

Arizona

Table with 13 columns for age groups and 13 rows for race/sex categories in Arizona. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Kentucky

Table with 13 columns representing age groups and one column for 'Age not stated'. Rows include 'All races, both sexes', 'Male', 'Female', 'White, both sexes', 'Male', 'Female', 'Black, both sexes', 'Male', 'Female', 'American Indian 1/, both sexes', 'Male', 'Female', 'Asian or Pacific Islander 2/, both sexes', 'Male', 'Female'.

Louisiana

Table with 13 columns representing age groups and one column for 'Age not stated'. Rows include 'All races, both sexes', 'Male', 'Female', 'White, both sexes', 'Male', 'Female', 'Black, both sexes', 'Male', 'Female', 'American Indian 1/, both sexes', 'Male', 'Female', 'Asian or Pacific Islander 2/, both sexes', 'Male', 'Female'.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Maine

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maine. All cells contain dashes (-).

Maryland

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maryland. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Massachusetts

Table with 13 columns (Age groups) and 13 rows (Race categories) for Massachusetts. All values are dashes (-).

Michigan

Table with 13 columns (Age groups) and 13 rows (Race categories) for Michigan. Values are dashes (-) or 1.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Nebraska

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified acute lower respiratory infections (J22,U04)

Oregon

All races, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	-	-	-	-	-	1	1	2	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	1	1	-	-	-	-
White, both sexes.....	-	-	-	-	-	1	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Texas

Table with 13 columns (Age groups) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Age groups) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Vermont

Table with 13 columns for age groups and one for 'Age not stated'. Rows include 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with 'Male' and 'Female' sub-rows.

Virginia

Table with 13 columns for age groups and one for 'Age not stated'. Rows include 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with 'Male' and 'Female' sub-rows.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified acute lower respiratory infections (J22,U04)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of upper respiratory tract (J30-J39)

Arkansas

All races, both sexes.....	2	1	-	-	-	-	-	1	-	-	-	-
Male.....	2	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes.....	1	-	-	-	-	-	-	1	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes.....	1	1	-	-	-	-	-	-	-	-	-	-
Male.....	1	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	2	2	1	1	1	-	7	8	2	-	-	-
Male.....	2	1	1	-	1	-	4	3	2	-	-	-
Female.....	-	1	-	1	-	-	3	5	-	-	-	-
White, both sexes.....	2	1	1	1	1	-	6	5	2	-	-	-
Male.....	2	-	1	-	1	-	3	2	2	-	-	-
Female.....	-	1	-	1	-	-	3	3	-	-	-	-
Black, both sexes.....	-	1	-	-	-	-	1	-	-	-	-	-
Male.....	-	1	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	3	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of upper respiratory tract (J30-J39)

Colorado

All races, both sexes.....	1	1	-	-	-	-	1	1	2	-	-	-
Male.....	1	1	-	-	-	-	1	-	2	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes.....	1	1	-	-	-	-	1	1	2	-	-	-
Male.....	1	1	-	-	-	-	1	-	2	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Connecticut

All races, both sexes.....	1	-	-	-	-	1	1	1	-	-	-	-
Male.....	1	-	-	-	-	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes.....	1	-	-	-	-	-	1	1	-	-	-	-
Male.....	1	-	-	-	-	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other diseases of upper respiratory tract (J30-J39)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
Arkansas												
All races, both sexes.....	1,656	4	-	1	1	2	-	2	1	2	15	25
Male.....	857	3	-	-	1	1	-	2	-	1	6	14
Female.....	799	1	-	1	-	1	-	-	1	1	9	11
White, both sexes.....	1,558	1	-	-	1	-	-	2	1	2	14	23
Male.....	813	-	-	-	1	-	-	2	-	1	6	14
Female.....	745	1	-	-	-	-	-	-	1	1	8	9
Black, both sexes.....	94	3	-	1	-	2	-	-	-	-	1	2
Male.....	42	3	-	-	-	1	-	-	-	-	-	-
Female.....	52	-	-	1	-	1	-	-	-	-	1	2
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	12,532	3	5	6	4	7	7	16	5	20	49	135
Male.....	5,876	2	4	4	4	5	5	7	2	10	23	62
Female.....	6,656	1	1	2	-	2	2	9	3	10	26	73
White, both sexes.....	11,104	3	1	6	2	6	5	9	1	12	35	101
Male.....	5,071	2	1	4	2	4	4	3	1	7	17	48
Female.....	6,033	1	-	2	-	2	1	6	-	5	18	53
Black, both sexes.....	650	-	2	-	2	-	2	6	3	6	10	25
Male.....	321	-	1	-	2	-	1	3	1	3	3	9
Female.....	329	-	1	-	-	-	1	3	2	3	7	16
American Indian 1/, both sexes.....	71	-	-	-	-	-	-	-	-	-	1	2
Male.....	37	-	-	-	-	-	-	-	-	-	1	1
Female.....	34	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	707	-	2	-	-	1	-	1	1	2	3	7
Male.....	447	-	2	-	-	1	-	1	-	-	2	4
Female.....	260	-	-	-	-	-	-	-	1	2	1	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic lower respiratory diseases (J40-J47)												
New Mexico												
All races, both sexes.....	884	-	-	-	1	1	-	-	3	2	4	7
Male.....	441	-	-	-	1	-	-	-	2	1	3	1
Female.....	443	-	-	-	-	1	-	-	1	1	1	6
White, both sexes.....	860	-	-	-	1	1	-	-	3	1	4	7
Male.....	428	-	-	-	1	-	-	-	2	-	3	1
Female.....	432	-	-	-	-	1	-	-	1	1	1	6
Black, both sexes.....	7	-	-	-	-	-	-	-	-	-	-	-
Male.....	6	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	14	-	-	-	-	-	-	-	-	1	-	-
Male.....	6	-	-	-	-	-	-	-	-	1	-	-
Female.....	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	6,561	1	2	4	3	2	11	13	11	13	33	75
Male.....	2,922	1	1	1	2	1	5	12	7	6	18	40
Female.....	3,639	-	1	3	1	1	6	1	4	7	15	35
White, both sexes.....	5,962	-	1	2	-	1	8	8	7	9	21	57
Male.....	2,621	-	-	1	-	-	4	7	6	3	12	33
Female.....	3,341	-	1	1	-	1	4	1	1	6	9	24
Black, both sexes.....	500	1	1	2	3	1	2	5	4	3	10	17
Male.....	243	1	1	-	2	1	1	5	1	2	5	7
Female.....	257	-	-	2	1	-	1	-	3	1	5	10
American Indian 1/, both sexes.....	13	-	-	-	-	-	-	-	-	-	1	1
Male.....	7	-	-	-	-	-	-	-	-	-	1	-
Female.....	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	86	-	-	-	-	-	1	-	-	1	1	-
Male.....	51	-	-	-	-	-	-	-	-	1	-	-
Female.....	35	-	-	-	-	-	1	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85-89	90-94	95-99	100 years and over	Age not stated
--	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	--------------------	----------------

Chronic lower respiratory diseases (J40-J47)

United States

All races, both sexes.....	2,756	4,641	8,136	11,975	16,689	22,758	25,283	19,724	9,810	2,936	387	6
Male.....	1,422	2,478	4,255	6,207	8,472	11,303	12,136	8,729	3,737	927	87	6
Female.....	1,334	2,163	3,881	5,768	8,217	11,455	13,147	10,995	6,073	2,009	300	-
White, both sexes.....	2,293	4,037	7,375	10,944	15,445	21,215	23,771	18,643	9,256	2,734	343	5
Male.....	1,209	2,161	3,795	5,624	7,775	10,419	11,316	8,195	3,474	846	74	5
Female.....	1,084	1,876	3,580	5,320	7,670	10,796	12,455	10,448	5,782	1,888	269	-
Black, both sexes.....	411	525	653	871	1,026	1,213	1,127	775	399	147	34	-
Male.....	189	275	397	490	571	686	610	359	180	57	10	-
Female.....	222	250	256	381	455	527	517	416	219	90	24	-
American Indian 1/, both sexes.....	31	41	59	85	88	89	96	50	22	13	1	-
Male.....	14	17	32	44	48	41	45	21	12	4	-	-
Female.....	17	24	27	41	40	48	51	29	10	9	1	-
Asian or Pacific Islander 2/, both sexes.....	21	38	49	75	130	241	289	256	133	42	9	1
Male.....	10	25	31	49	78	157	165	154	71	20	3	1
Female.....	11	13	18	26	52	84	124	102	62	22	6	-

Alabama

All races, both sexes.....	65	90	216	300	372	467	462	307	135	48	5	-
Male.....	39	45	126	167	199	240	218	143	50	19	1	-
Female.....	26	45	90	133	173	227	244	164	85	29	4	-
White, both sexes.....	56	76	191	273	327	423	425	282	122	39	3	-
Male.....	36	38	108	155	174	214	197	130	46	13	-	-
Female.....	20	38	83	118	153	209	228	152	76	26	3	-
Black, both sexes.....	9	14	24	27	45	43	37	23	13	8	2	-
Male.....	3	7	17	12	25	25	21	13	4	6	1	-
Female.....	6	7	7	15	20	18	16	10	9	2	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	1	-	2	-	-	-	-
Male.....	-	-	1	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Alaska

All races, both sexes.....	7	6	17	21	29	40	27	15	4	2	-	-
Male.....	3	1	10	10	14	16	17	8	1	1	-	-
Female.....	4	5	7	11	15	24	10	7	3	1	-	-
White, both sexes.....	5	4	13	15	20	35	24	10	4	2	-	-
Male.....	2	1	8	7	8	14	15	5	1	1	-	-
Female.....	3	3	5	8	12	21	9	5	3	1	-	-
Black, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	1	4	6	9	4	3	3	-	-	-	-
Male.....	1	-	2	3	6	1	2	1	-	-	-	-
Female.....	1	1	2	3	3	3	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	1	-	2	-	-	-	-
Male.....	-	-	-	-	-	1	-	2	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	49	84	172	236	381	458	548	436	205	64	6	-
Male.....	24	41	72	127	203	234	261	209	83	29	1	-
Female.....	25	43	100	109	178	224	287	227	122	35	5	-
White, both sexes.....	45	83	169	230	371	448	537	430	203	63	5	-
Male.....	23	41	71	123	199	228	258	207	82	28	1	-
Female.....	22	42	98	107	172	220	279	223	121	35	4	-
Black, both sexes.....	3	1	2	2	5	6	5	3	-	-	1	-
Male.....	1	-	1	1	1	2	1	-	-	-	-	-
Female.....	2	1	1	1	4	4	4	3	-	-	1	-
American Indian 1/, both sexes.....	1	-	1	4	3	3	5	2	1	1	-	-
Male.....	-	-	-	3	2	3	2	1	1	1	-	-
Female.....	1	-	1	1	1	-	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	2	1	1	1	1	-	-	-
Male.....	-	-	-	-	1	1	-	1	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Arkansas

All races, both sexes.....	45	65	129	197	222	324	271	210	94	38	8	-
Male.....	26	33	82	104	115	176	143	101	34	13	2	-
Female.....	19	32	47	93	107	148	128	109	60	25	6	-
White, both sexes.....	31	62	123	185	211	313	259	201	86	36	7	-
Male.....	19	33	77	100	109	172	136	98	30	13	2	-
Female.....	12	29	46	85	102	141	123	103	56	23	5	-
Black, both sexes.....	14	3	6	11	10	9	12	9	8	2	1	-
Male.....	7	-	5	4	5	3	7	3	4	-	-	-
Female.....	7	3	1	7	5	6	5	6	4	2	1	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	2	-	-	-	-	-	-
Male.....	-	-	-	-	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

California

All races, both sexes.....	269	414	693	1,006	1,472	2,160	2,555	2,193	1,140	326	47	-
Male.....	130	244	334	513	721	1,069	1,189	988	435	118	7	-
Female.....	139	170	359	493	751	1,091	1,366	1,205	705	208	40	-
White, both sexes.....	217	338	605	887	1,313	1,929	2,311	1,975	1,019	290	39	-
Male.....	108	201	284	446	631	924	1,040	866	368	105	5	-
Female.....	109	137	321	441	682	1,005	1,271	1,109	651	185	34	-
Black, both sexes.....	32	55	64	76	91	95	84	53	30	10	4	-
Male.....	14	30	34	36	45	55	50	16	16	1	1	-
Female.....	18	25	30	40	46	40	34	37	14	9	3	-
American Indian 1/, both sexes.....	6	7	5	7	15	6	8	8	4	2	-	-
Male.....	2	4	2	4	9	1	5	5	3	-	-	-
Female.....	4	3	3	3	6	5	3	3	1	2	-	-
Asian or Pacific Islander 2/, both sexes.....	14	14	19	36	53	130	152	157	87	24	4	-
Male.....	6	9	14	27	36	89	94	101	48	12	1	-
Female.....	8	5	5	9	17	41	58	56	39	12	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Florida

All races, both sexes.....	178	329	561	813	1,198	1,617	1,912	1,532	810	226	31	-
Male.....	103	188	309	412	626	812	908	644	292	71	9	-
Female.....	75	141	252	401	572	805	1,004	888	518	155	22	-
White, both sexes.....	155	291	518	756	1,135	1,546	1,842	1,491	781	216	30	-
Male.....	94	167	280	382	591	764	869	625	280	65	8	-
Female.....	61	124	238	374	544	782	973	866	501	151	22	-
Black, both sexes.....	21	35	40	56	56	66	64	37	28	9	1	-
Male.....	7	19	28	30	34	46	35	17	12	6	1	-
Female.....	14	16	12	26	22	20	29	20	16	3	-	-
American Indian 1/, both sexes.....	2	-	2	1	2	1	3	-	-	-	-	-
Male.....	2	-	1	-	-	-	2	-	-	-	-	-
Female.....	-	-	1	1	2	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	3	1	-	5	4	3	4	1	1	-	-
Male.....	-	2	-	-	1	2	2	2	-	-	-	-
Female.....	-	1	1	-	4	2	1	2	1	1	-	-

Georgia

All races, both sexes.....	90	158	259	363	497	553	641	452	200	70	5	-
Male.....	49	81	145	201	246	271	304	214	69	23	1	-
Female.....	41	77	114	162	251	282	337	238	131	47	4	-
White, both sexes.....	70	126	221	310	435	501	582	411	184	58	3	-
Male.....	37	63	124	165	209	237	268	196	62	17	-	-
Female.....	33	63	97	145	226	264	314	215	122	41	3	-
Black, both sexes.....	20	32	38	51	57	48	56	40	15	11	2	-
Male.....	12	18	21	35	33	31	35	18	6	5	1	-
Female.....	8	14	17	16	24	17	21	22	9	6	1	-
American Indian 1/, both sexes.....	-	-	-	-	2	2	1	-	-	-	-	-
Male.....	-	-	-	-	1	2	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	3	2	2	1	1	1	-	-
Male.....	-	-	-	1	3	1	1	-	1	1	-	-
Female.....	-	-	-	1	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Hawaii

All races, both sexes.....	5	13	21	22	30	44	78	39	16	11	2	1
Male.....	3	7	14	11	18	22	42	16	4	3	2	1
Female.....	2	6	7	11	12	22	36	23	12	8	-	-
White, both sexes.....	3	5	9	9	16	17	29	14	8	6	-	1
Male.....	2	2	5	3	10	7	17	4	2	1	-	1
Female.....	1	3	4	6	6	10	12	10	6	5	-	-
Black, both sexes.....	-	-	-	-	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	8	12	13	14	26	49	24	8	5	2	-
Male.....	1	5	9	8	8	15	25	12	2	2	2	-
Female.....	1	3	3	5	6	11	24	12	6	3	-	-

Idaho

All races, both sexes.....	19	21	38	59	90	117	151	95	50	14	2	-
Male.....	13	10	19	37	50	61	71	46	24	3	1	-
Female.....	6	11	19	22	40	56	80	49	26	11	1	-
White, both sexes.....	19	21	36	59	89	116	150	94	50	14	2	-
Male.....	13	10	19	37	50	61	70	46	24	3	1	-
Female.....	6	11	17	22	39	55	80	48	26	11	1	-
Black, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	1	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Illinois

All races, both sexes.....	102	157	279	394	605	897	924	756	400	104	12	-
Male.....	55	80	155	219	305	448	429	316	123	30	2	-
Female.....	47	77	124	175	300	449	495	440	277	74	10	-
White, both sexes.....	72	118	239	338	539	823	857	700	373	97	12	-
Male.....	40	61	128	185	272	407	390	293	114	29	2	-
Female.....	32	57	111	153	267	416	467	407	259	68	10	-
Black, both sexes.....	30	39	38	55	63	66	59	52	22	6	-	-
Male.....	15	19	25	33	32	36	32	20	8	1	-	-
Female.....	15	20	13	22	31	30	27	32	14	5	-	-
American Indian 1/, both sexes.....	-	-	-	1	1	-	2	-	-	-	-	-
Male.....	-	-	-	1	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	-	2	8	6	4	5	1	-	-
Male.....	-	-	2	-	1	5	6	3	1	-	-	-
Female.....	-	-	-	-	1	3	-	1	4	1	-	-

Indiana

All races, both sexes.....	74	128	217	331	460	548	642	463	198	52	16	-
Male.....	35	60	122	169	241	278	326	213	76	12	5	-
Female.....	39	68	95	162	219	270	316	250	122	40	11	-
White, both sexes.....	71	117	213	323	448	527	622	439	191	52	16	-
Male.....	34	53	120	165	236	268	313	202	72	12	5	-
Female.....	37	64	93	158	212	259	309	237	119	40	11	-
Black, both sexes.....	3	11	4	8	11	21	20	23	7	-	-	-
Male.....	1	7	2	4	5	10	13	11	4	-	-	-
Female.....	2	4	2	4	6	11	7	12	3	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Iowa

All races, both sexes.....	29	49	97	119	210	295	341	283	152	49	4	-
Male.....	18	28	52	53	116	158	180	156	67	16	-	-
Female.....	11	21	45	66	94	137	161	127	85	33	4	-
White, both sexes.....	29	48	95	118	206	291	340	283	152	49	4	-
Male.....	18	27	50	52	114	156	179	156	67	16	-	-
Female.....	11	21	45	66	92	135	161	127	85	33	4	-
Black, both sexes.....	-	1	2	1	4	4	1	-	-	-	-	-
Male.....	-	1	2	1	2	2	1	-	-	-	-	-
Female.....	-	-	-	-	2	2	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Kansas

All races, both sexes.....	24	52	78	127	187	273	327	221	115	39	3	-
Male.....	12	23	47	69	99	135	169	109	49	11	-	-
Female.....	12	29	31	58	88	138	158	112	66	28	3	-
White, both sexes.....	23	46	73	123	180	258	319	215	114	38	3	-
Male.....	12	20	43	66	96	125	166	106	48	11	-	-
Female.....	11	26	30	57	84	133	153	109	66	27	3	-
Black, both sexes.....	1	3	3	3	2	9	4	3	1	1	-	-
Male.....	-	2	3	3	1	7	2	1	1	-	-	-
Female.....	1	1	-	-	1	2	2	2	-	1	-	-
American Indian 1/, both sexes.....	-	3	2	1	3	6	4	3	-	-	-	-
Male.....	-	1	1	-	2	3	1	2	-	-	-	-
Female.....	-	2	1	1	1	3	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Maine

All races, both sexes.....	12	28	54	72	102	118	131	127	56	20	1	-
Male.....	2	18	23	43	51	60	58	52	18	7	-	-
Female.....	10	10	31	29	51	58	73	75	38	13	1	-
White, both sexes.....	12	28	53	67	101	118	131	127	55	20	1	-
Male.....	2	18	23	41	51	60	58	52	18	7	-	-
Female.....	10	10	30	26	50	58	73	75	37	13	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	4	1	-	-	-	1	-	-	-
Male.....	-	-	-	2	-	-	-	-	-	-	-	-
Female.....	-	-	1	2	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	32	58	98	141	222	342	404	333	173	46	4	-
Male.....	15	24	48	71	123	154	156	134	59	13	2	-
Female.....	17	34	50	70	99	188	248	199	114	33	2	-
White, both sexes.....	25	43	87	118	188	290	360	301	155	43	3	-
Male.....	12	18	38	57	106	130	135	123	52	12	2	-
Female.....	13	25	49	61	82	160	225	178	103	31	1	-
Black, both sexes.....	7	15	11	22	32	48	38	32	18	2	1	-
Male.....	3	6	10	13	16	21	16	11	7	1	-	-
Female.....	4	9	1	9	16	27	22	21	11	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	2	3	6	-	-	-	-	-
Male.....	-	-	-	1	1	3	5	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Massachusetts

All races, both sexes.....	25	79	119	204	272	367	491	431	235	70	10	-
Male.....	7	45	54	103	103	165	219	182	71	21	1	-
Female.....	18	34	65	101	169	202	272	249	164	49	9	-
White, both sexes.....	21	77	111	201	264	356	483	418	233	69	9	-
Male.....	7	45	49	101	101	159	217	174	70	20	1	-
Female.....	14	32	62	100	163	197	266	244	163	49	8	-
Black, both sexes.....	4	1	8	3	5	8	7	6	1	-	-	-
Male.....	-	-	5	2	-	4	2	4	-	-	-	-
Female.....	4	1	3	1	5	4	5	2	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	1	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	2	3	1	6	1	1	1	-
Male.....	-	-	-	-	1	2	-	4	1	1	-	-
Female.....	-	1	-	-	1	1	1	2	-	-	1	-

Michigan

All races, both sexes.....	109	171	274	407	583	836	960	703	359	106	9	-
Male.....	57	99	140	201	304	417	447	334	150	32	3	-
Female.....	52	72	134	206	279	419	513	369	209	74	6	-
White, both sexes.....	86	146	248	363	531	778	896	670	333	100	8	-
Male.....	42	87	123	177	269	389	412	320	138	31	3	-
Female.....	44	59	125	186	262	389	484	350	195	69	5	-
Black, both sexes.....	22	22	25	35	47	54	58	32	24	5	1	-
Male.....	14	11	16	20	32	26	33	14	11	1	-	-
Female.....	8	11	9	15	15	28	25	18	13	4	1	-
American Indian 1/, both sexes.....	1	2	-	7	3	4	3	1	1	1	-	-
Male.....	1	1	-	3	1	2	1	-	-	-	-	-
Female.....	-	1	-	4	2	2	2	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	2	2	-	3	-	1	-	-	-
Male.....	-	-	1	1	2	-	1	-	1	-	-	-
Female.....	-	1	-	1	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Missouri

All races, both sexes.....	81	113	214	314	415	526	559	462	228	70	13	1
Male.....	45	62	111	175	200	275	279	203	99	24	1	1
Female.....	36	51	103	139	215	251	280	259	129	46	12	-
White, both sexes.....	67	101	197	289	401	504	535	447	221	67	11	1
Male.....	37	59	101	161	192	264	270	198	96	22	-	1
Female.....	30	42	96	128	209	240	265	249	125	45	11	-
Black, both sexes.....	14	12	17	24	12	21	21	14	7	3	2	-
Male.....	8	3	10	13	8	10	7	5	3	2	1	-
Female.....	6	9	7	11	4	11	14	9	4	1	1	-
American Indian 1/, both sexes.....	-	-	-	1	1	1	1	-	-	-	-	-
Male.....	-	-	-	1	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	2	1	-	-	-	-
Male.....	-	-	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	1	-	-	-	-

Montana

All races, both sexes.....	12	15	33	67	95	117	115	83	45	13	2	-
Male.....	6	9	18	34	50	56	65	37	18	2	-	-
Female.....	6	6	15	33	45	61	50	46	27	11	2	-
White, both sexes.....	11	14	28	65	90	110	108	82	44	13	2	-
Male.....	5	9	16	33	48	51	63	37	18	2	-	-
Female.....	6	5	12	32	42	59	45	45	26	11	2	-
Black, both sexes.....	-	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	5	2	5	6	5	1	1	-	-	-
Male.....	1	-	2	1	2	4	2	-	-	-	-	-
Female.....	-	1	3	1	3	2	3	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Nebraska

All races, both sexes.....	22	19	43	89	101	169	181	156	81	33	4	1
Male.....	11	13	21	47	54	88	92	78	43	16	1	1
Female.....	11	6	22	42	47	81	89	78	38	17	3	-
White, both sexes.....	20	17	41	85	99	166	178	154	80	32	4	1
Male.....	10	11	20	45	53	87	91	77	43	15	1	1
Female.....	10	6	21	40	46	79	87	77	37	17	3	-
Black, both sexes.....	-	2	2	3	2	-	1	2	1	1	-	-
Male.....	-	2	1	2	1	-	-	1	-	1	-	-
Female.....	-	-	1	1	1	-	1	1	1	-	-	-
American Indian 1/, both sexes.....	2	-	-	1	-	1	2	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Nevada

All races, both sexes.....	18	29	70	115	144	204	199	154	78	13	-	-
Male.....	7	19	34	62	81	98	87	52	22	5	-	-
Female.....	11	10	36	53	63	106	112	102	56	8	-	-
White, both sexes.....	18	29	67	107	138	196	193	149	72	13	-	-
Male.....	7	19	32	57	75	92	84	50	20	5	-	-
Female.....	11	10	35	50	63	104	109	99	52	8	-	-
Black, both sexes.....	-	-	3	6	5	4	6	3	2	-	-	-
Male.....	-	-	2	4	5	2	3	2	1	-	-	-
Female.....	-	-	1	2	-	2	3	1	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	1	2	-	-	1	-	-	-
Male.....	-	-	-	1	1	2	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	2	-	2	3	-	-	-
Male.....	-	-	-	-	-	2	-	-	1	-	-	-
Female.....	-	-	-	1	-	-	-	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

New Hampshire

All races, both sexes.....	12	20	45	45	82	117	116	101	40	20	1	-
Male.....	6	8	22	25	32	64	48	45	11	6	-	-
Female.....	6	12	23	20	50	53	68	56	29	14	1	-
White, both sexes.....	12	20	45	45	82	117	116	101	40	20	1	-
Male.....	6	8	22	25	32	64	48	45	11	6	-	-
Female.....	6	12	23	20	50	53	68	56	29	14	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	36	104	149	247	317	503	624	566	288	83	12	-
Male.....	17	50	77	115	152	232	270	229	103	21	-	-
Female.....	19	54	72	132	165	271	354	337	185	62	12	-
White, both sexes.....	24	84	133	216	279	467	589	528	275	74	11	-
Male.....	15	43	66	95	134	213	254	209	95	20	-	-
Female.....	9	41	67	121	145	254	335	319	180	54	11	-
Black, both sexes.....	11	19	16	29	36	33	32	30	12	9	1	-
Male.....	2	6	11	20	17	17	16	15	7	1	-	-
Female.....	9	13	5	9	19	16	16	15	5	8	1	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	2	1	2	3	8	1	-	-	-
Male.....	-	1	-	-	-	2	-	5	1	-	-	-
Female.....	1	-	-	2	1	-	3	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)												
New Mexico												
All races, both sexes.....	19	32	44	73	108	172	189	128	72	25	4	-
Male.....	7	18	23	37	58	87	102	58	32	10	1	-
Female.....	12	14	21	36	50	85	87	70	40	15	3	-
White, both sexes.....	18	31	43	73	106	169	184	126	69	21	3	-
Male.....	7	18	22	37	56	85	99	58	30	8	1	-
Female.....	11	13	21	36	50	84	85	68	39	13	2	-
Black, both sexes.....	1	-	1	-	1	2	1	-	1	-	-	-
Male.....	-	-	1	-	1	2	1	-	1	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	3	1	2	4	1	-
Male.....	-	-	-	-	1	-	1	-	1	2	-	-
Female.....	-	-	-	-	-	1	2	1	1	2	1	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	1	-	-	-	-
New York												
All races, both sexes.....	133	201	362	511	746	1,162	1,320	1,097	623	206	32	-
Male.....	65	105	196	244	344	528	611	457	223	46	9	-
Female.....	68	96	166	267	402	634	709	640	400	160	23	-
White, both sexes.....	98	172	310	459	685	1,070	1,222	1,020	590	192	30	-
Male.....	51	93	163	218	312	482	562	419	205	42	8	-
Female.....	47	79	147	241	373	588	660	601	385	150	22	-
Black, both sexes.....	33	28	47	46	48	75	86	60	16	10	2	-
Male.....	13	11	29	22	23	38	42	29	7	2	1	-
Female.....	20	17	18	24	25	37	44	31	9	8	1	-
American Indian 1/, both sexes.....	1	-	2	-	2	3	2	1	-	-	-	-
Male.....	1	-	2	-	1	1	1	-	-	-	-	-
Female.....	-	-	-	-	1	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	3	6	11	14	10	16	17	4	-	-
Male.....	-	1	2	4	8	7	6	9	11	2	-	-
Female.....	1	-	1	2	3	7	4	7	6	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Ohio

All races, both sexes.....	162	251	400	625	830	1,192	1,324	949	433	131	17	-
Male.....	78	130	209	324	426	571	630	384	153	43	7	-
Female.....	84	121	191	301	404	621	694	565	280	88	10	-
White, both sexes.....	143	226	368	564	771	1,110	1,249	903	410	124	15	-
Male.....	69	117	185	293	387	531	590	369	144	40	6	-
Female.....	74	109	183	271	384	579	659	534	266	84	9	-
Black, both sexes.....	19	23	32	60	59	81	72	45	23	7	2	-
Male.....	9	13	24	31	39	39	38	14	9	3	1	-
Female.....	10	10	8	29	20	42	34	31	14	4	1	-
American Indian 1/, both sexes.....	-	1	-	1	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	1	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	1	1	1	-	-	-	-
Male.....	-	-	-	-	-	1	1	1	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-

Oklahoma

All races, both sexes.....	65	118	184	256	355	466	404	298	137	36	3	-
Male.....	27	65	84	141	184	235	210	152	56	14	-	-
Female.....	38	53	100	115	171	231	194	146	81	22	3	-
White, both sexes.....	53	109	166	237	330	435	380	284	128	32	3	-
Male.....	22	63	72	127	171	222	195	146	52	11	-	-
Female.....	31	46	94	110	159	213	185	138	76	21	3	-
Black, both sexes.....	5	1	9	7	11	11	8	5	4	3	-	-
Male.....	3	-	7	5	6	5	6	1	2	2	-	-
Female.....	2	1	2	2	5	6	2	4	2	1	-	-
American Indian 1/, both sexes.....	7	8	8	12	14	20	15	9	5	1	-	-
Male.....	2	2	5	9	7	8	9	5	2	1	-	-
Female.....	5	6	3	3	7	12	6	4	3	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Oregon

All races, both sexes.....	49	67	145	175	240	325	385	305	136	29	2	-
Male.....	34	39	71	86	117	166	182	139	60	7	-	-
Female.....	15	28	74	89	123	159	203	166	76	22	2	-
White, both sexes.....	47	65	137	171	235	318	373	302	133	29	2	-
Male.....	34	37	67	84	113	162	177	136	57	7	-	-
Female.....	13	28	70	87	122	156	196	166	76	22	2	-
Black, both sexes.....	1	-	2	1	3	3	1	1	1	-	-	-
Male.....	-	-	2	-	3	1	-	1	1	-	-	-
Female.....	1	-	-	1	-	2	1	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	5	3	2	3	8	-	2	-	-	-
Male.....	-	1	2	2	1	2	4	-	2	-	-	-
Female.....	1	-	3	1	1	1	4	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	-	-	1	3	2	-	-	-	-
Male.....	-	1	-	-	-	1	1	2	-	-	-	-
Female.....	-	-	1	-	-	-	2	-	-	-	-	-

Pennsylvania

All races, both sexes.....	103	182	332	481	716	1,061	1,308	1,049	545	158	21	1
Male.....	57	104	176	260	338	528	633	446	198	50	6	1
Female.....	46	78	156	221	378	533	675	603	347	108	15	-
White, both sexes.....	84	155	300	441	656	1,005	1,258	1,012	525	153	21	-
Male.....	49	87	159	239	311	498	602	430	191	49	6	-
Female.....	35	68	141	202	345	507	656	582	334	104	15	-
Black, both sexes.....	19	26	32	40	59	56	44	35	19	4	-	-
Male.....	8	17	17	21	27	30	26	14	7	1	-	-
Female.....	11	9	15	19	32	26	18	21	12	3	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	6	2	1	1	-	1
Male.....	-	-	-	-	-	-	5	2	-	-	-	1
Female.....	-	-	-	-	1	-	1	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Rhode Island

All races, both sexes.....	5	13	32	20	50	67	92	84	35	15	3	-
Male.....	2	6	17	8	28	27	47	33	13	3	-	-
Female.....	3	7	15	12	22	40	45	51	22	12	3	-
White, both sexes.....	5	13	32	19	49	63	90	83	34	15	3	-
Male.....	2	6	17	8	27	23	46	33	13	3	-	-
Female.....	3	7	15	11	22	40	44	50	21	12	3	-
Black, both sexes.....	-	-	-	-	1	4	1	-	1	-	-	-
Male.....	-	-	-	-	1	4	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

South Carolina

All races, both sexes.....	58	92	177	230	323	376	375	219	97	27	5	1
Male.....	26	48	100	118	172	203	193	97	36	8	1	1
Female.....	32	44	77	112	151	173	182	122	61	19	4	-
White, both sexes.....	42	72	152	200	284	336	336	196	89	21	4	1
Male.....	16	39	85	103	151	182	175	84	32	5	1	1
Female.....	26	33	67	97	133	154	161	112	57	16	3	-
Black, both sexes.....	16	20	25	28	39	40	38	23	8	6	1	-
Male.....	10	9	15	15	21	21	18	13	4	3	-	-
Female.....	6	11	10	13	18	19	20	10	4	3	1	-
American Indian 1/, both sexes.....	-	-	-	2	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

South Dakota

All races, both sexes.....	1	8	26	44	56	83	85	80	46	16	3	-
Male.....	-	6	13	21	33	47	54	46	28	7	1	-
Female.....	1	2	13	23	23	36	31	34	18	9	2	-
White, both sexes.....	-	8	24	41	55	81	81	77	45	16	3	-
Male.....	-	6	12	20	32	46	53	46	28	7	1	-
Female.....	-	2	12	21	23	35	28	31	17	9	2	-
Black, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	2	3	1	2	3	3	1	-	-	-
Male.....	-	-	1	1	1	1	-	-	-	-	-	-
Female.....	1	-	1	2	-	1	3	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	102	159	275	352	442	549	581	396	170	45	8	-
Male.....	58	75	147	193	204	277	302	168	70	13	3	-
Female.....	44	84	128	159	238	272	279	228	100	32	5	-
White, both sexes.....	80	144	252	328	412	515	544	372	156	42	6	-
Male.....	48	68	135	179	190	256	280	158	62	11	3	-
Female.....	32	76	117	149	222	259	264	214	94	31	3	-
Black, both sexes.....	22	14	23	21	29	33	37	23	14	3	2	-
Male.....	10	6	12	12	14	21	22	9	8	2	-	-
Female.....	12	8	11	9	15	12	15	14	6	1	2	-
American Indian 1/, both sexes.....	-	1	-	1	-	-	-	-	-	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	1	1	-	1	-	-	-	-
Male.....	-	-	-	1	-	-	-	1	-	-	-	-
Female.....	-	-	-	1	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Texas

All races, both sexes.....	179	300	572	844	1,178	1,500	1,487	1,144	514	159	16	1
Male.....	96	166	289	445	615	768	753	534	218	52	2	1
Female.....	83	134	283	399	563	732	734	610	296	107	14	-
White, both sexes.....	150	270	523	791	1,090	1,380	1,402	1,082	490	144	15	1
Male.....	78	151	262	416	562	702	710	492	204	46	2	1
Female.....	72	119	261	375	528	678	692	590	286	98	13	-
Black, both sexes.....	27	27	45	48	80	105	68	57	22	13	1	-
Male.....	17	14	26	26	48	62	38	39	13	6	-	-
Female.....	10	13	19	22	32	43	30	18	9	7	1	-
American Indian 1/, both sexes.....	1	2	-	2	2	6	6	1	-	-	-	-
Male.....	-	-	-	-	2	-	2	-	-	-	-	-
Female.....	1	2	-	2	-	6	4	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	4	3	6	9	11	4	2	2	-	-
Male.....	1	1	1	3	3	4	3	3	1	-	-	-
Female.....	-	-	3	-	3	5	8	1	1	2	-	-

Utah

All races, both sexes.....	13	40	40	57	107	107	114	78	31	6	1	-
Male.....	7	16	24	35	67	60	68	39	16	2	-	-
Female.....	6	24	16	22	40	47	46	39	15	4	1	-
White, both sexes.....	13	40	38	53	104	105	112	77	30	5	1	-
Male.....	7	16	24	33	65	58	66	39	15	1	-	-
Female.....	6	24	14	20	39	47	46	38	15	4	1	-
Black, both sexes.....	-	-	-	1	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	3	-	1	1	-	-	-	-	-
Male.....	-	-	-	2	-	1	1	-	-	-	-	-
Female.....	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	3	1	-	1	1	1	-	-
Male.....	-	-	-	-	2	1	-	-	1	1	-	-
Female.....	-	-	1	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Vermont

All races, both sexes.....	7	8	21	29	48	50	56	56	28	7	1	-
Male.....	3	4	12	9	24	21	32	28	11	-	-	-
Female.....	4	4	9	20	24	29	24	28	17	7	1	-
White, both sexes.....	7	8	21	29	48	49	56	56	28	7	1	-
Male.....	3	4	12	9	24	20	32	28	11	-	-	-
Female.....	4	4	9	20	24	29	24	28	17	7	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	49	91	174	274	350	547	549	422	199	52	7	-
Male.....	25	43	80	132	176	264	232	167	65	15	-	-
Female.....	24	48	94	142	174	283	317	255	134	37	7	-
White, both sexes.....	39	66	149	239	315	492	493	396	178	43	6	-
Male.....	21	29	67	112	160	224	205	155	59	12	-	-
Female.....	18	37	82	127	155	268	288	241	119	31	6	-
Black, both sexes.....	10	24	23	34	32	50	53	25	21	9	1	-
Male.....	4	13	13	19	15	36	27	11	6	3	-	-
Female.....	6	11	10	15	17	14	26	14	15	6	1	-
American Indian 1/, both sexes.....	-	-	1	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	2	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	1	3	3	3	1	-	-	-	-
Male.....	-	1	-	1	1	2	-	1	-	-	-	-
Female.....	-	-	1	-	2	1	3	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Washington

All races, both sexes.....	54	108	173	271	407	473	480	435	188	58	5	-
Male.....	26	52	100	135	205	230	219	184	76	20	3	-
Female.....	28	56	73	136	202	243	261	251	112	38	2	-
White, both sexes.....	50	95	161	257	387	446	465	424	186	57	4	-
Male.....	24	43	93	126	193	211	212	180	75	19	3	-
Female.....	26	52	68	131	194	235	253	244	111	38	1	-
Black, both sexes.....	2	7	6	8	5	8	3	-	1	-	-	-
Male.....	-	3	3	4	2	5	2	-	1	-	-	-
Female.....	2	4	3	4	3	3	1	-	-	-	-	-
American Indian 1/, both sexes.....	2	3	6	4	5	4	1	4	-	-	-	-
Male.....	2	3	4	3	3	3	-	2	-	-	-	-
Female.....	-	-	2	1	2	1	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	3	-	2	10	15	11	7	1	1	1	-
Male.....	-	3	-	2	7	11	5	2	-	1	-	-
Female.....	-	-	-	-	3	4	6	5	1	-	1	-

West Virginia

All races, both sexes.....	29	63	98	150	190	229	262	177	87	17	2	-
Male.....	17	35	54	81	87	106	132	85	38	6	-	-
Female.....	12	28	44	69	103	123	130	92	49	11	2	-
White, both sexes.....	28	61	97	150	187	227	260	175	86	16	2	-
Male.....	16	35	53	81	87	104	131	83	37	6	-	-
Female.....	12	26	44	69	100	123	129	92	49	10	2	-
Black, both sexes.....	1	2	1	-	2	2	2	2	1	1	-	-
Male.....	1	-	1	-	-	2	1	2	1	-	-	-
Female.....	-	2	-	-	2	-	1	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Arkansas

All races, both sexes.....	1,648	4	-	1	1	2	-	2	1	2	15	25
Male.....	855	3	-	-	1	1	-	2	-	1	6	14
Female.....	793	1	-	1	-	1	-	-	1	1	9	11
White, both sexes.....	1,550	1	-	-	1	-	-	2	1	2	14	23
Male.....	811	-	-	-	1	-	-	2	-	1	6	14
Female.....	739	1	-	-	-	-	-	-	1	1	8	9
Black, both sexes.....	94	3	-	1	-	2	-	-	-	-	1	2
Male.....	42	3	-	-	-	1	-	-	-	-	-	-
Female.....	52	-	-	1	-	1	-	-	-	-	1	2
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	12,419	3	5	6	4	7	7	16	5	20	49	133
Male.....	5,842	2	4	4	4	5	5	7	2	10	23	62
Female.....	6,577	1	1	2	-	2	2	9	3	10	26	71
White, both sexes.....	11,013	3	1	6	2	6	5	9	1	12	35	99
Male.....	5,044	2	1	4	2	4	4	3	1	7	17	48
Female.....	5,969	1	-	2	-	2	1	6	-	5	18	51
Black, both sexes.....	645	-	2	-	2	-	2	6	3	6	10	25
Male.....	320	-	1	-	2	-	1	3	1	3	3	9
Female.....	325	-	1	-	-	-	1	3	2	3	7	16
American Indian 1/, both sexes.....	71	-	-	-	-	-	-	-	-	-	1	2
Male.....	37	-	-	-	-	-	-	-	-	-	1	1
Female.....	34	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	690	-	2	-	-	1	-	1	1	2	3	7
Male.....	441	-	2	-	-	1	-	1	-	-	2	4
Female.....	249	-	-	-	-	-	-	-	1	2	1	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
New Mexico												
All races, both sexes.....	881	-	-	-	1	1	-	-	3	2	4	7
Male.....	441	-	-	-	1	-	-	-	2	1	3	1
Female.....	440	-	-	-	-	1	-	-	1	1	1	6
White, both sexes.....	857	-	-	-	1	1	-	-	3	1	4	7
Male.....	428	-	-	-	1	-	-	-	2	-	3	1
Female.....	429	-	-	-	-	1	-	-	1	1	1	6
Black, both sexes.....	7	-	-	-	-	-	-	-	-	-	-	-
Male.....	6	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	14	-	-	-	-	-	-	-	-	1	-	-
Male.....	6	-	-	-	-	-	-	-	-	1	-	-
Female.....	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	6,521	1	2	4	3	2	11	12	11	13	33	74
Male.....	2,911	1	1	1	2	1	5	11	7	6	18	40
Female.....	3,610	-	1	3	1	1	6	1	4	7	15	34
White, both sexes.....	5,924	-	1	2	-	1	8	7	7	9	21	56
Male.....	2,610	-	-	1	-	-	4	6	6	3	12	33
Female.....	3,314	-	1	1	-	1	4	1	1	6	9	23
Black, both sexes.....	499	1	1	2	3	1	2	5	4	3	10	17
Male.....	243	1	1	-	2	1	1	5	1	2	5	7
Female.....	256	-	-	2	1	-	1	-	3	1	5	10
American Indian 1/, both sexes.....	13	-	-	-	-	-	-	-	-	-	1	1
Male.....	7	-	-	-	-	-	-	-	-	-	1	-
Female.....	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	85	-	-	-	-	-	1	-	-	1	1	-
Male.....	51	-	-	-	-	-	-	-	-	1	-	-
Female.....	34	-	-	-	-	-	1	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

United States

All races, both sexes.....	2,746	4,623	8,111	11,910	16,602	22,622	25,063	19,544	9,700	2,913	387	6
Male.....	1,418	2,472	4,246	6,183	8,453	11,261	12,085	8,697	3,714	921	87	6
Female.....	1,328	2,151	3,865	5,727	8,149	11,361	12,978	10,847	5,986	1,992	300	-
White, both sexes.....	2,286	4,022	7,351	10,888	15,364	21,085	23,562	18,469	9,149	2,712	343	5
Male.....	1,205	2,155	3,786	5,606	7,758	10,380	11,267	8,164	3,453	840	74	5
Female.....	1,081	1,867	3,565	5,282	7,606	10,705	12,295	10,305	5,696	1,872	269	-
Black, both sexes.....	409	522	653	868	1,024	1,213	1,126	774	396	146	34	-
Male.....	189	275	397	488	570	686	610	359	178	57	10	-
Female.....	220	247	256	380	454	527	516	415	218	89	24	-
American Indian 1/, both sexes.....	31	41	59	84	87	89	95	50	22	13	1	-
Male.....	14	17	32	43	48	41	45	21	12	4	-	-
Female.....	17	24	27	41	39	48	50	29	10	9	1	-
Asian or Pacific Islander 2/, both sexes.....	20	38	48	70	127	235	280	251	133	42	9	1
Male.....	10	25	31	46	77	154	163	153	71	20	3	1
Female.....	10	13	17	24	50	81	117	98	62	22	6	-

Alabama

All races, both sexes.....	65	90	216	300	369	464	459	305	131	47	5	-
Male.....	39	45	126	167	198	238	217	143	49	18	1	-
Female.....	26	45	90	133	171	226	242	162	82	29	4	-
White, both sexes.....	56	76	191	273	324	420	422	280	118	38	3	-
Male.....	36	38	108	155	173	212	196	130	45	12	-	-
Female.....	20	38	83	118	151	208	226	150	73	26	3	-
Black, both sexes.....	9	14	24	27	45	43	37	23	13	8	2	-
Male.....	3	7	17	12	25	25	21	13	4	6	1	-
Female.....	6	7	7	15	20	18	16	10	9	2	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	1	-	2	-	-	-	-
Male.....	-	-	1	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Alaska

All races, both sexes.....	7	5	17	21	28	40	27	15	4	2	-	-
Male.....	3	-	10	10	14	16	17	8	1	1	-	-
Female.....	4	5	7	11	14	24	10	7	3	1	-	-
White, both sexes.....	5	3	13	15	20	35	24	10	4	2	-	-
Male.....	2	-	8	7	8	14	15	5	1	1	-	-
Female.....	3	3	5	8	12	21	9	5	3	1	-	-
Black, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	1	4	6	8	4	3	3	-	-	-	-
Male.....	1	-	2	3	6	1	2	1	-	-	-	-
Female.....	1	1	2	3	2	3	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	1	-	2	-	-	-	-
Male.....	-	-	-	-	-	1	-	2	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	49	83	171	235	378	455	543	430	205	64	6	-
Male.....	24	40	72	126	203	233	260	209	83	29	1	-
Female.....	25	43	99	109	175	222	283	221	122	35	5	-
White, both sexes.....	45	82	168	230	368	445	532	424	203	63	5	-
Male.....	23	40	71	123	199	227	257	207	82	28	1	-
Female.....	22	42	97	107	169	218	275	217	121	35	4	-
Black, both sexes.....	3	1	2	2	5	6	5	3	-	-	1	-
Male.....	1	-	1	1	1	2	1	-	-	-	-	-
Female.....	2	1	1	1	4	4	4	3	-	-	1	-
American Indian 1/, both sexes.....	1	-	1	3	3	3	5	2	1	1	-	-
Male.....	-	-	-	2	2	3	2	1	1	1	-	-
Female.....	1	-	1	1	1	-	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	2	1	1	1	1	-	-	-
Male.....	-	-	-	-	1	1	-	1	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Arkansas

All races, both sexes.....	45	64	129	196	222	324	269	208	92	38	8	-
Male.....	26	32	82	103	115	176	143	101	34	13	2	-
Female.....	19	32	47	93	107	148	126	107	58	25	6	-
White, both sexes.....	31	61	123	184	211	313	257	199	84	36	7	-
Male.....	19	32	77	99	109	172	136	98	30	13	2	-
Female.....	12	29	46	85	102	141	121	101	54	23	5	-
Black, both sexes.....	14	3	6	11	10	9	12	9	8	2	1	-
Male.....	7	-	5	4	5	3	7	3	4	-	-	-
Female.....	7	3	1	7	5	6	5	6	4	2	1	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	2	-	-	-	-	-	-
Male.....	-	-	-	-	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

California

All races, both sexes.....	266	413	691	995	1,464	2,144	2,531	2,162	1,128	323	47	-
Male.....	128	244	333	506	720	1,063	1,184	980	432	117	7	-
Female.....	138	169	358	489	744	1,081	1,347	1,182	696	206	40	-
White, both sexes.....	215	338	603	880	1,308	1,917	2,290	1,948	1,009	287	39	-
Male.....	106	201	283	441	631	920	1,035	859	366	104	5	-
Female.....	109	137	320	439	677	997	1,255	1,089	643	183	34	-
Black, both sexes.....	32	54	64	75	90	95	84	53	28	10	4	-
Male.....	14	30	34	36	45	55	50	16	15	1	1	-
Female.....	18	24	30	39	45	40	34	37	13	9	3	-
American Indian 1/, both sexes.....	6	7	5	7	15	6	8	8	4	2	-	-
Male.....	2	4	2	4	9	1	5	5	3	-	-	-
Female.....	4	3	3	3	6	5	3	3	1	2	-	-
Asian or Pacific Islander 2/, both sexes.....	13	14	19	33	51	126	149	153	87	24	4	-
Male.....	6	9	14	25	35	87	94	100	48	12	1	-
Female.....	7	5	5	8	16	39	55	53	39	12	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Florida

All races, both sexes.....	178	329	560	805	1,193	1,607	1,892	1,514	801	226	31	-
Male.....	103	188	309	410	624	809	902	642	290	71	9	-
Female.....	75	141	251	395	569	798	990	872	511	155	22	-
White, both sexes.....	155	291	517	748	1,130	1,536	1,822	1,473	772	216	30	-
Male.....	94	167	280	380	589	761	863	623	278	65	8	-
Female.....	61	124	237	368	541	775	959	850	494	151	22	-
Black, both sexes.....	21	35	40	56	56	66	64	37	28	9	1	-
Male.....	7	19	28	30	34	46	35	17	12	6	1	-
Female.....	14	16	12	26	22	20	29	20	16	3	-	-
American Indian 1/, both sexes.....	2	-	2	1	2	1	3	-	-	-	-	-
Male.....	2	-	1	-	-	-	2	-	-	-	-	-
Female.....	-	-	1	1	2	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	3	1	-	5	4	3	4	1	1	-	-
Male.....	-	2	-	-	1	2	2	2	-	-	-	-
Female.....	-	1	1	-	4	2	1	2	1	1	-	-

Georgia

All races, both sexes.....	89	156	256	363	493	551	638	448	197	70	5	-
Male.....	48	80	144	201	246	270	304	214	69	23	1	-
Female.....	41	76	112	162	247	281	334	234	128	47	4	-
White, both sexes.....	69	124	218	310	431	499	579	407	181	58	3	-
Male.....	36	62	123	165	209	236	268	196	62	17	-	-
Female.....	33	62	95	145	222	263	311	211	119	41	3	-
Black, both sexes.....	20	32	38	51	57	48	56	40	15	11	2	-
Male.....	12	18	21	35	33	31	35	18	6	5	1	-
Female.....	8	14	17	16	24	17	21	22	9	6	1	-
American Indian 1/, both sexes.....	-	-	-	-	2	2	1	-	-	-	-	-
Male.....	-	-	-	-	1	2	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	3	2	2	1	1	1	-	-
Male.....	-	-	-	1	3	1	1	-	1	1	-	-
Female.....	-	-	-	1	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Hawaii

All races, both sexes.....	5	13	21	21	29	44	74	39	16	11	2	1
Male.....	3	7	14	10	18	22	41	16	4	3	2	1
Female.....	2	6	7	11	11	22	33	23	12	8	-	-
White, both sexes.....	3	5	9	9	16	17	29	14	8	6	-	1
Male.....	2	2	5	3	10	7	17	4	2	1	-	1
Female.....	1	3	4	6	6	10	12	10	6	5	-	-
Black, both sexes.....	-	-	-	-	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	2	8	12	12	13	26	45	24	8	5	2	-
Male.....	1	5	9	7	8	15	24	12	2	2	2	-
Female.....	1	3	3	5	5	11	21	12	6	3	-	-

Idaho

All races, both sexes.....	19	21	38	59	90	115	149	93	50	14	2	-
Male.....	13	10	19	37	50	60	71	45	24	3	1	-
Female.....	6	11	19	22	40	55	78	48	26	11	1	-
White, both sexes.....	19	21	36	59	89	114	148	92	50	14	2	-
Male.....	13	10	19	37	50	60	70	45	24	3	1	-
Female.....	6	11	17	22	39	54	78	47	26	11	1	-
Black, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	1	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	1	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Illinois

All races, both sexes.....	102	156	276	393	603	892	914	745	395	104	12	-
Male.....	55	80	152	219	305	446	428	314	121	30	2	-
Female.....	47	76	124	174	298	446	486	431	274	74	10	-
White, both sexes.....	72	117	236	337	537	819	847	689	368	97	12	-
Male.....	40	61	125	185	272	406	389	291	112	29	2	-
Female.....	32	56	111	152	265	413	458	398	256	68	10	-
Black, both sexes.....	30	39	38	55	63	66	59	52	22	6	-	-
Male.....	15	19	25	33	32	36	32	20	8	1	-	-
Female.....	15	20	13	22	31	30	27	32	14	5	-	-
American Indian 1/, both sexes.....	-	-	-	1	1	-	2	-	-	-	-	-
Male.....	-	-	-	1	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	-	2	7	6	4	5	1	-	-
Male.....	-	-	2	-	1	4	6	3	1	-	-	-
Female.....	-	-	-	-	1	3	-	1	4	1	-	-

Indiana

All races, both sexes.....	74	128	217	331	456	545	638	455	196	52	16	-
Male.....	35	60	122	169	239	277	325	210	76	12	5	-
Female.....	39	68	95	162	217	268	313	245	120	40	11	-
White, both sexes.....	71	117	213	323	444	524	618	432	189	52	16	-
Male.....	34	53	120	165	234	267	312	199	72	12	5	-
Female.....	37	64	93	158	210	257	306	233	117	40	11	-
Black, both sexes.....	3	11	4	8	11	21	20	22	7	-	-	-
Male.....	1	7	2	4	5	10	13	11	4	-	-	-
Female.....	2	4	2	4	6	11	7	11	3	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Iowa

All races, both sexes.....	29	49	96	117	208	294	336	283	150	49	4	-
Male.....	18	28	52	52	116	157	179	156	67	16	-	-
Female.....	11	21	44	65	92	137	157	127	83	33	4	-
White, both sexes.....	29	48	94	116	204	290	335	283	150	49	4	-
Male.....	18	27	50	51	114	155	178	156	67	16	-	-
Female.....	11	21	44	65	90	135	157	127	83	33	4	-
Black, both sexes.....	-	1	2	1	4	4	1	-	-	-	-	-
Male.....	-	1	2	1	2	2	1	-	-	-	-	-
Female.....	-	-	-	-	2	2	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Kansas

All races, both sexes.....	24	52	78	127	185	272	324	218	115	39	3	-
Male.....	12	23	47	69	98	134	169	109	49	11	-	-
Female.....	12	29	31	58	87	138	155	109	66	28	3	-
White, both sexes.....	23	46	73	123	178	257	316	212	114	38	3	-
Male.....	12	20	43	66	95	124	166	106	48	11	-	-
Female.....	11	26	30	57	83	133	150	106	66	27	3	-
Black, both sexes.....	1	3	3	3	2	9	4	3	1	1	-	-
Male.....	-	2	3	3	1	7	2	1	1	-	-	-
Female.....	1	1	-	-	1	2	2	2	-	1	-	-
American Indian 1/, both sexes.....	-	3	2	1	3	6	4	3	-	-	-	-
Male.....	-	1	1	-	2	3	1	2	-	-	-	-
Female.....	-	2	1	1	1	3	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Maine

All races, both sexes.....	11	28	54	72	101	118	131	126	56	20	1	-
Male.....	2	18	23	43	50	60	58	52	18	7	-	-
Female.....	9	10	31	29	51	58	73	74	38	13	1	-
White, both sexes.....	11	28	53	67	100	118	131	126	55	20	1	-
Male.....	2	18	23	41	50	60	58	52	18	7	-	-
Female.....	9	10	30	26	50	58	73	74	37	13	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	4	1	-	-	-	1	-	-	-
Male.....	-	-	-	2	-	-	-	-	-	-	-	-
Female.....	-	-	1	2	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	32	58	98	140	220	340	402	331	171	43	4	-
Male.....	15	24	48	71	121	153	155	133	58	13	2	-
Female.....	17	34	50	69	99	187	247	198	113	30	2	-
White, both sexes.....	25	43	87	117	187	288	358	299	153	40	3	-
Male.....	12	18	38	57	105	129	134	122	51	12	2	-
Female.....	13	25	49	60	82	159	224	177	102	28	1	-
Black, both sexes.....	7	15	11	22	31	48	38	32	18	2	1	-
Male.....	3	6	10	13	15	21	16	11	7	1	-	-
Female.....	4	9	1	9	16	27	22	21	11	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	2	3	6	-	-	-	-	-
Male.....	-	-	-	1	1	3	5	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Massachusetts

All races, both sexes.....	24	79	118	204	267	366	486	424	231	70	10	-
Male.....	7	45	54	103	103	165	217	182	70	21	1	-
Female.....	17	34	64	101	164	201	269	242	161	49	9	-
White, both sexes.....	21	77	110	201	259	355	478	412	229	69	9	-
Male.....	7	45	49	101	101	159	215	174	69	20	1	-
Female.....	14	32	61	100	158	196	263	238	160	49	8	-
Black, both sexes.....	3	1	8	3	5	8	7	6	1	-	-	-
Male.....	-	-	5	2	-	4	2	4	-	-	-	-
Female.....	3	1	3	1	5	4	5	2	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	1	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	2	3	1	5	1	1	1	-
Male.....	-	-	-	-	1	2	-	4	1	1	-	-
Female.....	-	1	-	-	1	1	1	1	-	-	1	-

Michigan

All races, both sexes.....	109	171	273	406	579	831	955	700	356	106	9	-
Male.....	57	99	139	201	303	416	445	333	149	32	3	-
Female.....	52	72	134	205	276	415	510	367	207	74	6	-
White, both sexes.....	86	146	247	362	527	773	891	667	330	100	8	-
Male.....	42	87	122	177	268	388	410	319	137	31	3	-
Female.....	44	59	125	185	259	385	481	348	193	69	5	-
Black, both sexes.....	22	22	25	35	47	54	58	32	24	5	1	-
Male.....	14	11	16	20	32	26	33	14	11	1	-	-
Female.....	8	11	9	15	15	28	25	18	13	4	1	-
American Indian 1/, both sexes.....	1	2	-	7	3	4	3	1	1	1	-	-
Male.....	1	1	-	3	1	2	1	-	-	-	-	-
Female.....	-	1	-	4	2	2	2	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	2	2	-	3	-	1	-	-	-
Male.....	-	-	1	1	2	-	1	-	1	-	-	-
Female.....	-	1	-	1	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Missouri

All races, both sexes.....	81	113	214	313	414	525	555	459	223	69	13	1
Male.....	45	62	111	174	200	275	279	203	98	24	1	1
Female.....	36	51	103	139	214	250	276	256	125	45	12	-
White, both sexes.....	67	101	197	289	400	503	531	444	216	66	11	1
Male.....	37	59	101	161	192	264	270	198	95	22	-	1
Female.....	30	42	96	128	208	239	261	246	121	44	11	-
Black, both sexes.....	14	12	17	23	12	21	21	14	7	3	2	-
Male.....	8	3	10	12	8	10	7	5	3	2	1	-
Female.....	6	9	7	11	4	11	14	9	4	1	1	-
American Indian 1/, both sexes.....	-	-	-	1	1	1	1	-	-	-	-	-
Male.....	-	-	-	1	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	2	1	-	-	-	-
Male.....	-	-	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	1	-	-	-	-

Montana

All races, both sexes.....	12	15	33	67	95	117	115	82	44	13	2	-
Male.....	6	9	18	34	50	56	65	37	18	2	-	-
Female.....	6	6	15	33	45	61	50	45	26	11	2	-
White, both sexes.....	11	14	28	65	90	110	108	81	43	13	2	-
Male.....	5	9	16	33	48	51	63	37	18	2	-	-
Female.....	6	5	12	32	42	59	45	44	25	11	2	-
Black, both sexes.....	-	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	5	2	5	6	5	1	1	-	-	-
Male.....	1	-	2	1	2	4	2	-	-	-	-	-
Female.....	-	1	3	1	3	2	3	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Nebraska

All races, both sexes.....	22	19	43	88	100	167	179	156	78	33	4	1
Male.....	11	13	21	47	54	86	91	78	43	16	1	1
Female.....	11	6	22	41	46	81	88	78	35	17	3	-
White, both sexes.....	20	17	41	84	98	164	176	154	77	32	4	1
Male.....	10	11	20	45	53	85	90	77	43	15	1	1
Female.....	10	6	21	39	45	79	86	77	34	17	3	-
Black, both sexes.....	-	2	2	3	2	-	1	2	1	1	-	-
Male.....	-	2	1	2	1	-	-	1	-	1	-	-
Female.....	-	-	1	1	1	-	1	1	1	-	-	-
American Indian 1/, both sexes.....	2	-	-	1	-	1	2	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Nevada

All races, both sexes.....	18	29	70	115	144	204	196	154	78	13	-	-
Male.....	7	19	34	62	81	98	86	52	22	5	-	-
Female.....	11	10	36	53	63	106	110	102	56	8	-	-
White, both sexes.....	18	29	67	107	138	196	190	149	72	13	-	-
Male.....	7	19	32	57	75	92	83	50	20	5	-	-
Female.....	11	10	35	50	63	104	107	99	52	8	-	-
Black, both sexes.....	-	-	3	6	5	4	6	3	2	-	-	-
Male.....	-	-	2	4	5	2	3	2	1	-	-	-
Female.....	-	-	1	2	-	2	3	1	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	1	2	-	-	1	-	-	-
Male.....	-	-	-	1	1	2	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	2	-	2	3	-	-	-
Male.....	-	-	-	-	-	2	-	-	1	-	-	-
Female.....	-	-	-	1	-	-	-	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

New Hampshire

All races, both sexes.....	12	20	45	45	82	117	114	101	40	20	1	-
Male.....	6	8	22	25	32	64	47	45	11	6	-	-
Female.....	6	12	23	20	50	53	67	56	29	14	1	-
White, both sexes.....	12	20	45	45	82	117	114	101	40	20	1	-
Male.....	6	8	22	25	32	64	47	45	11	6	-	-
Female.....	6	12	23	20	50	53	67	56	29	14	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	36	104	149	246	316	502	615	563	286	83	12	-
Male.....	17	50	77	115	152	232	267	228	103	21	-	-
Female.....	19	54	72	131	164	270	348	335	183	62	12	-
White, both sexes.....	24	84	133	215	278	466	580	525	273	74	11	-
Male.....	15	43	66	95	134	213	251	208	95	20	-	-
Female.....	9	41	67	120	144	253	329	317	178	54	11	-
Black, both sexes.....	11	19	16	29	36	33	32	30	12	9	1	-
Male.....	2	6	11	20	17	17	16	15	7	1	-	-
Female.....	9	13	5	9	19	16	16	15	5	8	1	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	2	1	2	3	8	1	-	-	-
Male.....	-	1	-	-	-	2	-	5	1	-	-	-
Female.....	1	-	-	2	1	-	3	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

New Mexico

All races, both sexes.....	18	31	44	73	108	172	188	128	72	25	4	-
Male.....	7	18	23	37	58	87	102	58	32	10	1	-
Female.....	11	13	21	36	50	85	86	70	40	15	3	-
White, both sexes.....	17	30	43	73	106	169	183	126	69	21	3	-
Male.....	7	18	22	37	56	85	99	58	30	8	1	-
Female.....	10	12	21	36	50	84	84	68	39	13	2	-
Black, both sexes.....	1	-	1	-	1	2	1	-	1	-	-	-
Male.....	-	-	1	-	1	2	1	-	1	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	3	1	2	4	1	-
Male.....	-	-	-	-	1	-	1	-	1	2	-	-
Female.....	-	-	-	-	-	1	2	1	1	2	1	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	1	-	-	-	-

New York

All races, both sexes.....	132	199	360	510	745	1,155	1,311	1,088	619	204	32	-
Male.....	65	105	196	244	344	526	609	454	221	45	9	-
Female.....	67	94	164	266	401	629	702	634	398	159	23	-
White, both sexes.....	98	170	308	459	684	1,063	1,213	1,011	586	190	30	-
Male.....	51	93	163	218	312	480	560	416	203	41	8	-
Female.....	47	77	145	241	372	583	653	595	383	149	22	-
Black, both sexes.....	32	28	47	46	48	75	86	60	16	10	2	-
Male.....	13	11	29	22	23	38	42	29	7	2	1	-
Female.....	19	17	18	24	25	37	44	31	9	8	1	-
American Indian 1/, both sexes.....	1	-	2	-	2	3	2	1	-	-	-	-
Male.....	1	-	2	-	1	1	1	-	-	-	-	-
Female.....	-	-	-	-	1	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	3	5	11	14	10	16	17	4	-	-
Male.....	-	1	2	4	8	7	6	9	11	2	-	-
Female.....	1	-	1	1	3	7	4	7	6	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Ohio

All races, both sexes.....	161	250	398	624	828	1,181	1,314	944	428	127	17	-
Male.....	78	130	208	323	425	570	627	384	152	42	7	-
Female.....	83	120	190	301	403	611	687	560	276	85	10	-
White, both sexes.....	142	226	366	563	769	1,099	1,239	898	405	120	15	-
Male.....	69	117	184	292	386	530	587	369	143	39	6	-
Female.....	73	109	182	271	383	569	652	529	262	81	9	-
Black, both sexes.....	19	22	32	60	59	81	72	45	23	7	2	-
Male.....	9	13	24	31	39	39	38	14	9	3	1	-
Female.....	10	9	8	29	20	42	34	31	14	4	1	-
American Indian 1/, both sexes.....	-	1	-	1	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	1	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	1	1	1	-	-	-	-
Male.....	-	-	-	-	-	1	1	1	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-

Oklahoma

All races, both sexes.....	65	118	184	254	355	462	400	297	137	36	3	-
Male.....	27	65	84	141	184	235	208	152	56	14	-	-
Female.....	38	53	100	113	171	227	192	145	81	22	3	-
White, both sexes.....	53	109	166	235	330	431	377	283	128	32	3	-
Male.....	22	63	72	127	171	222	193	146	52	11	-	-
Female.....	31	46	94	108	159	209	184	137	76	21	3	-
Black, both sexes.....	5	1	9	7	11	11	8	5	4	3	-	-
Male.....	3	-	7	5	6	5	6	1	2	2	-	-
Female.....	2	1	2	2	5	6	2	4	2	1	-	-
American Indian 1/, both sexes.....	7	8	8	12	14	20	14	9	5	1	-	-
Male.....	2	2	5	9	7	8	9	5	2	1	-	-
Female.....	5	6	3	3	7	12	5	4	3	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Oregon

All races, both sexes.....	49	67	145	175	238	324	380	303	134	29	2	-
Male.....	34	39	71	86	117	166	182	139	59	7	-	-
Female.....	15	28	74	89	121	158	198	164	75	22	2	-
White, both sexes.....	47	65	137	171	233	317	368	300	131	29	2	-
Male.....	34	37	67	84	113	162	177	136	56	7	-	-
Female.....	13	28	70	87	120	155	191	164	75	22	2	-
Black, both sexes.....	1	-	2	1	3	3	1	1	1	-	-	-
Male.....	-	-	2	-	3	1	-	1	1	-	-	-
Female.....	1	-	-	1	-	2	1	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	5	3	2	3	8	-	2	-	-	-
Male.....	-	1	2	2	1	2	4	-	2	-	-	-
Female.....	1	-	3	1	1	1	4	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	-	-	1	3	2	-	-	-	-
Male.....	-	1	-	-	-	1	1	2	-	-	-	-
Female.....	-	-	1	-	-	-	2	-	-	-	-	-

Pennsylvania

All races, both sexes.....	103	181	331	479	711	1,051	1,291	1,041	536	155	21	1
Male.....	57	104	176	259	337	525	630	445	195	49	6	1
Female.....	46	77	155	220	374	526	661	596	341	106	15	-
White, both sexes.....	84	154	299	439	651	995	1,241	1,004	516	150	21	-
Male.....	49	87	159	238	310	495	599	429	188	48	6	-
Female.....	35	67	140	201	341	500	642	575	328	102	15	-
Black, both sexes.....	19	26	32	40	59	56	44	35	19	4	-	-
Male.....	8	17	17	21	27	30	26	14	7	1	-	-
Female.....	11	9	15	19	32	26	18	21	12	3	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	6	2	1	1	-	1
Male.....	-	-	-	-	-	-	5	2	-	-	-	1
Female.....	-	-	-	-	1	-	1	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Rhode Island

All races, both sexes.....	5	12	32	19	50	66	92	82	35	15	3	-
Male.....	2	6	17	8	28	27	47	32	13	3	-	-
Female.....	3	6	15	11	22	39	45	50	22	12	3	-
White, both sexes.....	5	12	32	18	49	62	90	81	34	15	3	-
Male.....	2	6	17	8	27	23	46	32	13	3	-	-
Female.....	3	6	15	10	22	39	44	49	21	12	3	-
Black, both sexes.....	-	-	-	-	1	4	1	-	1	-	-	-
Male.....	-	-	-	-	1	4	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

South Carolina

All races, both sexes.....	58	91	177	229	322	372	371	216	96	27	5	1
Male.....	26	48	100	118	172	201	192	97	36	8	1	1
Female.....	32	43	77	111	150	171	179	119	60	19	4	-
White, both sexes.....	42	72	152	199	283	332	332	193	88	21	4	1
Male.....	16	39	85	103	151	180	174	84	32	5	1	1
Female.....	26	33	67	96	132	152	158	109	56	16	3	-
Black, both sexes.....	16	19	25	28	39	40	38	23	8	6	1	-
Male.....	10	9	15	15	21	21	18	13	4	3	-	-
Female.....	6	10	10	13	18	19	20	10	4	3	1	-
American Indian 1/, both sexes.....	-	-	-	2	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

South Dakota

All races, both sexes.....	1	8	26	44	56	83	84	78	45	16	3	-
Male.....	-	6	13	21	33	47	54	44	28	7	1	-
Female.....	1	2	13	23	23	36	30	34	17	9	2	-
White, both sexes.....	-	8	24	41	55	81	80	75	44	16	3	-
Male.....	-	6	12	20	32	46	53	44	28	7	1	-
Female.....	-	2	12	21	23	35	27	31	16	9	2	-
Black, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	2	3	1	2	3	3	1	-	-	-
Male.....	-	-	1	1	1	1	-	-	-	-	-	-
Female.....	1	-	1	2	-	1	3	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	102	159	273	350	439	542	579	393	167	45	8	-
Male.....	58	75	147	193	203	275	302	168	70	13	3	-
Female.....	44	84	126	157	236	267	277	225	97	32	5	-
White, both sexes.....	80	144	250	326	409	508	542	369	153	42	6	-
Male.....	48	68	135	179	189	254	280	158	62	11	3	-
Female.....	32	76	115	147	220	254	262	211	91	31	3	-
Black, both sexes.....	22	14	23	21	29	33	37	23	14	3	2	-
Male.....	10	6	12	12	14	21	22	9	8	2	-	-
Female.....	12	8	11	9	15	12	15	14	6	1	2	-
American Indian 1/, both sexes.....	-	1	-	1	-	-	-	-	-	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	1	1	-	1	-	-	-	-
Male.....	-	-	-	1	-	-	-	1	-	-	-	-
Female.....	-	-	-	1	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Texas

All races, both sexes.....	179	299	570	836	1,169	1,488	1,474	1,137	506	158	16	1
Male.....	96	165	289	444	613	765	750	533	218	52	2	1
Female.....	83	134	281	392	556	723	724	604	288	106	14	-
White, both sexes.....	150	269	522	783	1,081	1,369	1,390	1,075	482	143	15	1
Male.....	78	150	262	415	560	699	707	491	204	46	2	1
Female.....	72	119	260	368	521	670	683	584	278	97	13	-
Black, both sexes.....	27	27	45	48	80	105	68	57	22	13	1	-
Male.....	17	14	26	26	48	62	38	39	13	6	-	-
Female.....	10	13	19	22	32	43	30	18	9	7	1	-
American Indian 1/, both sexes.....	1	2	-	2	2	6	6	1	-	-	-	-
Male.....	-	-	-	-	2	-	2	-	-	-	-	-
Female.....	1	2	-	2	-	6	4	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	3	3	6	8	10	4	2	2	-	-
Male.....	1	1	1	3	3	4	3	3	1	-	-	-
Female.....	-	-	2	-	3	4	7	1	1	2	-	-

Utah

All races, both sexes.....	13	40	40	57	107	107	111	77	31	6	1	-
Male.....	7	16	24	35	67	60	68	39	16	2	-	-
Female.....	6	24	16	22	40	47	43	38	15	4	1	-
White, both sexes.....	13	40	38	53	104	105	109	76	30	5	1	-
Male.....	7	16	24	33	65	58	66	39	15	1	-	-
Female.....	6	24	14	20	39	47	43	37	15	4	1	-
Black, both sexes.....	-	-	-	1	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	3	-	1	1	-	-	-	-	-
Male.....	-	-	-	2	-	1	1	-	-	-	-	-
Female.....	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	3	1	-	1	1	1	-	-
Male.....	-	-	-	-	2	1	-	-	1	1	-	-
Female.....	-	-	1	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Vermont

All races, both sexes.....	7	8	21	29	48	49	56	56	28	7	1	-
Male.....	3	4	12	9	24	21	32	28	11	-	-	-
Female.....	4	4	9	20	24	28	24	28	17	7	1	-
White, both sexes.....	7	8	21	29	48	48	56	56	28	7	1	-
Male.....	3	4	12	9	24	20	32	28	11	-	-	-
Female.....	4	4	9	20	24	28	24	28	17	7	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	49	90	174	273	349	545	540	412	196	52	7	-
Male.....	25	42	80	132	176	263	232	165	64	15	-	-
Female.....	24	48	94	141	173	282	308	247	132	37	7	-
White, both sexes.....	39	65	149	238	314	490	485	386	175	43	6	-
Male.....	21	28	67	112	160	223	205	153	58	12	-	-
Female.....	18	37	82	126	154	267	280	233	117	31	6	-
Black, both sexes.....	10	24	23	34	32	50	52	25	21	9	1	-
Male.....	4	13	13	19	15	36	27	11	6	3	-	-
Female.....	6	11	10	15	17	14	25	14	15	6	1	-
American Indian 1/, both sexes.....	-	-	1	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	2	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	1	3	3	3	1	-	-	-	-
Male.....	-	1	-	1	1	2	-	1	-	-	-	-
Female.....	-	-	1	-	2	1	3	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)

Washington

All races, both sexes.....	54	108	173	265	402	473	477	432	186	57	5	-
Male.....	26	52	100	132	204	230	219	182	76	20	3	-
Female.....	28	56	73	133	198	243	258	250	110	37	2	-
White, both sexes.....	50	95	161	252	382	446	462	421	184	56	4	-
Male.....	24	43	93	124	192	211	212	178	75	19	3	-
Female.....	26	52	68	128	190	235	250	243	109	37	1	-
Black, both sexes.....	2	7	6	7	5	8	3	-	1	-	-	-
Male.....	-	3	3	3	2	5	2	-	1	-	-	-
Female.....	2	4	3	4	3	3	1	-	-	-	-	-
American Indian 1/, both sexes.....	2	3	6	4	5	4	1	4	-	-	-	-
Male.....	2	3	4	3	3	3	-	2	-	-	-	-
Female.....	-	-	2	1	2	1	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	3	-	2	10	15	11	7	1	1	1	-
Male.....	-	3	-	2	7	11	5	2	-	1	-	-
Female.....	-	-	-	-	3	4	6	5	1	-	1	-

West Virginia

All races, both sexes.....	29	62	98	147	190	226	261	177	87	17	2	-
Male.....	17	35	54	78	87	106	132	85	38	6	-	-
Female.....	12	27	44	69	103	120	129	92	49	11	2	-
White, both sexes.....	28	60	97	147	187	224	259	175	86	16	2	-
Male.....	16	35	53	78	87	104	131	83	37	6	-	-
Female.....	12	25	44	69	100	120	128	92	49	10	2	-
Black, both sexes.....	1	2	1	-	2	2	2	2	1	1	-	-
Male.....	1	-	1	-	-	2	1	2	1	-	-	-
Female.....	-	2	-	-	2	-	1	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Bronchitis, not specified as acute or chronic (J40)

Alaska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Bronchitis, not specified as acute or chronic (J40)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchitis, not specified as acute or chronic (J40)

Arkansas

All races, both sexes.....	-	2	1	1	-	-	2	3	-	1	-	-
Male.....	-	1	-	-	-	-	-	1	-	-	-	-
Female.....	-	1	1	1	-	-	2	2	-	1	-	-
White, both sexes.....	-	2	1	1	-	-	2	3	-	-	-	-
Male.....	-	1	-	-	-	-	-	1	-	-	-	-
Female.....	-	1	1	1	-	-	2	2	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	2	2	2	3	2	3	10	5	3	8	1	-
Male.....	-	-	1	1	-	2	3	3	-	2	-	-
Female.....	2	2	1	2	2	1	7	2	3	6	1	-
White, both sexes.....	2	1	-	2	2	2	10	5	3	6	-	-
Male.....	-	-	-	1	-	1	3	3	-	1	-	-
Female.....	2	1	-	1	2	1	7	2	3	5	-	-
Black, both sexes.....	-	-	1	1	-	-	-	-	-	-	1	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	1	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	-	-	1	-	-	-	2	-	-
Male.....	-	-	1	-	-	1	-	-	-	1	-	-
Female.....	-	1	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Chronic bronchitis (J41-J42)

Alaska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Chronic bronchitis (J41-J42)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Chronic bronchitis (J41-J42)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

Chronic bronchitis (J41-J42)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Chronic bronchitis (J41-J42)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Chronic bronchitis (J41-J42)

Arkansas

All races, both sexes.....	-	-	-	-	1	1	2	2	-	-	-	-
Male.....	-	-	-	-	1	1	-	2	-	-	-	-
Female.....	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes.....	-	-	-	-	1	1	2	2	-	-	-	-
Male.....	-	-	-	-	1	1	-	2	-	-	-	-
Female.....	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	-	-	1	2	1	3	7	9	4	-	-	-
Male.....	-	-	-	1	-	3	4	4	3	-	-	-
Female.....	-	-	1	1	1	-	3	5	1	-	-	-
White, both sexes.....	-	-	1	2	1	3	7	8	4	-	-	-
Male.....	-	-	-	1	-	3	4	4	3	-	-	-
Female.....	-	-	1	1	1	-	3	4	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)												
Arkansas												
All races, both sexes.....	7	8	27	33	36	57	27	13	4	2	-	-
Male.....	3	5	18	19	19	31	15	9	3	1	-	-
Female.....	4	3	9	14	17	26	12	4	1	1	-	-
White, both sexes.....	7	8	25	32	34	56	25	13	4	2	-	-
Male.....	3	5	17	19	18	31	13	9	3	1	-	-
Female.....	4	3	8	13	16	25	12	4	1	1	-	-
Black, both sexes.....	-	-	2	-	1	1	2	-	-	-	-	-
Male.....	-	-	1	-	-	-	2	-	-	-	-	-
Female.....	-	-	1	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	32	47	66	123	153	220	236	197	87	18	4	-
Male.....	23	32	31	72	79	104	116	97	31	6	1	-
Female.....	9	15	35	51	74	116	120	100	56	12	3	-
White, both sexes.....	29	41	55	109	137	192	214	182	80	18	3	-
Male.....	21	27	25	62	72	83	103	85	26	6	1	-
Female.....	8	14	30	47	65	109	111	97	54	12	2	-
Black, both sexes.....	2	4	10	13	12	11	9	1	1	-	1	-
Male.....	1	3	5	9	6	9	4	1	1	-	-	-
Female.....	1	1	5	4	6	2	5	-	-	-	1	-
American Indian 1/, both sexes.....	1	-	-	-	-	1	-	2	1	-	-	-
Male.....	1	-	-	-	-	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	1	1	4	16	13	12	5	-	-	-
Male.....	-	2	1	1	1	12	9	10	3	-	-	-
Female.....	-	-	-	-	3	4	4	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Emphysema (J43)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Emphysema (J43)

Massachusetts

All races, both sexes.....	3	16	20	22	34	41	56	32	19	7	1	-
Male.....	1	11	10	10	14	23	25	11	5	2	-	-
Female.....	2	5	10	12	20	18	31	21	14	5	1	-
White, both sexes.....	3	16	19	22	34	40	55	31	19	7	1	-
Male.....	1	11	9	10	14	22	25	10	5	2	-	-
Female.....	2	5	10	12	20	18	30	21	14	5	1	-
Black, both sexes.....	-	-	1	-	-	1	1	1	-	-	-	-
Male.....	-	-	1	-	-	1	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Michigan

All races, both sexes.....	22	22	32	61	74	125	122	80	24	8	1	-
Male.....	12	12	19	27	37	69	60	33	13	3	-	-
Female.....	10	10	13	34	37	56	62	47	11	5	1	-
White, both sexes.....	17	21	29	56	64	113	113	77	21	8	-	-
Male.....	9	12	17	22	29	62	55	33	11	3	-	-
Female.....	8	9	12	34	35	51	58	44	10	5	-	-
Black, both sexes.....	5	-	3	4	10	11	6	3	3	-	1	-
Male.....	3	-	2	4	8	7	4	-	2	-	-	-
Female.....	2	-	1	-	2	4	2	3	1	-	1	-
American Indian 1/, both sexes.....	-	-	-	1	-	1	1	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	1	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Emphysema (J43)

Nebraska

All races, both sexes.....	1	2	4	13	11	15	12	9	7	1	-	1
Male.....	1	2	1	8	3	8	6	3	3	1	-	1
Female.....	-	-	3	5	8	7	6	6	4	-	-	-
White, both sexes.....	1	2	4	13	11	15	12	9	7	1	-	1
Male.....	1	2	1	8	3	8	6	3	3	1	-	1
Female.....	-	-	3	5	8	7	6	6	4	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	1	5	10	4	13	20	12	6	4	1	-	-
Male.....	-	2	6	2	7	13	5	2	2	-	-	-
Female.....	1	3	4	2	6	7	7	4	2	1	-	-
White, both sexes.....	1	5	10	3	13	20	11	5	4	1	-	-
Male.....	-	2	6	1	7	13	5	2	2	-	-	-
Female.....	1	3	4	2	6	7	6	3	2	1	-	-
Black, both sexes.....	-	-	-	1	-	-	1	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Emphysema (J43)

New Hampshire

All races, both sexes.....	2	3	8	3	13	10	10	17	3	4	1	-
Male.....	1	1	3	2	2	7	4	9	-	2	-	-
Female.....	1	2	5	1	11	3	6	8	3	2	1	-
White, both sexes.....	2	3	8	3	13	10	10	17	3	4	1	-
Male.....	1	1	3	2	2	7	4	9	-	2	-	-
Female.....	1	2	5	1	11	3	6	8	3	2	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	4	14	24	35	59	78	79	68	34	7	-	-
Male.....	1	7	14	15	29	37	32	31	13	2	-	-
Female.....	3	7	10	20	30	41	47	37	21	5	-	-
White, both sexes.....	3	13	22	29	53	73	79	66	34	7	-	-
Male.....	1	6	12	11	26	36	32	29	13	2	-	-
Female.....	2	7	10	18	27	37	47	37	21	5	-	-
Black, both sexes.....	1	-	2	6	5	5	-	2	-	-	-	-
Male.....	-	-	2	4	3	1	-	2	-	-	-	-
Female.....	1	-	-	2	2	4	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	1	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Emphysema (J43)

New Mexico

All races, both sexes.....	2	5	3	7	8	10	20	11	5	1	-	-
Male.....	1	3	2	2	3	6	8	5	2	-	-	-
Female.....	1	2	1	5	5	4	12	6	3	1	-	-
White, both sexes.....	2	5	3	7	8	10	20	10	5	1	-	-
Male.....	1	3	2	2	3	6	8	5	2	-	-	-
Female.....	1	2	1	5	5	4	12	5	3	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	18	26	39	54	90	132	111	91	40	12	3	-
Male.....	10	12	21	23	40	65	55	33	18	2	-	-
Female.....	8	14	18	31	50	67	56	58	22	10	3	-
White, both sexes.....	12	20	33	51	83	124	100	84	40	12	2	-
Male.....	7	9	16	22	36	60	51	29	18	2	-	-
Female.....	5	11	17	29	47	64	49	55	22	10	2	-
Black, both sexes.....	6	6	5	1	2	6	9	7	-	-	1	-
Male.....	3	3	4	-	2	4	2	4	-	-	-	-
Female.....	3	3	1	1	-	2	7	3	-	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	2	4	1	1	-	-	-	-	-
Male.....	-	-	1	1	2	1	1	-	-	-	-	-
Female.....	-	-	-	1	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Emphysema (J43)

Oregon

All races, both sexes.....	7	5	23	17	30	42	58	37	15	6	1	-
Male.....	6	1	13	5	13	25	24	19	6	2	-	-
Female.....	1	4	10	12	17	17	34	18	9	4	1	-
White, both sexes.....	7	5	21	17	30	41	58	37	15	6	1	-
Male.....	6	1	12	5	13	24	24	19	6	2	-	-
Female.....	1	4	9	12	17	17	34	18	9	4	1	-
Black, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	8	21	40	51	95	104	136	62	42	8	1	-
Male.....	4	10	29	33	50	56	60	29	16	4	-	-
Female.....	4	11	11	18	45	48	76	33	26	4	1	-
White, both sexes.....	6	17	37	48	89	98	129	61	39	8	1	-
Male.....	3	7	27	30	46	52	57	28	14	4	-	-
Female.....	3	10	10	18	43	46	72	33	25	4	1	-
Black, both sexes.....	2	4	3	3	5	6	6	1	3	-	-	-
Male.....	1	3	2	3	4	4	2	1	2	-	-	-
Female.....	1	1	1	-	1	2	4	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Emphysema (J43)

South Dakota

All races, both sexes.....	-	1	2	8	3	8	12	7	4	3	-	-
Male.....	-	1	2	4	1	6	10	3	2	1	-	-
Female.....	-	-	-	4	2	2	2	4	2	2	-	-
White, both sexes.....	-	1	1	8	3	8	12	7	4	3	-	-
Male.....	-	1	1	4	1	6	10	3	2	1	-	-
Female.....	-	-	-	4	2	2	2	4	2	2	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	15	17	48	42	55	50	60	19	9	2	-	-
Male.....	14	12	31	23	24	28	29	7	4	1	-	-
Female.....	1	5	17	19	31	22	31	12	5	1	-	-
White, both sexes.....	14	16	44	40	52	47	58	18	9	2	-	-
Male.....	13	12	28	21	24	26	28	6	4	1	-	-
Female.....	1	4	16	19	28	21	30	12	5	1	-	-
Black, both sexes.....	1	1	4	2	2	3	2	1	-	-	-	-
Male.....	1	-	3	2	-	2	1	1	-	-	-	-
Female.....	-	1	1	-	2	1	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Emphysema (J43)

Texas

All races, both sexes.....	21	43	68	107	129	165	142	82	41	5	2	-
Male.....	13	28	41	68	75	86	78	42	14	2	-	-
Female.....	8	15	27	39	54	79	64	40	27	3	2	-
White, both sexes.....	21	40	62	102	123	148	134	77	37	4	2	-
Male.....	13	27	37	68	71	75	73	39	12	1	-	-
Female.....	8	13	25	34	52	73	61	38	25	3	2	-
Black, both sexes.....	-	3	5	4	5	16	5	5	4	1	-	-
Male.....	-	1	3	-	3	10	4	3	2	1	-	-
Female.....	-	2	2	4	2	6	1	2	2	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	1	1	1	-	-	-	-	-
Male.....	-	-	1	-	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Utah

All races, both sexes.....	4	5	3	6	12	8	12	6	3	1	-	-
Male.....	3	2	2	5	7	6	9	3	2	1	-	-
Female.....	1	3	1	1	5	2	3	3	1	-	-	-
White, both sexes.....	4	5	3	6	11	8	12	6	3	1	-	-
Male.....	3	2	2	5	7	6	9	3	2	1	-	-
Female.....	1	3	1	1	4	2	3	3	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

United States

All races, both sexes.....	2,133	3,755	6,876	10,264	14,454	19,888	22,212	17,519	8,660	2,600	331	3
Male.....	1,105	2,013	3,586	5,322	7,356	9,881	10,750	7,830	3,359	840	80	3
Female.....	1,028	1,742	3,290	4,942	7,098	10,007	11,462	9,689	5,301	1,760	251	-
White, both sexes.....	1,834	3,311	6,281	9,418	13,398	18,567	20,917	16,575	8,170	2,426	295	3
Male.....	964	1,777	3,226	4,846	6,752	9,127	10,040	7,359	3,125	769	68	3
Female.....	870	1,534	3,055	4,572	6,646	9,440	10,877	9,216	5,045	1,657	227	-
Black, both sexes.....	267	384	513	719	877	1,048	983	689	351	126	27	-
Male.....	127	206	312	400	493	590	538	320	159	50	10	-
Female.....	140	178	201	319	384	458	445	369	192	76	17	-
American Indian 1/, both sexes.....	22	37	46	71	83	82	88	40	20	12	1	-
Male.....	9	16	26	37	47	39	43	17	11	3	-	-
Female.....	13	21	20	34	36	43	45	23	9	9	1	-
Asian or Pacific Islander 2/, both sexes.....	10	23	36	56	96	191	224	215	119	36	8	-
Male.....	5	14	22	39	64	125	129	134	64	18	2	-
Female.....	5	9	14	17	32	66	95	81	55	18	6	-

Alabama

All races, both sexes.....	57	69	186	268	332	422	413	276	116	42	5	-
Male.....	34	33	110	152	181	218	190	130	46	17	1	-
Female.....	23	36	76	116	151	204	223	146	70	25	4	-
White, both sexes.....	50	60	163	247	293	382	379	254	106	34	3	-
Male.....	31	28	92	142	160	195	171	119	42	12	-	-
Female.....	19	32	71	105	133	187	208	135	64	22	3	-
Black, both sexes.....	7	9	22	21	39	39	34	20	10	7	2	-
Male.....	3	5	17	10	21	22	19	11	4	5	1	-
Female.....	4	4	5	11	18	17	15	9	6	2	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	1	-	2	-	-	-	-
Male.....	-	-	1	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

Alaska

All races, both sexes.....	4	5	14	17	23	34	24	13	4	2	-	-
Male.....	2	-	7	9	13	12	17	7	1	1	-	-
Female.....	2	5	7	8	10	22	7	6	3	1	-	-
White, both sexes.....	3	3	11	11	15	31	21	8	4	2	-	-
Male.....	2	-	6	6	7	11	15	4	1	1	-	-
Female.....	1	3	5	5	8	20	6	4	3	1	-	-
Black, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	3	6	8	3	3	3	-	-	-	-
Male.....	-	-	1	3	6	1	2	1	-	-	-	-
Female.....	1	1	2	3	2	2	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	2	-	-	-	-
Male.....	-	-	-	-	-	-	-	2	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	39	72	158	208	323	396	477	383	176	57	6	-
Male.....	19	33	67	110	174	206	232	186	72	27	1	-
Female.....	20	39	91	98	149	190	245	197	104	30	5	-
White, both sexes.....	36	71	156	204	314	388	468	378	174	56	5	-
Male.....	18	33	66	108	170	201	229	184	71	26	1	-
Female.....	18	38	90	96	144	187	239	194	103	30	4	-
Black, both sexes.....	2	1	2	2	4	5	3	2	-	-	1	-
Male.....	1	-	1	1	1	2	1	-	-	-	-	-
Female.....	1	1	1	1	3	3	2	2	-	-	1	-
American Indian 1/, both sexes.....	1	-	-	2	3	2	5	2	1	1	-	-
Male.....	-	-	-	1	2	2	2	1	1	1	-	-
Female.....	1	-	-	1	1	-	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	1	1	1	1	-	-	-
Male.....	-	-	-	-	1	1	-	1	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic obstructive pulmonary disease (J44)

Arkansas

All races, both sexes.....	30	51	98	156	183	262	232	184	85	34	8	-
Male.....	21	23	62	80	94	142	123	89	31	12	2	-
Female.....	9	28	36	76	89	120	109	95	54	22	6	-
White, both sexes.....	22	48	95	146	175	252	223	175	78	33	7	-
Male.....	14	23	59	77	89	138	118	86	27	12	2	-
Female.....	8	25	36	69	86	114	105	89	51	21	5	-
Black, both sexes.....	8	3	3	10	8	8	9	9	7	1	1	-
Male.....	7	-	3	3	5	3	5	3	4	-	-	-
Female.....	1	3	-	7	3	5	4	6	3	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

California

All races, both sexes.....	200	328	594	851	1,280	1,883	2,233	1,908	1,005	283	40	-
Male.....	94	194	290	428	628	943	1,045	867	390	107	5	-
Female.....	106	134	304	423	652	940	1,188	1,041	615	176	35	-
White, both sexes.....	166	272	527	756	1,149	1,693	2,028	1,720	897	252	34	-
Male.....	79	165	251	375	550	826	917	760	330	96	3	-
Female.....	87	107	276	381	599	867	1,111	960	567	156	31	-
Black, both sexes.....	24	41	47	59	73	83	75	50	27	10	2	-
Male.....	11	21	27	26	38	46	46	15	14	1	1	-
Female.....	13	20	20	33	35	37	29	35	13	9	1	-
American Indian 1/, both sexes.....	3	6	5	7	15	5	8	6	3	2	-	-
Male.....	-	3	2	4	9	1	5	4	2	-	-	-
Female.....	3	3	3	3	6	4	3	2	1	2	-	-
Asian or Pacific Islander 2/, both sexes.....	7	9	15	29	43	102	122	132	78	19	4	-
Male.....	4	5	10	23	31	70	77	88	44	10	1	-
Female.....	3	4	5	6	12	32	45	44	34	9	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

Florida

All races, both sexes.....	146	285	480	691	1,045	1,412	1,697	1,373	709	207	27	-
Male.....	87	169	265	347	550	716	812	576	263	64	9	-
Female.....	59	116	215	344	495	696	885	797	446	143	18	-
White, both sexes.....	127	254	446	644	993	1,354	1,642	1,335	683	199	26	-
Male.....	80	149	243	323	519	675	780	558	252	59	8	-
Female.....	47	105	203	321	474	679	862	777	431	140	18	-
Black, both sexes.....	17	29	32	46	47	56	50	34	25	8	1	-
Male.....	5	18	21	24	30	41	29	16	11	5	1	-
Female.....	12	11	11	22	17	15	21	18	14	3	-	-
American Indian 1/, both sexes.....	2	-	1	1	2	1	3	-	-	-	-	-
Male.....	2	-	1	-	-	-	2	-	-	-	-	-
Female.....	-	-	-	1	2	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	1	-	3	1	2	4	1	-	-	-
Male.....	-	2	-	-	1	-	1	2	-	-	-	-
Female.....	-	-	1	-	2	1	1	2	1	-	-	-

Georgia

All races, both sexes.....	76	121	206	315	431	488	561	413	174	65	4	-
Male.....	40	66	111	171	206	237	271	197	61	19	1	-
Female.....	36	55	95	144	225	251	290	216	113	46	3	-
White, both sexes.....	61	100	177	270	379	442	513	375	162	54	2	-
Male.....	31	55	95	141	176	207	242	181	56	14	-	-
Female.....	30	45	82	129	203	235	271	194	106	40	2	-
Black, both sexes.....	15	21	29	43	48	43	46	37	11	10	2	-
Male.....	9	11	16	29	27	28	29	16	4	4	1	-
Female.....	6	10	13	14	21	15	17	21	7	6	1	-
American Indian 1/, both sexes.....	-	-	-	-	2	2	1	-	-	-	-	-
Male.....	-	-	-	-	1	2	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	2	1	1	1	1	1	-	-
Male.....	-	-	-	1	2	-	-	-	1	1	-	-
Female.....	-	-	-	1	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

Illinois

All races, both sexes.....	79	124	228	332	512	776	796	659	356	86	10	-
Male.....	47	59	126	187	256	380	371	285	110	27	1	-
Female.....	32	65	102	145	256	396	425	374	246	59	9	-
White, both sexes.....	63	93	198	290	458	713	734	614	331	79	10	-
Male.....	37	44	108	162	230	344	336	267	101	26	1	-
Female.....	26	49	90	128	228	369	398	347	230	53	9	-
Black, both sexes.....	16	31	29	41	53	57	54	42	20	6	-	-
Male.....	10	15	17	24	26	32	28	16	8	1	-	-
Female.....	6	16	12	17	27	25	26	26	12	5	-	-
American Indian 1/, both sexes.....	-	-	-	1	1	-	2	-	-	-	-	-
Male.....	-	-	-	1	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	6	6	3	5	1	-	-
Male.....	-	-	1	-	-	4	6	2	1	-	-	-
Female.....	-	-	-	-	-	2	-	1	4	1	-	-

Indiana

All races, both sexes.....	61	96	180	283	397	465	568	413	177	48	12	-
Male.....	29	46	102	150	208	230	291	192	68	12	4	-
Female.....	32	50	78	133	189	235	277	221	109	36	8	-
White, both sexes.....	59	92	176	277	385	451	550	392	172	48	12	-
Male.....	28	44	100	146	203	223	279	182	66	12	4	-
Female.....	31	48	76	131	182	228	271	210	106	36	8	-
Black, both sexes.....	2	4	4	6	11	14	18	20	5	-	-	-
Male.....	1	2	2	4	5	7	12	10	2	-	-	-
Female.....	1	2	2	2	6	7	6	10	3	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic obstructive pulmonary disease (J44)

Iowa

All races, both sexes.....	24	41	88	103	186	270	309	264	135	47	3	-
Male.....	14	23	47	46	102	142	161	146	61	15	-	-
Female.....	10	18	41	57	84	128	148	118	74	32	3	-
White, both sexes.....	24	40	86	102	182	266	308	264	135	47	3	-
Male.....	14	22	45	45	100	140	160	146	61	15	-	-
Female.....	10	18	41	57	82	126	148	118	74	32	3	-
Black, both sexes.....	-	1	2	1	4	4	1	-	-	-	-	-
Male.....	-	1	2	1	2	2	1	-	-	-	-	-
Female.....	-	-	-	-	2	2	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Kansas

All races, both sexes.....	18	44	72	111	163	240	288	195	107	31	2	-
Male.....	8	20	44	60	89	123	149	98	47	10	-	-
Female.....	10	24	28	51	74	117	139	97	60	21	2	-
White, both sexes.....	18	38	67	108	159	226	280	190	106	31	2	-
Male.....	8	17	40	58	86	114	146	95	46	10	-	-
Female.....	10	21	27	50	73	112	134	95	60	21	2	-
Black, both sexes.....	-	3	3	2	1	8	4	3	1	-	-	-
Male.....	-	2	3	2	1	6	2	1	1	-	-	-
Female.....	-	1	-	-	-	2	2	2	-	-	-	-
American Indian 1/, both sexes.....	-	3	2	1	2	6	4	2	-	-	-	-
Male.....	-	1	1	-	2	3	1	2	-	-	-	-
Female.....	-	2	1	1	-	3	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

Maine

All races, both sexes.....	9	21	51	66	96	110	122	119	53	19	1	-
Male.....	2	14	21	37	47	56	54	50	15	7	-	-
Female.....	7	7	30	29	49	54	68	69	38	12	1	-
White, both sexes.....	9	21	50	61	95	110	122	119	52	19	1	-
Male.....	2	14	21	35	47	56	54	50	15	7	-	-
Female.....	7	7	29	26	48	54	68	69	37	12	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	4	1	-	-	-	1	-	-	-
Male.....	-	-	-	2	-	-	-	-	-	-	-	-
Female.....	-	-	1	2	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	21	42	83	119	190	292	349	297	149	37	3	-
Male.....	10	14	41	60	98	130	127	119	50	12	2	-
Female.....	11	28	42	59	92	162	222	178	99	25	1	-
White, both sexes.....	18	34	74	102	165	248	311	268	132	34	2	-
Male.....	9	11	33	49	87	111	111	109	43	11	2	-
Female.....	9	23	41	53	78	137	200	159	89	23	-	-
Black, both sexes.....	3	8	9	17	23	42	34	29	17	2	1	-
Male.....	1	3	8	11	10	18	13	10	7	1	-	-
Female.....	2	5	1	6	13	24	21	19	10	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	1	4	-	-	-	-	-
Male.....	-	-	-	-	1	1	3	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

Missouri

All races, both sexes.....	69	93	184	272	366	465	502	422	205	66	12	1
Male.....	37	52	99	147	175	245	248	186	90	23	1	1
Female.....	32	41	85	125	191	220	254	236	115	43	11	-
White, both sexes.....	60	83	169	251	355	445	480	408	199	64	10	1
Male.....	33	50	89	136	167	234	239	181	88	22	-	1
Female.....	27	33	80	115	188	211	241	227	111	42	10	-
Black, both sexes.....	9	10	15	20	11	19	19	13	6	2	2	-
Male.....	4	2	10	10	8	10	7	5	2	1	1	-
Female.....	5	8	5	10	3	9	12	8	4	1	1	-
American Indian 1/, both sexes.....	-	-	-	1	-	1	1	-	-	-	-	-
Male.....	-	-	-	1	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	2	1	-	-	-	-
Male.....	-	-	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Montana

All races, both sexes.....	11	14	25	58	86	110	99	74	40	11	2	-
Male.....	5	8	14	31	45	54	55	34	16	2	-	-
Female.....	6	6	11	27	41	56	44	40	24	9	2	-
White, both sexes.....	10	13	21	56	81	103	92	73	39	11	2	-
Male.....	4	8	12	30	43	49	53	34	16	2	-	-
Female.....	6	5	9	26	38	54	39	39	23	9	2	-
Black, both sexes.....	-	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	4	2	5	6	5	1	1	-	-	-
Male.....	1	-	2	1	2	4	2	-	-	-	-	-
Female.....	-	1	2	1	3	2	3	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

Nebraska

All races, both sexes.....	18	16	35	75	88	150	166	141	64	32	4	-
Male.....	9	10	19	39	50	77	84	74	36	15	1	-
Female.....	9	6	16	36	38	73	82	67	28	17	3	-
White, both sexes.....	16	14	34	71	86	147	163	139	63	31	4	-
Male.....	8	8	18	37	49	76	83	73	36	14	1	-
Female.....	8	6	16	34	37	71	80	66	27	17	3	-
Black, both sexes.....	-	2	1	3	2	-	1	2	1	1	-	-
Male.....	-	2	1	2	1	-	-	1	-	1	-	-
Female.....	-	-	-	1	1	-	1	1	1	-	-	-
American Indian 1/, both sexes.....	2	-	-	1	-	1	2	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	-	-	1	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Nevada

All races, both sexes.....	13	24	59	111	131	182	181	146	74	12	-	-
Male.....	5	17	28	60	74	84	80	48	20	5	-	-
Female.....	8	7	31	51	57	98	101	98	54	7	-	-
White, both sexes.....	13	24	56	104	125	174	176	142	68	12	-	-
Male.....	5	17	26	56	68	78	77	46	18	5	-	-
Female.....	8	7	30	48	57	96	99	96	50	7	-	-
Black, both sexes.....	-	-	3	5	5	4	5	3	2	-	-	-
Male.....	-	-	2	3	5	2	3	2	1	-	-	-
Female.....	-	-	1	2	-	2	2	1	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	1	2	-	-	1	-	-	-
Male.....	-	-	-	1	1	2	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	2	-	1	3	-	-	-
Male.....	-	-	-	-	-	2	-	-	1	-	-	-
Female.....	-	-	-	1	-	-	-	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

New Hampshire

All races, both sexes.....	9	16	34	42	69	106	101	82	36	14	-	-
Male.....	5	7	18	23	30	56	42	36	10	3	-	-
Female.....	4	9	16	19	39	50	59	46	26	11	-	-
White, both sexes.....	9	16	34	42	69	106	101	82	36	14	-	-
Male.....	5	7	18	23	30	56	42	36	10	3	-	-
Female.....	4	9	16	19	39	50	59	46	26	11	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	26	79	118	201	253	417	527	481	245	74	10	-
Male.....	16	40	62	93	122	194	233	193	88	19	-	-
Female.....	10	39	56	108	131	223	294	288	157	55	10	-
White, both sexes.....	21	63	106	181	221	386	492	446	233	67	10	-
Male.....	14	34	54	80	107	176	217	175	81	18	-	-
Female.....	7	29	52	101	114	210	275	271	152	49	10	-
Black, both sexes.....	4	16	12	20	31	28	32	28	12	7	-	-
Male.....	2	6	8	13	14	16	16	13	7	1	-	-
Female.....	2	10	4	7	17	12	16	15	5	6	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	2	3	7	-	-	-	-
Male.....	-	-	-	-	-	2	-	5	-	-	-	-
Female.....	1	-	-	-	-	-	3	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

New Mexico

All races, both sexes.....	14	20	41	64	98	158	166	111	66	23	4	-
Male.....	5	13	21	34	54	80	93	52	30	9	1	-
Female.....	9	7	20	30	44	78	73	59	36	14	3	-
White, both sexes.....	14	19	40	64	96	155	161	110	63	20	3	-
Male.....	5	13	20	34	52	78	90	52	28	8	1	-
Female.....	9	6	20	30	44	77	71	58	35	12	2	-
Black, both sexes.....	-	-	1	-	1	2	1	-	1	-	-	-
Male.....	-	-	1	-	1	2	1	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	3	-	2	3	1	-
Male.....	-	-	-	-	1	-	1	-	1	1	-	-
Female.....	-	-	-	-	-	1	2	-	1	2	1	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	1	-	-	-	-

New York

All races, both sexes.....	92	157	304	441	645	1,005	1,177	973	558	183	27	-
Male.....	47	87	165	211	300	456	547	415	197	42	9	-
Female.....	45	70	139	230	345	549	630	558	361	141	18	-
White, both sexes.....	72	140	267	398	594	926	1,094	909	526	171	26	-
Male.....	39	80	143	191	273	416	504	384	179	38	8	-
Female.....	33	60	124	207	321	510	590	525	347	133	18	-
Black, both sexes.....	19	17	34	40	43	65	73	48	15	8	1	-
Male.....	7	7	20	17	20	34	38	22	7	2	1	-
Female.....	12	10	14	23	23	31	35	26	8	6	-	-
American Indian 1/, both sexes.....	1	-	1	-	1	2	1	1	-	-	-	-
Male.....	1	-	1	-	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	3	7	12	9	15	17	4	-	-
Male.....	-	-	1	3	6	5	5	9	11	2	-	-
Female.....	-	-	1	-	1	7	4	6	6	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

Ohio

All races, both sexes.....	131	210	332	528	707	1,017	1,147	831	378	116	13	-
Male.....	59	109	180	277	361	480	557	333	138	39	7	-
Female.....	72	101	152	251	346	537	590	498	240	77	6	-
White, both sexes.....	114	189	311	479	655	949	1,084	787	358	109	11	-
Male.....	51	98	163	250	328	449	524	318	132	36	6	-
Female.....	63	91	148	229	327	500	560	469	226	73	5	-
Black, both sexes.....	17	19	21	48	52	67	62	43	20	7	2	-
Male.....	8	11	17	27	33	30	32	14	6	3	1	-
Female.....	9	8	4	21	19	37	30	29	14	4	1	-
American Indian 1/, both sexes.....	-	1	-	1	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	1	-	1	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-

Oklahoma

All races, both sexes.....	59	96	165	222	314	422	361	271	125	29	3	-
Male.....	25	53	76	125	161	212	187	140	52	11	-	-
Female.....	34	43	89	97	153	210	174	131	73	18	3	-
White, both sexes.....	49	88	149	205	291	393	339	259	117	25	3	-
Male.....	21	51	65	112	148	200	173	135	48	8	-	-
Female.....	28	37	84	93	143	193	166	124	69	17	3	-
Black, both sexes.....	5	1	8	7	9	10	7	5	4	3	-	-
Male.....	3	-	6	5	6	4	5	1	2	2	-	-
Female.....	2	1	2	2	3	6	2	4	2	1	-	-
American Indian 1/, both sexes.....	5	7	7	10	14	19	14	7	4	1	-	-
Male.....	1	2	5	8	7	8	9	4	2	1	-	-
Female.....	4	5	2	2	7	11	5	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

Oregon

All races, both sexes.....	36	59	118	149	206	270	314	261	110	19	1	-
Male.....	26	36	57	78	103	136	156	120	51	5	-	-
Female.....	10	23	61	71	103	134	158	141	59	14	1	-
White, both sexes.....	35	57	113	147	202	264	302	258	107	19	1	-
Male.....	26	34	54	77	100	133	151	117	48	5	-	-
Female.....	9	23	59	70	102	131	151	141	59	14	1	-
Black, both sexes.....	-	-	1	1	2	3	1	1	1	-	-	-
Male.....	-	-	1	-	2	1	-	1	1	-	-	-
Female.....	-	-	-	1	-	2	1	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	4	1	2	2	8	-	2	-	-	-
Male.....	-	1	2	1	1	1	4	-	2	-	-	-
Female.....	1	-	2	-	1	1	4	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	1	3	2	-	-	-	-
Male.....	-	1	-	-	-	1	1	2	-	-	-	-
Female.....	-	-	-	-	-	-	2	-	-	-	-	-

Pennsylvania

All races, both sexes.....	83	148	280	418	608	940	1,139	956	475	135	20	-
Male.....	49	89	144	223	285	468	564	407	176	44	6	-
Female.....	34	59	136	195	323	472	575	549	299	91	14	-
White, both sexes.....	68	129	253	383	555	890	1,096	921	460	131	20	-
Male.....	43	75	130	205	262	442	536	392	171	43	6	-
Female.....	25	54	123	178	293	448	560	529	289	88	14	-
Black, both sexes.....	15	18	27	35	53	50	38	33	15	3	-	-
Male.....	6	14	14	18	23	26	24	13	5	1	-	-
Female.....	9	4	13	17	30	24	14	20	10	2	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	5	2	-	1	-	-
Male.....	-	-	-	-	-	-	4	2	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

South Dakota

All races, both sexes.....	1	7	21	35	52	73	69	70	40	12	2	-
Male.....	-	5	10	17	32	40	44	41	25	6	1	-
Female.....	1	2	11	18	20	33	25	29	15	6	1	-
White, both sexes.....	-	7	20	32	51	71	65	67	39	12	2	-
Male.....	-	5	10	16	31	39	43	41	25	6	1	-
Female.....	-	2	10	16	20	32	22	26	14	6	1	-
Black, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	1	3	1	2	3	3	1	-	-	-
Male.....	-	-	-	1	1	1	-	-	-	-	-	-
Female.....	1	-	1	2	-	1	3	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	74	134	219	303	377	485	514	365	152	39	7	-
Male.....	40	62	115	167	178	246	272	160	65	12	2	-
Female.....	34	72	104	136	199	239	242	205	87	27	5	-
White, both sexes.....	61	125	202	283	352	454	481	344	139	37	5	-
Male.....	34	55	107	156	165	227	252	151	57	10	2	-
Female.....	27	70	95	127	187	227	229	193	82	27	3	-
Black, both sexes.....	13	8	17	18	25	30	33	20	13	2	2	-
Male.....	6	6	8	10	13	19	20	8	8	2	-	-
Female.....	7	2	9	8	12	11	13	12	5	-	2	-
American Indian 1/, both sexes.....	-	1	-	1	-	-	-	-	-	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	1	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic obstructive pulmonary disease (J44)												
Texas												
All races, both sexes.....	135	241	481	713	1,026	1,308	1,304	1,026	451	151	11	1
Male.....	73	129	240	371	532	675	661	483	201	50	1	1
Female.....	62	112	241	342	494	633	643	543	250	101	10	-
White, both sexes.....	115	219	443	666	947	1,208	1,232	971	432	137	10	1
Male.....	60	118	218	343	483	620	624	445	190	45	1	1
Female.....	55	101	225	323	464	588	608	526	242	92	9	-
Black, both sexes.....	20	20	37	43	72	88	61	51	18	12	1	-
Male.....	13	11	22	25	45	52	33	36	11	5	-	-
Female.....	7	9	15	18	27	36	28	15	7	7	1	-
American Indian 1/, both sexes.....	-	2	-	1	2	6	3	1	-	-	-	-
Male.....	-	-	-	-	2	-	1	-	-	-	-	-
Female.....	-	2	-	1	-	6	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	3	5	6	8	3	1	2	-	-
Male.....	-	-	-	3	2	3	3	2	-	-	-	-
Female.....	-	-	1	-	3	3	5	1	1	2	-	-
Utah												
All races, both sexes.....	8	34	36	50	92	96	94	70	26	5	1	-
Male.....	4	14	22	30	60	52	57	36	12	1	-	-
Female.....	4	20	14	20	32	44	37	34	14	4	1	-
White, both sexes.....	8	34	35	47	90	94	92	69	25	4	1	-
Male.....	4	14	22	28	58	50	55	36	11	-	-	-
Female.....	4	20	13	19	32	44	37	33	14	4	1	-
Black, both sexes.....	-	-	-	1	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	2	-	1	1	-	-	-	-	-
Male.....	-	-	-	2	-	1	1	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	1	-	1	1	1	-	-
Male.....	-	-	-	-	2	1	-	-	1	1	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

Vermont

All races, both sexes.....	4	6	15	20	46	43	50	53	25	7	1	-
Male.....	1	2	9	8	23	18	29	26	11	-	-	-
Female.....	3	4	6	12	23	25	21	27	14	7	1	-
White, both sexes.....	4	6	15	20	46	42	50	53	25	7	1	-
Male.....	1	2	9	8	23	17	29	26	11	-	-	-
Female.....	3	4	6	12	23	25	21	27	14	7	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	39	72	149	223	301	479	470	366	171	47	6	-
Male.....	20	32	66	111	152	233	205	147	53	15	-	-
Female.....	19	40	83	112	149	246	265	219	118	32	6	-
White, both sexes.....	34	57	130	196	272	430	423	344	152	40	5	-
Male.....	18	24	56	95	140	198	180	137	47	12	-	-
Female.....	16	33	74	101	132	232	243	207	105	28	5	-
Black, both sexes.....	5	15	18	26	26	45	45	21	19	7	1	-
Male.....	2	8	10	15	11	31	25	9	6	3	-	-
Female.....	3	7	8	11	15	14	20	12	13	4	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	2	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	1	3	2	2	1	-	-	-	-
Male.....	-	-	-	1	1	2	-	1	-	-	-	-
Female.....	-	-	1	-	2	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other chronic obstructive pulmonary disease (J44)

Washington

All races, both sexes.....	43	85	146	231	346	423	421	387	165	53	5	-
Male.....	22	41	82	113	180	204	200	166	68	19	3	-
Female.....	21	44	64	118	166	219	221	221	97	34	2	-
White, both sexes.....	40	73	137	220	330	399	407	377	163	52	4	-
Male.....	20	33	78	106	170	187	193	162	67	18	3	-
Female.....	20	40	59	114	160	212	214	215	96	34	1	-
Black, both sexes.....	1	6	4	6	5	8	3	-	1	-	-	-
Male.....	-	2	1	3	2	5	2	-	1	-	-	-
Female.....	1	4	3	3	3	3	1	-	-	-	-	-
American Indian 1/, both sexes.....	2	3	5	3	4	4	1	3	-	-	-	-
Male.....	2	3	3	2	2	3	-	2	-	-	-	-
Female.....	-	-	2	1	2	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	3	-	2	7	12	10	7	1	1	1	-
Male.....	-	3	-	2	6	9	5	2	-	1	-	-
Female.....	-	-	-	-	1	3	5	5	1	-	1	-

West Virginia

All races, both sexes.....	25	52	85	133	173	206	233	163	81	16	2	-
Male.....	15	30	47	72	80	98	120	77	35	6	-	-
Female.....	10	22	38	61	93	108	113	86	46	10	2	-
White, both sexes.....	25	50	85	133	170	204	231	161	80	15	2	-
Male.....	15	30	47	72	80	96	119	75	34	6	-	-
Female.....	10	20	38	61	90	108	112	86	46	9	2	-
Black, both sexes.....	-	2	-	-	2	2	2	2	1	1	-	-
Male.....	-	-	-	-	-	2	1	2	1	-	-	-
Female.....	-	2	-	-	2	-	1	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
Arkansas												
All races, both sexes.....	55	-	-	-	-	2	-	2	1	-	5	3
Male.....	25	-	-	-	-	1	-	2	-	-	3	-
Female.....	30	-	-	-	-	1	-	-	1	-	2	3
White, both sexes.....	40	-	-	-	-	-	-	2	1	-	4	2
Male.....	22	-	-	-	-	-	-	2	-	-	3	-
Female.....	18	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes.....	15	-	-	-	-	2	-	-	-	-	1	1
Male.....	3	-	-	-	-	1	-	-	-	-	-	-
Female.....	12	-	-	-	-	1	-	-	-	-	1	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	419	-	4	6	4	7	6	9	4	15	21	35
Male.....	157	-	3	4	4	5	5	4	1	8	6	13
Female.....	262	-	1	2	-	2	1	5	3	7	15	22
White, both sexes.....	287	-	1	6	2	6	4	5	-	8	14	20
Male.....	99	-	1	4	2	4	4	2	-	5	5	8
Female.....	188	-	-	2	-	2	-	3	-	3	9	12
Black, both sexes.....	65	-	2	-	2	-	2	3	3	5	6	11
Male.....	25	-	1	-	2	-	1	1	1	3	1	3
Female.....	40	-	1	-	-	-	1	2	2	2	5	8
American Indian 1/, both sexes.....	3	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	64	-	1	-	-	1	-	1	1	2	1	4
Male.....	31	-	1	-	-	1	-	1	-	-	-	2
Female.....	33	-	-	-	-	-	-	-	1	2	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
New Mexico												
All races, both sexes.....	28	-	-	-	-	1	-	-	2	2	1	1
Male.....	12	-	-	-	-	-	-	-	2	1	1	-
Female.....	16	-	-	-	-	1	-	-	-	1	-	1
White, both sexes.....	25	-	-	-	-	1	-	-	2	1	1	1
Male.....	10	-	-	-	-	-	-	-	2	-	1	-
Female.....	15	-	-	-	-	1	-	-	-	1	-	1
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	-	1	-	-
Male.....	2	-	-	-	-	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	237	-	1	4	3	2	10	9	10	9	15	22
Male.....	101	-	1	1	2	1	4	9	7	3	6	12
Female.....	136	-	-	3	1	1	6	-	3	6	9	10
White, both sexes.....	153	-	-	2	-	1	7	5	7	5	9	11
Male.....	59	-	-	1	-	-	3	5	6	-	4	7
Female.....	94	-	-	1	-	1	4	-	1	5	5	4
Black, both sexes.....	77	-	1	2	3	1	2	4	3	3	6	11
Male.....	38	-	1	-	2	1	1	4	1	2	2	5
Female.....	39	-	-	2	1	-	1	-	2	1	4	6
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6	-	-	-	-	-	1	-	-	1	-	-
Male.....	3	-	-	-	-	-	-	-	-	1	-	-
Female.....	3	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Asthma (J45-J46)

Arkansas

All races, both sexes.....	8	3	3	6	2	4	6	6	3	1	-	-
Male.....	2	3	2	4	1	2	5	-	-	-	-	-
Female.....	6	-	1	2	1	2	1	6	3	1	-	-
White, both sexes.....	2	3	2	5	1	4	5	6	2	1	-	-
Male.....	2	3	1	3	1	2	5	-	-	-	-	-
Female.....	-	-	1	2	-	2	-	6	2	1	-	-
Black, both sexes.....	6	-	1	1	1	-	1	-	1	-	-	-
Male.....	-	-	1	1	-	-	-	-	-	-	-	-
Female.....	6	-	-	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	32	36	28	16	28	35	45	43	29	14	2	-
Male.....	11	18	11	4	13	11	16	9	8	2	1	-
Female.....	21	18	17	12	15	24	29	34	21	12	1	-
White, both sexes.....	18	24	20	11	19	27	31	33	25	11	2	-
Male.....	6	9	7	2	9	7	8	7	7	1	1	-
Female.....	12	15	13	9	10	20	23	26	18	10	1	-
Black, both sexes.....	6	9	6	2	5	1	-	2	-	-	-	-
Male.....	2	6	2	1	1	-	-	-	-	-	-	-
Female.....	4	3	4	1	4	1	-	2	-	-	-	-
American Indian 1/, both sexes.....	2	1	-	-	-	-	-	-	-	-	-	-
Male.....	1	1	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6	2	2	3	4	7	14	8	4	3	-	-
Male.....	2	2	2	1	3	4	8	2	1	1	-	-
Female.....	4	-	-	2	1	3	6	6	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Hawaii												
All races, both sexes.....	2	3	2	2	1	1	4	4	1	-	1	1
Male.....	1	2	1	1	-	-	-	3	-	-	1	1
Female.....	1	1	1	1	1	1	4	1	1	-	-	-
White, both sexes.....	1	-	1	1	-	-	1	-	-	-	-	1
Male.....	-	-	-	1	-	-	-	-	-	-	-	1
Female.....	1	-	1	-	-	-	1	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	1	3	1	1	1	1	3	4	1	-	1	-
Male.....	1	2	1	-	-	-	-	3	-	-	1	-
Female.....	-	1	-	1	1	1	3	1	1	-	-	-
Idaho												
All races, both sexes.....	1	-	2	1	2	1	4	5	-	2	-	-
Male.....	1	-	-	-	1	1	2	1	-	-	-	-
Female.....	-	-	2	1	1	-	2	4	-	2	-	-
White, both sexes.....	1	-	2	1	2	1	4	4	-	2	-	-
Male.....	1	-	-	-	1	1	2	1	-	-	-	-
Female.....	-	-	2	1	1	-	2	3	-	2	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Asthma (J45-J46)

Iowa

All races, both sexes.....	1	3	-	4	3	1	5	2	4	1	1	-
Male.....	1	2	-	1	-	-	4	1	1	1	-	-
Female.....	-	1	-	3	3	1	1	1	3	-	1	-
White, both sexes.....	1	3	-	4	3	1	5	2	4	1	1	-
Male.....	1	2	-	1	-	-	4	1	1	1	-	-
Female.....	-	1	-	3	3	1	1	1	3	-	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Kansas

All races, both sexes.....	1	1	-	2	4	2	2	2	2	4	1	-
Male.....	1	-	-	1	-	1	1	-	-	-	-	-
Female.....	-	1	-	1	4	1	1	2	2	4	1	-
White, both sexes.....	1	1	-	1	3	2	2	2	2	3	1	-
Male.....	1	-	-	-	-	1	1	-	-	-	-	-
Female.....	-	1	-	1	3	1	1	2	2	3	1	-
Black, both sexes.....	-	-	-	1	-	-	-	-	-	1	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Asthma (J45-J46)

Massachusetts

All races, both sexes.....	2	3	3	2	4	3	6	6	5	2	1	-
Male.....	-	1	3	2	1	2	2	1	1	1	-	-
Female.....	2	2	-	-	3	1	4	5	4	1	1	-
White, both sexes.....	-	3	3	1	3	3	6	6	5	2	1	-
Male.....	-	1	3	1	1	2	2	1	1	1	-	-
Female.....	-	2	-	-	2	1	4	5	4	1	1	-
Black, both sexes.....	2	-	-	1	1	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Michigan

All races, both sexes.....	15	10	6	7	8	11	11	6	8	3	-	-
Male.....	8	5	2	-	1	1	4	2	-	1	-	-
Female.....	7	5	4	7	7	10	7	4	8	2	-	-
White, both sexes.....	9	6	4	6	7	11	8	6	7	3	-	-
Male.....	5	4	2	-	1	1	4	2	-	1	-	-
Female.....	4	2	2	6	6	10	4	4	7	2	-	-
Black, both sexes.....	6	4	2	1	1	-	2	-	1	-	-	-
Male.....	3	1	-	-	-	-	-	-	-	-	-	-
Female.....	3	3	2	1	1	-	2	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
New Hampshire												
All races, both sexes.....	1	-	3	-	-	1	1	2	1	2	-	-
Male.....	-	-	1	-	-	1	-	-	1	1	-	-
Female.....	1	-	2	-	-	-	1	2	-	1	-	-
White, both sexes.....	1	-	3	-	-	1	1	2	1	2	-	-
Male.....	-	-	1	-	-	1	-	-	1	1	-	-
Female.....	1	-	2	-	-	-	1	2	-	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes.....	6	10	6	8	3	5	5	9	5	2	2	-
Male.....	-	2	1	5	1	-	1	3	1	-	-	-
Female.....	6	8	5	3	2	5	4	6	4	2	2	-
White, both sexes.....	-	7	5	3	3	5	5	8	4	-	1	-
Male.....	-	2	-	2	1	-	1	3	-	-	-	-
Female.....	-	5	5	1	2	5	4	5	4	-	1	-
Black, both sexes.....	6	3	1	3	-	-	-	-	-	2	1	-
Male.....	-	-	1	3	-	-	-	-	-	-	-	-
Female.....	6	3	-	-	-	-	-	-	-	2	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	-	-	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	2	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Asthma (J45-J46)

New Mexico

All races, both sexes.....	2	6	-	1	1	4	2	3	1	1	-	-
Male.....	1	2	-	-	1	1	1	1	-	1	-	-
Female.....	1	4	-	1	-	3	1	2	1	-	-	-
White, both sexes.....	1	6	-	1	1	4	2	3	1	-	-	-
Male.....	1	2	-	-	1	1	1	1	-	-	-	-
Female.....	-	4	-	1	-	3	1	2	1	-	-	-
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	1	-
Male.....	-	-	-	-	-	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	21	16	17	13	8	15	20	19	16	5	2	-
Male.....	8	6	10	10	3	4	7	4	3	-	-	-
Female.....	13	10	7	3	5	11	13	15	13	5	2	-
White, both sexes.....	13	10	8	8	5	11	16	14	15	4	2	-
Male.....	5	4	4	5	2	3	5	2	3	-	-	-
Female.....	8	6	4	3	3	8	11	12	12	4	2	-
Black, both sexes.....	7	5	8	5	3	3	4	4	1	1	-	-
Male.....	3	1	5	5	1	-	2	2	-	-	-	-
Female.....	4	4	3	-	2	3	2	2	1	1	-	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	-	-	1	-	1	-	-	-	-
Male.....	-	1	-	-	-	1	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Asthma (J45-J46)

Oregon

All races, both sexes.....	6	2	4	6	2	10	8	3	8	3	-	-
Male.....	2	1	1	2	1	4	2	-	2	-	-	-
Female.....	4	1	3	4	1	6	6	3	6	3	-	-
White, both sexes.....	5	2	3	5	1	10	8	3	8	3	-	-
Male.....	2	1	1	2	-	4	2	-	2	-	-	-
Female.....	3	1	2	3	1	6	6	3	6	3	-	-
Black, both sexes.....	1	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	12	10	10	9	6	6	11	15	9	6	-	1
Male.....	4	4	2	3	2	1	3	3	-	-	-	1
Female.....	8	6	8	6	4	5	8	12	9	6	-	-
White, both sexes.....	10	6	8	7	5	6	11	14	7	6	-	-
Male.....	3	4	1	3	2	1	3	3	-	-	-	-
Female.....	7	2	7	4	3	5	8	11	7	6	-	-
Black, both sexes.....	2	4	2	2	1	-	-	1	1	-	-	-
Male.....	1	-	1	-	-	-	-	-	-	-	-	-
Female.....	1	4	1	2	1	-	-	1	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	1
Male.....	-	-	-	-	-	-	-	-	-	-	-	1
Female.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Asthma (J45-J46)

Rhode Island

All races, both sexes.....	-	-	1	-	1	-	2	2	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	1	-	1	-	2	2	-	-	-	-
White, both sexes.....	-	-	1	-	1	-	2	2	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	1	-	1	-	2	2	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

South Carolina

All races, both sexes.....	6	6	5	4	4	4	4	9	-	4	-	-
Male.....	3	2	2	2	1	3	-	-	-	1	-	-
Female.....	3	4	3	2	3	1	4	9	-	3	-	-
White, both sexes.....	3	2	4	2	1	2	2	6	-	2	-	-
Male.....	-	-	1	1	-	1	-	-	-	-	-	-
Female.....	3	2	3	1	1	1	2	6	-	2	-	-
Black, both sexes.....	3	4	1	1	3	2	1	3	-	2	-	-
Male.....	3	2	1	1	1	2	-	-	-	1	-	-
Female.....	-	2	-	-	2	-	1	3	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)												
Vermont												
All races, both sexes.....	1	-	-	-	-	1	1	1	2	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	1	1	2	-	-	-
White, both sexes.....	1	-	-	-	-	1	1	1	2	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	1	1	2	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes.....	4	6	2	9	5	8	15	4	3	3	-	-
Male.....	2	3	1	1	2	1	2	-	1	-	-	-
Female.....	2	3	1	8	3	7	13	4	2	3	-	-
White, both sexes.....	1	-	1	6	2	6	9	4	3	2	-	-
Male.....	1	-	-	1	1	-	1	-	1	-	-	-
Female.....	-	-	1	5	1	6	8	4	2	2	-	-
Black, both sexes.....	3	6	1	3	3	1	5	-	-	1	-	-
Male.....	1	3	1	-	1	1	1	-	-	-	-	-
Female.....	2	3	-	3	2	-	4	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Bronchiectasis (J47)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Bronchiectasis (J47)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Bronchiectasis (J47)

Wisconsin

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Bronchiectasis (J47)

Arkansas

All races, both sexes.....	-	1	-	1	-	-	2	2	2	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	2	2	2	-	-	-
White, both sexes.....	-	1	-	1	-	-	2	2	2	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	2	2	2	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	3	1	2	11	8	16	24	31	12	3	-	-
Male.....	2	-	1	7	1	6	5	8	3	1	-	-
Female.....	1	1	1	4	7	10	19	23	9	2	-	-
White, both sexes.....	2	-	2	7	5	12	21	27	10	3	-	-
Male.....	2	-	1	5	-	4	5	7	2	1	-	-
Female.....	-	-	1	2	5	8	16	20	8	2	-	-
Black, both sexes.....	-	1	-	1	1	-	-	-	2	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	1	-	1	1	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	3	2	4	3	4	-	-	-	-
Male.....	-	-	-	2	1	2	-	1	-	-	-	-
Female.....	1	-	-	1	1	2	3	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lung diseases due to external agents (J60-J70)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Lung diseases due to external agents (J60-J70)

Alaska

All races, both sexes.....	2	-	-	2	2	6	4	1	4	-	-	-
Male.....	-	-	-	-	2	4	2	-	-	-	-	-
Female.....	2	-	-	2	-	2	2	1	4	-	-	-
White, both sexes.....	1	-	-	-	2	3	4	1	3	-	-	-
Male.....	-	-	-	-	2	2	2	-	-	-	-	-
Female.....	1	-	-	-	-	1	2	1	3	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	1	-	3	-	-	1	-	-	-
Male.....	-	-	-	-	-	2	-	-	-	-	-	-
Female.....	1	-	-	1	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	2	5	12	8	21	37	44	57	32	22	3	-
Male.....	1	2	8	5	15	26	31	34	19	13	1	-
Female.....	1	3	4	3	6	11	13	23	13	9	2	-
White, both sexes.....	2	4	10	8	20	34	43	53	31	19	3	-
Male.....	1	2	8	5	15	24	31	32	18	12	1	-
Female.....	1	2	2	3	5	10	12	21	13	7	2	-
Black, both sexes.....	-	-	1	-	-	2	1	1	-	1	-	-
Male.....	-	-	-	-	-	1	-	1	-	-	-	-
Female.....	-	-	1	-	-	1	1	-	-	1	-	-
American Indian 1/, both sexes.....	-	1	1	-	1	-	-	2	1	2	-	-
Male.....	-	-	-	-	-	-	-	1	1	1	-	-
Female.....	-	1	1	-	1	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Lung diseases due to external agents (J60-J70)

Arkansas

All races, both sexes.....	2	4	8	6	16	25	41	57	29	15	1	-
Male.....	2	2	4	4	12	19	20	32	13	3	-	-
Female.....	-	2	4	2	4	6	21	25	16	12	1	-
White, both sexes.....	2	3	6	5	14	23	39	49	27	12	1	-
Male.....	2	2	3	3	10	17	19	28	13	2	-	-
Female.....	-	1	3	2	4	6	20	21	14	10	1	-
Black, both sexes.....	-	1	2	1	2	2	2	8	2	3	-	-
Male.....	-	-	1	1	2	2	1	4	-	1	-	-
Female.....	-	1	1	-	-	-	1	4	2	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	18	26	23	39	67	108	177	215	164	61	15	-
Male.....	11	18	13	21	44	66	102	124	77	21	6	-
Female.....	7	8	10	18	23	42	75	91	87	40	9	-
White, both sexes.....	15	20	21	35	57	96	159	194	152	54	10	-
Male.....	8	13	12	18	36	64	94	114	70	17	3	-
Female.....	7	7	9	17	21	32	65	80	82	37	7	-
Black, both sexes.....	2	4	2	1	6	5	10	8	4	3	2	-
Male.....	2	3	1	1	4	-	4	5	2	1	1	-
Female.....	-	1	1	-	2	5	6	3	2	2	1	-
American Indian 1/, both sexes.....	-	1	-	1	-	2	-	1	-	-	-	-
Male.....	-	1	-	1	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	2	4	5	8	12	8	4	3	-
Male.....	1	1	-	1	4	1	4	5	5	3	2	-
Female.....	-	-	-	1	-	4	4	7	3	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Lung diseases due to external agents (J60-J70)

Colorado

All races, both sexes.....	5	7	11	4	25	23	50	70	54	20	3	-
Male.....	3	7	8	2	18	17	31	40	26	7	1	-
Female.....	2	-	3	2	7	6	19	30	28	13	2	-
White, both sexes.....	5	7	9	4	25	22	49	70	54	19	3	-
Male.....	3	7	7	2	18	16	30	40	26	7	1	-
Female.....	2	-	2	2	7	6	19	30	28	12	2	-
Black, both sexes.....	-	-	2	-	-	-	-	-	-	1	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Connecticut

All races, both sexes.....	4	4	11	15	19	31	61	108	83	25	8	-
Male.....	3	4	6	11	12	21	37	55	41	9	1	-
Female.....	1	-	5	4	7	10	24	53	42	16	7	-
White, both sexes.....	4	4	10	13	19	30	59	104	82	24	8	-
Male.....	3	4	6	9	12	20	36	54	41	9	1	-
Female.....	1	-	4	4	7	10	23	50	41	15	7	-
Black, both sexes.....	-	-	1	2	-	-	1	4	-	1	-	-
Male.....	-	-	-	2	-	-	1	1	-	-	-	-
Female.....	-	-	1	-	-	-	-	3	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	-	1	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Lung diseases due to external agents (J60-J70)

Kentucky

All races, both sexes.....	7	7	13	13	31	47	77	83	70	25	7	-
Male.....	5	3	7	7	23	25	42	38	33	6	1	-
Female.....	2	4	6	6	8	22	35	45	37	19	6	-
White, both sexes.....	5	7	10	13	30	46	68	81	65	25	6	-
Male.....	4	3	5	7	22	24	39	37	30	6	1	-
Female.....	1	4	5	6	8	22	29	44	35	19	5	-
Black, both sexes.....	2	-	3	-	1	1	9	2	5	-	1	-
Male.....	1	-	2	-	1	1	3	1	3	-	-	-
Female.....	1	-	1	-	-	-	6	1	2	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Louisiana

All races, both sexes.....	4	9	7	15	14	46	47	55	31	14	2	-
Male.....	4	8	3	8	9	23	27	31	15	8	-	-
Female.....	-	1	4	7	5	23	20	24	16	6	2	-
White, both sexes.....	3	5	3	9	11	40	40	42	26	11	2	-
Male.....	3	5	1	4	7	19	25	24	14	7	-	-
Female.....	-	-	2	5	4	21	15	18	12	4	2	-
Black, both sexes.....	1	4	4	6	3	6	6	12	5	3	-	-
Male.....	1	3	2	4	2	4	2	6	1	1	-	-
Female.....	-	1	2	2	1	2	4	6	4	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Lung diseases due to external agents (J60-J70)

Maine

All races, both sexes.....	2	3	4	6	7	12	18	25	19	7	2	-
Male.....	1	3	2	6	5	9	12	12	9	3	-	-
Female.....	1	-	2	-	2	3	6	13	10	4	2	-
White, both sexes.....	2	3	4	6	7	12	18	25	19	7	2	-
Male.....	1	3	2	6	5	9	12	12	9	3	-	-
Female.....	1	-	2	-	2	3	6	13	10	4	2	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	8	11	11	14	23	33	80	104	68	20	1	-
Male.....	7	9	7	6	14	22	45	54	37	11	-	-
Female.....	1	2	4	8	9	11	35	50	31	9	1	-
White, both sexes.....	6	6	9	8	15	26	60	91	59	19	1	-
Male.....	6	5	6	3	8	20	33	48	34	11	-	-
Female.....	-	1	3	5	7	6	27	43	25	8	1	-
Black, both sexes.....	2	5	2	6	8	7	19	12	9	1	-	-
Male.....	1	4	1	3	6	2	11	6	3	-	-	-
Female.....	1	1	1	3	2	5	8	6	6	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Lung diseases due to external agents (J60-J70)

Massachusetts

All races, both sexes.....	4	14	16	15	36	67	97	133	99	33	6	-
Male.....	4	10	9	13	21	43	63	74	49	10	2	-
Female.....	-	4	7	2	15	24	34	59	50	23	4	-
White, both sexes.....	4	14	16	13	35	65	92	129	94	33	6	-
Male.....	4	10	9	11	20	41	59	73	44	10	2	-
Female.....	-	4	7	2	15	24	33	56	50	23	4	-
Black, both sexes.....	-	-	-	2	1	1	4	3	1	-	-	-
Male.....	-	-	-	2	1	1	3	1	1	-	-	-
Female.....	-	-	-	-	-	-	1	2	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	1	4	-	-	-
Male.....	-	-	-	-	-	1	1	-	4	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Michigan

All races, both sexes.....	10	14	16	26	36	68	108	116	90	41	10	-
Male.....	8	8	4	17	24	43	57	64	48	13	3	-
Female.....	2	6	12	9	12	25	51	52	42	28	7	-
White, both sexes.....	7	9	11	24	32	58	93	101	85	38	7	-
Male.....	7	6	3	17	22	37	51	56	44	12	3	-
Female.....	-	3	8	7	10	21	42	45	41	26	4	-
Black, both sexes.....	3	5	5	2	3	9	13	12	4	3	3	-
Male.....	1	2	1	-	2	6	5	5	3	1	-	-
Female.....	2	3	4	2	1	3	8	7	1	2	3	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	1	2	2	1	-	-	-
Male.....	-	-	-	-	-	-	1	2	1	-	-	-
Female.....	-	-	-	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Lung diseases due to external agents (J60-J70)

New Hampshire

All races, both sexes.....	-	1	4	5	7	3	7	10	18	4	-	-
Male.....	-	1	1	2	4	-	7	1	11	1	-	-
Female.....	-	-	3	3	3	3	-	9	7	3	-	-
White, both sexes.....	-	1	4	5	7	3	7	10	18	4	-	-
Male.....	-	1	1	2	4	-	7	1	11	1	-	-
Female.....	-	-	3	3	3	3	-	9	7	3	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	4	11	13	17	33	75	112	150	113	47	5	-
Male.....	3	5	8	14	22	47	58	80	56	19	1	-
Female.....	1	6	5	3	11	28	54	70	57	28	4	-
White, both sexes.....	2	7	12	17	27	63	108	142	106	43	4	-
Male.....	2	4	8	14	18	37	57	78	53	18	1	-
Female.....	-	3	4	3	9	26	51	64	53	25	3	-
Black, both sexes.....	2	4	1	-	6	12	4	7	3	4	1	-
Male.....	1	1	-	-	4	10	1	2	1	1	-	-
Female.....	1	3	1	-	2	2	3	5	2	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	4	-	-	-
Male.....	-	-	-	-	-	-	-	-	2	-	-	-
Female.....	-	-	-	-	-	-	-	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Lung diseases due to external agents (J60-J70)

New Mexico

All races, both sexes.....	2	4	2	6	6	10	21	15	8	5	2	-
Male.....	1	3	1	3	4	7	10	7	4	2	2	-
Female.....	1	1	1	3	2	3	11	8	4	3	-	-
White, both sexes.....	1	4	1	6	4	9	19	14	8	4	1	-
Male.....	1	3	1	3	3	7	9	7	4	2	1	-
Female.....	-	1	-	3	1	2	10	7	4	2	-	-
Black, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	2	-	2	1	-	1	1	-
Male.....	-	-	-	-	1	-	1	-	-	-	1	-
Female.....	-	-	1	-	1	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	9	16	20	22	45	82	160	201	121	62	13	-
Male.....	6	10	10	14	30	51	86	110	63	22	3	-
Female.....	3	6	10	8	15	31	74	91	58	40	10	-
White, both sexes.....	8	13	17	19	39	79	151	190	118	60	11	-
Male.....	5	10	8	12	25	49	83	106	61	21	3	-
Female.....	3	3	9	7	14	30	68	84	57	39	8	-
Black, both sexes.....	1	3	3	2	6	2	7	11	3	2	2	-
Male.....	1	-	2	2	5	1	2	4	2	1	-	-
Female.....	-	3	1	-	1	1	5	7	1	1	2	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	2	-	-	-	-	-
Male.....	-	-	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lung diseases due to external agents (J60-J70)												
Oregon												
All races, both sexes.....	4	7	5	10	14	21	35	56	37	16	3	-
Male.....	2	6	5	8	7	15	17	33	18	8	-	-
Female.....	2	1	-	2	7	6	18	23	19	8	3	-
White, both sexes.....	4	7	5	9	11	21	34	54	36	16	3	-
Male.....	2	6	5	8	5	15	16	33	17	8	-	-
Female.....	2	1	-	1	6	6	18	21	19	8	3	-
Black, both sexes.....	-	-	-	-	1	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	-	1	1	1	-	-	-
Male.....	-	-	-	-	2	-	1	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes.....	11	19	37	50	74	132	243	328	205	73	8	-
Male.....	5	12	19	32	49	84	133	198	95	25	3	-
Female.....	6	7	18	18	25	48	110	130	110	48	5	-
White, both sexes.....	7	18	36	39	68	122	227	313	192	68	7	-
Male.....	4	12	18	23	45	78	123	190	90	24	3	-
Female.....	3	6	18	16	23	44	104	123	102	44	4	-
Black, both sexes.....	4	1	1	11	6	9	16	14	11	5	1	-
Male.....	1	-	1	9	4	5	10	8	4	1	-	-
Female.....	3	1	-	2	2	4	6	6	7	4	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	1	2	-	-	-
Male.....	-	-	-	-	-	1	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Lung diseases due to external agents (J60-J70)

Texas

All races, both sexes.....	21	39	42	46	77	140	232	256	195	76	19	-
Male.....	11	21	23	28	46	95	113	144	88	19	4	-
Female.....	10	18	19	18	31	45	119	112	107	57	15	-
White, both sexes.....	20	30	37	39	65	129	214	241	185	67	19	-
Male.....	10	15	20	24	38	86	102	137	84	18	4	-
Female.....	10	15	17	15	27	43	112	104	101	49	15	-
Black, both sexes.....	1	8	4	7	12	10	17	11	9	9	-	-
Male.....	1	5	2	4	8	8	10	5	3	1	-	-
Female.....	-	3	2	3	4	2	7	6	6	8	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	-	-	1	1	3	1	-	-	-
Male.....	-	1	1	-	-	1	1	1	1	-	-	-
Female.....	-	-	-	-	-	-	-	2	-	-	-	-

Utah

All races, both sexes.....	4	3	5	2	5	9	18	24	22	3	-	-
Male.....	4	2	4	2	4	4	8	10	14	1	-	-
Female.....	-	1	1	-	1	5	10	14	8	2	-	-
White, both sexes.....	4	3	5	2	5	9	18	23	22	3	-	-
Male.....	4	2	4	2	4	4	8	10	14	1	-	-
Female.....	-	1	1	-	1	5	10	13	8	2	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pneumoconioses and chemical effects (J60-J66,J68)

New Mexico

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonitis due to solids and liquids (J69)

Alaska

All races, both sexes.....	2	-	-	2	2	6	4	1	4	-	-	-
Male.....	-	-	-	-	2	4	2	-	-	-	-	-
Female.....	2	-	-	2	-	2	2	1	4	-	-	-
White, both sexes.....	1	-	-	-	2	3	4	1	3	-	-	-
Male.....	-	-	-	-	2	2	2	-	-	-	-	-
Female.....	1	-	-	-	-	1	2	1	3	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	1	-	3	-	-	1	-	-	-
Male.....	-	-	-	-	-	2	-	-	-	-	-	-
Female.....	1	-	-	1	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	2	5	12	7	15	36	38	55	31	22	3	-
Male.....	1	2	8	4	9	25	25	32	18	13	1	-
Female.....	1	3	4	3	6	11	13	23	13	9	2	-
White, both sexes.....	2	4	10	7	14	33	37	51	30	19	3	-
Male.....	1	2	8	4	9	23	25	30	17	12	1	-
Female.....	1	2	2	3	5	10	12	21	13	7	2	-
Black, both sexes.....	-	-	1	-	-	2	1	1	-	1	-	-
Male.....	-	-	-	-	-	1	-	1	-	-	-	-
Female.....	-	-	1	-	-	1	1	-	-	1	-	-
American Indian 1/, both sexes.....	-	1	1	-	1	-	-	2	1	2	-	-
Male.....	-	-	-	-	-	-	-	1	1	1	-	-
Female.....	-	1	1	-	1	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to solids and liquids (J69)

Arkansas

All races, both sexes.....	2	4	7	4	15	22	39	57	29	15	1	-
Male.....	2	2	3	2	11	16	19	32	13	3	-	-
Female.....	-	2	4	2	4	6	20	25	16	12	1	-
White, both sexes.....	2	3	5	4	13	21	37	49	27	12	1	-
Male.....	2	2	2	2	9	15	18	28	13	2	-	-
Female.....	-	1	3	2	4	6	19	21	14	10	1	-
Black, both sexes.....	-	1	2	-	2	1	2	8	2	3	-	-
Male.....	-	-	1	-	2	1	1	4	-	1	-	-
Female.....	-	1	1	-	-	-	1	4	2	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	18	23	18	37	62	97	164	209	159	61	15	-
Male.....	11	16	10	19	41	55	90	118	74	21	6	-
Female.....	7	7	8	18	21	42	74	91	85	40	9	-
White, both sexes.....	15	17	16	33	52	86	146	188	147	54	10	-
Male.....	8	11	9	16	33	54	82	108	67	17	3	-
Female.....	7	6	7	17	19	32	64	80	80	37	7	-
Black, both sexes.....	2	4	2	1	6	5	10	8	4	3	2	-
Male.....	2	3	1	1	4	-	4	5	2	1	1	-
Female.....	-	1	1	-	2	5	6	3	2	2	1	-
American Indian 1/, both sexes.....	-	1	-	1	-	1	-	1	-	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	2	4	5	8	12	8	4	3	-
Male.....	1	1	-	1	4	1	4	5	5	3	2	-
Female.....	-	-	-	1	-	4	4	7	3	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to solids and liquids (J69)

Colorado

All races, both sexes.....	5	7	10	3	21	19	45	69	52	19	3	-
Male.....	3	7	7	2	14	13	27	39	24	6	1	-
Female.....	2	-	3	1	7	6	18	30	28	13	2	-
White, both sexes.....	5	7	8	3	21	18	44	69	52	18	3	-
Male.....	3	7	6	2	14	12	26	39	24	6	1	-
Female.....	2	-	2	1	7	6	18	30	28	12	2	-
Black, both sexes.....	-	-	2	-	-	-	-	-	-	1	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Connecticut

All races, both sexes.....	4	4	11	15	19	30	60	108	83	25	8	-
Male.....	3	4	6	11	12	20	36	55	41	9	1	-
Female.....	1	-	5	4	7	10	24	53	42	16	7	-
White, both sexes.....	4	4	10	13	19	29	58	104	82	24	8	-
Male.....	3	4	6	9	12	19	35	54	41	9	1	-
Female.....	1	-	4	4	7	10	23	50	41	15	7	-
Black, both sexes.....	-	-	1	2	-	-	1	4	-	1	-	-
Male.....	-	-	-	2	-	-	1	1	-	-	-	-
Female.....	-	-	1	-	-	-	-	3	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	-	1	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to solids and liquids (J69)

Kentucky

All races, both sexes.....	5	6	12	9	28	42	68	75	67	25	7	-
Male.....	3	2	6	5	20	20	33	30	30	6	1	-
Female.....	2	4	6	4	8	22	35	45	37	19	6	-
White, both sexes.....	3	6	9	9	27	41	60	73	62	25	6	-
Male.....	2	2	4	5	19	19	31	29	27	6	1	-
Female.....	1	4	5	4	8	22	29	44	35	19	5	-
Black, both sexes.....	2	-	3	-	1	1	8	2	5	-	1	-
Male.....	1	-	2	-	1	1	2	1	3	-	-	-
Female.....	1	-	1	-	-	-	6	1	2	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Louisiana

All races, both sexes.....	4	9	7	13	14	43	45	53	30	14	2	-
Male.....	4	8	3	6	9	20	25	29	14	8	-	-
Female.....	-	1	4	7	5	23	20	24	16	6	2	-
White, both sexes.....	3	5	3	8	11	37	38	40	25	11	2	-
Male.....	3	5	1	3	7	16	23	22	13	7	-	-
Female.....	-	-	2	5	4	21	15	18	12	4	2	-
Black, both sexes.....	1	4	4	5	3	6	6	12	5	3	-	-
Male.....	1	3	2	3	2	4	2	6	1	1	-	-
Female.....	-	1	2	2	1	2	4	6	4	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to solids and liquids (J69)

Maine

All races, both sexes.....	2	2	4	4	3	11	15	21	17	7	2	-
Male.....	1	2	2	4	2	8	9	8	7	3	-	-
Female.....	1	-	2	-	1	3	6	13	10	4	2	-
White, both sexes.....	2	2	4	4	3	11	15	21	17	7	2	-
Male.....	1	2	2	4	2	8	9	8	7	3	-	-
Female.....	1	-	2	-	1	3	6	13	10	4	2	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	8	11	10	14	21	31	74	103	66	20	1	-
Male.....	7	9	7	6	13	20	40	54	35	11	-	-
Female.....	1	2	3	8	8	11	34	49	31	9	1	-
White, both sexes.....	6	6	9	8	13	24	54	90	57	19	1	-
Male.....	6	5	6	3	7	18	28	48	32	11	-	-
Female.....	-	1	3	5	6	6	26	42	25	8	1	-
Black, both sexes.....	2	5	1	6	8	7	19	12	9	1	-	-
Male.....	1	4	1	3	6	2	11	6	3	-	-	-
Female.....	1	1	-	3	2	5	8	6	6	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to solids and liquids (J69)

Massachusetts

All races, both sexes.....	3	14	16	15	34	65	90	130	98	33	6	-
Male.....	3	10	9	13	19	41	57	71	48	10	2	-
Female.....	-	4	7	2	15	24	33	59	50	23	4	-
White, both sexes.....	3	14	16	13	33	63	85	126	93	33	6	-
Male.....	3	10	9	11	18	39	53	70	43	10	2	-
Female.....	-	4	7	2	15	24	32	56	50	23	4	-
Black, both sexes.....	-	-	-	2	1	1	4	3	1	-	-	-
Male.....	-	-	-	2	1	1	3	1	1	-	-	-
Female.....	-	-	-	-	-	-	1	2	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	1	4	-	-	-
Male.....	-	-	-	-	-	1	1	-	4	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Michigan

All races, both sexes.....	10	14	16	24	33	61	106	113	89	40	10	-
Male.....	8	8	4	16	21	37	55	61	47	13	3	-
Female.....	2	6	12	8	12	24	51	52	42	27	7	-
White, both sexes.....	7	9	11	23	29	52	91	98	84	37	7	-
Male.....	7	6	3	16	19	31	49	53	43	12	3	-
Female.....	-	3	8	7	10	21	42	45	41	25	4	-
Black, both sexes.....	3	5	5	1	3	8	13	12	4	3	3	-
Male.....	1	2	1	-	2	6	5	5	3	1	-	-
Female.....	2	3	4	1	1	2	8	7	1	2	3	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	1	2	2	1	-	-	-
Male.....	-	-	-	-	-	-	1	2	1	-	-	-
Female.....	-	-	-	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to solids and liquids (J69)

New Hampshire

All races, both sexes.....	-	1	4	5	7	3	7	10	18	4	-	-
Male.....	-	1	1	2	4	-	7	1	11	1	-	-
Female.....	-	-	3	3	3	3	-	9	7	3	-	-
White, both sexes.....	-	1	4	5	7	3	7	10	18	4	-	-
Male.....	-	1	1	2	4	-	7	1	11	1	-	-
Female.....	-	-	3	3	3	3	-	9	7	3	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	4	11	10	17	29	70	103	142	107	46	5	-
Male.....	3	5	5	14	18	42	50	74	53	18	1	-
Female.....	1	6	5	3	11	28	53	68	54	28	4	-
White, both sexes.....	2	7	9	17	23	60	99	134	100	42	4	-
Male.....	2	4	5	14	14	34	49	72	50	17	1	-
Female.....	-	3	4	3	9	26	50	62	50	25	3	-
Black, both sexes.....	2	4	1	-	6	10	4	7	3	4	1	-
Male.....	1	1	-	-	4	8	1	2	1	1	-	-
Female.....	1	3	1	-	2	2	3	5	2	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	4	-	-	-
Male.....	-	-	-	-	-	-	-	-	2	-	-	-
Female.....	-	-	-	-	-	-	-	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
New Mexico												
All races, both sexes.....	2	4	2	5	5	10	20	15	7	5	2	-
Male.....	1	3	1	2	3	7	9	7	3	2	2	-
Female.....	1	1	1	3	2	3	11	8	4	3	-	-
White, both sexes.....	1	4	1	5	3	9	18	14	7	4	1	-
Male.....	1	3	1	2	2	7	8	7	3	2	1	-
Female.....	-	1	-	3	1	2	10	7	4	2	-	-
Black, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	2	-	2	1	-	1	1	-
Male.....	-	-	-	-	1	-	1	-	-	-	1	-
Female.....	-	-	1	-	1	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	9	16	18	21	41	72	151	190	118	60	13	-
Male.....	6	10	9	13	26	43	78	100	60	21	3	-
Female.....	3	6	9	8	15	29	73	90	58	39	10	-
White, both sexes.....	8	13	16	18	36	70	142	180	115	58	11	-
Male.....	5	10	8	11	22	42	75	97	58	20	3	-
Female.....	3	3	8	7	14	28	67	83	57	38	8	-
Black, both sexes.....	1	3	2	2	5	1	7	10	3	2	2	-
Male.....	1	-	1	2	4	-	2	3	2	1	-	-
Female.....	-	3	1	-	1	1	5	7	1	1	2	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	2	-	-	-	-	-
Male.....	-	-	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)												
Oregon												
All races, both sexes.....	4	7	4	8	13	18	32	53	36	15	3	-
Male.....	2	6	4	6	6	12	14	30	17	7	-	-
Female.....	2	1	-	2	7	6	18	23	19	8	3	-
White, both sexes.....	4	7	4	7	10	18	31	51	35	15	3	-
Male.....	2	6	4	6	4	12	13	30	16	7	-	-
Female.....	2	1	-	1	6	6	18	21	19	8	3	-
Black, both sexes.....	-	-	-	-	1	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	-	1	1	1	-	-	-
Male.....	-	-	-	-	2	-	1	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes.....	11	17	33	45	69	112	215	283	194	71	8	-
Male.....	5	10	15	27	45	65	106	153	84	23	3	-
Female.....	6	7	18	18	24	47	109	130	110	48	5	-
White, both sexes.....	7	16	32	35	63	103	200	268	181	66	7	-
Male.....	4	10	14	19	41	60	96	145	79	22	3	-
Female.....	3	6	18	16	22	43	104	123	102	44	4	-
Black, both sexes.....	4	1	1	10	6	8	15	14	11	5	1	-
Male.....	1	-	1	8	4	4	10	8	4	1	-	-
Female.....	3	1	-	2	2	4	5	6	7	4	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	1	2	-	-	-
Male.....	-	-	-	-	-	1	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Pneumonitis due to solids and liquids (J69)

Rhode Island

Table with 13 columns (Age groups) and 13 rows (Race categories) for Rhode Island. Rows include All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.

South Carolina

Table with 13 columns (Age groups) and 13 rows (Race categories) for South Carolina. Rows include All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to solids and liquids (J69)

Texas

All races, both sexes.....	19	36	40	39	75	127	225	246	191	76	19	-
Male.....	9	18	21	23	44	83	106	136	84	19	4	-
Female.....	10	18	19	16	31	44	119	110	107	57	15	-
White, both sexes.....	18	27	35	32	63	117	210	231	181	67	19	-
Male.....	8	12	18	19	36	75	98	129	80	18	4	-
Female.....	10	15	17	13	27	42	112	102	101	49	15	-
Black, both sexes.....	1	8	4	7	12	9	14	11	9	9	-	-
Male.....	1	5	2	4	8	7	7	5	3	1	-	-
Female.....	-	3	2	3	4	2	7	6	6	8	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	-	-	1	1	3	1	-	-	-
Male.....	-	1	1	-	-	1	1	1	1	-	-	-
Female.....	-	-	-	-	-	-	-	2	-	-	-	-

Utah

All races, both sexes.....	4	1	4	1	3	9	15	23	22	3	-	-
Male.....	4	-	3	1	2	4	5	9	14	1	-	-
Female.....	-	1	1	-	1	5	10	14	8	2	-	-
White, both sexes.....	4	1	4	1	3	9	15	22	22	3	-	-
Male.....	4	-	3	1	2	4	5	9	14	1	-	-
Female.....	-	1	1	-	1	5	10	13	8	2	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Pneumonitis due to solids and liquids (J69)

Washington

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.

West Virginia

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to food and vomit (J69.0)

Alaska

All races, both sexes.....	2	-	-	2	2	6	4	1	4	-	-	-
Male.....	-	-	-	-	2	4	2	-	-	-	-	-
Female.....	2	-	-	2	-	2	2	1	4	-	-	-
White, both sexes.....	1	-	-	-	2	3	4	1	3	-	-	-
Male.....	-	-	-	-	2	2	2	-	-	-	-	-
Female.....	1	-	-	-	-	1	2	1	3	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	1	-	3	-	-	1	-	-	-
Male.....	-	-	-	-	-	2	-	-	-	-	-	-
Female.....	1	-	-	1	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	2	5	12	7	15	36	38	55	31	22	3	-
Male.....	1	2	8	4	9	25	25	32	18	13	1	-
Female.....	1	3	4	3	6	11	13	23	13	9	2	-
White, both sexes.....	2	4	10	7	14	33	37	51	30	19	3	-
Male.....	1	2	8	4	9	23	25	30	17	12	1	-
Female.....	1	2	2	3	5	10	12	21	13	7	2	-
Black, both sexes.....	-	-	1	-	-	2	1	1	-	1	-	-
Male.....	-	-	-	-	-	1	-	1	-	-	-	-
Female.....	-	-	1	-	-	1	1	-	-	1	-	-
American Indian 1/, both sexes.....	-	1	1	-	1	-	-	2	1	2	-	-
Male.....	-	-	-	-	-	-	-	1	1	1	-	-
Female.....	-	1	1	-	1	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to food and vomit (J69.0)

Arkansas

All races, both sexes.....	2	4	7	4	15	22	39	57	29	15	1	-
Male.....	2	2	3	2	11	16	19	32	13	3	-	-
Female.....	-	2	4	2	4	6	20	25	16	12	1	-
White, both sexes.....	2	3	5	4	13	21	37	49	27	12	1	-
Male.....	2	2	2	2	9	15	18	28	13	2	-	-
Female.....	-	1	3	2	4	6	19	21	14	10	1	-
Black, both sexes.....	-	1	2	-	2	1	2	8	2	3	-	-
Male.....	-	-	1	-	2	1	1	4	-	1	-	-
Female.....	-	1	1	-	-	-	1	4	2	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	18	23	18	37	62	97	164	209	159	61	15	-
Male.....	11	16	10	19	41	55	90	118	74	21	6	-
Female.....	7	7	8	18	21	42	74	91	85	40	9	-
White, both sexes.....	15	17	16	33	52	86	146	188	147	54	10	-
Male.....	8	11	9	16	33	54	82	108	67	17	3	-
Female.....	7	6	7	17	19	32	64	80	80	37	7	-
Black, both sexes.....	2	4	2	1	6	5	10	8	4	3	2	-
Male.....	2	3	1	1	4	-	4	5	2	1	1	-
Female.....	-	1	1	-	2	5	6	3	2	2	1	-
American Indian 1/, both sexes.....	-	1	-	1	-	1	-	1	-	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	2	4	5	8	12	8	4	3	-
Male.....	1	1	-	1	4	1	4	5	5	3	2	-
Female.....	-	-	-	1	-	4	4	7	3	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to food and vomit (J69.0)

Colorado

All races, both sexes.....	5	7	10	3	21	19	44	69	52	19	3	-
Male.....	3	7	7	2	14	13	26	39	24	6	1	-
Female.....	2	-	3	1	7	6	18	30	28	13	2	-
White, both sexes.....	5	7	8	3	21	18	43	69	52	18	3	-
Male.....	3	7	6	2	14	12	25	39	24	6	1	-
Female.....	2	-	2	1	7	6	18	30	28	12	2	-
Black, both sexes.....	-	-	2	-	-	-	-	-	-	1	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Connecticut

All races, both sexes.....	4	4	11	15	19	30	60	108	83	25	8	-
Male.....	3	4	6	11	12	20	36	55	41	9	1	-
Female.....	1	-	5	4	7	10	24	53	42	16	7	-
White, both sexes.....	4	4	10	13	19	29	58	104	82	24	8	-
Male.....	3	4	6	9	12	19	35	54	41	9	1	-
Female.....	1	-	4	4	7	10	23	50	41	15	7	-
Black, both sexes.....	-	-	1	2	-	-	1	4	-	1	-	-
Male.....	-	-	-	2	-	-	1	1	-	-	-	-
Female.....	-	-	1	-	-	-	-	3	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	-	1	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to food and vomit (J69.0)

Kentucky

All races, both sexes.....	5	6	12	9	28	42	68	75	67	25	7	-
Male.....	3	2	6	5	20	20	33	30	30	6	1	-
Female.....	2	4	6	4	8	22	35	45	37	19	6	-
White, both sexes.....	3	6	9	9	27	41	60	73	62	25	6	-
Male.....	2	2	4	5	19	19	31	29	27	6	1	-
Female.....	1	4	5	4	8	22	29	44	35	19	5	-
Black, both sexes.....	2	-	3	-	1	1	8	2	5	-	1	-
Male.....	1	-	2	-	1	1	2	1	3	-	-	-
Female.....	1	-	1	-	-	-	6	1	2	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Louisiana

All races, both sexes.....	4	9	7	13	14	43	45	53	30	14	2	-
Male.....	4	8	3	6	9	20	25	29	14	8	-	-
Female.....	-	1	4	7	5	23	20	24	16	6	2	-
White, both sexes.....	3	5	3	8	11	37	38	40	25	11	2	-
Male.....	3	5	1	3	7	16	23	22	13	7	-	-
Female.....	-	-	2	5	4	21	15	18	12	4	2	-
Black, both sexes.....	1	4	4	5	3	6	6	12	5	3	-	-
Male.....	1	3	2	3	2	4	2	6	1	1	-	-
Female.....	-	1	2	2	1	2	4	6	4	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Maine												
All races, both sexes.....	2	2	4	4	3	11	15	21	17	7	2	-
Male.....	1	2	2	4	2	8	9	8	7	3	-	-
Female.....	1	-	2	-	1	3	6	13	10	4	2	-
White, both sexes.....	2	2	4	4	3	11	15	21	17	7	2	-
Male.....	1	2	2	4	2	8	9	8	7	3	-	-
Female.....	1	-	2	-	1	3	6	13	10	4	2	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes.....	8	11	10	14	21	31	74	103	66	20	1	-
Male.....	7	9	7	6	13	20	40	54	35	11	-	-
Female.....	1	2	3	8	8	11	34	49	31	9	1	-
White, both sexes.....	6	6	9	8	13	24	54	90	57	19	1	-
Male.....	6	5	6	3	7	18	28	48	32	11	-	-
Female.....	-	1	3	5	6	6	26	42	25	8	1	-
Black, both sexes.....	2	5	1	6	8	7	19	12	9	1	-	-
Male.....	1	4	1	3	6	2	11	6	3	-	-	-
Female.....	1	1	-	3	2	5	8	6	6	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pneumonitis due to food and vomit (J69.0)

Massachusetts

All races, both sexes.....	3	14	16	15	34	65	89	130	98	33	6	-
Male.....	3	10	9	13	19	41	56	71	48	10	2	-
Female.....	-	4	7	2	15	24	33	59	50	23	4	-
White, both sexes.....	3	14	16	13	33	63	84	126	93	33	6	-
Male.....	3	10	9	11	18	39	52	70	43	10	2	-
Female.....	-	4	7	2	15	24	32	56	50	23	4	-
Black, both sexes.....	-	-	-	2	1	1	4	3	1	-	-	-
Male.....	-	-	-	2	1	1	3	1	1	-	-	-
Female.....	-	-	-	-	-	-	1	2	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	1	4	-	-	-
Male.....	-	-	-	-	-	1	1	-	4	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Michigan

All races, both sexes.....	10	14	16	24	33	61	106	113	89	40	10	-
Male.....	8	8	4	16	21	37	55	61	47	13	3	-
Female.....	2	6	12	8	12	24	51	52	42	27	7	-
White, both sexes.....	7	9	11	23	29	52	91	98	84	37	7	-
Male.....	7	6	3	16	19	31	49	53	43	12	3	-
Female.....	-	3	8	7	10	21	42	45	41	25	4	-
Black, both sexes.....	3	5	5	1	3	8	13	12	4	3	3	-
Male.....	1	2	1	-	2	6	5	5	3	1	-	-
Female.....	2	3	4	1	1	2	8	7	1	2	3	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	1	2	2	1	-	-	-
Male.....	-	-	-	-	-	-	1	2	1	-	-	-
Female.....	-	-	-	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
New Hampshire												
All races, both sexes.....	-	1	4	5	7	3	7	10	18	4	-	-
Male.....	-	1	1	2	4	-	7	1	11	1	-	-
Female.....	-	-	3	3	3	3	-	9	7	3	-	-
White, both sexes.....	-	1	4	5	7	3	7	10	18	4	-	-
Male.....	-	1	1	2	4	-	7	1	11	1	-	-
Female.....	-	-	3	3	3	3	-	9	7	3	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes.....	4	11	10	17	29	70	103	142	107	46	5	-
Male.....	3	5	5	14	18	42	50	74	53	18	1	-
Female.....	1	6	5	3	11	28	53	68	54	28	4	-
White, both sexes.....	2	7	9	17	23	60	99	134	100	42	4	-
Male.....	2	4	5	14	14	34	49	72	50	17	1	-
Female.....	-	3	4	3	9	26	50	62	50	25	3	-
Black, both sexes.....	2	4	1	-	6	10	4	7	3	4	1	-
Male.....	1	1	-	-	4	8	1	2	1	1	-	-
Female.....	1	3	1	-	2	2	3	5	2	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	4	-	-	-
Male.....	-	-	-	-	-	-	-	-	2	-	-	-
Female.....	-	-	-	-	-	-	-	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
New Mexico												
All races, both sexes.....	2	4	2	5	5	10	20	15	7	5	2	-
Male.....	1	3	1	2	3	7	9	7	3	2	2	-
Female.....	1	1	1	3	2	3	11	8	4	3	-	-
White, both sexes.....	1	4	1	5	3	9	18	14	7	4	1	-
Male.....	1	3	1	2	2	7	8	7	3	2	1	-
Female.....	-	1	-	3	1	2	10	7	4	2	-	-
Black, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	2	-	2	1	-	1	1	-
Male.....	-	-	-	-	1	-	1	-	-	-	1	-
Female.....	-	-	1	-	1	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	9	16	18	21	41	72	151	190	118	60	13	-
Male.....	6	10	9	13	26	43	78	100	60	21	3	-
Female.....	3	6	9	8	15	29	73	90	58	39	10	-
White, both sexes.....	8	13	16	18	36	70	142	180	115	58	11	-
Male.....	5	10	8	11	22	42	75	97	58	20	3	-
Female.....	3	3	8	7	14	28	67	83	57	38	8	-
Black, both sexes.....	1	3	2	2	5	1	7	10	3	2	2	-
Male.....	1	-	1	2	4	-	2	3	2	1	-	-
Female.....	-	3	1	-	1	1	5	7	1	1	2	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	2	-	-	-	-	-
Male.....	-	-	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Oregon												
All races, both sexes.....	4	7	4	8	13	18	32	53	36	15	3	-
Male.....	2	6	4	6	6	12	14	30	17	7	-	-
Female.....	2	1	-	2	7	6	18	23	19	8	3	-
White, both sexes.....	4	7	4	7	10	18	31	51	35	15	3	-
Male.....	2	6	4	6	4	12	13	30	16	7	-	-
Female.....	2	1	-	1	6	6	18	21	19	8	3	-
Black, both sexes.....	-	-	-	-	1	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	-	1	1	1	-	-	-
Male.....	-	-	-	-	2	-	1	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes.....	11	17	33	45	69	112	215	282	194	71	8	-
Male.....	5	10	15	27	45	65	106	152	84	23	3	-
Female.....	6	7	18	18	24	47	109	130	110	48	5	-
White, both sexes.....	7	16	32	35	63	103	200	267	181	66	7	-
Male.....	4	10	14	19	41	60	96	144	79	22	3	-
Female.....	3	6	18	16	22	43	104	123	102	44	4	-
Black, both sexes.....	4	1	1	10	6	8	15	14	11	5	1	-
Male.....	1	-	1	8	4	4	10	8	4	1	-	-
Female.....	3	1	-	2	2	4	5	6	7	4	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	1	2	-	-	-
Male.....	-	-	-	-	-	1	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to food and vomit (J69.0)												
Texas												
All races, both sexes.....	19	36	40	39	75	127	225	246	191	76	19	-
Male.....	9	18	21	23	44	83	106	136	84	19	4	-
Female.....	10	18	19	16	31	44	119	110	107	57	15	-
White, both sexes.....	18	27	35	32	63	117	210	231	181	67	19	-
Male.....	8	12	18	19	36	75	98	129	80	18	4	-
Female.....	10	15	17	13	27	42	112	102	101	49	15	-
Black, both sexes.....	1	8	4	7	12	9	14	11	9	9	-	-
Male.....	1	5	2	4	8	7	7	5	3	1	-	-
Female.....	-	3	2	3	4	2	7	6	6	8	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	-	-	1	1	3	1	-	-	-
Male.....	-	1	1	-	-	1	1	1	1	-	-	-
Female.....	-	-	-	-	-	-	-	2	-	-	-	-
Utah												
All races, both sexes.....	4	1	4	1	3	9	15	23	22	3	-	-
Male.....	4	-	3	1	2	4	5	9	14	1	-	-
Female.....	-	1	1	-	1	5	10	14	8	2	-	-
White, both sexes.....	4	1	4	1	3	9	15	22	22	3	-	-
Male.....	4	-	3	1	2	4	5	9	14	1	-	-
Female.....	-	1	1	-	1	5	10	13	8	2	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Delaware

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Alaska

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Kentucky

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Oregon

Table with 13 columns (Age groups) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Age groups) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other pneumonitis due to solids and liquids (J69.1-J69.8)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other lung diseases due to external agents (J67,J70)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other lung diseases due to external agents (J67,J70)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other lung diseases due to external agents (J67,J70)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other lung diseases due to external agents (J67,J70)

North Carolina

Table with 13 columns representing age groups and 13 rows representing race and sex categories for North Carolina. Data values are mostly dashes, with '1' appearing for 'All races, both sexes' and 'Asian or Pacific Islander 2/, both sexes'.

North Dakota

Table with 13 columns representing age groups and 13 rows representing race and sex categories for North Dakota. All data values are dashes.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other lung diseases due to external agents (J67,J70)

Washington

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other lung diseases due to external agents (J67,J70)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other lung diseases due to external agents (J67,J70)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Suppurative and necrotic conditions of lower respiratory tract (J85-J86)

Arkansas

All races, both sexes.....	1	-	1	1	1	1	1	1	-	-	-	-
Male.....	1	-	1	1	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	1	-	-	-	-
White, both sexes.....	1	-	1	1	1	1	1	1	-	-	-	-
Male.....	1	-	1	1	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	1	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	4	13	2	3	4	8	9	4	3	-	-	-
Male.....	3	10	1	2	3	4	6	2	3	-	-	-
Female.....	1	3	1	1	1	4	3	2	-	-	-	-
White, both sexes.....	2	9	1	2	2	6	9	4	3	-	-	-
Male.....	1	7	1	1	2	2	6	2	3	-	-	-
Female.....	1	2	-	1	-	4	3	2	-	-	-	-
Black, both sexes.....	2	4	-	-	1	2	-	-	-	-	-	-
Male.....	2	3	-	-	1	2	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	1	1	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Suppurative and necrotic conditions of lower respiratory tract (J85-J86)

New Hampshire

All races, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	2	-	-	4	2	1	3	3	-	-	-	-
Male.....	1	-	-	3	1	1	2	-	-	-	-	-
Female.....	1	-	-	1	1	-	1	3	-	-	-	-
White, both sexes.....	2	-	-	3	2	1	3	2	-	-	-	-
Male.....	1	-	-	3	1	1	2	-	-	-	-	-
Female.....	1	-	-	-	1	-	1	2	-	-	-	-
Black, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Suppurative and necrotic conditions of lower respiratory tract (J85-J86)

New Mexico

All races, both sexes.....	1	2	-	-	2	-	1	-	-	-	-	-
Male.....	1	2	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
White, both sexes.....	1	2	-	-	1	-	1	-	-	-	-	-
Male.....	1	2	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	3	5	3	1	6	2	7	2	-	-	-	-
Male.....	3	4	1	-	3	1	4	1	-	-	-	-
Female.....	-	1	2	1	3	1	3	1	-	-	-	-
White, both sexes.....	2	5	3	-	3	2	6	2	-	-	-	-
Male.....	2	4	1	-	1	1	4	1	-	-	-	-
Female.....	-	1	2	-	2	1	2	1	-	-	-	-
Black, both sexes.....	1	-	-	1	1	-	1	-	-	-	-	-
Male.....	1	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Pleural effusion and plaque (J90,J92)

Nebraska

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Pleural effusion and plaque (J90,J92)

Rhode Island

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pleural effusion and plaque (J90,J92)

Arkansas

All races, both sexes.....	-	1	-	1	-	2	2	4	1	2	-	-
Male.....	-	-	-	1	-	1	2	1	-	1	-	-
Female.....	-	1	-	-	-	1	-	3	1	1	-	-
White, both sexes.....	-	1	-	1	-	2	2	4	1	2	-	-
Male.....	-	-	-	1	-	1	2	1	-	1	-	-
Female.....	-	1	-	-	-	1	-	3	1	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	4	4	1	2	6	10	19	13	10	5	1	-
Male.....	3	2	1	1	1	5	5	5	5	2	-	-
Female.....	1	2	-	1	5	5	14	11	5	3	1	-
White, both sexes.....	4	3	1	2	6	9	16	10	9	5	1	-
Male.....	3	1	1	1	1	4	4	1	5	2	-	-
Female.....	1	2	-	1	5	5	12	9	4	3	1	-
Black, both sexes.....	-	1	-	-	-	-	2	1	1	-	-	-
Male.....	-	1	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	2	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pleural effusion and plaque (J90,J92)

Colorado

All races, both sexes.....	-	1	1	-	1	3	4	1	3	-	-	-
Male.....	-	-	1	-	1	2	2	1	2	-	-	-
Female.....	-	1	-	-	-	1	2	-	1	-	-	-
White, both sexes.....	-	-	1	-	1	3	4	1	3	-	-	-
Male.....	-	-	1	-	1	2	2	1	2	-	-	-
Female.....	-	-	-	-	-	1	2	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Connecticut

All races, both sexes.....	-	-	-	-	2	1	4	3	4	2	-	-
Male.....	-	-	-	-	1	-	1	1	-	2	-	-
Female.....	-	-	-	-	1	1	3	2	4	-	-	-
White, both sexes.....	-	-	-	-	2	1	3	3	4	2	-	-
Male.....	-	-	-	-	1	-	1	1	-	2	-	-
Female.....	-	-	-	-	1	1	2	2	4	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pleural effusion and plaque (J90,J92)

Florida

All races, both sexes.....	-	5	2	2	5	9	9	13	10	2	-	-
Male.....	-	3	1	1	4	4	6	7	-	1	-	-
Female.....	-	2	1	1	1	5	3	6	10	1	-	-
White, both sexes.....	-	4	2	1	4	9	9	13	7	2	-	-
Male.....	-	2	1	-	3	4	6	7	-	1	-	-
Female.....	-	2	1	1	1	5	3	6	7	1	-	-
Black, both sexes.....	-	1	-	1	1	-	-	-	2	-	-	-
Male.....	-	1	-	1	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-

Georgia

All races, both sexes.....	1	1	2	1	7	6	6	4	5	1	-	-
Male.....	1	-	1	1	3	4	2	2	1	-	-	-
Female.....	-	1	1	-	4	2	4	2	4	1	-	-
White, both sexes.....	1	1	2	-	4	5	5	4	4	-	-	-
Male.....	1	-	1	-	1	4	2	2	1	-	-	-
Female.....	-	1	1	-	3	1	3	2	3	-	-	-
Black, both sexes.....	-	-	-	-	2	1	1	-	1	1	-	-
Male.....	-	-	-	-	2	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	1	-	1	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	1	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Pleural effusion and plaque (J90,J92)

New Hampshire

All races, both sexes.....	-	-	1	-	1	1	1	2	1	-	-	-
Male.....	-	-	-	-	1	1	-	1	-	-	-	-
Female.....	-	-	1	-	-	-	1	1	1	-	-	-
White, both sexes.....	-	-	1	-	1	1	1	2	1	-	-	-
Male.....	-	-	-	-	1	1	-	1	-	-	-	-
Female.....	-	-	1	-	-	-	1	1	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	2	-	2	2	4	8	12	11	7	3	2	-
Male.....	1	-	-	2	2	2	6	2	2	1	-	-
Female.....	1	-	2	-	2	6	6	9	5	2	2	-
White, both sexes.....	1	-	2	-	2	8	11	11	6	3	2	-
Male.....	1	-	-	-	1	2	5	2	2	1	-	-
Female.....	-	-	2	-	1	6	6	9	4	2	2	-
Black, both sexes.....	1	-	-	2	2	-	1	-	-	-	-	-
Male.....	-	-	-	2	1	-	1	-	-	-	-	-
Female.....	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases of respiratory system (J80-J84,J93-J98)

Arkansas

All races, both sexes.....	283	1	-	-	1	-	2	1	1	3	3	7
Male.....	144	-	-	-	1	-	-	1	1	3	-	2
Female.....	139	1	-	-	-	-	2	-	-	-	3	5
White, both sexes.....	259	1	-	-	1	-	2	1	-	3	1	6
Male.....	132	-	-	-	1	-	-	1	-	3	-	2
Female.....	127	1	-	-	-	-	2	-	-	-	1	4
Black, both sexes.....	23	-	-	-	-	-	-	-	1	-	2	1
Male.....	12	-	-	-	-	-	-	-	1	-	-	-
Female.....	11	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	2,007	19	2	2	2	4	3	6	6	12	24	28
Male.....	1,032	14	1	1	2	2	-	5	3	1	14	23
Female.....	975	5	1	1	-	2	3	1	3	11	10	5
White, both sexes.....	1,736	14	1	1	1	3	2	5	5	7	19	20
Male.....	893	11	-	-	1	2	-	5	2	1	12	17
Female.....	843	3	1	1	-	1	2	-	3	6	7	3
Black, both sexes.....	88	2	1	1	-	1	-	1	-	3	2	6
Male.....	41	1	1	1	-	-	-	-	-	-	-	4
Female.....	47	1	-	-	-	1	-	1	-	3	2	2
American Indian 1/, both sexes.....	19	-	-	-	-	-	-	-	-	1	1	-
Male.....	7	-	-	-	-	-	-	-	-	-	-	-
Female.....	12	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes.....	164	3	-	-	1	-	1	-	1	1	2	2
Male.....	91	2	-	-	1	-	-	-	1	-	2	2
Female.....	73	1	-	-	-	-	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

Alaska

All races, both sexes.....	1	1	3	10	7	11	6	2	-	2	-	-
Male.....	1	1	3	6	5	6	2	1	-	-	-	-
Female.....	-	-	-	4	2	5	4	1	-	2	-	-
White, both sexes.....	-	1	2	7	6	8	5	2	-	2	-	-
Male.....	-	1	2	5	4	5	1	1	-	-	-	-
Female.....	-	-	-	2	2	3	4	1	-	2	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	2	1	3	1	-	-	-	-	-
Male.....	-	-	1	1	1	1	1	-	-	-	-	-
Female.....	-	-	-	1	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	1	-	-	1	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	13	17	29	43	48	70	57	63	23	17	-	-
Male.....	8	11	18	26	25	38	26	29	10	6	-	-
Female.....	5	6	11	17	23	32	31	34	13	11	-	-
White, both sexes.....	13	17	26	38	41	59	54	59	21	16	-	-
Male.....	8	11	17	24	20	34	26	26	10	6	-	-
Female.....	5	6	9	14	21	25	28	33	11	10	-	-
Black, both sexes.....	-	-	-	1	-	2	-	-	1	-	-	-
Male.....	-	-	-	1	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	2	4	5	9	3	3	-	1	-	-
Male.....	-	-	-	1	4	3	-	2	-	-	-	-
Female.....	-	-	2	3	1	6	3	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	1	-	2	-	-	1	1	-	-	-
Male.....	-	-	1	-	1	-	-	1	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

Arkansas

All races, both sexes.....	13	12	15	28	32	51	56	34	17	6	-	-
Male.....	6	7	9	15	20	31	28	10	7	3	-	-
Female.....	7	5	6	13	12	20	28	24	10	3	-	-
White, both sexes.....	8	10	14	25	30	49	54	33	16	5	-	-
Male.....	3	6	8	14	19	31	27	9	6	2	-	-
Female.....	5	4	6	11	11	18	27	24	10	3	-	-
Black, both sexes.....	5	2	1	2	2	2	2	1	1	1	-	-
Male.....	3	1	1	1	1	-	1	1	1	1	-	-
Female.....	2	1	-	1	1	2	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	59	81	112	173	244	306	376	343	149	45	11	-
Male.....	30	48	66	101	131	169	179	162	61	13	6	-
Female.....	29	33	46	72	113	137	197	181	88	32	5	-
White, both sexes.....	47	67	94	147	209	275	325	304	139	41	10	-
Male.....	22	41	53	86	115	155	154	145	55	10	6	-
Female.....	25	26	41	61	94	120	171	159	84	31	4	-
Black, both sexes.....	8	7	9	10	11	8	8	7	2	-	1	-
Male.....	5	4	6	5	5	4	-	3	2	-	-	-
Female.....	3	3	3	5	6	4	8	4	-	-	1	-
American Indian 1/, both sexes.....	1	2	-	1	4	1	3	3	2	-	-	-
Male.....	-	1	-	1	1	-	1	2	1	-	-	-
Female.....	1	1	-	-	3	1	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	5	9	15	20	22	40	29	6	4	-	-
Male.....	3	2	7	9	10	10	24	12	3	3	-	-
Female.....	-	3	2	6	10	12	16	17	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases of respiratory system (J80-J84,J93-J98)

Florida

All races, both sexes.....	38	46	91	130	183	285	315	254	122	40	6	-
Male.....	16	27	50	83	107	159	180	129	52	9	2	-
Female.....	22	19	41	47	76	126	135	125	70	31	4	-
White, both sexes.....	29	38	81	116	169	272	306	241	118	38	4	-
Male.....	14	22	44	75	99	152	174	123	51	8	1	-
Female.....	15	16	37	41	70	120	132	118	67	30	3	-
Black, both sexes.....	9	7	10	12	13	10	7	13	4	2	2	-
Male.....	2	4	6	8	7	5	5	6	1	1	1	-
Female.....	7	3	4	4	6	5	2	7	3	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	2	1	3	2	-	-	-	-	-
Male.....	-	1	-	-	1	2	1	-	-	-	-	-
Female.....	-	-	-	2	-	1	1	-	-	-	-	-

Georgia

All races, both sexes.....	30	54	78	87	141	138	182	127	83	27	4	-
Male.....	16	32	41	43	76	67	77	47	25	7	2	-
Female.....	14	22	37	44	65	71	105	80	58	20	2	-
White, both sexes.....	19	38	53	74	111	115	154	103	72	23	2	-
Male.....	10	25	28	37	63	58	64	39	21	7	1	-
Female.....	9	13	25	37	48	57	90	64	51	16	1	-
Black, both sexes.....	11	16	25	12	28	21	26	22	11	4	2	-
Male.....	6	7	13	6	11	8	11	8	4	-	1	-
Female.....	5	9	12	6	17	13	15	14	7	4	1	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	2	2	2	-	-	-	-
Male.....	-	-	-	-	2	1	2	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

Maine

All races, both sexes.....	4	4	11	11	22	27	34	38	18	3	2	-
Male.....	1	3	5	7	9	15	24	18	6	1	1	-
Female.....	3	1	6	4	13	12	10	20	12	2	1	-
White, both sexes.....	3	4	11	11	22	27	34	38	18	3	2	-
Male.....	1	3	5	7	9	15	24	18	6	1	1	-
Female.....	2	1	6	4	13	12	10	20	12	2	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	10	17	25	31	44	62	64	63	42	10	1	-
Male.....	8	9	16	16	23	32	35	27	16	3	-	-
Female.....	2	8	9	15	21	30	29	36	26	7	1	-
White, both sexes.....	8	13	21	22	35	49	55	59	40	9	1	-
Male.....	6	6	14	12	19	28	30	23	16	2	-	-
Female.....	2	7	7	10	16	21	25	36	24	7	1	-
Black, both sexes.....	2	3	4	8	7	12	8	3	2	1	-	-
Male.....	2	2	2	3	2	4	4	3	-	1	-	-
Female.....	-	1	2	5	5	8	4	-	2	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	1	-	-	-	-	-	-	-
Male.....	-	-	-	1	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	1	1	1	1	-	-	-	-
Male.....	-	1	-	-	1	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

Massachusetts

All races, both sexes.....	8	23	34	42	67	98	138	116	68	29	7	-
Male.....	1	9	21	21	34	54	65	52	25	6	1	-
Female.....	7	14	13	21	33	44	73	64	43	23	6	-
White, both sexes.....	6	21	33	40	64	95	137	115	66	28	6	-
Male.....	1	8	20	19	33	52	64	52	25	6	1	-
Female.....	5	13	13	21	31	43	73	63	41	22	5	-
Black, both sexes.....	2	2	-	2	3	2	-	1	1	1	1	-
Male.....	-	1	-	2	1	1	-	-	-	-	-	-
Female.....	2	1	-	-	2	1	-	1	1	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	1	1	-	1	-	-	-
Male.....	-	-	1	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-

Michigan

All races, both sexes.....	24	33	45	78	95	137	151	137	57	18	3	-
Male.....	14	18	27	43	61	69	71	55	27	6	2	-
Female.....	10	15	18	35	34	68	80	82	30	12	1	-
White, both sexes.....	17	26	41	63	83	127	142	129	52	17	3	-
Male.....	10	15	26	36	53	66	68	52	24	6	2	-
Female.....	7	11	15	27	30	61	74	77	28	11	1	-
Black, both sexes.....	7	7	4	13	10	7	7	8	3	1	-	-
Male.....	4	3	1	5	6	1	2	3	2	-	-	-
Female.....	3	4	3	8	4	6	5	5	1	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	1	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	1	3	1	-	2	-	-	-
Male.....	-	-	-	2	1	2	1	-	1	-	-	-
Female.....	-	-	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

Minnesota

All races, both sexes.....	8	15	28	31	42	60	79	84	45	12	4	-
Male.....	5	10	18	17	27	36	49	45	14	5	1	-
Female.....	3	5	10	14	15	24	30	39	31	7	3	-
White, both sexes.....	6	14	26	28	39	60	78	84	45	12	4	-
Male.....	3	10	18	15	26	36	48	45	14	5	1	-
Female.....	3	4	8	13	13	24	30	39	31	7	3	-
Black, both sexes.....	-	1	1	1	1	-	-	-	-	-	-	-
Male.....	-	-	-	1	1	-	-	-	-	-	-	-
Female.....	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	-	1	-	1	-	1	-	-	-	-	-
Male.....	2	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	1	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	1	-	-	-	-	-	-	-

Mississippi

All races, both sexes.....	10	18	25	33	44	48	59	42	30	12	1	-
Male.....	6	8	14	21	26	19	24	18	5	4	-	-
Female.....	4	10	11	12	18	29	35	24	25	8	1	-
White, both sexes.....	7	12	17	20	32	37	48	38	24	9	1	-
Male.....	3	5	10	14	21	14	20	18	3	4	-	-
Female.....	4	7	7	6	11	23	28	20	21	5	1	-
Black, both sexes.....	3	6	7	13	9	11	11	4	6	3	-	-
Male.....	3	3	3	7	4	5	4	-	2	-	-	-
Female.....	-	3	4	6	5	6	7	4	4	3	-	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	3	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

Nebraska

All races, both sexes.....	9	5	12	13	9	36	43	32	16	5	2	-
Male.....	6	3	3	7	4	26	18	12	4	1	-	-
Female.....	3	2	9	6	5	10	25	20	12	4	2	-
White, both sexes.....	9	5	11	12	7	36	42	32	16	4	2	-
Male.....	6	3	2	7	3	26	17	12	4	1	-	-
Female.....	3	2	9	5	4	10	25	20	12	3	2	-
Black, both sexes.....	-	-	1	1	2	-	1	-	-	1	-	-
Male.....	-	-	1	-	1	-	1	-	-	-	-	-
Female.....	-	-	-	1	1	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	8	9	15	15	21	32	32	18	7	1	-	-
Male.....	5	7	4	11	13	17	19	5	4	1	-	-
Female.....	3	2	11	4	8	15	13	13	3	-	-	-
White, both sexes.....	6	8	15	11	18	27	31	17	7	-	-	-
Male.....	4	7	4	8	10	15	18	5	4	-	-	-
Female.....	2	1	11	3	8	12	13	12	3	-	-	-
Black, both sexes.....	1	-	-	2	1	1	-	1	-	1	-	-
Male.....	-	-	-	2	1	1	-	-	-	1	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	1	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	2	2	3	1	-	-	-	-	-
Male.....	-	-	-	1	2	1	1	-	-	-	-	-
Female.....	-	1	-	1	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

New Hampshire

All races, both sexes.....	3	2	9	13	10	26	21	13	10	4	2	-
Male.....	2	1	4	13	5	14	10	6	3	1	-	-
Female.....	1	1	5	-	5	12	11	7	7	3	2	-
White, both sexes.....	3	2	9	13	10	26	21	13	10	4	2	-
Male.....	2	1	4	13	5	14	10	6	3	1	-	-
Female.....	1	1	5	-	5	12	11	7	7	3	2	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	18	19	32	51	86	107	172	115	75	24	2	-
Male.....	14	13	14	27	54	55	91	50	24	7	1	-
Female.....	4	6	18	24	32	52	81	65	51	17	1	-
White, both sexes.....	11	16	26	44	77	104	157	111	72	22	2	-
Male.....	10	12	11	22	50	53	82	48	24	6	1	-
Female.....	1	4	15	22	27	51	75	63	48	16	1	-
Black, both sexes.....	6	2	4	4	6	3	11	3	2	2	-	-
Male.....	4	1	1	2	2	2	6	2	-	1	-	-
Female.....	2	1	3	2	4	1	5	1	2	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	2	3	3	-	3	1	1	-	-	-
Male.....	-	-	2	3	2	-	2	-	-	-	-	-
Female.....	1	1	-	-	1	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

New Mexico

All races, both sexes.....	2	4	12	20	26	35	29	38	10	4	1	-
Male.....	2	2	9	13	15	19	14	20	5	2	-	-
Female.....	-	2	3	7	11	16	15	18	5	2	1	-
White, both sexes.....	2	4	9	14	23	32	29	28	10	3	1	-
Male.....	2	2	7	9	13	18	14	16	5	2	-	-
Female.....	-	2	2	5	10	14	15	12	5	1	1	-
Black, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	3	6	2	3	-	10	-	1	-	-
Male.....	-	-	2	4	1	1	-	4	-	-	-	-
Female.....	-	-	1	2	1	2	-	6	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	32	58	79	102	139	197	247	198	87	30	7	-
Male.....	22	31	40	47	75	99	119	78	24	7	-	-
Female.....	10	27	39	55	64	98	128	120	63	23	7	-
White, both sexes.....	26	49	64	89	120	178	232	185	82	29	5	-
Male.....	17	25	32	40	68	91	110	76	23	6	-	-
Female.....	9	24	32	49	52	87	122	109	59	23	5	-
Black, both sexes.....	5	8	14	11	15	15	11	10	5	1	2	-
Male.....	4	6	7	6	4	7	6	2	1	1	-	-
Female.....	1	2	7	5	11	8	5	8	4	-	2	-
American Indian 1/, both sexes.....	-	-	1	1	-	1	-	-	-	-	-	-
Male.....	-	-	1	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	1	4	3	4	3	-	-	-	-
Male.....	1	-	-	-	3	1	3	-	-	-	-	-
Female.....	-	1	-	1	1	2	1	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

Ohio

All races, both sexes.....	28	38	60	110	117	170	208	184	97	18	2	-
Male.....	12	20	30	52	67	84	111	81	38	5	1	-
Female.....	16	18	30	58	50	86	97	103	59	13	1	-
White, both sexes.....	21	33	52	103	100	156	196	171	93	18	2	-
Male.....	11	18	25	49	57	78	105	74	38	5	1	-
Female.....	10	15	27	54	43	78	91	97	55	13	1	-
Black, both sexes.....	6	4	7	7	16	11	11	13	2	-	-	-
Male.....	1	1	5	3	9	4	5	7	-	-	-	-
Female.....	5	3	2	4	7	7	6	6	2	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	1	-	1	3	1	-	2	-	-	-
Male.....	-	1	-	-	1	2	1	-	-	-	-	-
Female.....	1	-	1	-	-	1	-	-	2	-	-	-

Oklahoma

All races, both sexes.....	8	18	26	26	52	53	46	41	27	4	1	-
Male.....	4	10	18	8	25	34	18	13	9	1	-	-
Female.....	4	8	8	18	27	19	28	28	18	3	1	-
White, both sexes.....	2	11	21	20	47	51	43	38	26	4	1	-
Male.....	1	6	14	6	21	32	17	13	9	1	-	-
Female.....	1	5	7	14	26	19	26	25	17	3	1	-
Black, both sexes.....	3	3	2	2	2	-	-	-	1	-	-	-
Male.....	-	1	2	1	2	-	-	-	-	-	-	-
Female.....	3	2	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	2	4	3	4	3	2	1	3	-	-	-	-
Male.....	2	3	2	1	2	2	-	-	-	-	-	-
Female.....	-	1	1	3	1	-	1	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	2	-	-	-	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

Oregon

All races, both sexes.....	7	18	19	34	39	58	42	57	32	13	1	-
Male.....	3	10	13	17	29	36	22	26	7	3	-	-
Female.....	4	8	6	17	10	22	20	31	25	10	1	-
White, both sexes.....	6	15	16	31	38	55	40	55	32	12	1	-
Male.....	3	9	11	16	28	35	22	26	7	2	-	-
Female.....	3	6	5	15	10	20	18	29	25	10	1	-
Black, both sexes.....	-	2	-	-	-	-	-	1	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	1	-	-
Female.....	-	2	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	1	1	3	3	-	-	-	1	-	-	-	-
Male.....	-	1	2	1	-	-	-	-	-	-	-	-
Female.....	1	-	1	2	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	3	2	-	-	-	-	-
Male.....	-	-	-	-	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	2	2	-	-	-	-	-

Pennsylvania

All races, both sexes.....	42	62	100	89	168	265	313	268	168	42	7	-
Male.....	18	35	59	50	88	134	167	118	50	10	1	-
Female.....	24	27	41	39	80	131	146	150	118	32	6	-
White, both sexes.....	29	52	89	80	148	237	301	253	159	39	6	-
Male.....	13	30	54	45	80	120	161	117	48	9	1	-
Female.....	16	22	35	35	68	117	140	136	111	30	5	-
Black, both sexes.....	13	10	8	8	19	23	10	13	8	2	-	-
Male.....	5	5	3	4	7	13	5	-	2	-	-	-
Female.....	8	5	5	4	12	10	5	13	6	2	-	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	1	1	5	2	2	1	1	1	-
Male.....	-	-	1	1	1	1	1	1	-	1	-	-
Female.....	-	-	1	-	-	4	1	1	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

Texas

All races, both sexes.....	65	106	137	162	194	256	311	226	113	43	5	-
Male.....	35	50	71	93	108	134	140	94	46	10	2	-
Female.....	30	56	66	69	86	122	171	132	67	33	3	-
White, both sexes.....	57	84	122	144	179	227	291	208	106	41	3	-
Male.....	29	39	64	86	104	115	130	88	43	10	2	-
Female.....	28	45	58	58	75	112	161	120	63	31	1	-
Black, both sexes.....	8	19	11	14	12	20	15	18	6	1	2	-
Male.....	6	10	5	4	4	14	8	6	3	-	-	-
Female.....	2	9	6	10	8	6	7	12	3	1	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	3	4	4	3	9	4	-	1	1	-	-
Male.....	-	1	2	3	-	5	2	-	-	-	-	-
Female.....	-	2	2	1	3	4	2	-	1	1	-	-

Utah

All races, both sexes.....	1	10	8	18	17	25	35	21	10	3	-	-
Male.....	1	7	6	10	11	16	24	14	6	-	-	-
Female.....	-	3	2	8	6	9	11	7	4	3	-	-
White, both sexes.....	1	9	8	18	17	25	35	20	10	3	-	-
Male.....	1	6	6	10	11	16	24	14	6	-	-	-
Female.....	-	3	2	8	6	9	11	6	4	3	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All other diseases of respiratory system (J80-J84,J93-J98)

Vermont

All races, both sexes.....	-	3	3	2	2	5	15	17	3	2	-	-
Male.....	-	-	3	1	1	3	9	6	2	-	-	-
Female.....	-	3	-	1	1	2	6	11	1	2	-	-
White, both sexes.....	-	3	3	2	2	5	15	17	3	2	-	-
Male.....	-	-	3	1	1	3	9	6	2	-	-	-
Female.....	-	3	-	1	1	2	6	11	1	2	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	27	30	57	63	101	128	132	117	50	25	4	-
Male.....	19	14	31	33	56	73	67	52	17	8	1	-
Female.....	8	16	26	30	45	55	65	65	33	17	3	-
White, both sexes.....	19	22	42	55	81	109	114	107	46	24	3	-
Male.....	14	13	23	31	42	66	60	47	15	8	-	-
Female.....	5	9	19	24	39	43	54	60	31	16	3	-
Black, both sexes.....	8	8	14	8	18	16	13	9	3	1	-	-
Male.....	5	1	8	2	13	6	4	4	2	-	-	-
Female.....	3	7	6	6	5	10	9	5	1	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	2	3	5	1	1	-	1	-
Male.....	-	-	-	-	1	1	3	1	-	-	1	-
Female.....	-	-	1	-	1	2	2	-	1	-	-	-

