

**Subset of alphabetical index to diseases and nature of injury for use with perinatal conditions  
(P00-P96)**

Conditions arising in the perinatal period

*Note* - Conditions arising in the perinatal period, even though death or morbidity occurs later, should, as far as possible, be coded to chapter XVI, which takes precedence over chapters containing codes for diseases by their anatomical site.

These exclude:

- Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)
- Endocrine, nutritional and metabolic diseases (E00-E99)
- Injury, poisoning and certain other consequences of external causes (S00-T99)
- Neoplasms (C00-D48)
- Tetanus neonatorum (A33)

**A**

- ablatio, ablation
- - placentae (*see also* Abruptio placentae)
- - - affecting fetus or newborn P02.1 2a
- abnormal, abnormality, abnormalities - *see also* Anomaly
- - alphafetoprotein
- - - maternal, affecting fetus or newborn P00.8
- - amnion, amniotic fluid
- - - affecting fetus or newborn P02.9
- - anticoagulation
- - - newborn (transient) P61.6
- - cervix NEC, maternal (acquired) (congenital), in pregnancy or childbirth
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- - chorion
- - - affecting fetus or newborn P02.9
- - coagulation
- - - newborn, transient P61.6
- - fetus, fetal
- - - causing disproportion
- - - - affecting fetus or newborn P03.1
- - forces of labor
- - - affecting fetus or newborn P03.6
- - labor NEC
- - - affecting fetus or newborn P03.6
- - membranes (fetal)
- - - affecting fetus or newborn P02.9
- - - specified type NEC, affecting fetus or newborn P02.8
- - organs or tissues of maternal pelvis
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- - parturition
- - - affecting fetus or newborn P03.9
- - presentation (fetus) (*see also* Presentation, fetal, abnormal)
- - - before labor, affecting fetus or newborn P01.7

## ICD-10-Mortality Perinatal Subset - 2014

- - pulmonary
- - - function, newborn P28.8
- - - ventilation, newborn P28.8
- - umbilical cord
- - - affecting fetus or newborn P02.6
- - vagina, maternal (acquired) (congenital), in pregnancy or childbirth
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- - vulva and perineum, maternal (acquired) (congenital), in pregnancy or childbirth
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- ABO hemolytic disease (fetus or newborn) P55.1
- aborter, habitual or recurrent NEC
- - current abortion
- - - affecting fetus or newborn P01.8
- abortion (complete) (incomplete)
- - accidental
- - - fetus or newborn P01.8
- - fetus or newborn P96.4
- - - in fetal death file P01.8
- - habitual or recurrent
- - - with current abortion
- - - - fetus P01.8
- - idiopathic
- - - fetus or newborn P01.8
- - inevitable
- - - fetus or newborn P01.8
- - infected
- - - fetus or newborn P01.8
- - legal (induced)
- - - fetus P96.4
- - medical
- - - fetus P96.4
- - missed
- - - fetus or newborn P01.8
- - natural
- - - fetus or newborn P01.8
- - previous (habitual) (recurrent) (spontaneous)
- - - affecting fetus or newborn P01.8
- - recurrent
- - - fetus or newborn P01.8
- - retained
- - - fetus or newborn P01.8
- - septic
- - - fetus or newborn P01.8
- - spontaneous
- - - fetus P01.8
- - - threatened
- - - - affecting fetus or newborn P01.8
- - therapeutic
- - - fetus P96.4
- - threatened (spontaneous)
- - - affecting fetus or newborn P01.8
- - tubal
- - - fetus or newborn P01.4
- - unavoidable

## ICD-10-Mortality Perinatal Subset - 2014

- - - fetus or newborn P01.8
- - unintended
- - - fetus or newborn P01.8
- abruptio placentae
- - affecting fetus or newborn P02.1 2a
- abscess (embolic) (infective) (metastatic) (multiple) (perforated) (pyogenic) (septic)
- - abdomen, abdominal NEC
- - - cavity
- - - - newborn P78.1
- - - newborn P78.1
- - abdominopelvic
- - - newborn P78.1
- - breast (acute) (chronic) (nonpuerperal)
- - - newborn P39.0
- - diaphragm, diaphragmatic
- - - newborn P78.1
- - epiploon, epiploic
- - - newborn P78.1
- - extraperitoneal
- - - newborn P78.1
- - intra-abdominal (*see also* Abscess, Peritoneum)
- - - newborn P78.1
- - intraperitoneal
- - - newborn P78.1
- - kidney, maternal
- - - complicating pregnancy
- - - - affecting fetus or newborn P00.1
- - mesentery, mesenteric (*see also* Abscess, peritoneum)
- - - newborn P78.1
- - navel
- - - newborn P38
- - omentum (*see also* Abscess, peritoneum)
- - - newborn P78.1
- - pelvis, pelvic
- - - newborn P78.1
- - peritoneum, peritoneal (perforated) (ruptured)
- - - newborn P78.1
- - - pelvic
- - - - male
- - - - - newborn P78.1
- - pyemic - *see* Septicemia
- - - newborn (*see also* Septicemia, newborn) P36.9
- - retrocecal
- - - newborn P78.1
- - retroperitoneal
- - - newborn P78.1
- - subdiaphragmatic
- - - newborn P78.1
- - subhepatic
- - - newborn P78.1
- - subphrenic
- - - newborn P78.1
- - umbilicus
- - - newborn P38
- - viscera (multiple) NEC
- - - newborn P78.1

## ICD-10-Mortality Perinatal Subset - 2014

- absence, absent, (organ or part)(complete or partial)
- - amniotic fluid
- - - affecting fetus or newborn P01.2 2a
- - diastolic flow (umbilical) P02.2
- - fetal heart tone P95
- - umbilical
- - - diastolic flow P02.2
- absent or reverse end diastolic flow (AREDF) P02.2
- absorption
- - chemical
- - - through placenta (fetus or newborn) P04.9
- - - - environmental substance P04.6
- - - - nutritional substance P04.5
- - - - obstetric anesthetic or analgesic drug P04.0
- - drug NEC (*see also* Reaction, drug)
- - - addictive
- - - - through placenta (fetus or newborn) P04.4
- - - through placenta (fetus or newborn) P04.1
- - - - obstetric anesthetic or analgesic medication P04.0
- - maternal medication NEC through placenta (fetus or newborn) P04.1
- accident
- - birth - *see* Birth, injury
- - cord (umbilical) P02.6
- - during pregnancy, to mother
- - - affecting fetus or newborn P00.5
- acidosis (lactic) (respiratory)
- - fetal -*see* Distress, fetal
- - intrauterine -*see* Distress, fetal
- - metabolic NEC
- - - late, of newborn P74.0
- - - newborn -*see* Distress, fetal
- acrocyanosis
- - newborn P28.2
- action, heart
- - disorder
- - - newborn P29.1
- - irregular
- - - newborn P29.1
- addiction, maternal (*see also* Dependence)
- - alcohol, alcoholic (ethyl) (methyl) (wood)
- - - complicating pregnancy or childbirth
- - - - affecting fetus or newborn P04.3
- adhesions, adhesive (postinfective)
- - amnion to fetus
- - - affecting fetus or newborn P02.8
- - lung
- - - newborn P28.8
- - pulmonary
- - - newborn P28.8
- - uterus
- - - affecting fetus or newborn P03.8
- adiponecrosis neonatorum P83.8
- aeration lung imperfect, newborn P28.1
- albuminuria, albuminuric (acute) (chronic) (subacute) (*see also* Proteinuria)
- - pre-eclamptic (*see also* Pre-eclampsia)
- - - affecting fetus or newborn P00.0

## ICD-10-Mortality Perinatal Subset - 2014

- alcoholism (acute) (chronic)
- - complicating pregnancy or childbirth
- - - affecting fetus or newborn P04.3
- amnion, amniotic
- - fluid
- - - absent P01.2 2a
- - fluid index (AFI)
- - - less than 1.4 - *see* Oligohydramnios
- - - more than 3.8 - *see* Polyhydramnios
- amnionitis
- - affecting fetus or newborn P02.7
- amputation
- - any part of fetus, to facilitate delivery P03.8
- Anaerosis of newborn P28.8
- anasarca
- - fetus or newborn P83.2
- android pelvis, maternal
- - with disproportion (fetopelvic)
- - - affecting fetus or newborn P03.1
- anemia
- - congenital P61.4
- - - due to isoimmunization NEC P55.9
- - - following fetal blood loss P61.3
- - due to
- - - fetal blood loss P61.3
- - - prematurity P61.2
- - erythroblastic
- - - fetus or newborn (*see also* Disease, hemolytic) P55.9
- - fetus or newborn P61.4
- - - due to
- - - - ABO (antibodies) (isoimmunization) (maternal/fetal incompatibility) P55.1
- - - - Rh (antibodies) (isoimmunization) (maternal/fetal incompatibility) P55.0
- - - following fetal blood loss P61.3
- - hemolytic
- - - acute
- - - - fetus or newborn (*see also* Disease, hemolytic) P55.9
- - maternal, of or complicating pregnancy
- - - affecting fetus or newborn P00.8
- - newborn P61.4
- - - posthemorrhagic (fetal) P61.3
- - of prematurity P61.2
- - posthemorrhagic (chronic)
- - - newborn P61.3
- anhydramnios
- - affecting fetus or newborn P01.2 2a
- anomaly, anomalous (congenital) (unspecified type)
- - cervix, maternal
- - - in pregnancy or childbirth NEC
- - - - affecting fetus or newborn P03.8
- - - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- - cord P02.6
- - umbilical cord NEC P02.6
- - uterus, maternal
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8

## ICD-10-Mortality Perinatal Subset - 2014

- - - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- anoxemia (*see also* Anoxia)
- - newborn (*see also* Asphyxia, newborn) P21.9
- anoxia (*see also* Hypoxia)
- - cerebral
- - - newborn (*see also* Asphyxia, newborn) P21.9
- - fetal, fetus (*see also* Distress, fetal) P20-
- - newborn (*see also* Asphyxia, newborn) P21.9
- anteversion
- - uterus, uterine, maternal (cervix) (postinfectious) (postpartal, old)
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8
- - - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- anthropoid pelvis, maternal
- - with disproportion (fetopelvic)
- - - affecting fetus or newborn P03.1
- antibodies (blood group) (*see also* Incompatibility)
- - anti-D
- - - fetus or newborn P55.0
- - anti-Kell
- - - fetus or newborn P55.8
- anuria
- - newborn P96.0
- apgar (score)
- - low NEC, with asphyxia P21.9
- - 0-3 at 1 minute, with asphyxia P21.0
- - 4-7 at 1 minute, with asphyxia P21.1
- apnea, apneic (spells)
- - newborn NEC (obstructive) P28.4
- - - sleep (central) (primary) P28.3
- - prematurity (of) P28.4
- arrest, arrested
- - active phase of labor
- - - affecting fetus or newborn P03.6
- - cardiac
- - - fetal, fetus P29.8
- - - newborn P29.1
- - coronary
- - - infant P29.1
- - deep transverse
- - - affecting fetus or newborn P03.1
- - development or growth
- - - fetus P05.9
- - respiratory
- - - newborn P28.5 2a
- arrhythmia (cardiac) (ventricular) (*see also* Dysrhythmia)
- - extrasystolic
- - - newborn P29.1
- - fetal P29.8
- - newborn P29.1
- - sinus
- - - newborn P29.1
- - specified NEC
- - - newborn P29.1

## ICD-10-Mortality Perinatal Subset - 2014

- asphyxia, asphyxiation (by)
  - - antenatal (*see also* Distress, fetal) P20.9
  - - birth (*see also* Asphyxia, newborn) P21.9
  - - fetal (*see also* Distress, fetal) P20.9
  - - food or foreign body (in)
    - - - newborn P24.3
  - - intrauterine (*see also* Distress, fetal) P20.9
  - - - mucus (in)
    - - - newborn P24.1
  - - newborn P21.9
    - - - with 1-minute Apgar score
      - - - - low NEC P21.9
      - - - - 0-3 P21.0
      - - - - 4-7 P21.1
    - - - blue P21.1
    - - - livida P21.1
    - - - mild or moderate P21.1
    - - - pallida P21.0
    - - - severe P21.0
    - - - white P21.0
  - - perinatal (*see also* Asphyxia, newborn) P21.9
  - - postnatal (*see also* Asphyxia, newborn) P21.9
  - - prenatal (*see also* Distress, fetal) P20.9
- aspiration
  - - amniotic fluid (newborn) P24.1
  - - blood, newborn P24.2
  - - liquor (amnii) (newborn) P24.1
  - - meconium (newborn) P24.0
  - - milk (newborn) P24.3
  - - mucus
    - - - newborn P24.1
  - - newborn (massive) (syndrome) P24.9
  - - - meconium P24.0
  - - vernix caseosa (newborn) P24.8
- asthma, asthmatic (bronchial) (catarrh) (spasmodic)
  - - heart
    - - - fetal, fetus P29.8
    - - - newborn P29.0
  - - Rostan's
    - - - fetal, fetus P29.8
    - - - newborn P29.0
- asymmetry - *see also* Distortion
  - - pelvis with disproportion (fetopelvic)
    - - - affecting fetus or newborn P03.1
- atelectasis (massive) (partial) (pressure) (pulmonary)
  - - fetus or newborn (secondary) P28.1
    - - - due to resorption P28.1
    - - - partial P28.1
    - - - primary P28.0 2a
    - - - subtotal P28.0 2a
  - - primary (newborn) P28.0 2a
- atonia, atony, atonic
  - - congenital P94.2
  - - uterus, maternal, during labor
    - - - affecting fetus or newborn P03.6
- atresia, atretic
  - - cervix, maternal (acquired)

## ICD-10-Mortality Perinatal Subset - 2014

- - - in pregnancy or childbirth
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- atrophy, atrophic
- - lung (senile)
- - - newborn P28.8
- attack
- - cyanotic, newborn P28.2
- - respiration, respiratory
- - - newborn P28.8
- awareness of heart beat
- - fetal P20.9
- - newborn P29.1



**B**

- baby
- - floppy (syndrome) P94.2
- bacteremia
- - due to
- - - specified bacteria (*see also* Infection, by specified bacteria)
- - - - newborn - *see* Sepsis, newborn, by specified bacteria
- - newborn (any bacteria classifiable to A49- ) - *see* Sepsis, newborn, by specified bacteria
- - organism other than bacteria - *see* Septicemia
- - - newborn - *see* Infection, congenital, by organism
- bacteriuria, bacteruria
- - in pregnancy
- - - affecting fetus or newborn P00.1
- ballooning
- - membranes
- - - affecting fetus or newborn P01.0
- band(s)
- - amniotic
- - - umbilical cord P02.5
- Bandl's ring (contraction), complicating delivery
- - affecting fetus or newborn P03.6
- barotrauma
- - nontraumatic
- - - newborn P28.8
- beat(s)
- - atrial, premature
- - - newborn P29.1
- - ectopic
- - - newborn P29.1
- - escaped, heart
- - - newborn P29.1
- - premature
- - - atrial
- - - - newborn P29.1
- - - newborn P29.1
- - - nodal
- - - - newborn P29.1
- - - ventricular (supra)
- - - - newborn P29.1
- bicornate or bicornis maternal uterus
- - in pregnancy or childbirth
- - - affecting fetus or newborn P03.8
- bigeminal pulse
- - fetal P20.9
- - newborn P29.1
- birth
- - abnormal NEC, affecting fetus or newborn P03.9
- - delayed, fetus P03.8
- - difficult NEC, affecting fetus or newborn P03.9
- - forced NEC, affecting fetus or newborn P03.8
- - forceps, affecting fetus or newborn P03.2
- - immature (between 28 and 37 completed weeks) P07.3

## ICD-10-Mortality Perinatal Subset - 2014

- - - extremely (less than 28 completed weeks) P07.2
- - induced, affecting fetus or newborn P03.8
- - injury P15.9 2a
- - - basal ganglia P11.1 2a
- - - brachial plexus NEC P14.3
- - - brain (compression) (pressure) P11.2 2a
- - - - anoxic P21.9
- - - central nervous system NEC P11.9 2a
- - - cerebellum P11.1 2a
- - - cerebral hemorrhage P10.1
- - - external genitalia P15.5
- - - eye P15.3
- - - face P15.4
- - - fracture
- - - - bone P13.9
- - - - - specified NEC P13.8
- - - - clavicle P13.4
- - - - femur P13.2
- - - - humerus P13.3
- - - - long bone, except femur P13.3
- - - - radius and ulna P13.3
- - - - skull P13.0
- - - - spine P11.5
- - - - tibia and fibula P13.3
- - - intracranial P11.2 2a
- - - - laceration or hemorrhage P10.9
- - - - - specified NEC P10.8
- - - intraventricular hemorrhage P10.2
- - - laceration
- - - - brain P10.1
- - - - by scalpel P15.8
- - - - peripheral nerve P14.9
- - - liver P15.0
- - - meninges
- - - - brain P11.1 2a
- - - - spinal cord P11.5
- - - nerve
- - - - brachial plexus P14.3
- - - - cranial NEC (except facial) P11.4
- - - - facial P11.3
- - - - peripheral P14.9
- - - - phrenic (paralysis) P14.2
- - - penis P15.5
- - - scalp P12.9
- - - scalpel wound P15.8
- - - scrotum P15.5
- - - skull NEC P13.1
- - - - fracture P13.0
- - - - specified site or type NEC P15.8
- - - spinal cord P11.5
- - - spine P11.5
- - - spleen P15.1
- - - sternomastoid (hematoma) P15.2
- - - subarachnoid hemorrhage P10.3
- - - subcutaneous fat necrosis P15.6
- - - subdural hemorrhage P10.0

## ICD-10-Mortality Perinatal Subset - 2014

- - - tentorial tear P10.4
- - - testes P15.5
- - - vulva P15.5
- - instrumental NEC, affecting fetus or newborn P03.8
- - multiple, affecting fetus or newborn P01.5
- - palsy or paralysis, newborn, NEC (birth injury) P14.9
- - post-term (42 weeks or more) P08.2
- - precipitate, affecting fetus or newborn P03.5
- - premature (infant) P07.3
- - prolonged, affecting fetus or newborn P03.8
- - retarded, affecting fetus or newborn P03.8
- - shock, newborn P96.8
- - trauma - *see* Birth, injury
- - twin, affecting fetus or newborn P01.5
- - vacuum extractor, affecting fetus or newborn P03.3
- - ventouse, affecting fetus or newborn P03.3 - - weight - *see also* Weight
- - - low (between 1000 and 2499 grams) P07.1
- - - - extremely (999 grams or less) P07.0
- - - 4500 grams or more P08.0
- biventricular failure (heart)
- - fetal, fetus P29.8
- - newborn P29.0
- bleb(s)
- - lung (ruptured)
- - - fetus or newborn P25.8
- bleeding (*see also* Hemorrhage)
- - rectum, rectal
- - - newborn P54.2
- - umbilical stump P51.9
- - vagina, vaginal (abnormal)
- - - maternal
- - - - affecting fetus or newborn P02.1 2a
- - - newborn P54.6
- blighted ovum
- - affecting fetus or newborn P01.8
- blood
- - dyscrasia
- - - fetus or newborn P61.9
- - in
- - - amniotic fluid P02.1 2a
- born in toilet (*see also* Birth, precipitate fetus or newborn) P03.5
- brachycardia
- - fetal P20.9
- - newborn P29.1
- bradyarrhythmia, cardiac
- - newborn P29.1
- bradycardia (sinoatrial) (sinus) (vagal)
- - fetal -*see* Distress, fetal
- - newborn P29.1
- bradytachyarrhythmia
- - newborn P29.1
- bradytachycardia
- - newborn P29.1
- breech
- - delivery NEC
- - - affecting fetus or newborn P03.0

## ICD-10-Mortality Perinatal Subset - 2014

- - extraction
- - - affecting fetus or newborn P03.0
- - presentation
- - - affecting fetus or newborn P01.7
- - - before labor, affecting fetus or newborn P01.7
- - - during labor, affecting fetus or newborn P03.0
- bronze baby syndrome P83.8
- bruise (skin surface intact) (*see also* Contusion)
- - fetus or newborn P54.5
- - scalp, due to birth injury, newborn P12.3
- - umbilical cord
- - - affecting fetus or newborn P02.6
- bubbly lung syndrome P27.0
- bulging membranes P02.8
- bulla(e)
- - lung
- - - fetus or newborn P25.8

C

- calcification
  - - lung (active) (postinfectious)
    - - - newborn P28.8
  - - umbilical cord P02.6
- calculus, calculi, calculous
  - - lung
    - - - newborn P28.8
  - - pulmonary
    - - - newborn P28.8
- candidiasis, candidal
  - - congenital P37.5
    - - neonatal P37.5
- capsulitis (joint)
  - - hepatic
    - - - newborn P78.1
- caput
  - - succedaneum P12.8
- catastrophe, catastrophe
  - - cardiorespiratory
    - - - newborn P28.8
- caul over face (causing asphyxia) P21.9
- cavitation of lung (*see also* Tuberculosis, pulmonary)
  - - nontuberculous
    - - - newborn P28.8
- cecitis
  - - with perforation, peritonitis, or rupture
    - - - newborn P78.1
- cellulitis (diffuse) (with lymphangitis)
  - - navel
    - - - newborn P38
  - - pelvis, pelvic (chronic)
    - - - male
      - - - -newborn P78.1
  - - retroperitoneal
    - - - newborn P78.1
  - - umbilicus
    - - - newborn P38
- cephalocele, cephal(o)hemocele
  - - fetus or newborn P52.8
    - - - birth injury P10.8
- cephalematoma, cephalhematoma (calcified)
  - - fetus or newborn (birth injury) P12.0
- cervicitis, maternal (acute) (chronic) (nonvenereal) (subacute) (with ulceration)
  - - complicating pregnancy
    - - - affecting fetus or newborn P00.8
- cesarean
  - - operation or section NEC (*see also* Delivery, cesarean)
    - - - affecting fetus or newborn P03.4
    - - - post mortem, affecting fetus or newborn P01.6
- cessation
  - - cardiac

## ICD-10-Mortality Perinatal Subset - 2014

- - - newborn P29.0
- - cardiorespiratory
- - - newborn P29.0
- - respiratory
- - - newborn P28.5 2a
- chemotherapy (session) (for)
- - cancer
- - - maternal, affecting fetus or newborn P04.1
- chickenpox (*see also* Varicella)
- - congenital P35.8
- chignon
- - fetus or newborn (birth injury) (from vacuum extraction) P12.1
- choleperitoneum, choleperitonitis
- - newborn P78.1
- chorioamnionitis
- - fetus or newborn P02.7
- chorioangioma
- - affecting fetus or newborn P02.2
- chorioretinitis
- - in (due to)
- - - toxoplasmosis
- - - - congenital (active) P37.1† H32.0\*
- chylothorax (nonfilarial)
- - fetal, fetus P28.8
- cicatrix (adherent) (contracted) (painful) (vicious) (*see also* Scar)
- - lung
- - - newborn P28.8
- circulation
- - failure (peripheral)
- - - fetus or newborn P29.8
- - persistent (fetal)
- - - newborn P29.3
- circum nuchal cord (X2) (X3) (X4) (X5) P02.5
- cirrhosis, cirrhotic (hepatic)
- - liver (chronic) (fatty) (hepatolienal) (hypertrophic) (nodular) (nonalcoholic) (splenomegalic)
- - - congenital P78.8
- cleidotomy, fetus or newborn P03.8
- clot (blood)(*see also* Embolism)
- - placenta P02.2
- - umbilical cord P02.6
- clotting
- - disseminated
- - - intravascular
- - - - newborn P60
- coagulation
- - fetus or newborn P60
- - intravascular (diffuse) (disseminated) (*see also* Defibrination)
- - - fetus or newborn P60
- - umbilical cord P02.6
- coagulopathy (*see also* Defect, coagulation)
- - consumption
- - - newborn P60
- - disseminated
- - - newborn P60
- - intravascular
- - - - diffuse

## ICD-10-Mortality Perinatal Subset - 2014

- - - - newborn P60
- - - disseminated
- - - - newborn P60
- cold
- - injury syndrome (newborn) P80.0
- colibacillosis NEC
- - generalized
- - - newborn P36.8
- collapse
- - cardiocirculatory
- - - newborn P29.8
- - cardiopulmonary
- - - newborn P29.8
- - cardiorespiratory
- - - newborn P29.8
- - cardiovascular
- - - newborn P29.8
- - circulatory (peripheral)
- - - fetus or newborn P29.8
- - heart
- - - fetal, fetus P29.8
- - - newborn P29.0
- - respiratory
- - - newborn P28.8
- - vascular (peripheral)
- - - during labor and delivery
- - - - affecting fetus or newborn P03.8
- - - fetus or newborn P29.8
- coma
- - newborn P91.5
- complications (from) (of)
- - intrauterine
- - - procedure NEC, affecting fetus or newborn P96.5
- - maternal sedation during labor and delivery
- - - affecting fetus or newborn P04.0
- - umbilical cord
- - - affecting fetus or newborn P02.6
- compression
- - during birth (fetus or newborn) P15.9 2a
- - lung
- - - newborn P28.8
- - umbilical cord
- - - affecting fetus or newborn P02.5
- - - - with cord prolapse P02.4
- compromise, compromised
- - respiratory
- - - newborn P28.5 2a
- congestion, congestive (chronic) (passive)
- - facial, due to birth injury P15.4
- - respiratory
- - - newborn P28.8
- - umbilical cord P02.6
- conjunctivitis (in) (due to)
- - chlamydial
- - - neonatal P39.1
- - neonatal P39.1

## ICD-10-Mortality Perinatal Subset - 2014

- constriction (*see also* Stricture)
- - ring dystocia (uterus)
- - - affecting fetus or newborn P03.6
- - umbilical cord P02.5
- contraction, contracture, contracted (deformity) NEC
- - hourglass
- - - uterus (complicating delivery)
- - - - affecting fetus or newborn P03.6
- - pelvis, maternal (acquired) (general)
- - - with disproportion (fetopelvic)
- - - - affecting fetus or newborn P03.1
- - premature
- - - atrium
- - - - newborn P29.1
- - - auriculoventricular
- - - - newborn P29.1
- - - heart
- - - - newborn P29.1
- - - junctional
- - - - newborn P29.1
- - - newborn P29.1
- - - ventricular
- - - - newborn P29.1
- - ring (Bandl's) (complicating delivery)
- - - affecting fetus or newborn P03.6
- - uterus, maternal
- - - abnormal NEC
- - - - affecting fetus or newborn P03.6
- contusion (skin surface intact)
- - fetus or newborn P54.5
- - scalp
- - - due to birth injury P12.3
- convulsions (idiopathic) (*see also* Seizure(s))
- - newborn P90
- cord (umbilical)
- - accident P02.6
- - around any part of the body (tightly) (with compression) (wrapped)
- - - affecting fetus or newborn P02.5
- - body (around) P02.5
- - break, broken P50.1
- - coiled, coiling P02.6
- - - with constriction (obstruction) (occlusion) P02.5
- - cyst P02.6
- - decreased blood flow P02.6
- - dilated P02.6
- - flat P02.5
- - injury P02.6
- - long (excessive) P02.6
- - narrowing P02.6
- - obstructed P02.6
- - occlusion (by) P02.6
- - - amniotic band P02.5
- - - fetal, fetus P02.5
- - supercoiled P02.5
- - thin P02.6
- - tight P02.5



## ICD-10-Mortality Perinatal Subset - 2014

- - tightly wrapped P02.5
- - twisted P02.5
- coupled rhythm
- - fetal P20.9
- - newborn P29.1
- cranioclasia, fetus P03.8
- craniotabes (cause unknown)
- - neonatal P96.3
- craniotomy, fetus P03.8
- cretin, cretinism (congenital) (endemic) (nongoitrous) (sporadic)
- - pelvis, maternal, with disproportion (fetopelvic)
- - - affecting fetus or newborn P03.1
- cripple
- - pulmonary
- - - newborn P28.8
- cyst (colloid) (mucous) (retention) (simple)
- - air, lung
- - - newborn P28.8
- - bronchogenic (mediastinal) (sequestration)
- - - newborn P28.8
- - lung
- - - newborn P28.8
- - periventricular, acquired, newborn P91.1
- cystic
- - lung disease
- - - newborn P28.8
- cystitis, maternal (exudative) (hemorrhagic) (septic) (suppurative)
- - complicating pregnancy
- - - affecting fetus or newborn P00.1
- cystocele(-urethrocele), maternal
- - in pregnancy or childbirth
- - - affecting fetus or newborn P03.8
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1

D

- dacryocystitis (acute) (phlegmonous)
  - - neonatal P39.1
- damage
  - - brain (nontraumatic)
    - - - anoxic, hypoxic
      - - - - at birth P21.9
      - - - - fetal P20.9
      - - - - intrauterine P20.9
      - - - - newborn P21.9
    - - - due to birth injury P11.2 2a
      - - - ischemic
        - - - - newborn P91.0
  - - newborn P11.2 2a
- eye, birth injury P15.3
  - - lung
    - - - newborn P28.8
- deadborn fetus NEC - *see* Death, fetus
- death
  - - fetus, fetal (cause not stated) (intrauterine)
    - - - in fetal death file P95
    - - - in mortality file P96.9 2a
  - - infant - from intrauterine coil P00.7
  - - intrauterine (late)
    - - - fetus or newborn
      - - - - in fetal death file P95
      - - - - in mortality file P96.9 2a
  - - neonatal NEC P96.8
  - - obstetric, maternal (cause unknown)
    - - - affecting fetus or newborn P01.6
- debility (chronic) (general)
  - - congenital or neonatal NEC P96.9 2a
    - - pulmonary
      - - - newborn P28.8
- decapitation
  - - fetal (to facilitate delivery) P03.8
- deciduitis (acute), affecting fetus or newborn P00.8
- decompensation
  - - lung
    - - - newborn P28.8
  - - pulmonary
    - - - newborn P28.8
  - - respiratory
    - - - newborn P28.8
- decrease(d)
  - - blood
    - - - flow
      - - - - cord (umbilical)
        - - - - - affecting fetus or newborn P02.6
- defect, defective
  - - coagulation (factor) (*see also* Deficiency, factor)
    - - - antepartum with hemorrhage, maternal

## ICD-10-Mortality Perinatal Subset - 2014

- - - - affecting fetus or newborn P02.1 2a
- - - newborn, transient P61.6
- defibrination (syndrome)
- - antepartum, maternal
- - - affecting fetus or newborn P02.1 2a
- - fetus or newborn P60
- deficiency, deficient
- - amniotic fluid
- - - affecting fetus or newborn P01.2 2a
- - cardiac (*see also* Insufficiency, myocardial)
- - - advanced
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - grave
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - severe
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - coagulation
- - - antepartum, maternal
- - - - affecting fetus or newborn P02.1 2a
- - - newborn, transient P61.6
- - myocardial (*see also* Insufficiency, myocardial)
- - - advanced
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - grave
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - severe
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - surfactant (disease) (lung) P28.0 2a
- - vitamin K
- - - of newborn P53
- deformity
- - fetal
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- - lung (congenital)
- - - acquired
- - - - newborn P28.8
- - pelvis, pelvic, maternal (acquired) (bony)
- - - with disproportion (fetopelvic)
- - - - affecting fetus or newborn P03.1
- - soft parts, maternal organs or tissues (of pelvis)
- - - in pregnancy or childbirth NEC
- - - - affecting fetus or newborn P03.8
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- degeneration, degenerative (hyaline)
- - lung
- - - newborn P28.8
- - villi (maternal), affecting fetus or newborn P02.8
- dehiscence

## ICD-10-Mortality Perinatal Subset - 2014

- - uterine
- - - affecting fetus or newborn P03.8
- dehydration (hypernatremic)
- - newborn P74.1
- delay, delayed
- - birth or delivery NEC
- - - affecting fetus or newborn P03.8
- - closure, ductus arteriosus (Botalli) P29.3
- - delivery, second twin, triplet, etc.
- - - affecting fetus or newborn P03.8
- - growth P05.9
- - primary respiration (*see also* Asphyxia, newborn) P21.9
- delivery (single)
- - breech NEC
- - - affecting fetus or newborn P03.0
- - cesarean (for)
- - - affecting fetus or newborn P03.4
- - extremely rapid
- - - newborn P03.5
- - forceps
- - - affecting fetus or newborn P03.2
- - precipitate
- - - affecting fetus or newborn P03.5
- - premature or preterm NEC
- - - affecting fetus or newborn P07.3
- - - - previable P07.2
- - vacuum extractor NEC
- - - affecting fetus or newborn P03.3
- - ventouse NEC
- - - affecting fetus or newborn P03.3
- demise
- - fetal - *see* Death, fetal
- density
- - lung (nodular)
- - - newborn P28.8
- dependence, dependency
- - due to
- - - drug NEC
- - - - maternal, complicating pregnancy or childbirth
- - - - - affecting fetus or newborn P04.4
- - - - - withdrawal symptoms in newborn P96.1
- depolarization, premature
- - atrial
- - - newborn P29.1
- - junctional
- - - newborn P29.1
- - newborn P29.1
- - specified NEC
- - - newborn P29.1
- - ventricular
- - - newborn P29.1
- depression
- - central nervous system (CNS) NEC
- - - newborn P91.4
- - cerebral
- - - newborn P91.4

## ICD-10-Mortality Perinatal Subset - 2014

- - function
  - - - cardiac
    - - - - newborn P29.1
  - - perinatal P96.8
  - - respiration, respiratory
    - - - newborn P28.5 2a
  - - vital centers
    - - - newborn P91.4
- dehiscence
  - - uterine
    - - - affecting fetus or newborn P03.8
- destruction, destructive
  - - live fetus to facilitate delivery
    - - - fetus P03.8
- development
  - - arrested
    - - - fetus P05.9
  - - incomplete P05.9
    - - - organ or site not listed - *see* Hypoplasia, by site
- device
  - - contraceptive
    - - - intrauterine
      - - - - affecting fetus or newborn P00.8
- diabetes, diabetic (controlled) (familial) (mellitus) (on insulin) (severe) (uncontrolled)
  - - A1 or A2 P70.0
    - - arising in pregnancy, maternal
      - - - affecting fetus or newborn P70.0
      - - class A1 or A2 P70.0
    - - complicating pregnancy or childbirth, maternal
      - - - affecting fetus or newborn P70.1
      - - - arising in pregnancy
        - - - - affecting fetus or newborn P70.0
      - - - gestational
        - - - - affecting fetus or newborn P70.0
    - - congenital
      - - - age less than 28 days P70.2
      - - gestational
        - - - affecting fetus or newborn P70.0
      - - neonatal (transient) P70.2
  - diarrhea, diarrheal (disease) (infantile) (*see also* Enteritis, Enterocolitis)
    - - neonatal (noninfectious) P78.3
  - difficult, difficulty (in)
    - - birth, affecting fetus or newborn P03.9
  - feeding
    - - newborn P92.9
      - - - breast P92.5
      - - - specified NEC P92.8
  - respiratory
    - - - newborn P28.8
  - dilatation
    - - advanced (cervix) (uteri) - *see also* Incompetency, cervix
    - - cervix (uteri), maternal - *see also* Incompetency, cervix
      - - - affecting fetus or newborn P01.0
      - - - incomplete, poor, slow
        - - - - affecting fetus or newborn P03.6
    - - cord (umbilical) P02.6

## ICD-10-Mortality Perinatal Subset - 2014

- discordant
- - fetal growth P01.5
- disease, diseased - *see also* Syndrome
- - airway (*see also* Disease, respiratory)
- - - hyperactive
- - - - newborn P28.8
- - - reactive
- - - - newborn P28.8
- - - restrictive
- - - - newborn P28.8
- - brain
- - - anoxic
- - - - newborn (*see also* Asphyxia, newborn) P21.9
- - breast
- - - inflammatory
- - - - fetus or newborn P83.4
- - bronchopulmonary (chronic)
- - - newborn P27.8
- - cardiorespiratory
- - - newborn P96.8
- - cardiovascular
- - - fetus or newborn P29.9
- - - - specified NEC P29.8
- - cerebrovascular
- - - hemorrhagic
- - - - newborn P52.4
- - circulatory (system) NEC
- - - fetus or newborn P29.9
- - - maternal, affecting fetus or newborn P00.3
- - cystic (adventitial) (artery)
- - - lung
- - - - newborn P28.8
- - facial nerve (seventh)
- - - newborn (birth injury) P11.3
- - heart (organic)
- - - congenital
- - - - maternal, affecting fetus or newborn P00.3
- - - - rheumatic (chronic) (inactive) (old) (quiescent) (with chorea)
- - - - maternal, affecting fetus or newborn P00.3
- - hemolytic (fetus) (newborn) P55.9
- - - due to
- - - - incompatibility
- - - - ABO (blood group) P55.1
- - - - - blood (group) (Duffy) (K(ell)) (Kidd) (Lewis) (M) (S) NEC P55.8
- - - - - Rh (blood group) (factor) P55.0
- - - - Rh-negative mother P55.0
- - - specified type NEC P55.8
- - hemorrhagic
- - - fetus or newborn P53
- - hyaline (diffuse) (generalized)
- - - membrane (lung) (newborn) P22.0
- - infectious, infective
- - - congenital P37.9
- - - - specified NEC P37.8
- - - maternal, complicating pregnancy or childbirth
- - - - affecting fetus or newborn P00.2

## ICD-10-Mortality Perinatal Subset - 2014

- - lung (acute) (chronic) (end-stage)
- - - bullous (obstructive)
- - - - congenital P25.0
- - - - newborn P25.0
- - - - cystic
- - - - newborn P28.8
- - - - interstitial
- - - - newborn P28.8
- - - - newborn P28.9
- - - - chronic P27.9
- - - - obstructive (chronic)
- - - - newborn (acute) P28.8
- - - - polycystic
- - - - newborn P28.8
- - - - restrictive
- - - - newborn P28.8
- - maternal, unrelated to pregnancy NEC, affecting fetus or newborn P00.9
- - pelvis, pelvic, maternal
- - - inflammatory (female)
- - - - complicating pregnancy
- - - - - affecting fetus or newborn P00.8
- - - placenta
- - - affecting fetus or newborn P02.2
- - - polycystic
- - - lung or pulmonary
- - - - newborn P28.8
- - - renal, maternal (functional) (pelvis)
- - - - complicating pregnancy
- - - - - affecting fetus or newborn P00.1
- - - respiratory (tract) (*see also* Disease, airway)
- - - - chronic NEC
- - - - - fetus or newborn P27.9
- - - - - specified NEC P27.8
- - - - newborn P28.9
- - - - - specified type NEC P28.8
- - - - specified type NEC
- - - - - newborn P28.8
- - - viral, virus (*see also* Disease, by type of virus)
- - - - congenital P35.9
- - - - - specified NEC P35.8
- - disorder (of) - *see also* Disease
- - - amino-acid
- - - - neonatal, transitory P74.8
- - - coagulation (factor) (*see also* Defect, coagulation)
- - - - newborn, transient P61.6
- - - digestive (system)
- - - - fetus or newborn P78.9
- - - - - specified NEC P78.8
- - - feeding
- - - - newborn P92.9
- - - fetus or newborn P96.9 2a
- - - - specified NEC P96.8
- - - glomerular (in)
- - - - septicemia NEC
- - - - - newborn P36.9
- - - - - streptococcal

## ICD-10-Mortality Perinatal Subset - 2014

- - - - newborn P36.1
- - heart action
- - - newborn P29.1
- - hematological
- - - fetus or newborn P61.9
- - - - specified NEC P61.8
- - hemorrhagic NEC
- - - newborn P53
- - integument, fetus or newborn P83.9
- - - specified NEC P83.8
- - membranes or fluid, amniotic
- - - affecting fetus or newborn P02.9
- - muscle
- - - tone, newborn P94.9
- - - - specified NEC P94.8
- - seizure
- - - maternal, affecting fetus P00.0
- - - newborn P90
- - skin
- - - fetus or newborn P83.9
- - - - specified NEC P83.8
- - temperature regulation, fetus or newborn P81.9
- - - specified NEC P81.8
- - thyroid (gland)
- - - function NEC, neonatal, transitory P72.2
- - tubulo-interstitial (in)
- - - septicemia
- - - - newborn P36.9
- - - - streptococcal
- - - - - newborn P36.1
- disproportion (fetopelvic)
- - affecting fetus or newborn P03.1
- distortion (congenital)
- - lumbar spine, maternal
- - - with disproportion
- - - - affecting fetus or newborn P03.1
- distress
- - cardiac
- - - congenital P20.9
- - - newborn P29.8
- - cardiopulmonary
- - - newborn P96.8
- - cardiorespiratory
- - - newborn P96.8
- - circulatory
- - - newborn P96.8
- - fetal (syndrome) P20-
- - - first noted
- - - - before onset of labor P20.0
- - - - during labor and delivery P20.1
- - intrauterine -see Distress, fetal
- - respiratory
- - - adult, so stated
- - - - newborn P22.9
- - - newborn P22.9
- - - - specified NEC P22.8



## ICD-10-Mortality Perinatal Subset - 2014

- - - syndrome (idiopathic)
- - - - adult
- - - - newborn P22.0
- - - - newborn P22.0
- disturbance - *see also* Disease
- - cerebral
- - - status, newborn P91.9
- - - - specified NEC P91.8
- - electrolyte (*see also* Imbalance, electrolyte)
- - - newborn, transitory P74.4
- - - - potassium balance P74.3
- - - - sodium balance P74.2
- - - - specified type NEC P74.4
- - endocrine (gland)
- - - neonatal, transitory P72.9
- - - - specified NEC P72.8
- - feeding
- - - newborn P92.9
- - metabolism
- - - neonatal, transitory P74.9
- - - - calcium and magnesium P71.9
- - - - - specified type NEC P71.8
- - - - carbohydrate metabolism P70.9
- - - - - specified type NEC P70.8
- - - - specified NEC P74.8
- - potassium balance, newborn P74.3
- - rhythm, heart
- - - newborn P29.1
- - sinus node
- - - newborn P29.1
- - sodium balance, newborn P74.2
- - temperature
- - - regulation, newborn P81.9
- - - - specified NEC P81.8
- double
- - uterus, maternal
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8
- Duchenne's
- - paralysis
- - - birth injury P14.0
- dyscrasia
- - blood
- - - fetus or newborn P61.9
- - - - specified type NEC P61.8
- dysfunction
- - atrioventricular node
- - - newborn P29.1
- - sinus node
- - - newborn P29.1
- - uterus, complicating delivery
- - - affecting fetus or newborn P03.6
- dysmaturity (*see also* Immaturity) P05.0
- - pulmonary (newborn) (Wilson-Mikity) P27.0
- dysplasia - *see also* Anomaly
- - bronchopulmonary (perinatal) P27.1

## ICD-10-Mortality Perinatal Subset - 2014

- - - less than 18 years of age P27.1
- - lung (congenital) (not associated with short gestation)
- - - associated with short gestation P28.0 2a
- dyspnea (nocturnal) (paroxysmal)
- - newborn P22.8
- dysrhythmia (*see also* Arrhythmia)
- - cardiac
- - - newborn P29.1
- - newborn P29.1
- dystocia
- - affecting fetus or newborn P03.1
- - cervical, maternal (failure of cervical dilation)
- - - affecting fetus or newborn P03.6
- - contraction ring, maternal (failure of cervical dilation)
- - - affecting fetus or newborn P03.6
- - fetal, fetus
- - - affecting fetus or newborn P03.1
- - maternal
- - - affecting fetus or newborn P03.1
- - positional
- - - affecting fetus or newborn P03.1
- - shoulder (girdle)
- - - affecting fetus or newborn P03.1
- - uterine NEC, maternal
- - - affecting fetus or newborn P03.6

E

- ecchymosis (*see also* Hemorrhage)
- - fetus or newborn P54.5
- eclampsia, eclamptic (coma) (convulsions) (delirium) (with pre-existing or pregnancy- related hypertension) NEC
- - pregnancy
- - - affecting fetus or newborn P00.0
- ectopic, ectopia (congenital)
- - atrial beats
- - - newborn P29.1
- - beats
- - - atrial
- - - - newborn P29.1
- - - newborn P29.1
- - - ventricular
- - - - newborn P29.1
- - ventricular beats
- - - newborn P29.1
- edema, edematous
- - brain
- - - due to birth injury P11.0 2a
- - - fetus or newborn (anoxia or hypoxia) P52.4
- - - - birth injury P11.0 2a
- - fetus or newborn NEC, classifiable to R60- P83.3
- - newborn, classifiable to R60- P83.3
- - umbilical P02.6
- effect, adverse NEC
- - anesthesia
- - - in
- - - - labor and delivery
- - - - - affecting fetus or newborn P04.0
- effusion
- - cardiac
- - - advanced
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - fetal, fetus P29.8
- - - grave
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - newborn P29.0
- - - severe
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - myocardium, myocardial
- - - advanced
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - fetal, fetus P29.8
- - - grave
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - newborn P29.0

## ICD-10-Mortality Perinatal Subset - 2014

- - - severe
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - pericardium, pericardial (noninflammatory)
- - - fetus or newborn P29.8
- - pleura, pleurisy, pleuritic, pleuropericardial
- - - fetus or newborn P28.8
- elevated, elevation
- - maternal serum P00.8
- embarrassment
- - respiratory
- - - newborn (*see also* Distress, respiratory)
- embolism (septic)
- - air (any site) (traumatic)
- - - newborn NEC P25.8
- - amniotic fluid (pulmonary)
- - - affecting fetus or newborn P03.8
- embryotomy (to facilitate delivery)
- - fetus P03.8
- emphysema (atrophic) (chronic) (interlobular) (lung) (obstructive) (pulmonary) (senile)
- - congenital (interstitial) P25.0
- - fetus or newborn (interstitial) P25.0
- - interstitial
- - - congenital P25.0
- - - perinatal period P25.0
- - mediastinal
- - - fetus or newborn P25.2
- encephalitis
- - in (due to)
- - - toxoplasmosis (acquired)
- - - - congenital P37.1† G05.2\*
- encephalopathia hyperbilirubinemia, newborn P57.9
- - due to isoimmunization (conditions in P55-) P57.0
- encephalopathy (acute) (cerebral)
- - hyperbilirubinemic, newborn P57.9
- - - due to isoimmunization (conditions in P55-) P57.0
- - hypoxic – *see* Damage, brain, anoxic
- - - ischaemic of newborn P91.6
- - in (due to)
- - - birth injury P11.1 2a
- - ischemic
- - - anoxic
- - - - newborn P91.6
- - - hypoxic
- - - - newborn P91.6
- - - newborn P91.0
- endometritis (nonspecific) (purulent) (septic) (suppurative)
- - affecting fetus or newborn P00.8
- - maternal, complicating pregnancy
- - - affecting fetus or newborn P00.8
- endotoxemia
- - newborn P36.9
- endotoxiosis
- - newborn P36.9
- engorgement
- - breast

## ICD-10-Mortality Perinatal Subset - 2014

- - - newborn P83.4
- enlargement, enlarged - *see also* Hypertrophy
- - placenta P02.2
- entanglement
- - umbilical cord(s)
- - - affecting fetus or newborn P02.5
- enteritis (diarrheal) (hemorrhagic) (*see also* Diarrhea, Enterocolitis)
- - necrotizing of fetus or newborn P77
- enterocolitis (*see also* Diarrhea, Enteritis)
- - necrotizing (chronic)
- - - fetus or newborn P77
- entrapment
- - head, fetus or newborn P03.8
- Erb(-Duchenne) paralysis (birth injury) (newborn) P14.0
- Erb's
- - palsy, paralysis (brachial) (birth) (newborn) P14.0
- erythema, erythematous
- - neonatorum P83.8
- - - toxic P83.1
- - rash, newborn P83.8
- - toxic, toxicum NEC
- - - newborn P83.1
- erythroblastosis (fetalis) (newborn) P55.9
- - due to
- - - ABO (antibodies) (incompatibility) (isoimmunization) P55.1
- - - Rh (antibodies) (incompatibility) (isoimmunization) P55.0
- erythroderma (*see also* Erythema)
- - neonatorum P83.8
- evisceration
- - birth injury P15.8
- excess, excessive, excessively
- - large
- - - fetus or infant P08.0
- - long
- - - umbilical cord (entangled)
- - - - affecting fetus or newborn P02.5
- - short
- - - umbilical cord
- - - - affecting fetus or newborn P02.6
- exhaustion (physical NEC)
- - fetus or newborn P96.8
- - maternal, complicating delivery
- - - affecting fetus or newborn P03.8
- exsanguination - *see* Hemorrhage
- - fetal P50.9
- extraction
- - with hook (fetus) P03.8
- - breech NEC
- - - affecting fetus or newborn P03.0
- extrasystoles (supraventricular)
- - atrial
- - - newborn P29.1
- - auricular
- - - newborn P29.1
- - junctional
- - - newborn P29.1

**ICD-10-Mortality Perinatal Subset - 2014**

- - newborn P29.1
- - ventricular
- - - newborn P29.1

F

- face, facial
- - presentation, affecting fetus or newborn P01.7
- failure, failed
- - biventricular
- - - advanced
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - fetal, fetus P29.8
- - - grave
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - newborn P29.0
- - - severe
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - cardiac (*see also* Failure, heart)
- - - newborn P29.0
- - cardiopulmonary (acute) (chronic) (*see also* Failure, heart)
- - - newborn P29.0
- - cardiorenal (chronic)
- - - fetal, fetus P29.8
- - - newborn P29.0
- - cardiovascular (*see also* Failure, cardiac)
- - - newborn P29.0
- - - pulmonary
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - cervical dilatation in labor
- - - affecting fetus or newborn P03.6
- - congestive (*see also* Failure, heart, congestive)
- - - circulatory
- - - - advanced
- - - - - fetal, fetus P29.8
- - - - - newborn P29.0
- - - - fetal, fetus P29.8
- - - - grave
- - - - - fetal, fetus P29.8
- - - - - newborn P29.0
- - - - newborn P29.0
- - - - severe
- - - - - fetal, fetus P29.8
- - - - - newborn P29.0
- - descent of head (at term)
- - - affecting fetus or newborn P03.1
- - expansion, terminal respiratory units (newborn) (primary) P28.0 2a
- - fetal head to enter pelvic brim
- - - affecting fetus or newborn P03.1
- - forceps NEC (with subsequent delivery by cesarean section)
- - - affecting fetus or newborn P03.1
- - heart (acute) (sudden)
- - - biventricular

## ICD-10-Mortality Perinatal Subset - 2014

- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - congenital
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - congestive
- - - - advanced
- - - - - fetal, fetus P29.8
- - - - - newborn P29.0
- - - - biventricular
- - - - - advanced
- - - - - - fetal, fetus P29.8
- - - - - - newborn P29.0
- - - - - fetal, fetus P29.8
- - - - - grave
- - - - - fetal, fetus P29.8
- - - - - newborn P29.0
- - - - - newborn P29.0
- - - - - severe
- - - - - fetal, fetus P29.8
- - - - - newborn P29.0
- - - - decompensated
- - - - - advanced
- - - - - fetal, fetus P29.8
- - - - - newborn P29.0
- - - - - fetal, fetus P29.8
- - - - - grave
- - - - - fetal, fetus P29.8
- - - - - newborn P29.0
- - - - - newborn P29.0
- - - - - severe
- - - - - fetal, fetus P29.8
- - - - - newborn P29.0
- - - - fetal, fetus P29.8
- - - - grave
- - - - - fetal, fetus P29.8
- - - - - newborn P29.0
- - - - left
- - - - - newborn P29.0
- - - - newborn P29.0
- - - - severe
- - - - - fetal, fetus P29.8
- - - - - newborn P29.0
- - - decompensated
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - - fetal, fetus P29.8
- - - left (ventricular) (*see also* Failure, ventricular, left)
- - - - newborn P29.0
- - - newborn P29.0
- - kidney
- - - congenital P96.0
- - - diabetic (acute) (any stage) (chronic) (*see also* E10-E14 with fourth character .2)
- - - - congenital P70.2
- - lungs to expand, fetus or newborn P28.0 2a
- - myocardial, myocardium (*see also* Failure, heart)



## ICD-10-Mortality Perinatal Subset - 2014

- - - renal
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - pulmonary
- - - newborn P28.5 2a
- - pump
- - - fetal, fetus P29.8
- - - newborn P29.0
- - respiration, respiratory
- - - newborn P28.5 2a
- - to
- - - take fluid
- - - - newborn P74.1
- - trial of labor (with subsequent cesarean section)
- - - affecting fetus or newborn P03.1
- - vacuum extraction NEC (with subsequent cesarean section)
- - - affecting fetus or newborn P03.1
- - ventilatory
- - - newborn P28.5 2a
- - ventricular (*see also* Failure, heart)
- - - left
- - - - fetal, fetus P29.8
- - ventouse NEC (with subsequent cesarean section)
- - - affecting fetus or newborn P03.1
- - ventricular (*see also* Failure, heart)
- - - left
- - - - newborn P29.0
- - vital centers, fetus or newborn P91.8
- fecalith (impaction)
- - congenital P76.8
- feeding
- - problem
- - - newborn P92.9
- - - - specified NEC P92.8
- fever
- - newborn P81.9
- - - environmental P81.0
- fibrillation
- - atrial or auricular (established)
- - - newborn P29.1
- - cardiac
- - - newborn P29.1
- - heart
- - - newborn P29.1
- - newborn P29.1
- - ventricular
- - - newborn P29.1
- fibroid (tumor) (M8890), maternal
- - in pregnancy or childbirth
- - - affecting fetus or newborn P03.8
- - uterus
- - - affecting fetus or newborn P03.8
- fibromyoma (M8890), maternal
- - uterus (corpus)
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8

## ICD-10-Mortality Perinatal Subset - 2014

- fibrosis, fibrotic
  - - cystic (of pancreas)
  - - - lung, acquired
  - - - - newborn P28.8
  - - lung (atrophic) (capillary) (chronic) (confluent) (massive) (perialveolar) (peribronchial)
  - - - congenital P27.8
  - - perineum, maternal, in pregnancy or childbirth
  - - - causing obstructed labor
  - - - - affecting fetus or newborn P03.1
  - - pulmonary (*see also* Fibrosis, lung)
  - - - congenital P27.8
- fistula
  - - peritoneum
  - - - newborn P78.1
- fit
  - - newborn P90
- flail
  - - chest (*see also* Fracture, rib(s), multiple)
  - - - newborn (birth injury) P13.8
- flat - *see also* Anomaly, by site
  - - cord (umbilical) P02.5
  - - pelvis, maternal
    - - - with disproportion (fetopelvic)
    - - - - affecting fetus or newborn P03.1
- floppy baby syndrome (nonspecific) P94.2
- flutter
  - - atrial or auricular
    - - - newborn P29.1
  - - heart
    - - - atrial
      - - - - newborn P29.1
      - - - newborn P29.1
      - - - ventricular
        - - - - newborn P29.1
    - - - newborn P29.1
    - - ventricular
      - - - newborn P29.1
- forced birth or delivery NEC
  - - affecting fetus or newborn P03.8
- forceps delivery NEC
  - - affecting fetus or newborn P03.2
- fracture (abduction) (adduction) (avulsion) (comminuted) (compression) (dislocation) (oblique) (separation)
  - - bone
    - - - birth injury P13.9
  - - clavicle (acromial end) (interligamentous) (shaft)
    - - - birth injury P13.4
  - - femur, femoral
    - - - birth injury P13.2
  - - skull
    - - - birth injury P13.0
  - - vertebra, vertebral (back) (body) (column) (neural arch) (pedicle) (spinous process) (transverse process)
    - - - birth injury P11.5
- funisitis
  - - affecting fetus or newborn P02.7
- funnel, funneling
  - - cervix

## **ICD-10-Mortality Perinatal Subset - 2014**

- - - maternal, affecting fetus or newborn P01.0
- - pelvis, maternal (acquired)
- - - with disproportion (fetopelvic)
- - - - affecting fetus or newborn P03.1

G

- gallop rhythm
- - fetal P20.9
- - newborn P29.1
- ganglioneitis
- - geniculate
- - - newborn (birth injury) P11.3
- gastro-esophageal reflux in newborn P78.8
- gestation (period) -*see also* Pregnancy
- - ectopic -*see* Pregnancy, by site
- - less than 28 weeks P07.2
- - 28 weeks but less than 37 weeks P07.3
- - 42 or more completed weeks P08.2
- glaucoma
- - traumatic
- - - newborn (birth injury) P15.3
- glomerulonephritis (*see also* Nephritis)
- - in (due to)
- - - septicemia
- - - - newborn P36.9
- glycosuria
- - maternal, affecting fetus or newborn P00.8
- goiter (plunging) (substernal)
- - congenital (nontoxic)
- - - transitory, with normal functioning P72.0
- - neonatal NEC P72.0
- gonococcus, gonococcal (disease) (infection)
- - maternal, complicating pregnancy or childbirth
- - - affecting fetus or newborn P00.2
- gonorrhoea (acute) (chronic)
- - maternal, complicating pregnancy or childbirth
- - affecting fetus or newborn P00.2
- granuloma
- - umbilicus
- - - newborn P38
- Gray syndrome (newborn) P93
- Grey syndrome (newborn) P93

H

- heart beat
  - - abnormality
    - - - fetal P20.9
    - - - newborn P29.1
  - - awareness
    - - - fetal P20.9
    - - - newborn P29.1
- heavy-for-dates NEC (fetus or infant) P08.1 - - exceptionally (4500 g or more) P08.0
- hemangioma
  - - placental, affecting fetus or newborn P02.2
- hematemesis
  - - newborn, neonatal P54.0
    - - - due to swallowed maternal blood P78.2
- hematoma (traumatic) (skin surface intact) (*see also* Injury, superficial)
  - - arachnoid
    - - - fetus or newborn (nontraumatic) P52.5
      - - - - birth injury P10.3
  - - birth injury NEC (*see also* Birth, injury) P15.8
  - - brain (traumatic)
    - - - fetus or newborn NEC P52.4
      - - - - birth injury P10.1
  - - cerebellar - *see* Hematoma, brain
    - - - newborn P52.6
  - - face, birth injury P15.4
  - - intracranial - *see* Injury, intracranial
    - - - newborn P52.9
  - - liver (subcapsular) (nontraumatic)
    - - - birth injury P15.0
  - - penis (nontraumatic)
    - - - birth injury P15.5
  - - scrotum, superficial
    - - - birth injury P15.5
  - - spinal (cord) (meninges) (*see also* Injury, spinal cord, by region)
    - - - fetus or newborn (birth injury) P11.5
  - - spine (*see also* Injury, spine)
    - - - fetus or newborn (birth injury) P11.5
  - - sternocleidomastoid, birth injury P15.2
  - - sternomastoid, birth injury P15.2
  - - subarachnoid (traumatic)
    - - - fetus or newborn (nontraumatic) P52.5
      - - - - birth injury P10.3
  - - subdural (traumatic)
    - - - fetus or newborn (localized) P52.8
      - - - - birth injury P10.0
  - - superficial, fetus or newborn P54.5
  - - testis (nontraumatic)
    - - - birth injury P15.5
  - - umbilical cord, complicating delivery
    - - - affecting fetus or newborn P02.6
  - - vulva (nonobstetric) (nontraumatic)
    - - - fetus or newborn (birth injury) P15.5

## ICD-10-Mortality Perinatal Subset - 2014

- hematomyelia (central)
  - - fetus or newborn (birth injury) P11.5
- hematorachis, hematorrhachis
  - - fetus or newborn (birth injury) P11.5
- hemiplegia
  - - newborn NEC P91.8
  - - - birth injury P11.9 2a
- hemolysis
  - - autoimmune
    - - - newborn P55.9
  - - intravascular NEC
    - - - newborn P60
  - - neonatal (excessive) P58.8
- hemopericardium
  - - newborn P54.8
- hemoperitoneum
  - - infectious
    - - - newborn P78.1
- hemoptysis
  - - newborn P26.9
- hemorrhage, hemorrhagic
  - - accidental antepartum
    - - - affecting fetus or newborn P02.1 2a
  - - adrenal (capsule) (gland)
    - - - newborn P54.4
  - - alveolar
    - - - lung
      - - - - newborn P26.8
    - - - respiratory
      - - - - newborn P26.8
  - - antepartum (*see also* Hemorrhage, pregnancy)
    - - - affecting fetus or newborn P02.1 2a
  - - bowel
    - - - newborn P54.3
  - - brain (interparenchymal) (miliary) (nontraumatic) (petechial)
    - - - due to
      - - - - birth injury P10.1
      - - - fetus or newborn P52.4
      - - - - birth injury P10.1
    - - - stem (basilar) (ganglia) (ganglion)
      - - - - newborn P52.4
  - - cerebellar, cerebellum (nontraumatic)
    - - - fetus or newborn P52.6
  - - cerebral, cerebrum (*see also* Hemorrhage, intracerebral)
    - - - fetus or newborn (anoxic) P52.4
      - - - - birth injury P10.1
  - - complicating
    - - - delivery
      - - - - affecting fetus or newborn P02.1 2a
      - - - - associated with coagulation defect, maternal (afibrinognemia) hyperfibrinolysis) (hypofibrinogeneia)
        - - - - - affecting fetus or newborn P03.8
  - - - - due to
    - - - - - low-lying placenta
      - - - - - - affecting fetus or newborn P02.0
    - - - - - placenta previa

## ICD-10-Mortality Perinatal Subset - 2014

- - - - - affecting fetus or newborn P02.0
- - - - - premature separation of placenta (*see also* Abruptio placentae)
- - - - - affecting fetus or newborn P02.1 2a
- - - - - trauma
- - - - - affecting fetus or newborn P03.8
- - - - - uterine leiomyoma
- - - - - affecting fetus or newborn P03.8
- - conjunctiva
- - newborn P54.8
- - cord, newborn (stump) P51.9
- - cutaneous
- - fetus or newborn P54.5
- - disease
- - fetus or newborn P53
- - epicranial subaponeurotic (massive), birth injury P12.2
- - extradural (traumatic)
- - birth injury P10.8
- - fetus or newborn (anoxic) (nontraumatic) P52.8
- - fetal, fetus (specified sites NEC) P50.9
- - from
- - - cut end of co-twin's cord P50.5
- - - placenta P50.2
- - - umbilical cord P50.8
- - - ruptured P50.1
- - - vasa previa P50.0
- - into
- - - co-twin P50.3
- - - maternal circulation P50.4
- - - specified NEC P50.8
- - fetal-maternal P50.4
- - gastroenteric
- - newborn P54.3
- - gastrointestinal (tract)
- - newborn P54.3
- - internal (organs) NEC
- - newborn P54.8
- - intestine
- - newborn P54.3
- - intra-alveolar (lung), newborn P26.8
- - intracerebral (nontraumatic) (*see also* Hemorrhage, brain)
- - fetus or newborn P52.4
- - - birth injury P10.1
- - intracranial (nontraumatic)
- - birth injury P10.9
- - fetus or newborn P52.9
- - - specified NEC P52.8
- - intraventricular (*see also* Hemorrhage, brain)
- - fetus or newborn (nontraumatic) P52.3
- - - birth injury P10.2
- - - grade
- - - - 1 P52.0
- - - - 2 P52.1
- - - - 3 P52.2
- - - - 4 P52.3
- - lung
- - - newborn P26.9

## ICD-10-Mortality Perinatal Subset - 2014

- - - - massive P26.1
- - - - specified NEC P26.8
- - massive umbilical, newborn P51.0
- - maternal, gestational, affecting fetus or newborn P02.1 2a
- - maternal-fetal P50.4
- - mucous membrane NEC
- - - newborn P54.8
- - nasal turbinate
- - - newborn P54.8
- - navel, newborn P51.9
- - newborn P54.9
- - - specified NEC P54.8
- - nose
- - - newborn P54.8
- - placenta NEC
- - - affecting fetus or newborn P02.1 2a
- - - from surgical or instrumental damage
- - - - affecting fetus or newborn P02.1 2a
- - - previa
- - - - affecting fetus or newborn P02.0
- - posterior fossa (nontraumatic) (*see also* Hemorrhage, brain)
- - - fetus or newborn P52.6
- - pregnancy - *see also* Hemorrhage, antepartum
- - - due to
- - - - abruptio placentae
- - - - - affecting fetus or newborn P02.1 2a
- - - - afibrinogenemia, or other coagulation defect (conditions in category D65-D68)
- - - - - affecting fetus or newborn P02.1 2a
- - - - leiomyoma, uterus
- - - - - affecting fetus or newborn P02.1 2a
- - - - placenta previa
- - - - - affecting fetus or newborn P02.0
- - - - premature separation of placenta (normally implanted) - *see also* Abruptio placentae
- - - - - affecting fetus or newborn P02.1 2a
- - - - threatened abortion
- - - - - affecting fetus or newborn P02.1 2a
- - - - trauma, affecting fetus or newborn P02.1 2a
- - - early
- - - - affecting fetus or newborn P02.1 2a
- - pulmonary
- - - newborn P26.9
- - - - massive P26.1
- - - - specified NEC P26.8
- - rectum (sphincter)
- - - newborn P54.2
- - renal
- - - newborn P54.8
- - skin
- - - fetus or newborn P54.5
- - slipped umbilical ligature P51.8
- - spinal (cord)
- - - fetus or newborn (birth injury) P11.5
- - stomach
- - - newborn P54.3
- - subarachnoid (nontraumatic)
- - - fetus or newborn P52.5



## ICD-10-Mortality Perinatal Subset - 2014

- - - - birth injury P10.3
- - subchorionic
- - - affecting fetus or newborn P02.1 2a
- - subconjunctival
- - - birth injury P15.3
- - subdural (acute) (nontraumatic)
- - - birth injury P10.0
- - - fetus or newborn (anoxic) (hypoxic) P52.8
- - - - birth injury P10.0
- - subependymal
- - - fetus or newborn P52.0
- - - - with intraventricular extension P52.1
- - - - - and intracerebral extension P52.2
- - suprarenal (capsule) (gland)
- - - newborn P54.4
- - tentorium (traumatic) NEC
- - - fetus or newborn (birth injury) P10.4
- - tracheobronchial
- - - newborn P26.0
- - umbilicus, umbilical
- - - cord
- - - - after birth, newborn P51.9
- - - - fetus P50.8
- - - - - from ruptured cord P50.1
- - - newborn P51.9
- - - - massive P51.0
- - - - slipped ligature P51.8
- - - stump P51.9
- - unavoidable (ante partum) (due to placenta previa)
- - - affecting fetus or newborn P02.0
- - vagina (abnormal)
- - - newborn P54.6
- - vasa previa
- - - affecting fetus or newborn P50.0
- - viscera NEC
- - - newborn P54.8
- hemothorax (bacterial) (nontuberculous)
- - newborn P54.8
- hepatitis
- - fetus or newborn (giant cell) (idiopathic) P59.2
- - in (due to)
- - - toxoplasmosis
- - - - congenital (active) P37.1† K77.0\*
- - neonatal (giant cell) (idiopathic) (toxic) P59.2
- - viral, virus
- - - congenital P35.3
- hernia, hernial (acquired) (recurrent)
- - lung (subcutaneous)
- - - newborn P28.8
- herpes, herpetic
- - simplex
- - - congenital P35.2
- honeycomb lung
- - newborn P28.8
- hyaline membrane (disease) (lung) (pulmonary) (newborn) P22.0
- hydramnios

## ICD-10-Mortality Perinatal Subset - 2014

- - affecting fetus or newborn P01.3
- hydrocele (spermatic cord) (testis) (tunica vaginalis)
- - communicating
- - - congenital P83.5
- - congenital P83.5
- - fetus or newborn P83.5
- hydrocephalus (acquired) (external) (internal) (malignant) (recurrent)
- - causing disproportion
- - - affecting fetus or newborn P03.1
- - due to toxoplasmosis (congenital) P37.1
- hydrops
- - fetal(is) or newborn (idiopathic) P83.2
- - - due to
- - - - ABO isoimmunization P56.0
- - - - hemolytic disease NEC P56.9
- - - - isoimmunization (ABO) (Rh) P56.0
- - - - Rh incompatibility P56.0
- hydrothorax (double) (pleura)
- - fetus or newborn P28.8
- hyperbilirubinemia NEC
- - neonatal (transient) (*see also* Jaundice, fetus or newborn) P59.9
- - - of prematurity P59.0
- hypercoiled, hypercoiling
- - cord (umbilical) P02.5
- - umbilicus (cord) P02.5
- hyperemesis (*see also* Vomiting)
- - gravidarum, maternal (mild)
- - - affecting fetus or newborn P01.8
- hypermagnesemia
- - neonatal P71.8
- hypermaturity (fetus or newborn) P08.2
- hypertension, hypertensive (accelerated) (benign) (essential) (idiopathic) (malignant) (primary) (systemic)
- - maternal, complicating pregnancy or childbirth
- - - affecting fetus or newborn P00.0
- - newborn P29.2
- - - pulmonary (persistent) P29.3
- - pulmonary (artery)
- - - of newborn (persistent) P29.3
- hyperthermia (of unknown origin)
- - newborn, environmental P81.0
- hyperthyroidism (apathetic) (latent) (pre-adult) (recurrent)
- - neonatal (transitory) P72.1
- hypertony, hypertonia, hypertonicity
- - congenital P94.1
- - uterus, uterine (contractions) (complicating delivery)
- - - affecting fetus or newborn P03.6
- hypertrophy, hypertrophic (*see also* Enlargement)
- - breast
- - - fetus or newborn P83.4
- hypocalcemia
- - neonatal P71.1
- - - due to cow's milk P71.0
- - phosphate-loading (newborn) P71.1
- hypoglycemia (spontaneous)
- - in infant of diabetic mother P70.1
- - - gestational diabetes P70.0

## ICD-10-Mortality Perinatal Subset - 2014

- - maternal, affecting fetus or newborn P00.8
- - neonatal (transitory) P70.4
  - - - iatrogenic P70.3
  - - - maternal diabetes P70.1
  - - - gestational P70.0
- - transitory neonatal P70.4
- hypomagnesemia
  - - neonatal P71.2
- hypoparathyroidism
  - - neonatal, transitory P71.4
- hypoperfusion
  - - lung
    - - - newborn P28.8
- hypoplasia, hypoplastic
  - - lung (congenital) (lobe)
    - - - with immaturity, prematurity or low birth weight P28.0 2a
    - - - associated with short gestation P28.0 2a
  - - pulmonary
    - - - associated with short gestation P28.0 2a
- hypoprothrombinemia (congenital) (hereditary) (idiopathic)
  - - newborn, transient P61.6
- hypothermia (accidental) (due to)
  - - neonatal P80.9
    - - - environmental (mild) NEC P80.8
    - - - mild P80.8
    - - - severe (chronic) (cold injury syndrome) P80.0
    - - - specified NEC P80.8
- hypothyroidism (acquired)
  - - neonatal, transitory P72.2
- hypoxia - *see also* Anoxia
  - - fetal -*see* Distress, fetal
- - intrauterine P20.9
  - - - first noted
    - - - - before onset of labor P20.0
    - - - - during labor and delivery P20.1
  - - newborn (*see also* Asphyxia, newborn) P21.9
- hysterotomy, affecting fetus or newborn P03.8

I

- icterus (*see also* Jaundice)
- - neonatorum (*see also* Jaundice, fetus or newborn) P59.9
- ileus (bowel) (colon) (inhibitory) (intestine)
- - meconium
- - - meaning meconium plug (without cystic fibrosis) P76.0
- - newborn
- - - due to meconium (with cystic fibrosis)
- - - - meaning meconium plug (without cystic fibrosis) P76.0
- - - without cystic fibrosis P76.0
- - - transitory P76.1
- imbalance
- - electrolyte
- - - neonatal, transitory NEC P74.4
- - - - potassium P74.3
- - - - sodium P74.2
- immature - *see also* Immaturity
- - birth (28 completed weeks or more but less than 37 completed weeks) P07.3
- - - extremely (less than 28 completed weeks) P07.2
- - placenta P02.2
- - villa (maternal), affecting fetus or newborn P02.8
- immaturity (28 completed weeks or more but less than 37 completed weeks) (*see also* Prematurity) P07.3
- - extreme (less than 28 completed weeks) P07.2
- - fetal, fetus P07.3
- - fetus or infant light-for-dates -*see* Light-for-dates
- - gross P07.2
- - infant P07.3
- - newborn P07.3
- - organ or site NEC -*see* Hypoplasia
- - pulmonary, fetus or newborn P28.0 2a
- - respiratory P28.0 2a
- immunization
- - ABO (*see also* Isoimmunization, ABO)
- - - affecting management of pregnancy
- - - - in fetus or newborn P55.1
- impaction, impacted
- - shoulder
- - - affecting fetus or newborn P03.1
- imperfect
- - aeration, lung (newborn) NEC P28.1
- improperly tied umbilical cord (causing hemorrhage) P51.8
- inability
- - to breathe properly
- - - newborn P28.8
- inadequate, inadequacy
- - development
- - - fetus P05.9
- - - lungs
- - - - associated with short gestation P28.0 2a
- - pulmonary
- - - function
- - - - newborn P28.5 2a

## ICD-10-Mortality Perinatal Subset - 2014

- - - ventilation, newborn P28.5 2a
- - ventilation
- - - newborn P28.5 2a
- incompatibility
- - ABO
- - - fetus or newborn P55.1
- - blood (group) (Duffy) (K(ell)) (Kidd) (Lewis) (M) (S) NEC
- - - fetus or newborn P55.8
- - Rh (blood group) (factor)
- - - fetus or newborn P55.0
- incompatible with life (newborn) (nonviable) P07.2
- incompetency, incompetent
- - cervix, cervical (os), maternal
- - - in pregnancy
- - - - affecting fetus or newborn P01.0
- incomplete
- - expansion, lungs (newborn) NEC P28.1
- incoordinate, incoordination
- - uterus (action) (contractions) (complicating delivery)
- - - affecting fetus or newborn P03.6
- induction of labor, affecting fetus or newborn P03.8
- inertia
- - uterus, uterine during labor
- - - affecting fetus or newborn P03.6
- infancy, infantile, infantilism
- - genitalia, genitals, maternal (after puberty)
- - - in pregnancy or childbirth NEC
- - - - affecting fetus or newborn P03.8
- - pelvis, maternal
- - - with disproportion (fetopelvic)
- - - - affecting fetus or newborn P03.1
- infant(s) - *see also* Infancy
- - of diabetic mother (syndrome of) P70.1
- - - gestational diabetes P70.0
- infarct, infarction (of)
- - placenta (complicating pregnancy)
- - - affecting fetus or newborn P02.2
- - umbilical cord P02.6
- infection, infected (opportunistic) (*see also* Inflammation)
- - amniotic fluid, sac or cavity
- - - affecting fetus or newborn P02.7
- - *Candida (albicans) (tropicalis)* (*see also* Candidiasis)
- - - neonatal P37.5
- - *Citrobacter, Citobacter (braakii) (freundii) (koseri)*
- - - newborn P37.8
- - *Clostridium*, clostridium
- - - congenital P39.8
- - - *difficile*
- - - - septicemia
- - - - - newborn P36.5
- - - *perfringens*
- - - - septicemia
- - - - - newborn P36.5
- - congenital NEC P39.9
- - - bacterial P39.8
- - - *Candida (albicans)* P37.5

## ICD-10-Mortality Perinatal Subset - 2014

- - - clostridium, other than *Clostridium tetani* P39.8
- - - cytomegalovirus P35.1
- - - *Escherichia coli* P39.8
- - - - sepsis P36.4
- - - fungus P37.8
- - - hepatitis, viral P35.3
- - - herpes (simplex) P35.2
- - - infectious or parasitic disease P37.9
- - - - specified NEC P37.8
- - - listeriosis (disseminated) P37.2
- - - malaria NEC P37.4
- - - - falciparum P37.3
- - - mycotic P37.8
- - - parvovirus P35.8
- - - *Plasmodium falciparum* P37.3
- - - poliomyelitis P35.8
- - - polymicrobial P37.9
- - - rubella P35.0 2a
- - - salmonella P39.8
- - - skin P39.4
- - - streptococcal NEC P39.8
- - - - sepsis P36.1
- - - - - group B P36.0
- - - toxoplasmosis (acute) (chronic) (subacute) P37.1
- - - tuberculosis P37.0
- - - urinary (tract) P39.3
- - - vaccinia P35.8
- - - virus P35.9
- - - - specified type NEC P35.8
- - cytomegalovirus, cytomegaloviral
- - - congenital P35.1
- - *Enterobacter (cloacae)*
- - - newborn P39.8
- - *Escherichia (E.) coli* NEC
- - - congenital P39.8
- - - - sepsis P36.4
- - - generalized
- - - - newborn P36.4
- - fetus (*see also* Infection, congenital) P39.9
- - - intra-amniotic NEC P39.2
- - fungus NEC
- - - newborn P37.8
- - genital organ or tract, maternal
- - - complicating pregnancy
- - - - affecting fetus or newborn P00.8
- - hemolysis NEC
- - - newborn P36.9
- - herpes (simplex) (*see also* Herpes)
- - - congenital P35.2
- - intra-amniotic, fetus P39.2
- - intrauterine (complicating pregnancy)
- - - fetus or newborn P00.8
- - - specified infection NEC, fetus P39.2
- - kidney, maternal (cortex) (hematogenous)
- - - complicating pregnancy
- - - - affecting fetus or newborn P00.1

## ICD-10-Mortality Perinatal Subset - 2014

- - *Listeria monocytogenes* (*see also* Listeriosis)
- - - congenital P37.2
- - lung NEC
- - - atypical mycobacterium
- - - newborn P23.9
- - *Monilia* (*see also* Candidiasis)
- - - neonatal P37.5
- - navel
- - - newborn P38
- - newborn P39.9
- - - skin P39.4
- - - specified type NEC (*see also* Infection, congenital) P39.8
- - pelvis
- - - male
- - - - newborn P78.1
- - perinatal period P39.9
- - - specified type NEC P39.8
- - pneumococcus, pneumococcal NEC
- - - generalized (purulent)
- - - - with pneumonia
- - - - - newborn P23.6
- - - - newborn P36.1
- - polymicrobial
- - - newborn P37.9
- - respiratory (tract) NEC
- - - chronic
- - - - newborn P27.8
- - - fetus P28.8
- - - newborn, neonatal P28.8
- - rubella (*see also* Rubella)
- - - congenital P35.0 2a
- - *Salmonella* (*arizonae*) (*cholerae-suis*) (*enteritidis*) (*typhimurium*)
- - - congenital P39.8
- - skin (local) (staphylococcal) (streptococcal)
- - - newborn P39.4
- - staphylococcal NEC (hemolytic)
- - - newborn P39.8
- - streptococcal NEC (hemolytic)
- - - congenital P39.8
- - - - sepsis P36.1
- - - - - group B P36.0
- - umbilicus
- - - fetus or newborn P38
- - urinary (tract) NEC
- - - maternal, complicating pregnancy
- - - - affecting fetus or newborn P00.1
- - - newborn P39.3
- - virus NEC
- - - chest
- - - - newborn P28.8
- - widespread
- - - newborn P36.9
- inflammation, inflamed, inflammatory (with exudation) (*see also* Infection)
- - lung (acute) (*see also* Pneumonia)
- - - chronic
- - - - newborn P27.8

## ICD-10-Mortality Perinatal Subset - 2014

- - navel
- - - newborn P38
- - umbilicus
- - - newborn P38
- influenza (specific virus not identified)
- - maternal, affecting fetus or newborn P00.2
- inhalation
- - meconium (newborn) P24.0
- injury, injuries
- - birth (*see also* Birth injury) P15.9 2a
- - brachial plexus
- - - newborn P14.3
- - brain
- - - anoxic
- - - - at birth P21.9
- - - - newborn P21.9
- - childbirth (fetus or newborn) (*see also* Birth, injury) P15.9 2a
- - delivery (fetus or newborn) P15.9 2a
- - ischemic (*see also* Ischemia)
- - - hypoxic, newborn P21.9
- - maternal, during pregnancy, affecting fetus or newborn P00.5
- - nerve
- - - facial
- - - - newborn P11.3
- - scalp
- - - fetus or newborn (birth injury) P12.9
- - - - due to monitoring (electrode) (sampling incision) P12.4
- - - - specified NEC P12.8
- - skeleton, skeletal
- - - birth injury P13.9
- - - - specified part NEC P13.8
- insertion
- - cord (umbilical) lateral or velamentous (*see also* Placenta, abnormal)
- - - abnormal P02.2
- - - marginal P02.2
- - velamentous (cord) (*see also* Placenta, abnormal)
- - - affecting fetus or newborn P02.2
- inspissated bile syndrome (newborn) P59.1
- insufficiency, insufficient
- - cardiac (*see also* Insufficiency, myocardial)
- - - congestive
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - circulatory NEC
- - - fetus or newborn P29.8
- - heart (*see also* Insufficiency, myocardial)
- - - newborn P29.0
- - lung (*see also* Insufficiency, pulmonary)
- - - newborn P28.5 2a
- - myocardial, myocardium (acute) (chronic)
- - - - fetal, fetus P29.8
- - - - newborn P29.0
- - placental
- - - affecting fetus or newborn P02.2
- - pulmonary
- - - newborn P28.5 2a



## ICD-10-Mortality Perinatal Subset - 2014

- - respiratory (acute)
- - - newborn P28.5 2a
- - subendocardial
- - - fetal, fetus P29.8
- - - newborn P29.0
- - ventilation, ventilatory
- - - newborn P28.8
- insufficiently
- - tied umbilical cord P51.8
- interruption (of)
- - oxygen cycle
- - - newborn P28.8
- - respiration
- - - newborn P28.8
- intolerance
- - fetal NEC (to labor) P20.9
- intoxication NEC
- - drug
- - - newborn P93
- - septic (*see also* Shock, septic)
- - - general
- - - - newborn P36.9
- intrauterine contraceptive device (IUD)
- - affecting fetus or newborn P00.8
- irregular, irregularity
- - action, heart
- - - newborn P29.1
- irritable, irritability
- - cerebral, in newborn P91.3
- ischemia, ischemic
- - cerebral (chronic) (generalized)
- - - newborn P91.0
- - - prenatal P91.0
- - myocardium, myocardial
- - - transient, of newborn P29.4
- isoimmunization NEC (*see also* Incompatibility)
- - fetus or newborn P55.9
- - - with
- - - - hydrops fetalis P56.0
- - - - kernicterus P57.0
- - - ABO P55.1
- - - anti-Kell P55.8
- - - Rh P55.0
- - - specified type NEC P55.8

J

- jaundice (yellow)
- - breast-milk (inhibitor) P59.3
- - due to or associated with
- - - delayed conjugation P59.8
- - - - associated with preterm delivery P59.0
- - - preterm delivery P59.0
- - fetus or newborn (physiological) P59.9
- - - due to or associated with
- - - - ABO
- - - - - antibodies P55.1
- - - - - incompatibility, maternal/fetal P55.1
- - - - - isoimmunization P55.1
- - - - absence or deficiency of enzyme system for bilirubin conjugation (congenital) P59.8
- - - - bleeding P58.1
- - - - breast milk inhibitors to conjugation P59.3
- - - - bruising P58.0
- - - - delayed conjugation P59.8
- - - - - associated with preterm delivery P59.0
- - - - drugs or toxins
- - - - - given to newborn P58.4
- - - - - transmitted from mother P58.4
- - - - excessive hemolysis NEC P58.9
- - - - - specified type NEC P58.8
- - - - hepatocellular damage P59.2
- - - - hereditary hemolytic anemia P58.8
- - - - incompatibility, maternal/fetal NEC P55.9
- - - - infection P58.2
- - - - inspissated bile syndrome P59.1
- - - - isoimmunization NEC P55.9
- - - - polycythemia P58.3
- - - - preterm delivery P59.0
- - - - Rh
- - - - - antibodies P55.0
- - - - - incompatibility, maternal/fetal P55.0
- - - - - isoimmunization P55.0
- - - - swallowed maternal blood P58.5
- - - specified cause NEC P59.8
- - nuclear, newborn (*see also* Kernicterus of newborn) P57.9

**K**

- kernicterus
- - newborn P57.9
- - - due to isoimmunization (conditions in P55-) P57.0
- - - specified type NEC P57.8
- Klumpke(-Déjerine) palsy, paralysis (birth) (newborn) P14.1
- knot (true), umbilical cord
- - affecting fetus or newborn P02.5

L

- labor (*see also* Delivery)
- - abnormal NEC
- - - affecting fetus or newborn P03.6
- - arrested active phase
- - - affecting fetus or newborn P03.6
- - desultory
- - - affecting fetus or newborn P03.6
- - dyscoordinate
- - - affecting fetus or newborn P03.6
- - forced or induced, affecting fetus or newborn P03.8
- - hypertonic
- - - affecting fetus or newborn P03.6
- - hypotonic
- - - affecting fetus or newborn P03.6
- - incoordinate
- - - affecting fetus or newborn P03.6
- - irregular
- - - affecting fetus or newborn P03.6
- - obstructed
- - - affecting fetus or newborn P03.1
- - - by or due to
- - - - abnormal
- - - - - pelvis (bony)
- - - - - - affecting fetus or newborn P03.1
- - - - - presentation or position
- - - - - - affecting fetus or newborn P03.1
- - - - - size, fetus
- - - - - - affecting fetus or newborn P03.1
- - - - cephalopelvic disproportion (normally formed fetus)
- - - - - affecting fetus or newborn P03.1
- - - - disproportion, fetopelvic NEC
- - - - - affecting fetus or newborn P03.1
- - - - malpresentation
- - - - - affecting fetus or newborn P03.1
- - precipitate
- - - affecting fetus or newborn P03.5
- - premature or preterm
- - - affecting fetus or newborn (*see also* Prematurity) P07.3
- - prolonged or protracted
- - - affecting fetus or newborn P03.8
- laceration (*see also* Wound)
- - brain (any part) (cortex) (diffuse) (membrane)
- - - during birth P10.8
- - - - with hemorrhage P10.1
- - cerebral (diffuse)
- - - during birth P10.8
- - - - with hemorrhage P10.1
- - intracranial NEC
- - - birth injury P10.9
- - spinal cord (meninges) (*see also* Injury, spinal cord, by region)
- - - fetus or newborn (birth injury) P11.5

## ICD-10-Mortality Perinatal Subset - 2014

- lack of
  - - exchange of gases
    - - - newborn P28.5 2a
  - large-for-dates NEC (fetus or infant) P08.1
    - - exceptionally (4500 g or more) P08.0
  - laryngismus (stridulus)
    - - congenital P28.8
  - lateroversion
    - - uterus, uterine, maternal (cervix) (postinfectious) (postpartal, old)
      - - - in pregnancy or childbirth
        - - - - affecting fetus or newborn P03.8
  - leak, leakage
    - - air (bronchus) (intrathoracic) (lung) (pleural) (pulmonary) (thorax) - *see* Pneumothorax
      - - - newborn P25.1
  - lesion (nontraumatic)
    - - lung
      - - - newborn P28.8
    - - pulmonary
      - - - newborn P28.8
    - - vascular
      - - - umbilical cord, complicating delivery
        - - - - affecting fetus or newborn P02.6
  - leukomalacia, cerebral
    - - newborn P91.2
  - lie, abnormal (maternal care) (*see also* Presentation, fetal, abnormal)
    - - before labor, affecting fetus or newborn P01.7
  - light
    - - fetus or newborn, for gestational age P05.0
    - light-for-dates (infant) P05.0
      - - and small-for-dates P05.1
  - listeriosis, listerellosis
    - - congenital (disseminated) P37.2
      - - fetal P37.2
      - - neonatal (disseminated) P37.2
  - lithopedion - *see* Death, fetus
  - locked twins causing obstructed labor
    - - affecting fetus or newborn P03.1
  - long
    - - labor
      - - - affecting fetus or newborn P03.8
  - loss (of)
    - - fluid (acute)
      - - - fetus or newborn P74.1
  - low
    - - alphafetoprotein
      - - - maternal, affecting fetus or newborn P00.8
    - - amniotic fluid (volume) P01.2 2a
      - - - birthweight (2499 grams or less) (*see also* Weight) P07.1
        - - - - extreme (999 grams or less) P07.0
      - - - for gestational age P05.0
    - - cardiac
      - - - output
        - - - - newborn P29.8
    - - lung volume
      - - - newborn P28.8
  - lupus

## ICD-10-Mortality Perinatal Subset - 2014

- - erythematosus (discoïd) (local)
- - - systemic
- - - - maternal, affecting fetus or newborn P00.8
- luxation
- - eyeball (nontraumatic)
- - - birth injury P15.3
- lymphadenopathy (generalized)
- - due to toxoplasmosis
- - - congenital (acute) (chronic) (subacute) P37.1

M

- maceration
- - fetus (cause not stated) - *see* Death, fetus
- - newborn - *see* Death, fetus
- macrosomia P08.0
- malaria, malarial (fever)
- - congenital NEC P37.4
- - - falciparum P37.3
- male type pelvis, maternal
- - with disproportion (fetopelvic)
- - - affecting fetus or newborn P03.1
- malformation (congenital) - *see also* Anomaly
- - pelvic organs or tissues NEC, maternal
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- - umbilical cord NEC (complicating delivery)
- - - affecting fetus or newborn P02.6
- malnutrition
- - intrauterine or fetal P05.2
- - - light-for-dates P05.0
- - - small-for-dates P05.1
- - maternal, affecting fetus or newborn P00.4
- mastitis (acute) (infective) (nonpuerperal) (subacute)
- - infective
- - - newborn P39.0
- - neonatal (noninfective) P83.4
- - - infective P39.0
- maternal condition, affecting fetus or newborn P00.9
- - acidosis P00.8
- - acute yellow atrophy of liver P00.8
- - alcohol use P04.3
- - anemia P00.8
- - anesthesia or analgesia P04.0
- - antidiarrheal (syndrome) (antibody) P00.8
- - antiphospholipid syndrome P00.8
- - appendicitis P00.8
- - asthma P00.3
- - autoimmune disease P00.8
- - bipolar P00.8
- - bleeding
- - - vaginal P02.1 2a
- - blood loss (gestational) P02.1 2a
- - bulging membranes P02.8
- - cancer chemotherapy P04.1
- - cardiomyopathy P00.3
- - cerebral palsy P00.8
- - cesarean section, previous P03.8
- - cholestasis P00.8
- - chorioamnionitis P02.7
- - circulatory disease (conditions in I00-I99, Q20-Q28) P00.3

## ICD-10-Mortality Perinatal Subset - 2014

- - cirrhosis (biliary) P00.8
- - complication of pregnancy NEC P01.9
- - conditions NEC in
  - - - A00-B99 P00.2
  - - - E00-E07 P00.8
  - - - E10-E14 P70.1
  - - - E40-E64 P00.4
  - - - I00-I99 P00.3
  - - - J00-J06.9 P00.3
  - - - J09-J11 P00.2
  - - - J12.0-J98.9 P00.3
  - - - N00-N39 P00.1
  - - - O10-O11 P00.0
  - - - O13-O16 P00.0
  - - - Q20-Q28 P00.3
  - - - Q30-Q34 P00.3
  - - - S00-T79 P00.5
- - congenital heart disease (conditions in Q20-Q24) P00.3
- - cortical necrosis of kidney P00.1
- - cramps (abdominal) P01.8
- - Crohn's disease P00.8
- - cytotoxic drug P04.1
- - death P01.6
- - deficiency
  - - - protein S P00.8
- - degeneration
  - - - villa, villous P02.8
- - depression P00.8
- - diabetes mellitus (conditions in E10-E14) P70.1
  - - - class A1 or A2 P70.0
  - - - gestational P70.0
- - disease NEC P00.9
- - disseminated intravascular coagulation (DIC) P02.1 2a
- - drug abuse or addiction or dependence P04.4
- - drug use NEC P04.9
  - - - named drug
    - - - - listed under Addiction or Dependence P04.4
    - - - - not listed under Addiction or Dependence P04.8
- - early onset of delivery P07.3
- - eclampsia P00.0
- - edema (gestational) P00.8
- - elevated afp P00.8
- - elevated enzymes P00.8
- - elevated serum hcg P00.8
- - exposure to environmental chemical substances P04.6
- - factor V Leiden mutation P00.8
- - fatty
  - - - liver, acute P00.8
- - febrile P00.8
- - fibroid (uterus) P03.8
- - funneling of cervix P01.0
- - gastric banding P00.6
- - gastric bypass P00.6
- - genital tract infections NEC P00.8
- - glomerular diseases (conditions in N00-N08) P00.1
- - glucose intolerance P00.8



## ICD-10-Mortality Perinatal Subset - 2014

- - glycosuria P00.8
- - hemolytic-uremic syndrome P00.8
- - hemorrhage
- - - gestational P02.1 2a
- - - intestinal P00.8
- - - subchorionic P02.1 2a
- - hepatitis, malignant P00.8
- - - acute or subacute P00.2
- - heterozygote C677T (MTHFR) P00.8
- - high blood sugar P00.8
- - HIV positive P00.8
- - hypercoagulation (state) P00.8
- - hyperemesis (gravidarum) P01.8
- - hyperglycemia P00.8
- - hypertension (conditions in O10-O11, O13-O16) P00.0
- - hyperthyroidism P00.8
- - hypoglycemia P00.8
- - infection NEC P00.2
- - - chlamydia P00.2
- - - streptococcal B P00.2
- - infectious and parasitic diseases (conditions in A00-B99, J09-J11) P00.2
- - influenza P00.2
- - - manifest influenza in the infant P35.8
- - injury (conditions in S00-T79) P00.5
- - insufficiency
- - - cervical P01.0
- - intrauterine coil P00.8
- - ketoacidosis P00.8
- - kidney stone(s) P00.1
- - leukocytosis P00.8
- - lipoproteinemia (elevated) P00.8
- - lupus P00.8
- - malaria P00.2
- - - manifest malaria NEC in infant or fetus P37.4
- - - - falciparum P37.3
- - malnutrition P00.4
- - mental illness P00.8
- - mirror syndrome P01.8
- - MTHFR P00.8
- - myocardial infarction (history of) P00.3
- - nausea P01.8
- - necrosis of liver P00.8
- - neoplasm NEC P00.8
- - nephritis, nephrotic syndrome and nephrosis (conditions in N00-N08) P00.1
- - noxious influence transmitted via breast milk or placenta P04.9
- - - specified NEC P04.8
- - nutritional disorder (conditions in E40-E64) P00.4
- - obesity P00.8
- - operation (unrelated to current pregnancy) P00.6
- - parvovirus P00.2
- - pre-eclampsia P00.0
- - previous surgery, uterus or pelvic organs P03.8
- - proteinuria P00.1
- - pyelitis or pyelonephritis P00.1
- - renal disease or failure P00.1
- - respiratory disease (conditions in J00-J06, J12-J99, Q30-Q34) P00.3

## ICD-10-Mortality Perinatal Subset - 2014

- - rheumatic heart disease (chronic) (conditions in I05-I09) P00.3
- - rheumatoid arthritis P00.8
- - rubella (conditions in B06 ) P00.2
- - - manifest rubella in the infant or fetus P35.0 2a
- - seizure (disorder) P00.0
- - septate vagina P03.8
- - shock (hypovolemic) P00.8
- - short, shortened cervix P00.8
- - specified condition NEC P00.8
- - stenosis or stricture of vagina P03.8
- - surgery (unrelated to current pregnancy) P00.6
- - - to uterus or pelvic organs P03.8
- - syncope P00.8
- - syphilis (conditions in A50-A53) P00.2
- - thalassemia P00.8
- - thrombocytopenia P00.8
- - thrombophilia P00.8
- - thrombophlebitis P00.3
- - tobacco use P04.2
- - toxemia (of pregnancy) P00.0
- - toxoplasmosis (conditions in B58-) P00.2
- - - manifest toxoplasmosis (acute) (chronic) (subacute) in the infant or fetus P37.1
- - transmission of chemical substance through the placenta (*see also* Absorption, chemical, through placenta) P04.8
- - uremia P00.1
- - urinary tract conditions (conditions in N00-N39) P00.1
- - vaginal leaking (amniotic fluid) P01.1 2a
- - vaginosis, bacterial P00.8
- - vomiting (pernicious) (persistent) (vicious) P01.8
- - Von Willebrand's disease P00.8
- meconium P03.8
- - foul smelling, fetus or newborn P03.8
- - heavy, fetus or newborn P03.8
- - ileus, fetus or newborn
- - in liquor - *see also* Distress, fetal
- - light, fetus or newborn P03.8
- - obstruction, fetus or newborn P76.0
- - particulate, fetus or newborn P03.8
- - peritonitis P78.0
- - plug syndrome (newborn) NEC P76.0
- - stain
- - - fetus or newborn P20.9
- - thick, fetus or newborn P03.8
- melena
- - newborn, neonatal P54.1
- - - due to swallowed maternal blood P78.2
- membrane(s), membranous
- - thick (maternal), affecting fetus or newborn P02.8
- membranitis
- - affecting fetus or newborn P02.7
- meningoencephalitis (*see also* Encephalitis)
- - in (due to)
- - - toxoplasmosis
- - - - congenital P37.1† G05.2\*
- meningoencephalomyelitis (*see also* Meningoencephalitis)
- - due to
- - - toxoplasma or toxoplasmosis

## ICD-10-Mortality Perinatal Subset - 2014

- - - congenital P37.1† G05.2\*
- microcephalus, microcephalic, microcephaly
- - due to toxoplasmosis (congenital) P37.1
- Mikity-Wilson disease or syndrome P27.0
- molding, head (during birth) P13.1
- mole (pigmented) (M8720)
- - hydatid, hydatidiform (benign) (complicating pregnancy) (delivered) (undelivered) (M9100)
- - - fetus or newborn P02.2
- - - newborn P02.2
- moniliasis (*see also* Candidiasis)
- - neonatal P37.5
- mosaicism, mosaic (autosomal) (chromosomal)
- - placental (confined) (CPM) P02.2
- mucus
- - asphyxia or suffocation (*see also* Asphyxia, mucus)
- - - newborn P24.1
- - plug (*see also* Asphyxia, mucus)
- - - aspiration, newborn P24.1
- - - tracheobronchial
- - - - newborn P24.1
- multiple, multiplex
- - birth, affecting fetus or newborn P01.5
- myasthenia, myasthenic
- - gravis
- - - neonatal, transient P94.0

## ICD-10-Mortality Perinatal Subset - 2014

### N

- narrowing
  - - cord (umbilical) P02.6
- necrosis, necrotic (ischemic) (necrotizing)
  - - fat (generalized)
    - - - subcutaneous, due to birth injury P15.6
  - - kidney (bilateral)
    - - - tubular
      - - - - complicating pregnancy
        - - - - - affecting fetus or newborn P00.1
    - - liver (cell), maternal
      - - - complicating pregnancy or childbirth
        - - - - affecting fetus or newborn P00.8
  - - mesentery
    - - - fat
      - - - - newborn P78.1
  - - omentum
    - - - fat
      - - - - newborn P78.1
  - - peritoneum
    - - - fat
      - - - - newborn P78.1
  - - subcutaneous fat, fetus or newborn P83.8
  - - umbilical cord, affecting fetus or newborn P02.6
- neonatal
  - - abstinence syndrome P96.1
  - - esophageal reflux P78.8
- nephritis, nephritic, maternal
  - - complicating pregnancy
    - - - with secondary hypertension, pre-existing
      - - - - affecting fetus or newborn P00.0
      - - - - affecting fetus or newborn P00.1
- neuritis
  - - cranial nerve
    - - - seventh or facial
      - - - - newborn (birth injury) P11.3
  - - facial
    - - - newborn (birth injury) P11.3
- neutropenia, neutropenic (congenital) (cyclic) (drug-induced) (periodic) (primary) (splenic) (toxic)
  - - neonatal, transitory (isoimmune) (maternal transfer) P61.5
- no amniotic fluid P01.2 2a
- nodal rhythm
  - - newborn P29.1
- nodule(s), nodular
  - - lung (solitary)
    - - - newborn P28.8
- nonexpansion, lung (newborn) P28.0 2a
- nonviable P07.2
- nuchal cord
  - - affecting fetus or newborn P02.5

O

- obstetric trauma NEC (complicating delivery)
  - - affecting fetus or newborn P03.8
- obstruction, obstructed, obstructive
  - - airway
    - - - newborn P28.8
  - - intestine (mechanical) (paroxysmal) (postinfective)
    - - - newborn P76.9
      - - - - due to
        - - - - - fecaliths P76.8
        - - - - - inspissated milk P76.2
        - - - - - meconium (plug) P76.0
        - - - - - specified NEC P76.8
  - - labor (*see also* Labor, obstructed)
    - - - affecting fetus or newborn P03.1
  - - lung (acute)
    - - - newborn P28.8
  - - meconium plug, newborn P76.0
  - - respiratory (acute)
    - - - newborn P28.8
  - - umbilical cord P02.6
- occlusion, occluded
  - - lung
    - - - newborn P28.8
  - - respiratory, (acute)
    - - - newborn P28.8
  - - umbilical cord P02.6
- oligohydramnios
  - - affecting fetus or newborn P01.2 2a
- omphalitis (congenital) (newborn) (with mild hemorrhage) P38
- omphalorrhagia, newborn P51.9
- operation
  - - for delivery, affecting fetus or newborn (*see also* Delivery, by type, affecting fetus) P03.8
  - - maternal, unrelated to current delivery, affecting fetus or newborn P00.6
- ophthalmia
  - - neonatorum, newborn P39.1
- overfeeding
  - - newborn P92.4
- oversize fetus P08.1
  - - causing disproportion
    - - - affecting fetus or newborn P03.1
    - - - exceptionally large (more than 4500 g) P08.0

P

- palpitations (heart)
- - fetal P20.9
- - newborn P29.1
- palsy (*see also* Paralysis)
- - brachial plexus NEC
- - - fetus or newborn (birth injury) P14.3
- - Erb's P14.0
- - facial
- - - newborn (birth injury) P11.3
- - Klumpke(-Déjerine) P14.1
- papyraceous fetus - *see* Death, fetus
- paralysis, paralytic (complete) (incomplete)
- - birth injury P14.9
- - brachial plexus NEC
- - - birth injury P14.3
- - - newborn P14.3
- - Duchenne's
- - - birth injury P14.0
- - Erb(-Duchenne) (birth) (newborn) P14.0
- - facial (nerve)
- - - birth injury P11.3
- - - newborn P11.3
- - Klumpke(-Déjerine) (birth injury) (newborn) P14.1
- - nerve - *see also* Disorder, nerve
- - - birth injury P14.9
- - - facial
- - - - birth injury P11.3
- - - - newborn P11.3
- - - newborn (birth injury) P14.9
- - - phrenic (birth injury) P14.2
- - - radial
- - - - birth injury P14.3
- - - - newborn P14.3
- - - seventh or facial
- - - - newborn (birth injury) P11.3
- - newborn P91.8
- - - birth injury P11.9 2a
- - radial nerve
- - - birth injury P14.3
- paraplegia (lower)
- - newborn P91.8
- - - birth injury P11.9 2a
- patent - *see also* Imperfect, closure
- - cervix, maternal
- - - complicating pregnancy
- - - - affecting fetus or newborn P01.0
- pendulous
- - abdomen, maternal, in pregnancy or childbirth
- - - affecting fetus or newborn P03.8
- penetration - *see* Perforation
- - lung

## ICD-10-Mortality Perinatal Subset - 2014

- - - nontraumatic
- - - - newborn P28.8
- perforation, perforated (nontraumatic) (*see also* Rupture)
- - bowel
- - - fetus or newborn P78.0
- - colon
- - - fetus or newborn P78.0
- - ileum
- - - fetus or newborn P78.0
- - intestine, unspecified part
- - - fetus or newborn P78.0
- - - ulcerative NEC
- - - - fetus or newborn P78.0
- - jejunum, jejunal
- - - fetus or newborn P78.0
- - lung (traumatic)
- - - nontraumatic
- - - - newborn P28.8
- - rectum
- - - fetus or newborn P78.0
- - sigmoid (*see also* Perforation, colon)
- - - fetus or newborn P78.0
- peritonitis (adhesive) (fibrinous) (with effusion)
- - congenital NEC P78.1
- - meconium (newborn) P78.0
- - neonatal P78.1
- - - meconium P78.0
- persistence, persistent (congenital)
- - fetal
- - - circulation
- - - - newborn P29.3
- - occipitoposterior or transverse (position)
- - - affecting fetus or newborn P03.1
- - occipitotransverse (position)
- - - affecting fetus or newborn P03.1
- petechia, petechiae
- - fetus or newborn P54.5
- pithecoïd maternal pelvis
- - with disproportion (fetopelvic)
- - - affecting fetus or newborn P03.1
- placenta, placental
- - ablatio (*see also* Abruptio placentae)
- - - affecting fetus or newborn P02.1 2a
- - abnormal, abnormality NEC
- - - with hemorrhage
- - - - affecting fetus or newborn P02.1 2a
- - - affecting fetus or newborn P02.2
- - abruptio (*see also* Abruptio placentae)
- - - affecting fetus or newborn P02.1 2a
- - accreta
- - - affecting fetus or newborn P02.2
- - cyst (amniotic)
- - - affecting fetus or newborn P02.2
- - disease NEC
- - - affecting fetus or newborn P02.2
- - dysfunction

## ICD-10-Mortality Perinatal Subset - 2014

- - - affecting fetus or newborn P02.2
- - enlarged P02.2
- - immature P02.2
- - implantation, insertion (abnormal) (poor)
- - - affecting fetus or newborn P02.2
- - - low - *see* Placenta, previa
- - increta
- - - affecting fetus or newborn P02.2
- - infarction
- - - affecting fetus or newborn P02.2
- - insufficiency
- - - affecting fetus or newborn P02.2
- - marginal (hemorrhage) (rupture)
- - - affecting fetus or newborn P02.0
- - percreta
- - - affecting fetus or newborn P02.2
- - polyp
- - - affecting fetus or newborn P02.2
- - preterm P02.2
- - previa (central) (complete) (marginal) (partial) (total) (with hemorrhage)
- - - affecting fetus or newborn P02.0
- placentitis
- - affecting fetus or newborn P02.7
- platypeloid maternal pelvis
- - with disproportion (fetopelvic)
- - - affecting fetus or newborn P03.1
- plug
- - meconium (newborn), syndrome P76.0
- pneumatocele
- - newborn P28.8
- pneumomediastinum
- - congenital or perinatal P25.2
- - newborn P25.2
- pneumonia (acute) (community acquired) (double) (health care associated (HCAP)) (hemorrhagic) (lobe) (migratory) (nosocomial) (obstructive) (post obstructive) (primary) (purulent) (septic) (unresolved) (*see also* Pneumonitis)
- - adenoviral
- - - newborn P23.0
- - adynamic
- - - newborn P23.9
- - aspiration
- - - bacterial
- - - - newborn P23.6
- - - due to
- - - - aspiration of micro-organisms
- - - - - bacterial
- - - - - newborn P23.6
- - - - - specified type NEC
- - - - - newborn P23.6
- - - - - newborn P24.9
- - - - - specified organism NEC
- - - - - newborn P23.8
- - - - - viral
- - - - - newborn P23.0
- - - - - specified type NEC
- - - - - newborn P23.0
- - - - bacterial organisms



## ICD-10-Mortality Perinatal Subset - 2014

- - - - newborn P23.6
- - - newborn P24.9
- - - - meconium P24.0
- - atypical NEC
- - - newborn P23.9
- - atypical, virus
- - - newborn P23.0
- - B (*see also* Pneumonia, bacterial)
- - - newborn P23.6
- - bacillus
- - - newborn P23.6
- - - specified NEC
- - - - newborn P23.6
- - bacterial
- - - newborn P23.6
- - - specified NEC
- - - - newborn P23.6
- - bacterium anitratum
- - - newborn P23.6
- - broncho, bronchial (confluent) (croupous) (diffuse) (disseminated) (involving lobes) (lobar) (*see also* Pneumonia)
- - - bacterial
- - - - newborn P23.6
- - - - specified NEC
- - - - - newborn P23.6
- - - Eaton's agent
- - - - newborn P23.6
- - - *Escherichia coli* (*E. coli*)
- - - - newborn P23.4
- - - Friedländer's bacillus
- - - - newborn P23.6
- - - *Haemophilus influenzae*
- - - - newborn P23.6
- - - hypostatic
- - - - newborn P23.9
- - - *Klebsiella* (*pneumoniae*)
- - - - newborn P23.6
- - - *Mycoplasma* (*pneumoniae*)
- - - - newborn P23.6
- - - newborn P23.9
- - - pleuro-pneumonia-like-organisms (PPLO)
- - - - newborn P23.6
- - - pneumococcal
- - - - newborn P23.6
- - - *Proteus*
- - - - newborn P23.6
- - - *Pseudomonas*
- - - - newborn P23.5
- - - specified organism NEC
- - - - newborn P23.8
- - - staphylococcal
- - - - newborn P23.2
- - - streptococcal NEC
- - - - newborn P23.6
- - - viral, virus (*see also* Pneumonia, viral)
- - - - newborn P23.0
- - bronchostatic

## ICD-10-Mortality Perinatal Subset - 2014

- - - newborn P23.9
- - *Burkholderia cepacia*
- - - newborn P23.6
- - catarrhal (*see also* Pneumonia, broncho)
- - - newborn P23.9
- - chlamydial
- - - congenital P23.1
- - - newborn P23.1
- - congenital (infective) P23.9
- - - due to
- - - - bacterium NEC P23.6
- - - - *Candida* P37.5
- - - - *Chlamydia* P23.1
- - - - *Escherichia coli* P23.4
- - - - *Haemophilus influenzae* P23.6
- - - - infective organism NEC P23.8
- - - - *Klebsiella pneumoniae* P23.6
- - - - *Mycoplasma* P23.6
- - - - *Pseudomonas* P23.5
- - - - staphylococcus P23.2
- - - - streptococcus (except group B) P23.6
- - - - - group B P23.3
- - - - viral agent P23.0
- - - specified NEC P23.8
- - consolidating
- - - newborn P23.9
- - diffuse - *see* Pneumonia, broncho
- - - interstitial
- - - - newborn P28.8
- - Eaton's agent
- - - newborn P23.6
- - *Enterobacter*
- - - newborn P23.6
- - enterococcus, enterococcal
- - - newborn P23.6
- - *Escherichia coli (E.coli)*
- - - newborn P23.4
- - Friedländer's bacillus
- - - newborn P23.6
- - gram-negative NEC
- - - newborn P23.6
- - gram-positive NEC
- - - newborn P23.6
- - *Haemophilus influenzae* (broncho) (lobar)
- - - newborn P23.6
- - *Herellea*
- - - newborn P23.6
- - hypostatic (broncho) (lobar)
- - - newborn P23.9
- - in (due to)
- - - adenovirus
- - - - newborn P23.0
- - - *Chlamydia*
- - - - neonatal P23.1
- - - Eaton's agent
- - - - newborn P23.6

## ICD-10-Mortality Perinatal Subset - 2014

- - - *Enterobacter*
- - - - newborn P23.6
- - - *Escherichia coli (E. coli)*
- - - - newborn P23.4
- - - Friedländer's bacillus
- - - - newborn P23.6
- - - *Haemophilus influenzae (H. influenzae)*
- - - - newborn P23.6
- - - *Klebsiella (pneumoniae)*
- - - - newborn P23.6
- - - *Mycoplasma (pneumoniae)*
- - - - newborn P23.6
- - - parainfluenza virus
- - - - newborn P23.0
- - - pleuro-pneumonia-like-organism (PPLO)
- - - - newborn P23.6
- - - pneumococcus
- - - - newborn P23.6
- - - *Proteus*
- - - - newborn P23.6
- - - *Pseudomonas* NEC
- - - - newborn P23.5
- - - respiratory syncytial virus
- - - - newborn P23.0
- - - septicemia
- - - - newborn P36.9
- - - *Serratia marcescens*
- - - - newborn P23.6
- - - specified
- - - - bacterium NEC
- - - - - newborn P23.6
- - - - - organism NEC
- - - - - newborn P23.8
- - - staphylococcus
- - - - newborn P23.2
- - - *Streptococcus*
- - - - group B
- - - - - newborn P23.3
- - - - - newborn P23.6
- - - - *pneumoniae*
- - - - - newborn P23.6
- - - - specified NEC
- - - - - newborn P23.6
- - - interstitial
- - - - diffuse
- - - - - newborn P28.8
- - - - newborn P28.8
- - - - - chronic P27.8
- - - *Klebsiella (pneumoniae)*
- - - - newborn P23.6
- - - lobar (disseminated) (interstitial) (*see also* Pneumonia)
- - - - bacterial
- - - - - newborn P23.6
- - - - - specified NEC
- - - - - newborn P23.6
- - - *Escherichia coli (E. coli)*

## ICD-10-Mortality Perinatal Subset - 2014

- - - - newborn P23.4
- - - Friedländer's bacillus
- - - - newborn P23.6
- - - *Haemophilus influenzae*
- - - - newborn P23.6
- - - *Klebsiella (pneumoniae)*
- - - - newborn P23.6
- - - newborn P23.9
- - - pneumococcal
- - - - newborn P23.6
- - - *Proteus*
- - - - newborn P23.6
- - - *Pseudomonas*
- - - - newborn P23.5
- - - specified organism NEC
- - - - newborn P23.8
- - - staphylococcal
- - - - newborn P23.2
- - - streptococcal NEC
- - - - newborn P23.6
- - - *Streptococcus pneumoniae*
- - - - newborn P23.6
- - - meconium P24.0
- - - metastatic
- - - newborn P36.9
- - - MRSA (methicillin resistant *Staphylococcus aureus*)
- - - newborn P23.2
- - - *Mycoplasma (pneumoniae)*
- - - newborn P23.6
- - - neonatal (*see also* Pneumonia, congenital) P23.9
- - - aspiration P24.9
- - - orthostatic
- - - newborn P23.9
- - - parainfluenza virus
- - - newborn P23.0
- - - pleuro-pneumonia-like-organism (PPLO)
- - - newborn P23.6
- - - pneumococcal (broncho) (lobar)
- - - newborn P23.6
- - - postural
- - - newborn P23.9
- - - *Proteus*
- - - newborn P23.6
- - - *Pseudomonas*
- - - newborn P23.5
- - - respiratory syncytial virus
- - - newborn P23.0
- - - *S. pneumoniae*
- - - newborn P23.6
- - - *Serratia marcescens*
- - - newborn P23.6
- - - specified NEC
- - - bacterium NEC
- - - - newborn P23.6
- - - newborn P23.9
- - - organism NEC

## ICD-10-Mortality Perinatal Subset - 2014

- - - - newborn P23.8
- - - virus NEC
- - - - newborn P23.0
- - staphylococcal (broncho) (lobar)
- - - newborn P23.2
- - static, stasis
- - - newborn P23.9
- - streptococcal NEC (broncho) (lobar)
- - - group B
- - - - newborn P23.3
- - - newborn P23.6
- - *Streptococcus pneumoniae*
- - - newborn P23.6
- - viral, virus (broncho) (interstitial) (lobar)
- - - adenoviral
- - - - newborn P23.0
- - - congenital P23.0
- - - newborn P23.0
- - - parainfluenza
- - - - newborn P23.0
- - - respiratory syncytial
- - - - newborn P23.0
- - - specified NEC
- - - - newborn P23.0
- pneumonitis (acute) (primary) (*see also* Pneumonia)
- - due to
- - - toxoplasmosis
- - - - congenital P37.1+ J17.3\*
- - meconium P24.0
- - neonatal aspiration P24.9
- - rubella, congenital P35.0 2a
- pneumopathy NEC
- - newborn P28.9
- pneumopericardium
- - congenital P25.3
- - fetus or newborn P25.3
- pneumothorax
- - congenital P25.1
- - newborn P25.1
- - perinatal period P25.1
- - spontaneous NEC
- - - fetus or newborn P25.1
- poliomyelitis (anterior) (epidemic)
- - congenital P35.8
- polycystic (disease)
- - lung
- - - newborn P28.8
- polycythemia (primary) (rubra) (vera) M9950)
- - neonatorum P61.1
- polyhydramnios
- - affecting fetus or newborn P01.3
- polyp, polypus
- - cervix (uteri), maternal
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8
- - umbilical, newborn P83.6

## ICD-10-Mortality Perinatal Subset - 2014

- - uterus (body) (corpus) (mucous), maternal
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8
- poor
- - contractions, labor
- - - affecting fetus or newborn P03.6
- - fetal growth NEC P05.9
- postdates (pregnancy) (mother)
- - affecting fetus or newborn P08.2
- postmaturity, postmature (fetus or newborn) P08.2
- postterm (pregnancy)
- - infant P08.2
- precipitate labor or delivery
- - affecting fetus or newborn P03.5
- pre-eclampsia
- - affecting fetus or newborn P00.0
- pregnancy (single) (uterine)
- - abdominal (ectopic)
- - - affecting fetus or newborn P01.4
- - complicated by (*see also* Pregnancy, management, affected by)
- - - pelvic organs or tissues
- - - - affecting fetus or newborn P03.8
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- - cornual
- - - fetus or newborn P01.4
- - ectopic (ruptured)
- - - affecting fetus or newborn P01.4
- - molar NEC
- - - affecting fetus or newborn P01.8
- - multiple NEC
- - - affecting fetus or newborn P01.5
- - mural
- - - fetus or newborn P01.4
- - ovarian
- - - fetus or newborn P01.4
- - quadruplet
- - - affecting fetus or newborn P01.5
- - quintuplet
- - - affecting fetus or newborn P01.5
- - sextuplet
- - - affecting fetus or newborn P01.5
- - triplet
- - - affecting fetus or newborn P01.5
- - tubal (with abortion) (with rupture)
- - - affecting fetus or newborn P01.4
- - twin
- - - affecting fetus or newborn P01.5
- premature (*see also* Prematurity)
- - birth NEC P07.3
- - contraction
- - - atrial
- - - - newborn P29.1
- - - atrioventricular
- - - - newborn P29.1
- - - heart (extrasystole)

## ICD-10-Mortality Perinatal Subset - 2014

- - - - newborn P29.1
- - - junctional
- - - - newborn P29.1
- - - ventricular
- - - - newborn P29.1
- - delivery
- - - newborn NEC P07.3
- - infant NEC P07.3
- - - light-for-dates P05.0
- - labor
- - - newborn NEC P07.3
- - lungs P28.0 2a
- - rupture, membranes or amnion (*see also* Rupture, membranes, premature)
- - - affecting fetus or newborn P01.1 2a
- - ventricular systole
- - - newborn P29.1
- prematurity NEC (less than 37 completed weeks) (*see also* Immaturity) P07.3
- - extreme (less than 28 completed weeks) P07.2
- - gross P07.2
- - marked P07.2
- - severe P07.2
- presentation, fetal
- - abnormal
- - - before labor, affecting fetus or newborn P01.7
- - - causing obstructed labor
- - - - affecting fetus or newborn (any, except breech) P03.1
- - - - - breech P03.0
- - breech
- - - with external version before labor, affecting fetus or newborn P01.7
- - - before labor, affecting fetus or newborn P01.7
- - - affecting fetus or newborn NEC P01.7
- - - causing obstructed labor
- - - - fetus or newborn P03.0
- pressure
- - birth, fetus or newborn, NEC P15.9 2a
- - brain
- - - injury at birth NEC P11.1 2a
- - increased
- - - intracranial (benign)
- - - - injury at birth P11.0 2a
- preterm infant, newborn NEC P07.3
- previa
- - affecting fetus or newborn P02.0
- - placenta (with hemorrhage) (*see also* Placenta, previa)
- - - affecting fetus or newborn P02.0
- - vasa
- - - affecting fetus or newborn P02.6
- previable P07.2
- prior to viability P07.2
- problem (related to) (with) (*see also* Disease)
- - feeding
- - - newborn P92.9
- - - - breast P92.5
- - - - overfeeding P92.4
- - - - slow P92.2
- - - - specified NEC P92.8

## ICD-10-Mortality Perinatal Subset - 2014

- - - - underfeeding P92.3
- procedure (surgical)
- - maternal (unrelated to current delivery), affecting fetus or newborn P00.6
- - - nonsurgical (medical) P00.7
- prolapse, prolapsed
- - arm or hand
- - - in fetus or newborn P03.1
- - cord - *see* Prolapse, umbilical cord
- - fetal limb NEC
- - - in fetus or newborn P03.1
- - leg
- - - in fetus or newborn P03.1
- - membranes (amniotic sac)
- - - affecting fetus or newborn P01.0
- - umbilical cord
- - - affecting fetus or newborn P02.4
- - uterus (with prolapse of vagina)
- - - pregnant, affecting fetus or newborn P03.8
- prolonged
- - labor
- - - affecting fetus or newborn P03.8
- - uterine contractions in labor
- - - affecting fetus or newborn P03.6
- prominent ischial spine or sacral promontory
- - with disproportion (fetopelvic)
- - - affecting fetus or newborn P03.1
- proteinuria
- - pre-eclamptic (*see also* Pre-eclampsia)
- - - affecting fetus or newborn P00.0
- pseudocyst
- - lung
- - - newborn P28.8
- pseudomenses (newborn) P54.6
- pseudomenstruation (newborn) P54.6
- pseudoparalysis
- - atonic, congenital P94.2
- pseudosclerema, newborn P83.8
- pulmolithiasis
- - newborn P28.8
- pulse
- - alternating
- - - fetal P20.9
- - - newborn P29.1
- - bigeminal
- - - fetal P20.9
- - - newborn P29.1
- pulsus alternans or trigeminus
- - fetal P20.9
- - newborn P29.1
- purpura
- - thrombocytopenic (congenital) (hereditary) (*see also* Thrombocytopenia)
- - - neonatal, transitory P61.0
- pyelitis, maternal (congenital) (uremic) (*see also* Pyelonephritis)
- - complicating pregnancy
- - - affecting fetus or newborn P00.1
- pyelonephritis, maternal (*see also* Nephritis, tubulo-interstitial)



## ICD-10-Mortality Perinatal Subset - 2014

- - complicating pregnancy
- - - affecting fetus or newborn P00.1
- - in (due to)
- - - septicemia NEC
- - - - newborn P36.9
- pyoderma, pyodermia NEC
- - newborn P39.4
- pyrexia (of unknown origin)
- - newborn, environmentally-induced P81.0

Q

- quadriplegia
- - newborn NEC P11.9 2a
- quadruplet, affecting fetus or newborn P01.5 - quintuplet, affecting fetus or newborn P01.5 - rachitic
- - pelvis, maternal (late effect)
- - - with disproportion (fetopelvic)
- - - - affecting fetus or newborn P03.1

R

- radiology, maternal, affecting fetus or newborn P00.7
- rapid
  - - heart(beat)
    - - - fetal P20.9
    - - - newborn P29.1
  - - second stage (delivery)
    - - - affecting fetus or newborn P03.5
- R.D.S. (newborn) (see also Syndrome, respiratory distress) P22.0
- reaction - *see also* Disorder
  - - drug NEC
    - - - newborn P93
    - - - withdrawal
      - - - - infant of dependent mother P96.1
      - - - - newborn P96.1
- rectocele, maternal
  - - in pregnancy or childbirth
    - - - causing obstructed labor
      - - - - affecting fetus or newborn P03.1
- reduction
  - - fetus(es) (any number) (multiple) P01.5
- reflux
  - - esophageal
    - - - neonatal P78.8
    - - gastroesophageal
      - - - in newborn P78.8
- regurgitation
  - - food (*see also* Vomiting)
    - - - newborn P92.1
- respiration
  - - insufficient, or poor
    - - - newborn NEC P28.5 2a
- restriction (of)
  - - intrauterine growth P05.9
- retardation
  - - growth (*see also* Retardation, physical)
    - - - fetus P05.9
    - - intrauterine growth P05.9
      - - physical (child)
        - - - fetus P05.9
- retention, retained
  - - placenta (total) (with hemorrhage)
    - - - affecting fetus or newborn P02.8
  - - products of conception
    - - - early pregnancy (dead fetus)
      - - - - affecting fetus or newborn P01.8
      - - - following
        - - - - delivery (with hemorrhage)
          - - - - - affecting fetus or newborn P02.8
  - - secretions
    - - - airway
      - - - - newborn P28.8

## ICD-10-Mortality Perinatal Subset - 2014

- - - of bronchus
- - - - newborn P28.8
- - - of lungs
- - - - newborn P28.8
- - - pulmonary
- - - - newborn P28.8
- retraction
- - lung
- - - newborn P28.8
- - ring, uterus (Bandl's) (pathological)
- - - affecting fetus or newborn P03.6
- retroversion, retroverted
- - uterus, uterine, maternal
- - - in pregnancy or childbirth
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- reverse
- - end diastolic flow P02.2
- Rh (factor)
- - hemolytic disease (fetus or newborn) P55.0
- - incompatibility, immunization or sensitization
- - - fetus or newborn P55.0
- - negative mother affecting fetus or newborn P55.0
- rhythm
- - atrioventricular nodal
- - - newborn P29.1
- - disorder
- - - coronary sinus
- - - - newborn P29.1
- - - ectopic
- - - - newborn P29.1
- - - newborn P29.1
- - - nodal
- - - - newborn P29.1
- - escape
- - - newborn P29.1
- - heart, abnormal
- - - newborn P29.1
- - nodal
- - - newborn P29.1
- rigid, rigidity
- - cervix, maternal (uteri)
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- - pelvic floor, maternal
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- - perineum or vulva, maternal
- - - in pregnancy or childbirth
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- - vagina, maternal

## ICD-10-Mortality Perinatal Subset - 2014

- - - in pregnancy or childbirth
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.8
- ring(s)
- - Bandl's
- - - fetus or newborn P03.6
- - contraction, complicating delivery
- - - affecting fetus or newborn P03.6
- - retraction, uterus, pathological
- - - affecting fetus or newborn P03.6
- rotation
- - manual, affecting fetus or newborn P03.8
- rubella
- - congenital P35.0 2a
- - maternal
- - - affecting fetus or newborn P00.2
- - - - manifest rubella in infant P35.0 2a
- rumination
- - newborn P92.1
- rupture, ruptured (*see also* Perforation)
- - fontanelle P13.1
- - kidney
- - - birth injury P15.8
- - liver
- - - birth injury P15.0
- - marginal sinus (placental) (with hemorrhage)
- - - with placenta previa
- - - - affecting fetus or newborn P02.0
- - - affecting fetus or newborn P02.1 2a
- - membranes (spontaneous)
- - - artificial
- - - - delayed delivery following
- - - - - affecting fetus or newborn P01.1 2a
- - - delayed delivery following
- - - - affecting fetus or newborn P01.1 2a
- - - premature
- - - - affecting fetus or newborn P01.1 2a
- - spinal cord (*see also* Injury, spinal cord, by region)
- - - fetus or newborn (birth injury) P11.5
- - spleen (traumatic)
- - - birth injury P15.1
- - - congenital (birth injury) P15.1
- - umbilical cord, complicating delivery
- - - affecting fetus or newborn P50.1
- - uterus, maternal (traumatic)
- - - affecting fetus or newborn P03.8
- - during or after labor
- - - affecting fetus or newborn P03.8

S

- salpingo-oophoritis, maternal (purulent) (ruptured) (septic) (suppurative)
  - - complicating pregnancy
    - - - affecting fetus or newborn P00.8
  - - scar, scarring (*see also* Cicatrix)
    - - - cervix, maternal
      - - - - in pregnancy or childbirth
        - - - - - affecting fetus or newborn P03.8
    - - - due to
      - - - - previous cesarean section, complicating pregnancy or childbirth
        - - - - - affecting fetus or newborn P03.8
    - - - uterus, maternal
      - - - - in pregnancy or childbirth
        - - - - - affecting fetus or newborn P03.8
  - scleredema
    - - newborn P83.0
  - sclerema
    - - adiposum (newborn) P83.0
    - - edematosum (newborn) P83.0
    - - newborn P83.0
  - scleroderma, sclerodermia (diffuse) (generalized)
    - - newborn P83.8
  - scoliotic maternal pelvis with disproportion (fetopelvic)
    - - affecting fetus or newborn P03.1
  - section
    - - cesarean (*see also* Delivery, cesarean)
      - - - affecting fetus or newborn P03.4
      - - - postmortem, affecting fetus or newborn P01.6
      - - - previous, in pregnancy or childbirth
        - - - - affecting fetus or newborn P03.8
    - - seizure(s) (*see also* Convulsions)
      - - - newborn P90
    - - sepsis (generalized) (*see also* Septicemia)
      - - - bacterial, newborn P36.9
        - - - - due to
          - - - - - anaerobes NEC P36.5
          - - - - - *Escherichia coli* P36.4
          - - - - - *Staphylococcus* NEC P36.3
            - - - - - - *aureus* P36.2
          - - - - - streptococcus NEC P36.1
            - - - - - - group B P36.0
        - - - - specified type NEC P36.8
      - - - newborn NEC P36.9
        - - - - due to
          - - - - - anaerobes NEC P36.5
          - - - - - *Escherichia coli* P36.4
          - - - - - *Staphylococcus* NEC P36.3
            - - - - - - *aureus* P36.2
          - - - - - streptococcus NEC P36.1
            - - - - - - group B P36.0
        - - - - specified NEC P36.8
      - - - umbilical (newborn) (organism unspecified) P36-

## ICD-10-Mortality Perinatal Subset - 2014

- septic
  - - umbilical cord P38
  - septicemia, septicemic (generalized) (suppurative) (*see also* Sepsis)
    - - *Actinobacter*
      - - - newborn P36.8
    - - *Aerobacter aerogenes*
      - - - newborn P36.8
    - - anaerobic
      - - - newborn P36.5
    - - *Bacillus coli*
      - - - newborn P36.8
    - - bacteriodes
      - - - newborn P36.5
    - - *Citrobacter Citrobacter (braakii) (freundii) (koseri)*
      - - - newborn P36.8
    - - *Clostridium (difficile)*
      - - - newborn P36.5
    - - congenital P36.9
    - - cryptogenic
      - - - newborn P36.9
    - - *Enterobacter (aerogenes) (clocae)*
      - - - newborn P36.8
    - - enterococcus, enterococcal
      - - - newborn P36.1
    - - *Escherichia coli*
      - - - newborn P36.4
    - - Friedländers'
      - - - newborn P36.8
    - - fungus
      - - - newborn P37.8
    - - gangrenous
      - - - newborn P36.9
    - - gram-negative (organism)
      - - - anaerobic
        - - - - newborn P36.5
      - - - newborn P36.9
    - - gram-positive
      - - - newborn P36.8
    - - *Haemophilus influenzae*
      - - - newborn P36.8
    - - *Klebsiella (pneumoniae)*
      - - - newborn P36.8
    - - newborn NEC (*see also* Sepsis, newborn) P36.9
    - - pneumococcal
      - - - newborn P36.1
    - - *Proteus (mirabilis) (vulgaris)*
      - - - newborn P36.8
    - - *Providencia (species)*
      - - - newborn P36.5
    - - *Pseudomonas (aeruginosa)*
      - - - newborn P36.8
    - - *Serratia*
      - - - newborn P36.8
    - - specified bacterial organism NEC
      - - - newborn P36.8
    - - *Staphylococcus, staphylococcal*

## ICD-10-Mortality Perinatal Subset - 2014

- - - aureus
- - - - newborn P36.2
- - - coagulase-negative
- - - - newborn P36.3
- - - newborn P36.3
- - - specified NEC
- - - - newborn P36.3
- - Streptococcus, streptococcal
- - - agalactiae
- - - - newborn P36.0
- - - group
- - - - A
- - - - - newborn P36.1
- - - - B
- - - - - newborn P36.0
- - - - D
- - - - - newborn P36.1
- - - neonatal P36.1
- - - pneumoniae
- - - - newborn P36.1
- - - pyogenes
- - - - newborn P36.1
- - - specified NEC
- - - - newborn P36.1
- - viral
- - - newborn P35.8
- sextuplet, affecting fetus or newborn P01.5
- shock (acute)
- - birth (fetus or newborn) P96.8
- - lung
- - - newborn P22.0
- - septic (*see also* Septicemia)
- - - newborn P36.9
- - septicemic
- - - newborn P36.9
- short, shortening, shortness
- - cervix, maternal affecting fetus or newborn P01.0
- - cord (umbilical), complicating delivery
- - - affecting fetus or newborn P02.6
- - umbilical cord
- - - complicating delivery
- - - - affecting fetus or newborn P02.6
- shutdown
- - cardiac
- - - newborn P29.0
- - fetal, fetus P29.8
- - cardiorenal
- - - fetal, fetus P29.8
- - - newborn P29.0
- sick
- - sinus (syndrome)
- - - newborn P29.1
- sinus - *see also* Fistula
- - arrhythmia
- - - newborn P29.1
- - marginal, ruptured or bleeding



## ICD-10-Mortality Perinatal Subset - 2014

- - - with placenta previa
- - - - affecting fetus or newborn P02.0
- - - affecting fetus or newborn P02.1 2a
- sleep
- - apnea
- - - newborn P28.3
- slipped, slipping
- - ligature, umbilical P51.8
- - skin, fetus - see Maceration, fetus
- slow
- - feeding, newborn P92.2
- - fetal growth NEC P05.9
- - heart(beat)
- - - fetal P20.9
- - - newborn P29.1
- small(ness)
- - fetus or newborn for gestational age P05.1
- - pelvis, maternal
- - - with disproportion (fetopelvic)
- - - - affecting fetus or newborn P03.1
- small-and-light-for-dates (infant) P05.1
- small-for-dates (infant) P05.1
- snuffles (non-syphilitic)
- - newborn P28.8
- spasm(s), spastic, spasticity
- - cervix, maternal, complicating delivery
- - - affecting fetus or newborn P03.6
- - uterus, maternal
- - - complicating labor
- - - - affecting fetus or newborn P03.6
- spondylolisthesis, maternal (acquired)
- - with disproportion (fetopelvic)
- - - affecting fetus or newborn P03.1
- spondylolysis, maternal
- - lumbosacral region
- - - with disproportion (fetopelvic)
- - - - affecting fetus or newborn P03.1
- spondylosis, maternal
- - with
- - - disproportion (fetopelvic)
- - - - affecting fetus or newborn P03.1
- standstill
- - respiration, respiratory
- - - newborn P28.5 2a
- staphylococemia
- - newborn P36.3
- stenosis (cicatricial) - *see also* Stricture
- - cervix, cervical (canal), maternal
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- - umbilical cord P02.6
- - vagina, maternal
- - - in pregnancy or childbirth
- - - - affecting fetus or newborn P03.8

## ICD-10-Mortality Perinatal Subset - 2014

- - - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- stillbirth NEC - *see* Death, fetus
- strangulation, strangulated
- - umbilical cord - *see also* Compression, umbilical cord P02.5
- - - fetus or newborn P02.5
- stricture (*see also* Stenosis)
- - cervix, cervical (canal), maternal
- - - in pregnancy or childbirth
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- - umbilical cord P02.6
- stridor
- - congenital (larynx) NEC P28.8
- stroke (apoplectic) (brain) (paralytic)
- - hemorrhagic
- - - newborn P52.4
- surgery
- - previous, in pregnancy or childbirth
- - - cervix, maternal
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- - - pelvic soft tissues NEC, maternal
- - - - affecting fetus or newborn P03.8
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- - - uterus, maternal
- - - - causing obstructed labor
- - - - - affecting fetus or newborn P03.1
- suspended maternal uterus
- - in pregnancy or childbirth
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- syncope
- - bradycardia
- - - fetal P20.9
- - - newborn P29.1
- syndrome - *see also* Disease
- - aspiration, of newborn (massive) P24.9
- - - meconium P24.0
- - bronze baby P83.8
- - bubbly lung P27.0
- - cardiorespiratory distress (idiopathic), newborn P22.0
- - cold injury (newborn) P80.0
- - congenital
- - - rubella (manifest) P35.0 2a
- - defibrination
- - - fetus or newborn P60
- - drug withdrawal, infant of dependent mother P96.1
- - fetal
- - - transfusion P02.3
- - fetomaternal dysfunctional P02.2
- - floppy
- - - baby P94.2
- - gray (newborn) P93
- - HELLP (hemolysis, elevated liver enzymes and low platelet count)

## ICD-10-Mortality Perinatal Subset - 2014

- - - affecting fetus or newborn P00.0
- - idiopathic cardiorespiratory distress, newborn P22.0
- - infant of diabetic mother P70.1
- - - gestational diabetes P70.0
- - inflammatory response
- - - newborn P36.9
- - inspissated bile (newborn) P59.1
- - low
- - - cardiac
- - - - output
- - - - - newborn P29.8
- - lower radicular, newborn (birth injury) P14.8
- - meconium plug (newborn) P76.0
- - mirror (maternal) P01.8
- - neonatal abstinence P96.1
- - perfusion
- - - newborn P28.8
- - placental
- - - dysfunction
- - - - affecting fetus or newborn P02.2
- - - - insufficiency
- - - - affecting fetus or newborn P02.2
- - - transfusion (mother)
- - - - in fetus or newborn P02.3
- - pulmonary
- - - dysmaturity (Wilson-Mikity) P27.0
- - radicular NEC
- - - upper limbs, newborn (birth injury) P14.3
- - respiratory
- - - distress (idiopathic) (newborn) P22.0
- - - - adult
- - - - - newborn P22.0
- - - - newborn P22.0
- - sick sinus
- - - newborn P29.1
- - sinus node
- - - newborn P29.1
- - stiff lung
- - - newborn P28.8
- - tachycardia-bradycardia
- - - newborn P29.1
- - transfusion
- - - fetomaternal P50.4
- - twin (to twin) transfusion
- - - in fetus or newborn P02.3
- - wet lung
- - - with injury
- - - - newborn P22.1
- - withdrawal - *see* Withdrawal, state
- - - drug
- - - - infant of dependent mother P96.1
- - - - therapeutic use, newborn P96.2
- synechia (anterior) (iris) (posterior) (pupil)
- - uterus (maternal)
- - - affecting fetus or newborn P03.8
- syphilis, syphilitic, maternal (acquired)

## **ICD-10-Mortality Perinatal Subset - 2014**

- - complicating pregnancy or childbirth
- - - affecting fetus or newborn P00.2
- - maternal, affecting fetus or newborn P00.2

T

- tachycardia
  - - fetal - *see* Distress, fetal
  - - newborn, *classifiable to* R00.0 , I47- P29.1
  - - sick sinus
    - - - newborn P29.1
- tachypnea
  - - transitory, of newborn P22.1
- tamponade
  - - pulmonary
    - - - newborn P28.8
- tear, torn (traumatic) -*see also* Wound
  - - tentorial, at birth P10.4
- umbilical cord
  - - - affecting fetus or newborn P50.1
- temperature
  - - cold, trauma from
    - - - newborn P80.0
- tetany (due to)
  - - neonatal (without calcium or magnesium deficiency) P71.3
- thick, thickening
  - - membranes (maternal), affecting fetus or newborn P02.8
- threatened
  - - abortion
    - - - affecting fetus P01.8
  - - labor (*see also* Labor, false)
    - - - affecting fetus P01.8
  - - miscarriage
    - - - affecting fetus P01.8
  - - premature delivery, affecting fetus P01.8
- thrombocytopenia, thrombocytopenic
  - - neonatal, transitory P61.0
    - - - due to
      - - - - exchange transfusion P61.0
      - - - - idiopathic maternal thrombocytopenia P61.0
      - - - - isoimmunization P61.0
  - - transient neonatal P61.0
- thrombophlebitis, maternal
  - - antepartum (superficial)
    - - - affecting fetus or newborn P00.3
  - - pregnancy (superficial)
    - - - affecting fetus or newborn P00.3
- thrombosis, thrombotic (multiple) (progressive) (septic) (vein) (vessel)
  - - placenta P02.2
    - - - marginal vein P02.2
  - - umbilical cord (vessels)
    - - - affecting fetus or newborn P02.6
    - - - complicating delivery
      - - - - affecting fetus or newborn P02.6
- thyrotoxicosis
  - - neonatal P72.1
- tobacco (nicotine)

## ICD-10-Mortality Perinatal Subset - 2014

- - maternal use, affecting fetus or newborn P04.2
- Torsades de pointes syndrome
- - newborn P29.1
- torsion
- - umbilical cord in fetus or newborn P02.5
- torticollis (intermittent) (spastic)
- - due to birth injury P15.2
- toxemia
- - maternal (of pregnancy), affecting fetus or newborn P00.0
- - of pregnancy (*see also* Pre-eclampsia)
- - - affecting fetus or newborn P00.0
- toxoplasma, toxoplasmosis (acquired)
- - congenital (acute) (chronic) (subacute) P37.1
- - maternal, affecting fetus or newborn P00.2
- - - manifest toxoplasmosis in infant or fetus (acute) (chronic) (subacute) P37.1
- transfusion
- - placental (syndrome) (mother)
- - - in fetus or newborn P02.3
- - twin-to-twin
- - - fetus or newborn P02.3
- transverse
- - arrest (deep), in labor
- - - affecting fetus or newborn P03.1
- - lie
- - - before labor, affecting fetus or newborn P01.7
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- trauma, traumatism (*see also* Injury)
- - birth -*see* Birth, injury
- - maternal, during pregnancy, affecting fetus or newborn P00.5
- trigeminy
- - fetal P20.9
- - newborn P29.1
- triplet, affecting fetus or newborn P01.5
- tuberculosis, tubercular, tuberculous (caseous) (degeneration) (gangrene) (necrosis)
- - congenital P37.0
- - maternal, complicating pregnancy or childbirth
- - - affecting fetus or newborn P00.2
- tumor (M8000)
- - cervix, maternal, in pregnancy or childbirth
- - - affecting fetus or newborn P03.8
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- - ovary, maternal, in pregnancy or childbirth
- - - affecting fetus or newborn P03.8
- - pelvic, maternal, in pregnancy or childbirth
- - - affecting fetus or newborn P03.8
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- - placental
- - - affecting fetus or newborn P02.2
- - uterus (body), maternal, in pregnancy or childbirth
- - - affecting fetus or newborn P03.8
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- - vagina, maternal, in pregnancy or childbirth

## ICD-10-Mortality Perinatal Subset - 2014

- - - affecting fetus or newborn P03.8
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- - vulva or perineum, maternal, in pregnancy or childbirth
- - - affecting fetus or newborn P03.8
- - - causing obstructed labor
- - - - affecting fetus or newborn P03.1
- twin (pregnancy) (fetus or newborn) P01.5
- twist, twisted
- - umbilical cord in fetus or newborn P02.5
- tyrosinemia
- - newborn, transitory P74.5

U

- ulcer, ulcerated, ulcerating, ulceration, ulcerative
- - intestine, intestinal
- - - perforating
- - - - fetus or newborn P78.0
- - lung
- - - newborn P28.8
- - peptic (site unspecified)
- - - newborn P78.8
- - unable (*see also* Inability)
- - - to breathe properly
- - - newborn P28.8
- - underfeeding, newborn P92.3
- - underweight
- - - for gestational age P05.0
- - Underwood's disease P83.0
- - unknown cause of
- - - fetal death in fetal death file P95
- - unstable
- - - lie
- - - fetus or newborn, before labor P01.7
- - uremia, uremic (coma)
- - - congenital P96.0
- - - maternal NEC, affecting fetus or newborn P00.1
- - - newborn P96.0
- - urticaria
- - - neonatorum P83.8
- - use (of)
- - - harmful
- - - - patent medicines
- - - - - maternal, affecting fetus or newborn P04.1
- - - uveitis (anterior)
- - - - due to toxoplasmosis
- - - - - congenital P37.1† H22.0\*



V

- vaccinia (generalized) (localized)
  - - congenital P35.8
- vaginitis, maternal (acute)
  - - complicating pregnancy
    - - - affecting fetus or newborn P00.8
- varicella
  - - congenital P35.8
- varicose
  - - vein (ruptured)
    - - - umbilical cord, affecting fetus or newborn P02.6
- varix (ruptured)
  - - umbilical cord, affecting fetus or newborn P02.6
- vasa previa
  - - affecting fetus or newborn P02.6
  - - hemorrhage from, affecting fetus or newborn P50.0
- vasculitis
  - - chorionic
    - - - affecting fetus or newborn P02.7
- vasculopathy, necrotizing
  - - placental fetal P02.2
    - - thrombotic, fetal, specified NEC P02.2
- ventilator lung, newborn P27.8
- ventouse delivery NEC
  - - affecting fetus or newborn P03.3
- ventricle, ventricular
  - - escape
    - - - newborn P29.1
- version
  - - with extraction
    - - - affecting fetus or newborn P01.7
- villitis (placenta)
  - - affecting fetus or newborn P02.2
- viremia
  - - newborn P35.9
- vitality, lack of
  - - newborn P96.8
- vomiting (*see also* Hyperemesis)
  - - newborn P92.0

W

- wandering
- - pacemaker (heart)
- - - newborn P29.1
- weak, weakness
- - newborn P96.8
- weight
- - over 2 pounds (1000 gms) but not more than 5½ pounds (2499 gms) at birth P07.1
- - 2 pounds (999 gms) or under at birth P07.0
- - 10 pounds (4500 gms) or more at birth P08.0
- - 999 grams or less at birth (extremely low) P07.0
- - 1000-2499 grams at birth (low) P07.1
- wet lung
- - newborn P22.1
- wide cranial sutures, newborn P96.3
- Wilson-Mikity syndrome P27.0
- withdrawal
- - state, symptoms, syndrome
- - - newborn
- - - - correct therapeutic substance properly administered P96.2
- - - - infant of dependent mother P96.1
- - therapeutic substance, neonatal P96.2
- wound (cut) (laceration) (open) (penetrating (puncture wound) (with penetrating foreign body)
- - scalpel, fetus or newborn (birth injury) P15.8